

Blow-up phenomena for a $p(x)$ -biharmonic heat equation with variable exponent

ERHAN PIŞKİN*, GÜLISTAN BUTAKIN

ABSTRACT. In this paper, we deal with a $p(x)$ -biharmonic heat equation with variable exponent under Dirichlet boundary and initial condition. We prove the blow up of solutions under suitable conditions.

1. INTRODUCTION

Let $\Omega \subset R^n$ is a bounded domain with smooth boundary $\partial\Omega$. We are concerned with the following $p(x)$ -biharmonic heat equation, with variable exponent, of the form

$$(1) \quad \begin{cases} u_t + \Delta^2 u_t + \Delta_{p(x)}^2 u = |u|^{q(x)-2} u, & Q = \Omega \times (0, T), \\ u(x, t) = \Delta u(x, t) = 0, & \partial Q = \partial\Omega \times [0, T), \\ u(x, 0) = u_0(x), & \Omega, \end{cases}$$

where $\Delta_{p(x)}^2$ is the so-called the $p(x)$ -biharmonic operator and is defined by

$$\Delta_{p(x)}^2 u = \Delta \left(|\Delta u|^{p(x)-2} \Delta u \right).$$

The exponents $p(\cdot)$ and $q(\cdot)$ are given measurable functions on $\bar{\Omega}$ such that

$$(2) \quad 2 \leq p_- \leq p(x) \leq p_+ < q_- \leq q(x) \leq q_+ < p_*(x),$$

with

$$p_*(x) = \begin{cases} \frac{np(x)}{(n-p(x))_+}, & \text{if } p_+ < n, \\ +\infty, & \text{if } p_+ \geq n. \end{cases}$$

We also suppose that

$$(3) \quad |p(x) - p(y)| \leq \frac{A}{\log \left(\frac{1}{|x-y|} \right)}, \text{ for all } x, y \in \Omega \text{ with } |x-y| < \delta,$$

with $A > 0$, $0 < \delta < 1$ and

$$(4) \quad \text{ess inf } (p^*(x) - q(x)) > 0.$$

2020 *Mathematics Subject Classification*. Primary: 40B05; Secondary: 35B44, 35K05.

Key words and phrases. Blow up, heat equation, variable exponent.

Full paper. Received 9 January 2023, revised 18 August 2023, accepted 3 September 2023, available online 22 September 2023.

The following problem was considered by Alaoui et al. in [2]

$$(5) \quad u_t - \operatorname{div} \left(|\nabla u|^{m(x)-2} \nabla u \right) = |u|^{p(x)-2} u.$$

The authors proved the blow up of solutions. Later, Rahmoune [13] proved an upper bound for blow up time of solutions eq. (5).

Di et al. [5] considered the following pseudo-parabolic equation with variable exponent

$$(6) \quad u_t - \Delta u_t - \operatorname{div} \left(|\nabla u|^{m(x)-2} \nabla u \right) = |u|^{p(x)-2} u.$$

They proved an upper bound and lower bound for blow up time. Later, some authors studied blow up of solutions of the equation (6) (see [8, 15]).

Liu [9] studied the $p(x)$ -biharmonic heat equation

$$u_t + \Delta_{p(x)}^2 u = |u|^{q(x)-2} u.$$

The author proved the local existence and blow up of solutions. Ferreira et al. [7] considered the beam-equation with a strong damping and the $p(x)$ -biharmonic operator

$$u_{tt} + \Delta_{p(x)}^2 u - \Delta u_t + f(x, t, u_t) = g(x, t).$$

They proved the local and global existence of solutions. Some other researchers considered the parabolic-type equations with variable exponents (see [3, 10, 11]).

The problems with variable exponents arise in many branches of sciences such as electrorheological fluids, nonlinear elasticity theory and image processing [4, 6, 14].

Motivated by the above studies, in this paper, we consider the blow up of the solution (1) under some conditions.

The present paper is structured as follows. In Section 2, we state some results about the variable exponent $L^{p(x)}(\Omega)$ Lebesgue and $W^{m,p(x)}(\Omega)$ Sobolev spaces. In Section 3, the blow up phenomena will be proved.

2. PRELIMINARIES

We recall some well-known results about the Lebesgue spaces and Sobolev spaces with variable exponents (see [6, 12]).

Let $p : \Omega \rightarrow [1, \infty]$ be a measurable function, where Ω is a bounded domain of R^n . We define the Lebesgue space with variable exponent $p(\cdot)$ by

$$L^{p(x)}(\Omega) = \left\{ u : \Omega \rightarrow R, u \text{ is measurable and } \rho_{p(\cdot)}(\lambda u) < \infty, \text{ for some } \lambda > 0 \right\},$$

where

$$\rho_{p(\cdot)}(u) = \int_{\Omega} |u|^{p(x)} dx.$$

Also endowed with the Luxemburg-type norm

$$\|u\|_{p(x)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{u}{\lambda} \right|^{p(x)} dx \leq 1 \right\},$$

$L^{p(x)}(\Omega)$ is a Banach space.

The Sobolev space with variable exponent $W^{m,p(x)}(\Omega)$ is defined as

$$W^{m,p(x)}(\Omega) = \left\{ u \in L^{p(x)}(\Omega) : D^{\alpha}u \in L^{p(x)}(\Omega), |\alpha| \leq m \right\}.$$

Sobolev space with variable exponent is a Banach space with respect to the norm

$$\|u\|_{2,p(x)} = \|u\|_{p(x)} + \|\nabla u\|_{p(x)} + \|\Delta u\|_{p(x)}.$$

Lemma 1 ([2]). (i) *If (4) holds, then $\|u\|_{p(\cdot)} \leq C \|\nabla u\|_{p(\cdot)}$ for all $u \in W_0^{1,p(\cdot)}(\Omega)$, where Ω is bounded. In particular, the space $W_0^{1,p(\cdot)}(\Omega)$ has a norm given by $\|u\|_{1,p(\cdot)} = \|\nabla u\|_{p(\cdot)}$, for all $u \in W_0^{1,p(\cdot)}(\Omega)$.*

(ii) *If $p \in C(\overline{\Omega})$, $q : \Omega \rightarrow [1, \infty)$ is a measurable function and*

$$\text{ess inf}(p^*(x) - q(x)) > 0,$$

with $p^(x) = \frac{np(x)}{(n-p(x))_+}$ then*

$$W_0^{1,p(\cdot)}(\Omega) \hookrightarrow L^{q(\cdot)}(\Omega).$$

3. BLOW UP

In this part, we study blow-up result of solutions. We first state a local existence theorem [1].

Theorem 1. *For all $u_0 \in W_0^{1,p(\cdot)}(\Omega)$, there exists a number $T_0 \in (0, T]$ such that the problem (1) has a strong solution u on $[0, T_0]$ satisfying*

$$u \in C_w([0, T_0]; W_0^{1,p(\cdot)}(\Omega)) \cap C([0, T_0], L^{q(\cdot)}(\Omega)) \cap W^{1,2}(0, T_0; L^2(\Omega)).$$

Lemma 2.

$$E(t) = \int_{\Omega} \left(\frac{1}{p(x)} |\Delta u|^{p(x)} - \frac{1}{q(x)} |u|^{q(x)} \right) dx$$

is a nonincreasing function for $t \geq 0$ and

$$E'(t) \leq 0.$$

Proof. Multiplying u_t on two sides of the equation (1), and integrating by parts, we get

$$\int_{\Omega} u_t^2 dx + \int_{\Omega} \Delta u_t^2 dx + \frac{d}{dt} \int_{\Omega} \frac{1}{p(x)} |\Delta u|^{p(x)} dx = \frac{d}{dt} \int_{\Omega} \frac{1}{q(x)} |u|^{q(x)} dx.$$

We then define the energy by

$$E(t) = \int_{\Omega} \left(\frac{1}{p(x)} |\Delta u|^{p(x)} - \frac{1}{q(x)} |u|^{q(x)} \right) dx.$$

Clearly, we get

$$E'(t) = - \int_{\Omega} u_t^2 dx - \int_{\Omega} |\Delta u_t|^2 dx \leq 0.$$

Let $H(t) = -E(t)$. So, $H'(t) \geq 0$. □

Theorem 2. Let $u_0 \in W_0^{1,m(\cdot)}(\Omega)$ such that $\int_{\Omega} u_0^2 dx + \int_{\Omega} \Delta^2 u_0 dx > 0$ and

$$\int_{\Omega} \left(\frac{1}{p(x)} |\Delta u_0|^{p(x)} - \frac{1}{q(x)} |u_0|^{q(x)} \right) dx \geq 0.$$

Then

$$F(t) = \frac{1}{2} \left(\int_{\Omega} u^2 dx + \int_{\Omega} |\Delta u|^2 dx \right)$$

blows up in finite time $t^* < +\infty$.

Proof. By differentiating F with respect to t , we obtain

$$\begin{aligned} F'(t) &= \int_{\Omega} (uu_t + \Delta u \Delta u_t) dx \\ &= \int_{\Omega} \left[u \left(-\Delta^2 u_t + \operatorname{div} \left(|\Delta u|^{p(x)-2} \Delta u \right) + |u|^{q(x)-2} u \right) + \Delta u \Delta u_t \right] dx \\ &= \int_{\Omega} \left(|u|^{q(x)} - |\Delta u|^{p(x)} \right) dx \\ &= \int_{\Omega} q(x) \left(\frac{|u|^{q(x)}}{q(x)} - \frac{|\Delta u|^{p(x)}}{p(x)} \right) dx \\ &\quad + \int_{\Omega} q(x) \left(\frac{1}{p(x)} - \frac{1}{q(x)} \right) |\Delta u|^{p(x)} dx. \end{aligned}$$

Since $E'(t) \leq 0$, we get

$$\begin{aligned} \int_{\Omega} q(x) \left(\frac{|u|^{q(x)}}{q(x)} - \frac{|\Delta u|^{p(x)}}{p(x)} \right) dx &\geq \int_{\Omega} q(x) \left(\frac{|u_0|^{q(x)}}{q(x)} - \frac{|\Delta u_0|^{p(x)}}{p(x)} \right) dx \geq \\ &q_- \int_{\Omega} \left(\frac{|u_0|^{q(x)}}{q(x)} - \frac{|\Delta u_0|^{p(x)}}{p(x)} \right) dx \geq 0. \end{aligned}$$

We see

$$\begin{aligned} F'(t) &\geq \int_{\Omega} q_- \left[\frac{1}{p^+} - \frac{1}{q^-} \right] |\Delta u|^{p(x)} dx \\ &= C_0 \int_{\Omega} |\Delta u|^{p(x)}. \end{aligned}$$

We define the sets $\Omega_+ = \{x \in \Omega : |\Delta u| \geq 1\}$ and $\Omega_- = \{x \in \Omega : |\Delta u| < 1\}$. So,

$$\begin{aligned} F'(t) &\geq C_0 \left(\int_{\Omega_-} |\Delta u|^{p^+} + \int_{\Omega_+} |\Delta u|^{p^-} \right) \\ &\geq C_1 \left(\left(\int_{\Omega_-} |\Delta u|^2 dx \right)^{p_+/2} + \left(\int_{\Omega_+} |\Delta u|^2 dx \right)^{p_-/2} \right), \end{aligned}$$

using the fact that $\|\Delta u\|_2 \leq C \|\Delta u\|_r$, for all $r \geq 2$.

This implies that

$$(F'(t))^{2/p_+} \geq C_2 \int_{\Omega_-} |\Delta u|^2 dx$$

and similarly,

$$(F'(t))^{2/p_-} \geq C_3 \int_{\Omega_+} |\Delta u|^2 dx.$$

The Poincare inequality gives $\|\Delta u\|^2 \geq \lambda_1 \|u\|^2$, where λ_1 is the first eigenvalue of the problem

$$\begin{cases} \Delta^2 w + \lambda w = 0, & \text{in } \Omega, \\ w = 0, & \text{on } \partial\Omega. \end{cases}$$

Thus, we obtain

$$\begin{aligned} \|\Delta u\|^2 &= \frac{1}{1 + \lambda_1} \|\Delta u\|^2 + \frac{\lambda_1}{1 + \lambda_1} \|\Delta u\|^2 \\ &\geq \frac{\lambda_1}{1 + \lambda_1} \|u\|_{H_0^2}^2. \end{aligned}$$

Simple addition leads to

$$\begin{aligned} (F'(t))^{2/p_-} + (F'(t))^{2/p_+} &\geq (C_3 + C_2) \|\Delta u\|^2 \\ (7) \quad &\geq \frac{\lambda_1(C_3 + C_2)}{1 + \lambda_1} \|u\|_{H_0^2}^2 = C_4 F(t), \end{aligned}$$

or

$$(8) \quad (F'(t))^{2/p_-} \left(1 + (F'(t))^{2(\frac{1}{p_+} - \frac{1}{p_-})}\right) \geq C_4 F(t).$$

By (7) and the fact that $F(t) \geq F(0) > 0$ ($F'(t) \geq 0$), we have, for each $t > 0$, either

$$(F'(t))^{2/p_-} \geq \frac{C_4}{2} F(t) \geq \frac{C_4}{2} F(0)$$

or

$$(F'(t))^{2/p_+} \geq \frac{C_4}{2} F(t) \geq \frac{C_4}{2} F(0),$$

which gives in turn

$$F'(t) \geq C_5 (F(0))^{p_-/2}$$

or

$$F'(t) \geq C_6 (F(0))^{p_+/2}.$$

Hence $F'(t) \geq \alpha$, where $\alpha = \min\{C_5(F(0))^{p_-/2}, C_6(F(0))^{p_+/2}\}$.

Since $\frac{1}{p_+} - \frac{1}{p_-} \leq 0$, (8) yields

$$(F'(t))^{2/p_-} (1 + \alpha)^{2(\frac{1}{p_+} - \frac{1}{p_-})} \geq C_4 F(t), \quad \forall t \geq 0.$$

Consequently,

$$(9) \quad F'(t) \geq \beta F^{p_-/2}(t), \quad \forall t \geq 0.$$

A simple integration of (9) over $(0, t)$ then yields

$$F(t)^{1-\frac{p_-}{2}} \leq F(0)^{1-\frac{p_-}{2}} - \frac{p_- - 2}{2} \beta t,$$

which implies that

$$F(t) \geq \frac{1}{\left(F(0)^{1-\frac{p_-}{2}} - \frac{p_- - 2}{2} \beta t\right)^{\frac{2}{p_- - 2}}}.$$

This shows that F blows up in a time

$$t^* \leq \frac{2F(0)^{1-\frac{p_-}{2}}}{(p_- - 2)\beta}.$$

□

4. CONCLUSION

In this paper, we have studied a $p(x)$ -biharmonic heat equation with a variable. The blow up of solutions has been proved. Our result improves earlier results in the literature.

REFERENCES

- [1] G. Akagi, M. Ôtani, *Evolutions inclusions governed by subdifferentials in reflexive Banach spaces*, Journal of Evolution Equations, 4 (2004) 519–541.
- [2] M.K. Alaoui, S.A. Messaoudi, H.B. Khenous, *A blow up result for nonlinear generalized heat equation*, Computers & Mathematics with Applications, 68 (12) (2014), 1723–1732.
- [3] A. Rahmoune, B. Benabderrahmane, *Bounds for blow-up time in a semilinear parabolic problem with variable exponents*, Studia Universitatis Babeş-Bolyai Mathematica, 67 (2022), 181–188.
- [4] Y. Chen, S. Levine, M. Rao, Variable Exponent, *Linear Growth Functionals in Image Restoration*, SIAM journal on Applied Mathematics, 66 (2006) 1383–1406.
- [5] H. Di, Y. Shang, X. Peng, *Blow-up phenomena for a pseudo-parabolic equation with variable exponents*, Applied Mathematics Letters, 64 (2017) 67–73.
- [6] L. Diening, P. Hasto, P. Harjulehto, M. Ruzicka, *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer, 2011.
- [7] J. Ferreira, W. S. Panni, E. Pişkin, M. Shahrouzi, *Existence of beam-equation solutions with strong damping and $p(x)$ -biharmonic operator*, Mathematica Moravica, 26 (2) (2022) 123–145.
- [8] M. Liao, *Non-global existence of solutions to pseudo-parabolic equations with variable exponents and positive initial energy*, Comptes Rendus Mécanique, 347 (2019), 710–715.
- [9] Y. Liu, *Existence and blow-up of solutions to a parabolic equation with nonstandard growth conditions*, Bulletin of the Australian Mathematical Society, 99 (2) (2019), 242–249.
- [10] E. Pişkin, G. Butakin, *Existence and Decay of solutions for a parabolic-type Kirchhoff equation with variable exponents*, Journal of Mathematical Sciences and Modelling, 6 (1) (2023), 32–41.
- [11] E. Pişkin, Y. Dinç, C. Tuñç, *Lower and Upper Bounds for the Blow up Time for Generalized Heat Equations with Variable Exponents*, Palestine Journal of Mathematics, 10 (2) (2021), 601–608.
- [12] E. Pişkin, B. Okutmuşur, *An Introduction to Sobolev Spaces*, Bentham Science, 2021.
- [13] A. Rahmoune, *Bounds for blow-up time in a nonlinear generalized heat equation*, Applicable Analysis, 101 (6) (2022), 1871–1879.

- [14] M. Ruzicka, *Electrorheological Fluids: Modeling and Mathematical Theory*, Springer Science & Business Media, 2000.
- [15] X. Zhu, B. Guo, M. Liao, *Global existence and blow-up of weak solutions for a pseudo-parabolic equation with high initial energy*, Applied Mathematics Letters, 104 (3) (2020), Article ID: 106270.

ERHAN PIŞKİN

DİCLE UNIVERSITY

DEPARTMENT OF MATHEMATICS

DİYARBAKIR

TURKEY

E-mail address: episkin@dicle.edu.tr

GÜLISTAN BUTAKIN

DİCLE UNIVERSITY

INSTITUTE OF NATURAL AND APPLIED SCIENCES

DİYARBAKIR

TURKEY

E-mail address: gulistanbutakin@gmail.com