

Treatment seeking and antibiotic use for urinary tract infection symptoms in the time of COVID-19 in Tanzania and Uganda

Emmanuel Olamijuwon^{1*},
Katherine Keenan^{1*}, Martha F Mushi²,
Catherine Kansime³, Eveline T Konje²,
Mike Kesby¹, Stella Neema⁴,
Benon Asiiimwe⁵, Stephen E Mshana²,
Kathryn J Fredricks¹, Benjamin Sunday⁶,
Joel Bazira⁶, Alison Sandeman⁷,
Derek J Sloan⁷, Joseph R Mwangi²,
Wilber Sabiti⁷, Matthew TG Holden⁷;
CARE Consortium

¹School of Geography and Sustainable Development,
University of St Andrews, St Andrews, UK

²Catholic University of Health and Allied Sciences,
Mwanza, Tanzania

³School of Public Health, College of Health Sciences,
Makerere University Kampala, Uganda

⁴Department of Sociology and Anthropology, Makerere
University, Kampala, Uganda

⁵Department of Medical Microbiology, Makerere
University College of Health Sciences, Kampala, Uganda

⁶Department of Microbiology, Mbarara University of
Science and Technology, Mbarara, Uganda

⁷School of Medicine, University of St Andrews, St
Andrews, UK

*Joint first authorship.

Correspondence to:

Emmanuel Olamijuwon
Population and Health Research Group, School of Geography
and Sustainable Development, University of St Andrews
Irvine Building, St Andrews KY16 9AL
UK
e.olamijuwon@st-andrews.ac.uk

Katherine Keenan
Population and Health Research Group, School of Geography
and Sustainable Development, University of St Andrews
Irvine Building, St Andrews KY16 9AL
UK
katherine.keenan@st-andrews.ac.uk

Background There is still little empirical evidence on how the outbreak of coronavirus disease 2019 (COVID-19) and associated regulations may have disrupted care-seeking for non-COVID-19 conditions or affected antibiotic behaviours in low- and middle-income countries (LMICs). We aimed to investigate the differences in treatment-seeking behaviours and antibiotic use for urinary tract infection (UTI)-like symptoms before and during the pandemic at recruitment sites in two East African countries with different COVID-19 control policies: Mbarara, Uganda and Mwanza, Tanzania.

Methods In this repeated cross-sectional study, we used data from outpatients (pregnant adolescents aged >14 and adults aged >18) with UTI-like symptoms who visited health facilities in Mwanza, Tanzania and Mbarara, Uganda. We assessed the prevalence of self-reported behaviours (delays in care-seeking, providers visited, antibiotics taken) at three different time points, labelled as 'pre-COVID-19 phase' (February 2019 to February 2020), 'COVID-19 phase 1' (March 2020 to April 2020), and 'COVID-19 phase 2' (July 2021 to February 2022).

Results In both study sites, delays in care-seeking were less common during the pandemic than they were in the pre-COVID phase. Patients in Mwanza, Tanzania had shorter care-seeking pathways during the pandemic compared to before it, but this difference was not observed in Mbarara, Uganda. Health centres were the dominant sources of antibiotics in both settings. Over time, reported antibiotic use for UTI-like symptoms became more common in both settings. During the COVID-19 phases, there was a significant increase in self-reported use of antibiotics like metronidazole (<30% in the pre-COVID-19 phase to 40% in COVID phase 2) and doxycycline (30% in the pre-COVID-19 phase to 55% in COVID phase 2) that were not recommended for treating UTI-like symptoms in the National Treatment Guidelines in Mbarara, Uganda.

Conclusions There was no clear evidence that patients with UTI-like symptoms attending health care facilities had longer

or more complex treatment pathways despite strict government-led interventions related to COVID-19. However, antibiotic use increased over time, including some antibiotics not recommended for treating UTI, which has implications for future antimicrobial resistance.

Antimicrobial resistance (AMR) is a global public health issue which makes infections more costly to treat and common health care interventions such as surgeries potentially life-threatening [1–3]. The burden of drug-resistant infections is estimated to be highest in low- and middle-income regions, particularly Africa [4], where infectious disease accounts for a higher share of morbidity and mortality, and where health care infrastructures are under-resourced compared with high-income regions. It is estimated that by 2050, AMR could cause low-income countries to lose more than 5% of their gross domestic product (GDP) and push millions into poverty [5]. AMR in low- and middle-income countries (LMICs) is driven by multiple factors, ranging from inappropriate use of antibiotics (including over-prescription, over-use of antibiotics and non-adherence to clinical advice [6]) and unprescribed purchase thereof [7], a lack of affordable clinical diagnostics [8,9], environmental factors such as sanitation [10,11], as well as socio-economic drivers like health care costs for patients [12].

Researchers have debated whether the coronavirus disease 2019 (COVID-19) pandemic intersects with the ongoing ‘silent pandemic’ of AMR, especially in LMICs [13–15]. Some suggested that travel restrictions, school, workplace, and non-essential service closures, improved hygiene practices, and enhanced public health information dissemination during the pandemic could decrease infection rates [16], reduce antibiotics use, and slow AMR [15]. In several high-income countries, COVID-19-related restrictions were associated with a substantial reduction in community dispensing of antibiotics primarily used to treat respiratory infections [17–21]. However, disruptions to health care access, financial hardships associated with pandemic restrictions, the fear of catching COVID-19, and the associated stigma may have motivated individuals to seek care outside the formal health care facilities or to self-medicate with antibiotics [13,14,22,23]. In Australia, for example, COVID-19-induced psychological distress was positively related to antibiotic self-medication [24]. In Iran, the fear of catching COVID-19 resulted in increased postponement of dental visits and antibiotic self-medication compared to pre-pandemic periods [25]. In sub-Saharan Africa, attempts to understand the indirect impacts of the pandemic have focused on service delivery, covering areas such as human immunodeficiency virus (HIV), chronic care, maternal, neonatal, and child health [26–33], but with limited evidence on patients’ experiences of care seeking for infections that often require antibiotics. Existing studies have offered mixed findings depending on context and the type of health care service. For example, antenatal and child health care seeking was delayed and reduced in Uganda, and there were periodic shortages of amoxicillin [26]. Conversely, services such as postnatal and HIV care experienced no disruptions in services or reductions in patient numbers [27,34,35].

In this study, we aimed to investigate changes in patient treatment-seeking for urinary tract infection (UTI) during the outbreak of COVID-19 at two sites in East Africa: Mbarara, Uganda and Mwanza, Tanzania. Understanding treatment pathways and antibiotic use in these countries is particularly important because drug sellers commonly sell them for these illnesses without prescription [7,36]. Despite commonalities in health care structures in these neighbouring countries, there were marked differences in government-led COVID-19 related interventions and policies (Figure 1), thus making a valuable comparative case study. The Ugandan government implemented strict household lockdowns and restricted travel and activities for key periods post-2020, while restrictions in Tanzania have been less consistent. Based on other studies from the region [26,30], we anticipated that formal health care use would be delayed and reduced during COVID-19 periods. We also anticipated that government-led COVID-19 prevention efforts would contribute to an increase in self-treatment, use of drug sellers, and related inappropriate antibiotics use, compared with pre-COVID-19. Given that strict lockdowns were more common in Uganda, we anticipated more obvious effects on UTI treatment-seeking there than in Tanzania.

METHODS

Data

In this repeated cross-sectional study, we used data collected by the Holistic Approach to Unravelling Antibacterial Resistance (HATUA) and COVID-19 and Antibacterial Resistance in East-Africa (CARE) consortia, which use UTI – a commonly occurring bacterial infection – as a clinical prism to investigate the drivers of




Figure 1. Timeline of recruitment and COVID-19 restrictions at the two study sites. The shaded area and brown-coloured texts refer to the period of data collection at both study sites.

antibacterial resistance [37]. The consortia recruited patients presenting with UTI-like symptoms at health facilities (including government and private/faith-based-run health facilities) in Kenya, Tanzania, and Uganda. The facilities ranged from community health centres (levels 2–3) to regional referral/zonal referral hospitals (levels 4/5). In this study, we used a subset of data collected at health facilities in Mwanza, Tanzania, and Mbarara, Uganda over three time points between January 2019 and February 2022 and labeled these as ‘pre-COVID-19 phase’ (February 2019 to February 2020), ‘COVID-19 phase 1’ (March 2020 to April 2020), and ‘COVID-19 phase 2’ (July 2021 to February 2022) (Figure 1). Our data collection periods also coincided with the pre-COVID-19 period, the early months of the pandemic, and 15–22 months later. Using standardised recruitment protocols and tools, doctors identified patients with symptomatic UTI for inclusion in the study. Patients answered a questionnaire on treatment-seeking for UTI-like symptoms, antibiotic use practices, and socio-demographic characteristics. Our analysis sample comprised 3666 individuals recruited in three waves (2408 in Mwanza, Tanzania and 1258 in Mbarara, Uganda).

Variables

Socio-demographic characteristics

Age was categorised into young adults (14–34 years), middle-aged adults (35–54 years), and older adults (≥ 55 years). We coded sex as male/female and highest educational attainment into ‘no formal education,’ ‘primary,’ ‘secondary,’ and ‘tertiary.’

Treatment seeking

All patients responded to a structured questionnaire (detailed elsewhere [38]) on their treatment-seeking behaviours for their recent UTI-like symptoms. We first asked them how long it took them to seek care after the onset of their UTI-like symptoms and coded individuals as having ‘delayed treatment’ if they waited for two weeks or more and ‘no delay’ if two weeks or less. The patients provided details on the treatment steps before their current visit to the health facility, including the type of provider (government clinic, private clinic, drug shop or pharmacy, self-treated at home) and type of medication taken. Patients reported medications using a drug card/drug bag method [39], which we used to create a binary response variable indicating whether antibiotics had been taken before coming to the recruitment health facility.

Statistical analysis

We used descriptive statistics to summarise the patients' sociodemographic characteristics, treatment behaviours, and antibiotics most commonly taken before the current health care visit. We used (χ^2) tests of homogeneity to compare their demographic characteristics and behaviours patterns across the different study phases and sites. We used chord diagrams to visualise more detailed flows of patient treatment seeking. Chord diagrams are a visualisation method used in migration studies [40–44] and population health research [45,46] that represents connections between several nodes as curved arcs within a circle. Each node in the chord diagram is a treatment attempt with a certain type of provider (that is, government clinic, private clinic, drug shops/pharmacies, self-treatment) and accounts for whether antibiotics were taken. We produced diagrams using the 'circlize' package in R [47]. We also fitted multivariable logistic regression models to assess significant differences in antibiotic use behaviour while adjusting for sociodemographic characteristics, including age, sex, education, and investigated variations by interacting country/time period. From the regression coefficients, we generated predicted probabilities and compared these across the three periods and countries. Finally, we compared patterns and trends in the use of specific antibiotics across the periods and countries, paying particular attention to antibiotics recommended for use in National Treatment Guidelines for UTI [48,49]. All statistical analyses were done in R, version 4.2.1. (R Core Team, Vienna, Austria) within the RStudio environment, version 2022.12.0+353 (Posit, PBC, Boston, Massachusetts, USA) [50,51]. We considered $P < 0.05$ as statistically significant.

RESULTS

Socio-demographic profile of patients

The socio-demographic characteristics of patients varied across the study periods in both sites (Table 1). About 13% of the patients recruited during the pre-COVID-19 phase in Mwanza, Tanzania were aged >55,

Table 1. Descriptive statistics on patients recruited from both sites – presented as percentages

	Mwanza, Tanzania			P-value†	Mbarara, Uganda			P-value†
	Pre-COVID-19 phase (n = 1295)	COVID-19 phase 1 (n = 231)	COVID-19 phase 2 (n = 882)		Pre-COVID-19 phase (n = 506)	COVID-19 phase 1 (n = 170)	COVID-19 phase 2 (n = 582)	
Age				<0.001				0.009
Young adults (18–34)	58.4	42.0	48.1		53.2	64.7	51.4	
Middle-aged adults (35–54)	28.6	34.6	30.4		37.9	30.0	36.6	
Older adults (55+)	13.1	23.4	21.5		8.9	5.3	12.0	
Gender				<0.001				0.210
Male	17.3	32.0	27.8		20.0	25.3	23.7	
Female	82.7	68.0	72.2		80.0	74.7	76.3	
Highest educational attainment				<0.001				<0.001
No formal education	10.3	19.9	7.82		14.2	14.7	8.93	
Primary	57.7	44.6	61.2		57.1	47.1	65.1	
Secondary	25.9	24.2	24.9		21.5	22.9	19.9	
Tertiary	6.10	11.3	6.01		7.11	15.3	6.0	
Delayed seeking treatment				<0.001				<0.001
No delay/less than two weeks delay	67.2	73.2	87.2		51.6	67.6	82.0	
Delay of more than two weeks	32.8	26.8	12.8		48.4	32.4	18.0	
First thing tried				<0.001				<0.001
Came to this clinic today	35.4	38.5	67.7		26.1	35.9	28.5	
Government clinic	39.6	30.7	20.4		33.2	33.5	38.5	
Private clinic	12.8	12.6	5.1		27.7	18.8	27.8	
Drug shop or pharmacy	8.3	10.4	3.3		5.5	3.5	1.7	
Self-treatment	3.9	7.8	3.5		7.5	8.3	3.4	
Taken antibiotics anytime in the pathway*				<0.001				<0.001
No	58.1	40.8	46.0		41.7	31.2	19.2	
Yes	41.9	59.2	54.0		58.3	68.8	80.8	

*Sample excludes patients who came straight to the health facility to treat the reported symptoms.

† χ^2 tests of homogeneity used to examine whether the demographic characteristics and behaviour patterns of patients follow the distribution across the different study phases.

compared to about 23% and 22% of patients recruited in COVID-19 phases 1 and 2, respectively. In the same study site, the proportion of patients with both the lowest and highest educational attainment nearly doubled in the COVID-19 phase 1, compared to the pre-COVID-19 period; however, in COVID-19 phase 2, educational distribution had returned to levels similar to pre-COVID-19 patterns. We observed a similar pattern in Mbarara, Uganda: there were proportionally more highly educated patients in COVID-19 phase 1, but this returned to pre-COVID-19 levels in phase 2. In Mwanza, Tanzania, there were proportionally more males in the patient samples recruited during the pandemic than in the pre-pandemic period.

Treatment-seeking pathways

We also noted differences in the treatment behaviours of patients recruited across the study periods (Table 1). During the pre-COVID-19 period, about 33% of patients in Mwanza, Tanzania and 48% of patients in Mbarara, Uganda had delays of two weeks or more, while fewer patients had delays during COVID-19 phase 2 (13% Mbarara, Uganda and 18% Mwanza, Tanzania). In Mwanza, Tanzania, the proportion of patients who came directly to the clinic for their symptoms increased across the study phases, while the proportion of patients who had taken antibiotics prior to presenting at a clinic increased over the study periods in both study sites.

We visualised the sequence of steps taken according to types of providers and types of medicine across sites and recruitment periods in the chord diagram (Figure 2 and Figure S1 in the Online Supplementary Document) and the related percentages separately (Table S1–S2 in the Online Supplementary Document). Considering Mwanza, Tanzania only (Figure 2, Panels A–C), we found that treatment-seeking complexity declined over the three periods, evidenced by a declining percentage of patients who had taken multiple steps to treat their UTI symptoms and an increase in those coming straight to the clinic. We did not observe this trend in Mbarara, Uganda. Regardless of the site or period, most respondents went to govern-




Figure 2. Treatment-seeking pathways during COVID-19. The chords represent percentages of the total sample, so the wider the circle's segments, the higher the percentage of patients taking those steps. The outer gray ring distinguishes the different steps (steps 1–3, and last step). The second ring shows the relative share of patients who accessed treatment from that provider overall. Finally, the innermost coloured bars depict the share of patients moving from their current step to another one.




Figure 3. Predicted probabilities of taking ABs anytime in the pathway among patients who tried treating their UTI symptoms before coming to the recruitment clinic. PP – predicted probabilities, CI – confidence interval.

in the COVID-19 phases were more likely to have taken antibiotics for UTI-like symptoms prior to recruitment to the study (Figure 3). In Mbarara, Uganda, the adjusted predicted probability of taking antibiotics anytime in the pathway increased (53% pre-COVID-19 phase, 64% COVID-19 phase 1 and 78% COVID-19 phase 2). Similar increases were observed in Mwanza, Tanzania (38% pre-COVID-19 phase, 58% COVID-19 phase 1 and 56% during COVID-19 phase 2). Over time, there was a significant increase in the use of antibiotics not recommended for treating UTIs in national treatment guidelines. In Mbarara, Uganda, for example, metronidazole use increased from 30% to 53% and doxycycline from 28% to 40% between the pre-COVID-19 phase and COVID-19 phase 2. In Mwanza, Tanzania, the use of ampicillin increased from 3% to 7% and azithromycin from 4% to 8% between the pre-COVID-19 phase and COVID-19 phase 1 (Figure 4).

ment and private health centres as their first point of care. However, over time, there were differences in the share of patients choosing clinics over self-care. In Mwanza, Tanzania, the proportion of patients who self-treated at home doubled (from 4%–8%) between the pre-COVID-19 and COVID-19 phase 1 period, but declined to 4% by COVID-19 phase 2. In Mbarara, Uganda, however, the proportion of patients who had self-treated at home during COVID-19 phase 2 declined significantly to 3% from 8% during the pre-COVID-19 and COVID-19 phases.

Antibiotic use patterns before and during COVID-19

After adjusting for the socio-demographic characteristics at both sites, patients recruited




Figure 4. Distribution of patients by the type of antibiotics they took to treat their UTI symptoms. Mwanza, TZ: pre-COVID-19 phase (n=836); Mwanza, TZ: COVID-19 phase 1 (n=142); Mwanza, TZ: COVID-19 phase 2 (n=285); Mbarara, UG: pre-COVID-19 phase (n=374); Mbarara, UG: COVID-19 phase 1 (n=109); Mbarara, UG, COVID-19 phase 2 (n=416). P-values are derived from χ^2 tests.

DISCUSSION

In this study, we investigated changes in treatment-seeking behaviour for UTI-like symptoms between pre-COVID-19 and the pandemic period in Uganda and Tanzania. We observed that patients who visited the recruitment clinics during the pandemic were less likely to have delayed treatment seeking and overall took fewer treatment steps, particularly in Mwanza, Tanzania compared to Mbarara, Uganda. Reduced delays in seeking care during COVID-19 could have resulted from patients fearing going to other community providers like drug sellers, wanting to take more direct routes to care, or having fewer barriers than usual to seeking timely care at facilities. The finding is not completely consistent across outcomes and time periods. For example, we observed marginal increases in reported rates of self-treatment and use of drug sellers in Mwanza, Tanzania but not Mbarara, Uganda during the COVID-19 phase 1. However, the overall trend of treatment pathways becoming less complex over the pandemic periods is somewhat contrary to our expectations. This could be explained in several ways. Taking the data at face value, it could be that UTI care was one of the less disrupted services. While there were documented interruptions in antenatal care and sexual health services in Uganda in 2020 [26], other services such as postnatal care [26], maternal and child health services [52,53] were seemingly not interrupted, while studies from other parts of sub-Saharan Africa document stability in health care provision and consistent patient attendance [30,32,54]. Additionally, our findings could also have been affected by selection bias. By using data from health care facility attendees, we likely underestimate the extent of self-medication and use of other providers because self-mediators whose symptoms resolve easily do not appear in our sample.


Although this simplification of patient care-seeking might predict more successful treatment outcomes, the increase in overall antibiotic use during the COVID-19 periods at both sites suggests otherwise. As a very prevalent disease commonly treated with antibiotics, data on UTI might be suggestive of community-level antibiotic use and drug resistance. Given the common misuse of antibiotics highlighted previously [7], the observed increase in antibiotic use among UTI patients is a cause for concern and may be suggestive of increasing AMR.

We also observed a significant increase in the proportion of patients treating UTI symptoms with antibiotics, such as metronidazole and doxycycline, that were not recommended for treating UTI in either country [48,49,55]. This increase was more obvious in Mbarara, Uganda, which experienced stricter government-led restrictions than Mwanza, Tanzania, and where antibiotics are commonly sold on demand [36]. Further investigation is necessary to determine if these changes are related to the pandemic or represent more generalised shifts in the prevalence of UTI in the population or medication use over time. Whatever the underlying drivers, this has important implications for AMR, as it has been well established that longer duration and multiple courses of antibiotic treatment further drive AMR [56].

The study has some limitations. First, our COVID-19 data collection periods did not coincide exactly with severe lockdowns (Figure 1), so patients in our study may have felt fewer concerns about attending health care facilities and not faced serious barriers. Our data collection periods in UG and TZ were slightly different, which could inhibit comparability. Moreover, the COVID-19 phase 1 data collection period covered a few months, while the comparable data collection periods were longer. Although continuous data collection could have solved these issues, it was not practical due to resource constraints and COVID-19 restrictions. Treatment-seeking data were self-reported and could be subject to response bias, and confusion on the part of patients about symptoms and medicines. Finally, our study design was observational and thus did not allow any causal inference for us to attribute behavioural patterns to the COVID-19 restrictions.

CONCLUSIONS

Treatment pathways for UTI care were less complex during the pandemic compared to the pre-pandemic period. However, we found evidence of increased antibiotics use, especially of those antibiotics not recommended for the treatment of UTI symptoms. Given what is known about the sub-optimal dispensing and use of antibiotics, both could further accelerate AMR. Our results emphasise the need to further investigate factors driving antibiotic use and AMR in these settings and to ensure antibiotic stewardship is resilient to future disease outbreaks.



Acknowledgments: We would like to thank the CARE field teams in Mbarara, Uganda and Mwanza, Tanzania, for conducting the survey during the COVID-19 pandemic.

Ethics statement: The study received ethical approval from the University of St Andrews, UK (number MD15722, 31/08/21); National Institute for Medical Research, Tanzania (number 2831, updated 17/06/21); CUHAS/BMC Research Ethics and Review Committee (number CREC /266/2018, updated on 09/02/2021); Uganda National Council for Science and Technology (number HS1519ES, 22/07/21); and Makerere University, Uganda (number SBS-2021-31, 24/05/21). Participants were informed about the study aims and written informed consent was obtained from all patients prior to interview and sample collection.

Data availability: Data supporting the reported results are available upon reasonable request from the authors.

Funding: CARE: COVID-19 and Antimicrobial Resistance in East Africa – impact and response is a Global Effort on COVID-19 (GECO) Health Research Award (MR/V036157/1) funded by UK Research and Innovation (Medical Research Council) and the Department of Health and Social Care (National Institute for Health Research).

Authorship contributions: EO conceptualised the paper, analysed quantitative data, visualized the results and wrote the manuscript. KK conceptualised the paper, co-designed the study, supervised data analysis, wrote the manuscript, and had the final responsibility for the decision to submit for publication. MFM supervised data collection and reviewed the manuscript. CK supervised data collection and reviewed the manuscript. ETK supervised data analysis and reviewed the manuscript. MK co-designed the study and reviewed the manuscript. SN co-designed the study, supervised data collection, and reviewed the manuscript. BA co-designed the study, supervised data collection, and reviewed the manuscript. SEM co-designed the study, supervised data collection, and reviewed the manuscript. KJF reviewed the manuscript. BS supervised data collection and reviewed the manuscript. JB supervised data collection and reviewed the manuscript. AS coordinated the study and reviewed the manuscript. DJS co-designed the study, supervised data analysis, and reviewed the manuscript. JRM co-designed the study, supervised data collection, and reviewed the manuscript. WS co-designed the study and reviewed the manuscript. MTGH led the CARE Consortium and reviewed the manuscript.

Disclosure of interest: The authors completed the ICMJE Disclosure of Interest Form (available upon request from the corresponding author) and disclose no relevant interests.

Additional material

Online Supplementary Document

- O'Neill J. Antimicrobial Resistance: Tackling a crisis for the health and wealth of nations. UK: HM Government; 2014. Available: <https://wellcomecollection.org/works/rdpck35v>. Accessed: 10 January 2024.
- Velin L, Umutesi G, Riviello R, Muwanguzi M, Bebell LM, Yankurije M, et al. Surgical site infections and antimicrobial resistance after cesarean section delivery in rural Rwanda. *Ann Glob Health*. 2021;87:77. Medline:34430227 doi:10.5334/aogh.3413
- Lakoh S, Yi L, Sevalie S, Guo X, Adekanmbi O, Smalle IO, et al. Incidence and risk factors of surgical site infections and related antibiotic resistance in Freetown, Sierra Leone: a prospective cohort study. *Antimicrob Resist Infect Control*. 2022;11:39. Medline:35189952 doi:10.1186/s13756-022-01078-y
- Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*. 2022;399:629–55. Medline:35065702 doi:10.1016/S0140-6736(21)02724-0
- World Bank. Drug-Resistant Infections: A threat to our economic future. USA: World Bank; 2017. Available: <https://documents.worldbank.org/curated/en/323311493396993758/final-report>. Accessed: 10 January 2024.
- Sharma A, Singh A, Dar MA, Kaur RJ, Charan J, Iskandar K, et al. Menace of antimicrobial resistance in LMICs: Current surveillance practices and control measures to tackle hostility. *J Infect Public Health*. 2022;15:172–81. Medline:34972026 doi:10.1016/j.jiph.2021.12.008
- Ndaki PM, Mushi MF, Mwanga JR, Konje ET, Ntinginya NE, Mmbaga BT, et al. Dispensing Antibiotics without Prescription at Community Pharmacies and Accredited Drug Dispensing Outlets in Tanzania: A Cross-Sectional Study. *Antibiotics (Basel)*. 2021;10:1025. Medline:34439074 doi:10.3390/antibiotics10081025
- Pearson M, Chandler C. Knowing antimicrobial resistance in practice: a multi-country qualitative study with human and animal healthcare professionals. *Glob Health Action*. 2019;12:1599560. Medline:31294679 doi:10.1080/16549716.2019.1599560
- Gebretekle GB, Mariam DH, Abebe W, Amogne W, Tenna A, Fenta TG, et al. Opportunities and barriers to implementing antibiotic stewardship in low and middle-income countries: Lessons from a mixed-methods study in a tertiary care hospital in Ethiopia. *PLoS One*. 2018;13:e0208447. Medline:30571688 doi:10.1371/journal.pone.0208447
- Samreen, Malak HA, Abulreesh HH. Environmental antimicrobial resistance and its drivers: a potential threat to public health. *J Glob Antimicrob Resist*. 2021;27:101–11. Medline:34454098 doi:10.1016/j.jgar.2021.08.001
- Nadimpalli ML, Marks SJ, Montealegre MC, Gilman RH, Pajuelo MJ, Saito M, et al. Urban informal settlements as hotspots of antimicrobial resistance and the need to curb environmental transmission. *Nat Microbiol*. 2020;5:787–95. Medline:32467623 doi:10.1038/s41564-020-0722-0
- Green DL, Keenan K, Huque SI, Kesby M, Mushi MF, Kansime C, et al. The Role of Multidimensional Poverty in Antibiotic Misuse: A Study of Self-Medication and Non-Adherence in Kenya, Tanzania, and Uganda. *Lancet Glob Health*. 2023;11:e59–68. Medline:36521953 doi:10.1016/S2214-109X(22)00423-5

- 13 Arshad M, Mahmood SF, Khan M, Hasan R. Covid-19, misinformation, and antimicrobial resistance. *BMJ*. 2020;371:m4501. Medline:33234500 doi:10.1136/bmj.m4501
- 14 Knight GM, Glover RE, McQuaid CF, Oлару ID, Gallandat K, Leclerc QJ, et al. Antimicrobial resistance and covid-19: Intersections and implications. *eLife*. 2021;10:1–27. Medline:33588991 doi:10.7554/eLife.64139
- 15 Collignon P, Beggs JJ. CON: COVID-19 will not result in increased antimicrobial resistance prevalence. *JAC Antimicrob Resist*. 2020;2:dlaa051. Medline:34192249 doi:10.1093/jacamr/dlaa051
- 16 Amegah AK. Improving handwashing habits and household air quality in Africa after COVID-19. *Lancet Glob Health*. 2020;8:e1110–1. Medline:32827474 doi:10.1016/S2214-109X(20)30353-3
- 17 Gillies MB, Burgner DP, Ivancic L, Nassar N, Miller JE, Sullivan SG, et al. Changes in antibiotic prescribing following COVID-19 restrictions: Lessons for post-pandemic antibiotic stewardship. *Br J Clin Pharmacol*. 2022;88:1143–51. Medline:34405427 doi:10.1111/bcp.15000
- 18 Zhu N, Aylin P, Rawson T, Gilchrist M, Majeed A, Holmes A. Investigating the impact of COVID-19 on primary care antibiotic prescribing in North West London across two epidemic waves. *Clin Microbiol Infect*. 2021;27:762–8. Medline:33601010 doi:10.1016/j.cmi.2021.02.007
- 19 Malcolm W, Seaton RA, Haddock G, Baxter L, Thirlwell S, Russell P, et al. Impact of the COVID-19 pandemic on community antibiotic prescribing in Scotland. *JAC Antimicrob Resist*. 2020;2:dlaa105. Medline:34192254 doi:10.1093/jacamr/dlaa105
- 20 Selke Krulichová I, Selke GW, Bennie M, Hajiebrahimi M, Nyberg F, Fürst J, et al. Comparison of drug prescribing before and during the COVID-19 pandemic: A cross-national European study. *Pharmacoepidemiol Drug Saf*. 2022;31:1046–55. Medline:35791700 doi:10.1002/pds.5509
- 21 Gagliotti C, Buttazzi R, Ricchizzi E, Di Mario S, Tedeschi S, Moro ML. Community use of antibiotics during the COVID-19 lockdown. *Infect Dis (Lond)*. 2021;53:142–4. Medline:33073645 doi:10.1080/23744235.2020.1834139
- 22 Subramanya SH, Czyż DM, Acharya KP, Humphreys H. The potential impact of the COVID-19 pandemic on antimicrobial resistance and antibiotic stewardship. *Virusdisease*. 2021;32:330–7. Medline:34056051 doi:10.1007/s13337-021-00695-2
- 23 Ahmed SAKS, Ajisola M, Azeem K, Bakibinga P, Chen YF, Choudhury NN, et al. Impact of the societal response to COVID-19 on access to healthcare for non-COVID-19 health issues in slum communities of Bangladesh, Kenya, Nigeria and Pakistan: results of pre-COVID and COVID-19 lockdown stakeholder engagements. *BMJ Glob Health*. 2020;5:e003042. Medline:32819917 doi:10.1136/bmjgh-2020-003042
- 24 Zhang A, Hobman EV, De Barro P, Young A, Carter DJ, Byrne M, et al. Self-Medication with Antibiotics for Protection against COVID-19: The Role of Psychological Distress, Knowledge of, and Experiences with Antibiotics. *Antibiotics*. 2021;10:232. Medline:33668953 doi:10.3390/antibiotics10030232
- 25 Khami MR, Zadeh AG, Rahi D. A cross-sectional study on COVID-19-related changes in self-medication with antibiotics. *PLoS One*. 2022;17:e0269782. Medline:35700193 doi:10.1371/journal.pone.0269782
- 26 Burt JF, Ouma J, Lubyayi L, Amone A, Aol L, Sekikubo M, et al. Indirect effects of COVID-19 on maternal, neonatal, child, sexual and reproductive health services in Kampala, Uganda. *BMJ Glob Health*. 2021;6:e006102. Medline:34452941 doi:10.1136/bmjgh-2021-006102
- 27 Tessema GA, Kinfu Y, Dachew BA, Gebresilassie Tesema A, Assefa Y, Addis Alene K. The COVID-19 pandemic and health-care systems in Africa: a scoping review of preparedness, impact and response. *BMJ Glob Health*. 2021;6:e007179. Medline:34853031 doi:10.1136/bmjgh-2021-007179
- 28 Bell D, Hansen KS, Kiragga AN, Kambugu A, Kissa J, Mbonye AK. Predicting the Impact of COVID-19 and the Potential Impact of the Public Health Response on Disease Burden in Uganda. *Am J Trop Med Hyg*. 2020;103:1191–7. Medline:32705975 doi:10.4269/ajtmh.20-0546
- 29 Ammor YM, Kaiitouni ZI, Darfaoui M, Lalya I, Elomrani A, Khouchani M. Managing cancer patients during COVID-19 pandemic: a North African oncological center experience. *Pan Afr Med J*. 2020;35:144. Medline:33193959 doi:10.11604/pamj.supp.2020.35.2.24582
- 30 Abdela SG, Berhanu AB, Ferede LM, van Griensven J. Essential Healthcare Services in the Face of COVID-19 Prevention: Experiences from a Referral Hospital in Ethiopia. *Am J Trop Med Hyg*. 2020;103:1198–200. Medline:32762799 doi:10.4269/ajtmh.20-0464
- 31 Jensen C, McKerrow NH. Child health services during a COVID-19 outbreak in KwaZulu-Natal Province, South Africa. *S Afr Med J*. 2020;0:13185. Medline:33334393
- 32 Siedner MJ, Kraemer JD, Meyer MJ, Harling G, Mngomezulu T, Gabela P, et al. Access to primary healthcare during lockdown measures for COVID-19 in rural South Africa: an interrupted time series analysis. *BMJ Open*. 2020;10:e043763. Medline:33020109 doi:10.1136/bmjopen-2020-043763
- 33 Mohammed H, Oljira L, Roba KT, Yimer G, Fekadu A, Manyazewal T. Containment of COVID-19 in Ethiopia and implications for tuberculosis care and research. *Infect Dis Poverty*. 2020;9:131. Medline:32938497 doi:10.1186/s40249-020-00753-9
- 34 Enbiale W, Abdela SG, Seyum M, Hundie DB, Bogale KA, Tamirat KS, et al. Effect of the COVID-19 Pandemic Preparation and Response on Essential Health Services in Primary and Tertiary Healthcare Settings of Amhara Region, Ethiopia. *Am J Trop Med Hyg*. 2021;105:1240–6. Medline:34544048 doi:10.4269/ajtmh.21-0354
- 35 Desta AA, Woldearegay TW, Gebremeskel E, Alemayehu M, Getachew T, Gebregzabihier G, et al. Impacts of COVID-19 on essential health services in Tigray, Northern Ethiopia: A pre-post study. *PLoS One*. 2021;16:e0256330. Medline:34449773 doi:10.1371/journal.pone.0256330
- 36 Olamijuwon E, Konje E, Kansiiime C, Kesby M, Keenan K, Neema S, et al. Antibiotic dispensing practices during COVID-19 and implications for Antimicrobial Resistance (AMR): Parallel mystery client studies in Uganda and Tanzania. 2023;12:10.

- 37 Asimwe BB, Kiiru J, Mshana SE, Neema S, Keenan K, Kesby M, et al. Protocol for an interdisciplinary cross-sectional study investigating the social, biological and community-level drivers of antimicrobial resistance (AMR): Holistic Approach to Unravel Antibacterial Resistance in East Africa (HATUA). *BMJ Open*. 2021;11:e041418. Medline:34006022 doi:10.1136/bmjopen-2020-041418
- 38 Keenan K, Fredricks K, Al Ahad MA, Neema S, Mwanga J, Kesby M, et al. Unravelling patient pathways in the context of antibacterial resistance in East Africa. *BMC Infect Dis*. 2023;23:414. Medline:37337134 doi:10.1186/s12879-023-08392-9
- 39 Dixon J, MacPherson E, Manyau S, Nayiga S, Khine Zaw Y, Kayendeke M, et al. The 'Drug Bag' method: lessons from anthropological studies of antibiotic use in Africa and South-East Asia. *Glob Health Action*. 2019;12:1639388. Medline:31339473 doi:10.1080/16549716.2019.1639388
- 40 Sander N, Abel GJ, Bauer R, Schmidt J. Visualising migration flow data with circular plots. Vienna: Austrian Academy of Sciences, Vienna Institute of Demography; 2014. Available: <https://www.econstor.eu/handle/10419/97018>. Accessed: 10 January 2024.
- 41 Qi W, Abel GJ, Muttarak R, Liu S. Circular visualisation of China's internal migration flows 2010–2015. *Environment and Planning A: Economy and Space*. 2017;49:2432–6. doi:10.1177/0308518X17718375
- 42 Azose JJ, Raftery AE. Estimation of emigration, return migration, and transit migration between all pairs of countries. *Proc Natl Acad Sci U S A*. 2019;116:116–22. Medline:30584106 doi:10.1073/pnas.1722334116
- 43 Gomez CJ, Herman AC, Parigi P. Moving more, but closer: Mapping the growing regionalisation of global scientific mobility using ORCID. *J Informetrics*. 2020;14:101044. doi:10.1016/j.joi.2020.101044
- 44 Abel GJ, Cohen JE. Bilateral international migration flow estimates for 200 countries. *Scientific Data*. 2019;6:82. Medline:31209218 doi:10.1038/s41597-019-0089-3
- 45 Aris-Brosou S, Ibeh N, Noël J. Viral outbreaks involve destabilised evolutionary networks: evidence from Ebola, Influenza and Zika. *Sci Rep*. 2017;7:11881. Medline:28928377 doi:10.1038/s41598-017-12268-9
- 46 Pierzynski JA, Clegg JL, Sim J-A, Forrest CB, Robison LL, Hudson MM, et al. Patient-reported outcomes in paediatric cancer survivorship: a qualitative study to elicit the content from cancer survivors and caregivers. *BMJ Open*. 2020;10:e032414. Medline:32423926 doi:10.1136/bmjopen-2019-032414
- 47 Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize implements and enhances circular visualisation in R. *Bioinformatics*. 2014;30:2811–2. Medline:24930139 doi:10.1093/bioinformatics/btu393
- 48 Ugandan Ministry of Health. Uganda Clinical Guidelines 2016: National Guidelines for Management of Common Conditions. Uganda: Ministry of Health; 2016. Available: https://health.go.ug/sites/default/files/Uganda%20Clinical%20Guidelines%20202016_FINAL.pdf. Accessed: 10 January 2024.
- 49 Ministry of Health, Community Development, Gender, Elderly and Children. Standard Treatment Guidelines and National Essential Medicines List For Tanzania Mainland. Tanzania: Ministry of Health, Community Development, Gender, Elderly and Children; 2021.
- 50 R Core Team. R: A Language and Environment for Statistical Computing. 2022. Available: <https://www.r-project.org/>. Accessed: 14 December 2023.
- 51 R Studio Team. RStudio: Integrated Development Environment for R. 2020. Available: <http://www.rstudio.com/>. Accessed: 14 December 2023.
- 52 Pallangyo E, Nakate MG, Maina R, Fleming V. The impact of covid-19 on midwives' practice in Kenya, Uganda and Tanzania: A reflective account. *Midwifery*. 2020;89:102775. Medline:32526597 doi:10.1016/j.midw.2020.102775
- 53 Shayo EH, Khamis N, Id N, Mboera LEG, Id EN, Mangesho P, et al. The impacts of COVID-19 and its policy response on access and utilisation of maternal and child health services in Tanzania: A mixed methods study. *PLOS Glob Public Health*. 2023;3:e0001549. Medline:37172038 doi:10.1371/journal.pgph.0001549
- 54 Shikuku DN, Nyaake IK, Nyaga LN, Ameh CA. Early indirect impact of COVID-19 pandemic on utilisation and outcomes of reproductive, maternal, newborn, child and adolescent health services in Kenya: A cross-sectional study. *Afr J Reprod Health*. 2021;25:76–87. Medline:37585823
- 55 Ndaki PM, Mushi MF, Mwanga JR, Konje ET, Mugassa S, Manyiri MW, et al. Non-prescribed antibiotic dispensing practices for symptoms of urinary tract infection in community pharmacies and accredited drug dispensing outlets in Tanzania: a simulated clients approach. *BMC Prim Care*. 2022;23:287. Medline:36402956 doi:10.1186/s12875-022-01905-6
- 56 Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ*. 2010;340:c2096. Medline:20483949 doi:10.1136/bmj.c2096