



UNIVERSITY OF LEEDS

This is a repository copy of *On the importance of ryanodine receptor subunit cooperativity in the heart*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/206767/>

Version: Accepted Version

Article:

Zhang, X. and Colman, M.A. orcid.org/0000-0003-2817-8508 (2022) On the importance of ryanodine receptor subunit cooperativity in the heart. *Biophysical Journal*, 122 (1). pp. 6-8. ISSN 0006-3495

<https://doi.org/10.1016/j.bpj.2022.11.016>

© 2022, Elsevier. This is an author produced version of an article published in *Biophysical Journal*, made available under the terms of the Creative Commons Attribution License (CC BY-NC-ND), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

New and Notable: On the importance of ryanodine receptor subunit cooperativity in the heart

Xianwei Zhang¹ and Michael A Colman²

¹ Department of Pharmacology, UC Davis, USA

²School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, UK

Intracellular calcium (Ca^{2+}) homeostasis in cardiac myocytes is an important determinant of both contractile and electrophysiological function, being responsible for excitation contraction coupling (ECC) through the process of Ca^{2+} -induced- Ca^{2+} -release (CICR). Cardiomyocytes contain an intracellular structure which facilitates rapid and robust CICR: type 2 ryanodine receptors (RyRs), responsible for Ca^{2+} release from the intracellular Ca^{2+} store, the sarcoplasmic reticulum (SR), are distributed throughout the intracellular volume in clusters closely juxtaposed ($< 20\text{nm}$) with the sarcolemmal membrane and the L-type Ca^{2+} channels which reside therein, forming dyads (Scriven *et al.*, 2013). RyR dysfunction may directly impair CICR, and thus ECC, resulting in inhibited or inefficient cellular contraction; moreover, many experimental reports have indicated that a change of RyR characteristics is a major contributor to arrhythmia associated with various conditions, such as atrial fibrillation (Voigt *et al.*, 2012) and heart failure (Dobrev & Wehrens, 2014).

To understand the role of RyRs at multiple scales (i.e., from molecular- to organ-level), a physiologically accurate mathematical description of RyR kinetics in health and disease is needed. An ideal RyR model should not only replicate the experimental observations of RyR kinetics but also conduct robustly in the context of the multiple interacting components of the subcellular or cell-level computational models of Ca^{2+} homeostasis. Multiple well-established (2-, 3-, and 4-state) RyR Markov models have been commonly used to replicate RyR kinetics (i.e., opening, closing, inactivation, and refractoriness) and systolic/diastolic Ca^{2+} regulation (e.g., SR leak, Ca^{2+} sparks, and Ca^{2+} waves). All the models are designed to replicate part of the RyR kinetics and Ca^{2+} dynamic features, but may be parameterised for different purposes, e.g., maintaining stable Ca^{2+} homeostasis, or reproducing spontaneous Ca^{2+} sparks at appropriate frequency. Despite the fundamental importance of RyR dynamics for cellular function, the development of a mathematical description of the RyR which is both determined from experimental channel observations and simultaneously robustly reproduces the full range of physiological and pathophysiological behaviours remains a major challenge (Colman *et al.*, 2022).

A better understanding of structure-function relationships at the molecular scale may help to develop such a robust description. Combined with more high-resolution RyR structure images in the recent decade (Yuchi & Van Petegem, 2016), RyR conformations with ligand-binding and RyR-RyR interaction are better understood (Williams *et al.*, 2018). In addition, experimental data suggest that the neighboring subunit-subunit interactions in RyR tetrameric structure substantially regulate RyR opening under not only physiology but also pathological conditions (Yuchi & Van Petegem, 2016). However, such subunit interactions have yet to be systematically assessed in regards to impact on channel and cellular function. To explore these interactions, Greene, Lucko and Shiferaw in this issue of *Biophysical Journal* (Greene *et al.*, 2023) present a theoretical study which mathematically models the RyR with four explicit subunits, applied to study single channel dynamics and whole-cell homeostasis.

The theoretical and numerical analysis reveals that 1) the minimal number of open-state subunits for RyR opening, and 2) the balance of cooperative interactions between neighboring subunits, are both important for regulating RyR opening and closure, and thus stable but efficient Ca^{2+} cycling. A major feature of RyR kinetics which was required to be captured was the simultaneous ability to rapidly respond to sub-millisecond changes to Ca^{2+} (i.e., to be triggered on activation of the L-type Ca^{2+} channels to facilitate CICR), to robustly close during electrical excitation to end CICR, and to demonstrate stability (i.e., low noise) at resting (diastolic) Ca^{2+} levels. Such function necessitates a steep dependence on intracellular Ca^{2+} concentration, which could be a result of a minimum number of required subunits to be simultaneously open before a flux is permitted.

The authors found that in the case where three or all four of the subunits were required to be in the open state in order to permit flux, the RyR can be both reliably shut at low Ca^{2+} levels while being able to rapidly respond to high Ca^{2+} concentrations (i.e., be a robust Ca^{2+} sensor) within a relatively wide parameter space of subunit-subunit interaction strengths (Figure); if two subunits were required to be simultaneously open, then the RyR can still reproduce this function but within a much more restricted region of cooperativity parameter space; for just one subunit, it was not possible to simultaneously fulfil all of these criteria.

The study also assessed the dependence of Ca^{2+} spark dynamics on the relationship between cooperativity energy penalties for the open or closed states, revealing that only under appropriate balance of these parameters could robust and self-terminating triggered Ca^{2+} sparks be induced: favouring the open state led to long-lasting (i.e., not self-terminating) Ca^{2+} sparks, whereas favouring the closed state led to unreliable Ca^{2+} sensitivity (sparks were unable to be robustly initiated). The size of the region of this parameter space became narrowed as the number of required subunits was reduced from four to two, and when just one subunit was required, only long-lasting (not self-terminating) sparks could be induced. Finally, spontaneous Ca^{2+} spark frequency was also highly sensitive to both the number of required subunits and the cooperativity parameter space, demonstrating again that the requirement for fewer subunits led to less stable behaviour, reflected in a higher frequency of spontaneous sparks.

By considering simultaneously multiple parameter spaces and features of Ca^{2+} spark dynamics, this study provides valuable insight into the fundamental physiology of RyR structure-function relationships. This paper highlights the importance of keeping proper subunit-subunit interaction energy and the number of open-state subunits for RyR opening. Meanwhile, this paper shows the potential of building a pipeline to study RyR dynamics from protein structure to a thermodynamic-based theoretical model with biophysics properties, similar to recent models developed for other ion channels (e.g., Silva *et al.*, 2009; DeMarco *et al.*, 2021). Overall, this study lends new insight into the effects of RyR subunits' interaction on Ca^{2+} signaling and will benefit the field.

Multi-scale RyR structural changes (i.e., binding-site, subunit-subunit, RyR-RyR, RyR cluster) underly myriad pathophysiological conditions, such as alternans and triggered activity. Meanwhile, the RyR conformations are regulated by varying targets (e.g., EGTA, Ca, Caffeine-ATP) and clustering (e.g., RyR-RyR regulation and subcellular distribution) (Williams *et al.*, 2018; Galice *et al.*, 2018). To understand the effects of RyR structure changes on disease mechanisms, this model could be extended with disease-specific structure information and be integrated into the cardiac cell model with subcellular spatial details.

Future works could incorporate further RyR regulations such as Calsequestrin (Restrepo *et al.*, 2008), RyR-RyR interactions (Sobie *et al.*, 2002), cluster size and distribution (Sutanto *et al.*, 2018; Zhang *et al.*, 2022). Meanwhile, disease-induced changes in RyR properties (e.g., hyperphosphorylation and disrupted distribution) could also be studied under the same pipeline from structural information to Ca^{2+} cycling.

Figure




Figure. Illustration of the relationship between the minimum number of subunits required for release flux and the size of the parameter space of subunit cooperativity that fulfils robust RyR function.

Author contributions

All authors conceived, designed, drafted, and edited the manuscript, and prepared figures.

Declaration of interests

The authors declare no competing interests.

Funding

XZ is supported by an American Heart Association Predoctoral Fellowship (grant no. 20PRE35120465); MAC is supported by Medical Research Council, UK, Career Development Award (grant no. MR/V010050/1).

References

- Colman MA, Alvarez-Lacalle E, Echebarria B, Sato D, Sutanto H & Heijman J (2022). Multi-Scale Computational Modeling of Spatial Calcium Handling From Nanodomain to Whole-Heart: Overview and Perspectives. *Frontiers in Physiology*. Available at: <https://www.frontiersin.org/article/10.3389/fphys.2022.836622> [Accessed March 11, 2022].
- DeMarco KR, Yang P-C, Singh V, Furutani K, Dawson JRD, Jeng M-T, Fettinger JC, Bekker S, Ngo VA, Noskov SY, Yarov-Yarovoy V, Sack JT, Wulff H, Clancy CE & Vorobyov I (2021). Molecular determinants of pro-arrhythmia proclivity of d- and l-sotalol via a multi-scale modeling pipeline. *Journal of Molecular and Cellular Cardiology* **158**, 163–177.
- Dobrev D & Wehrens XHT (2014). Role of RyR2 Phosphorylation in Heart Failure and Arrhythmias. *Circulation Research* **114**, 1311–1319.
- Galice S, Xie Y, Yang Y, Sato D & Bers DM (2018). Size Matters: Ryanodine Receptor Cluster Size Affects Arrhythmogenic Sarcoplasmic Reticulum Calcium Release. *Journal of the American Heart Association* **7**, e008724.

- Greene D, Luchko T & Shiferaw Y (2023). The role of subunit cooperativity on ryanodine receptor 2 calcium signaling. *Biophys J* **122**, 215–229.
- Restrepo JG, Weiss JN & Karma A (2008). Calsequestrin-Mediated Mechanism for Cellular Calcium Transient Alternans. *Biophys J* **95**, 3767–3789.
- Scriven DRL, Asghari P & Moore EDW (2013). Microarchitecture of the dyad. *Cardiovasc Res* **98**, 169–176.
- Silva JR, Pan H, Wu D, Nekouzadeh A, Decker KF, Cui J, Baker NA, Sept D & Rudy Y (2009). A multiscale model linking ion-channel molecular dynamics and electrostatics to the cardiac action potential. *Proceedings of the National Academy of Sciences* **106**, 11102–11106.
- Sobie EA, Dilly KW, dos Santos Cruz J, Lederer WJ & Jafri MS (2002). Termination of cardiac Ca²⁺ sparks: an investigative mathematical model of calcium-induced calcium release. *Biophys J* **83**, 59–78.
- Sutanto H, van Sloun B, Schönleitner P, van Zandvoort MAMJ, Antoons G & Heijman J (2018). The Subcellular Distribution of Ryanodine Receptors and L-Type Ca²⁺ Channels Modulates Ca²⁺-Transient Properties and Spontaneous Ca²⁺-Release Events in Atrial Cardiomyocytes. *Frontiers in Physiology* **9**, 1108.
- Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, Sun Q, Wieland T, Ravens U, Nattel S, Wehrens XHT & Dobrev D (2012). Enhanced Sarcoplasmic Reticulum Ca²⁺ Leak and Increased Na⁺-Ca²⁺ Exchanger Function Underlie Delayed Afterdepolarizations in Patients With Chronic Atrial Fibrillation. *Circulation* **125**, 2059–2070.
- Williams AJ, Thomas NL & George CH (2018). The ryanodine receptor: advances in structure and organization. *Current Opinion in Physiology* **1**, 1–6.
- Yuchi Z & Van Petegem F (2016). Ryanodine receptors under the magnifying lens: Insights and limitations of cryo-electron microscopy and X-ray crystallography studies. *Cell Calcium* **59**, 209–227.
- Zhang X, Smith CER, Morotti S, Edwards AG, Sato D, Louch WE, Ni H & Grandi E (n.d.). Mechanisms of spontaneous Ca²⁺ release-mediated arrhythmia in a novel 3D human atrial myocyte model: II. Ca²⁺-handling protein variation. *The Journal of Physiology*, DOI: 10.1113/JP283602.