



Kronične nezarazne bolesti

Frekvencija srca i njena varijabilnost kod bolesnika s hipertenzijom

(The heart frequency and it's variability in hypertensive patients considering A/B type of behaviour and eight basic emotions and levels of anger expression)

Karlo Kožul (1), Ksenija Vidović (1), Helga Heinzelman-Kožul (2), Senka Samardžić (3), Mirta Kopic (4) and Miroslav Šram (5)

1 Department of Epidemiology, Institute of Public Health for the Osijek-Baranja County, Osijek, Croatia

2 Specialist family medicine practice

3 Department of Social Medicine, Institute of Public Health for the Osijek-Baranja County, Osijek, Croatia

4 Department of Ophthalmology, Clinical hospital Osijek, Osijek, Croatia

5 Clinic for Internal medicine, Clinical hospital Osijek, Osijek, Croatia

Sažetak

U ovom radu istraživali smo autonomnu regulaciju rada srca putem frekvencije srca i njene varijabilnosti kod 100 bolesnika s esencijalnom hipertenzijom i usporedili sa 100 ispitanika zdrave kontrolne grupe. Bolesnici s esencijalnom hipertenzijom imali su značajno kraći prosječni RR interval, odnosno bržu frekvenciju srca nego kontrolna grupa. Prosječna varijabilnost frekvencije srca je statistički značajno niža u hipertenzivnoj, nego u zdravoj kontrolnoj grupi. Na frekvenciju srca i njenu varijabilnost ukupno značajno utječu, pri mirnom i dubokom disanju: sistolički tlak, dijastolički tlak, masa lijevog ventrikula, dijastolička disfunkcija lijevog ventrikula, veličina lijevog atrija, trajanje hipertenzije, visina kolesterola i triglicerida. Najznačajnija negativna povezanost nađena je između varijabilnosti frekvencije srca i trajanja hipertenzije. U hipertenzivnih bolesnika i u kontrolnoj skupini češći je "A tip" nego "B tip" ponašanja. U hipertenzivnoj grupi statistički je značajan ukupan utjecaj svih nezavisnih varijabli po Bortneru na varijabilnost frekvencije srca pri mirnom disanju. Frekvencija srca i njena varijabilnost povezane su sa osam temeljnih emocija kodu hipertenzivnih ispitanika i kontrolne skupine. U grupi hipertenzivnih bolesnika varijabilnost frekvencije srca značajno je pozitivno povezana s emocionalnom dimenzijom "Inkorporacija", a negativno s emocionalnim dimenzijama "Zaštita", "Odbacivanje" i "Reprodukcija". Naši podaci pokazuju da bolesnici s dugotrajnom hipertenzijom imaju povišenu frekvenciju srca i reduciranu varijabilnost frekvencije srca, što su poznati čimbenici rizika za povišeni kardijalni mortalitet.

Abstract

The aim of this paper was to comparison between autonomic regulation of the heart rate and it's variability within 100 essential hypertensive patients and 100 examinees in healthy control group. Essential hypertensive patients had significantly shorter average RR interval, that is, faster heart frequency then the control group. Average variability of the heart frequency was statistically considerably lower in hypertensive than in the healthy control group. Systolic blood pressure, diastolic blood pressure, duration of hypertension, levels have considerate total influence on heart frequency and it's variability during calm and during deep breathing. The most distinctive negative connection was found between variability of the heart frequency and duration of hypertension. The type A behaviour within hypertensive patients and the control group was more common then type B. In the hypertensive group, total influence of all independent Bortner's variables on to variability of the heart frequency during calm breathing was statistically important. Within hypertensive patients and the control group, the heart frequency and it's variability were connected with eight basic emotions. In the group of hypertensive patients variability of the heart frequency is significantly positively connected with emotional dimension "Incorporation", but negatively with emotional dimensions "Protection", "Rejection" and "Reproduction". Our data shows that patients with long-term hypertension have increased heart frequency and reduced variability of the heart frequency, which are well-known risk factors for increased cardial mortality.

Key words: hypertensive patients, A/B type of behaviour, basic emotions

* Ovaj članak je originalno objavljen u časopisu *Collegium Antropologicum*, 33 (2009) 2: 409-416

Introduction

It is estimated that 15-20% of adult population in industrialized high-developed countries, as well as in Croatia¹⁷ in the world have high blood pressure¹⁶. For adults, hypertension is considered to be with systolic blood pressure equal or higher than 21.3 kPa, and/or diastolic equal or higher then 12.0kPa. Untreated hypertension increases appearance of heart failure, coronary heart disease, haemorrhagic and thrombotic stroke, renal insufficiency, aortal dissection and death^{2,3,4,5}. Diagnosed hypertension is psychological and socioeconomical burden to a patient because it almost always demands life-long therapy^{8,9,10}. In arterial hypertension, the heart, at first, shows signs of hyperkinetic syndrome. The heart frequency is normally determined by early diastolic depolarization of the nodus sinoatrialis (heart automatism), function of the automatic nervous system, body temperature, thyroid hormones, metabolism, chatecolamines and neurotransmitters²². Numerous studies confirmed that average heart frequency is considerably higher in hypertensive then in those normotensive individuals, however it's not significantly connected to hypertensive cardiomyopathy²⁸.

Emotional state, i.e. neurogenic factors, affect many functions of cardiovascular system. Heart frequency, blood pressure, peripheral blood vessels succumb to quick changes due to stimulus from autonomic nervous system^{32,33}. There are two types of behaviour, A and B³⁴. Fully developed typ A connotes readiness, impatience, intolerance, agresive and hostile attitude, competitive spirit and constant personal ambition. In other words, type A is emotional complex that can be attributed to individuals in constant struggle to accomplish as much as they can in the shortest of time, or they are in constant conflict with people from their surroundings. Unlike this, type B is mostly released of this features, has no conflict with time and people, therefore no hostility.

According to Plutchnik, there are eight basic or primal emotions which are foundation for reactions of adults. 3 positive emotions are: acceptance (incorporation), joy (reproduction), anticipation (exploration), and 5 negative are, disgust (rejection), sadness (deprivation), fear (protection), surprize (orientation) and anger (destruction). They have adaptational meaning for survival of individuals.

The purpose of our study was to analyze autonomic regulation of heart function in relation to type of behavior, eight basic emotions and Minesota scale anger through frequency and it's variability in hypertensive patients compared to healthy individuals.

Material and Methods

This study was performed at the cardiovascular infirmary of Internal Clinic in Clinical Hospital Osijek, with the approval of the hospital Ethical Committee. For the group of examinees we chose only those hypertensives in which we did not find detectable secondary reasons of high blood pressure and its complications. Hypertensive patients with angina pectoris, myocardial infarction, valvular heart disease, congestive heart disease, diabetes mellitus, renal insufficiency, obstructive lung disease and musculoskeletal disorder had been excluded from the research. One hundred examinees, 40 men of average age of 47.1 ± 8.6 years old and 60 women of average age of 50.7 ± 7.0 years old made the hypertensive group. A control group of subjects were matched by gender, age and marital status from the normotensive healthy individuals who visited Internal Clinic or Institute of Public Health Osijek for some other reason. Every examinee, gave detailed medical anamnesis, physical status was taken, blood pressure, height and weight were measured, also body mass index was calculated.

After laying down and resting for 15 minutes, 12 lead electrocardiograph and paper rhythm of 100 R-R intervals (variability of the heart frequency) has been recorded during calm slow breathing and deep breathing. R-R intervals have been measured in order, with a pair of compasses. The variability of R-R intervals (variability of heart frequency) was expressed by coefficient of variability (CV) for 100 R-R intervals during calm and during deep breathing. By definition, deep breathing is a state in which a patient breathes freely and as deep as possible, with a frequency of 6 inhales per minute.

Each examinee filled in Bortner's scale to determine a type of behaviour, A or B with a physician's help. Maximum score is 336 points. Total count of 169 to 336 points have individuals that belong to A type behaviour, while 168 or less points indicate a person with B type behaviour. Also an EMOTIONS PROFILE INDEX (EPI) according to Plutchik was made for every examinee. That is a personality test formed to give information about basic personality types and intrapersonal conflicts in life of every individual. It's composed of 12 terms which are joined in pairs in every possible combination. So, we get 62 tasks in which an examinee is "forced" to choose one term in pair which describes him the most. For example, "Is he more shy or dispirited?". Selection is marked in a sense of emotions which are implied in chosen word, for example, shyness implies fear, while consternation implies sadness. Rough results of emotional dimension are expressed through frequency table within adequate percentage for every emotional dimension. This test measures intensity, analogy and bipolarity of eight basic emotions. Intensity of an emotion is shown with a scale from 0 to 100%. Percentage beyond 60% is considered high, and one below 40% low.

An examinee's anger level control was tested with Minnesota multiphasal personal scale of anger expression. That is a question-form with 16 statements on which are answered with "yes" or "no". If the number of affirmative answers is lower, examinees are better in control of their anger. So, lower rating anger is represented with 0-1, middle rating 2-4, higher rating 5-16 affirmative answers.

Statistical analysis

Data are represented as average value and standard deviation. Statistical significance was determined by Student's t-test between groups, variance analysis and χ^2 test. Mutual cohesion between variables were determined by linear correlation coefficient ($r_{xy} = \frac{SD_{xy}}{SD_x SD_y}$). Cohesion strength in between of depended variable and independent variables was determined by multiple regression coefficient and beta coefficient. Statistical importance of results was defined on the level $p < 0,05$.

Results

1. HEART FREQUENCY AND ITS VARIABILITY DURING CALM AND DEEP BREATHING

Within the group of hypertensives, during calm breathing, average R-R interval was shorter, that was higher heart rate than within control group ($758,7 \pm 29,3$ ms opposite $824,9 \pm 38,4$ ms), which was of statistical importance (t-test = -4,031, df = 99, $p < 0,05$). Equally, average frequency variability was statistically distinctively lower in the group of hypertensives, than in the control group (t-test = -2,394, df = 99, $p < 0,05$).

During deep breathing R-R interval was shorter than in calm breathing. The heart frequency was higher in the group of hypertensives than in control group ($726,7 \pm 40,3$ ms opposite $779,0 \pm 49,5$ ms), what was also of statistical importance (t-test = -3,507, df = 99, $p < 0,05$). Equally, average frequency variability was statistically distinctively lower in the group of hypertensives, than in the control group (t-test = -2,095, df = 99, $p < 0,05$).

2. SOMATIC FACTORS AND THEIR CORRELATION WITH HEART FREQUENCY AND ITS VARIABILITY

An average weight within the group of hypertensives was $80,0 \pm 15,1$ kg, and within the control group was $70,5 \pm 11,6$ kg, which is of statistical importance (t-test = 4,97, df = 198, $p < 0,05$). An average height in hypertensives was $170,3 \pm 8,8$ cm, and within the control group was $170,3 \pm 8,5$ cm. Body mass index (BMI) in the group of hypertensives was $27,4 \pm 3,8$ kg/m², and in the control group was $24,2 \pm 3,1$ kg/m². That difference had statistical importance (t-test = 6,44, df = 198, $p < 0,05$).

75 examinees from the group of hypertensives and 51 examinees from the control group had positive family history for hypertension, which made statistically important difference ($\chi^2 = 11,35$, df=1, $p < 0,05$).

A connection between length of average R-R interval, during calm and during deep breathing and coefficient of variability (CV) of R-R interval (heart frequency variability) with altitude of systolic and diastolic blood pressure and duration of hypertension had been tested with multiple regression analysis in the group of hypertensives. We found that total influence of independent variables on R-R interval length in calm (multiple regression coefficient $R = 0,569$, $p < 0,05$), and in deep breathing (multiple regression coefficient $R = 0,558$, $p < 0,05$) was statistically important. Also, effect of all independent variables on coefficient of variability (CV) of the R-R interval (variability of the heart frequency) during calm breathing was statistically important (multiple regression coefficient $R = 0,989$, $p < 0,05$). Coefficient of variability (CV) of the R-R interval during deep breathing was also statistically important and negative connected with duration of hypertension (beta = -0,291, $p < 0,05$). A total influence also showed statistical significance (multiple regression coefficient $R = 0,975$, $p < 0,05$) (Table 1).

3. TYPES OF BEHAVIOUR AND THEIR CONNECTION WITH HEART FREQUENCY AND ITS VARIABILITY

In the group of hypertensives A type behaviour was present within 63 examinees, and in the control group within 69 examinees (no statistically important difference).

A connection between average R-R interval during calm and during deep breathing and coefficient of variability (CV) of R-R interval (variability of the heart frequency) and type of behaviour and every individual Bortner's item was examined with multiple regression analysis.

It shows that, within control group, there's statistically important total influence of all independent variables on to length of average R-R interval during deep breathing (multiple regression coefficient $R = 0,561$, $p < 0,05$). Statistically important influence on to multiple regression coefficient have items which assess "Tension" (beta = -0,508, $p < 0,05$), "Expression of emotions" (beta = -0,314, $p < 0,05$), total "Bortner" (beta = -0,200, $p < 0,05$) and "Impetuosity" (beta = 0,324, $p < 0,05$) (Table 2).

Within the group of hypertensives influence of all independent variables on to coefficient of variability (CV) of R-R interval during calm breathing is statistically important (multiple regression coefficient $R = 0,509$, $p < 0,05$) (Table 3).

4. PROFILE INDEX OF EIGHT BASIC EMOTIONS AND ITS CONNECTION TO HEART FREQUENCY AND ITS VARIABILITY

In the group of hypertensives, emotional dimensions like "Incorporation" and "Reproduction" showed relative intensity above average (more than 60%), while "Rejection" and "Destruction" were below average relative intensity (less than 40%). In the control group, emotional dimensions "Rejection" and "Destruction" were above, and "Rejection" and "Exploration" were below average. "Rejection" as an emotional dimension was statistically importantly lower in the group of hypertensives (t-test = -2,066, df = 99, $p < 0,05$), while an emotional dimension "Exploration" in hypertensives was statistically importantly higher than in the control group (t-test = 2,986, df = 99, $p < 0,05$). In between the group of hypertensives and the control group, there were no statistically

important difference in relative intensity of other emotional dimensions.

A connection between an average R-R interval, during calm and during deep breathing, and coefficient of variability (CV) of the R-R interval (variability of the heart frequency) and Plutchik's eight basic emotions was tested with multiple regression analysis.

In the control group, total influence of all independent variables on coefficient of variability (CV) of the R-R interval (variability of the heart frequency) during calm breathing (multiple regression coefficient $R=0,396$, $p<0,05$) was statistically important, to which emotional dimensions "Incorporation" ($\beta=1,575$, $p<0,05$), "Exploration" ($\beta=1,274$, $p<0,05$) and "Destruction" had the most important positive contribution, while emotional dimensions "Protection" ($\beta=-0,821$, $p<0,05$), and "Rejection" ($\beta=-0,802$, $p<0,05$) and "Reproduction" ($\beta=-1,031$, $p<0,05$) negatively contributed the most (Table 4).

In the group of hypertonics, statistically important was total influence of all independent variables on length of average R-R interval (multiple regression coefficient $R=0,484$, $p<0,05$) and coefficient of variability (CV) of R-R interval (multiple regression coefficient $R=0,429$, $p<0,05$) during deep breathing. CV of R-R interval was importantly positively connected to emotional dimension "Incorporation" ($\beta=1,198$, $p<0,05$), and importantly negatively connected to emotional dimensions "Protection" ($\beta=-1,077$, $p<0,05$), "Rejection" ($\beta=-1,096$, $p<0,05$) and "Reproduction" ($\beta=-1,509$, $p<0,05$) (Table 5).

Difference in anger level between the group of hypertonics and the control group was small and wasn't statistically important ($\chi^2 = 1,67$, $df=2$, $p>0.05$).

Discussion

In this study we studied autonomic regulation of the heart function, threw heart frequency and it's variability expressed by coefficient of variability in 100 sequential heart beats recorded by the EKG during calm and deep breathing, within 100 patients with essential hypertension and compared it to 100 healthy examinees in the control group. 75% of hipertonics had a positive family history of hypertension, which coincides with literature data^{55,56,57}.

A body waight and a body mass index (BMI) were considerably higher in hypertonics then in the control group. According to dana from literature, there's positive correlation between increased body waight and hypertension.

Many epidemiologic studies of essential hypertension showed increased sympathetic activity, especially in the early phase of hypertension^{63,64}. Heart frequency in hypertonics has positive connection with cardiovascular morbidity and mortality. With higher heart rate, cardiovascular morbidity and mortality increases^{65,66,67}. Patients with essential hypertension, in our research, had considerably higher heart frequency at rest and during deep respiration than the control group. R-R variability at rest and during deep respiration was considerably lower in hypertonics than in the control group, which goes in favour of increased sympathetic activity, while vagal activity is reduced. A smaller increase in RR variability was found in hypertensive patients than in the control group during deep breathing.

A coefficient of variability calculated from standard deviation of the RR interval is useful clinical indicator of cardial parasympathetic activity^{70,71}. A mechanism of decreased parasympathetic activity in the essential hypertension is connected with abnormal autonomic stimulation of normal sinal node or abnormal function of the beta-receptors. A beta-adrenergic hypersensitivity syndrome has been described⁷².

A variability of heart frequency in hypertonics with left ventricle hypertrophy was significantly reduced considering the control group, which is already confirmed in some earlier studies^{30,73,74}. A number of hypertonics with heart changes in our research was considerably lower than the number of those hypertonics without heart changes. Systolic pressure, diastolic pressure, duration of hypertension, have total considerable influence on the heart frequency and it's variability. The most significant negative connection is found between variability of the heart frequency and duration of hypertension. Some recent studies showed that low variability of the heart frequency is a predictor of cardial events such as myocardial infarction, rapid progression of atherosclerosis and death after myocardial failure^{75,76,77}.

In hypertonics and in the control group "A" type behaviour was prevailing, but difference was not statistically important. That result could be explained with socioeconomic milieu and present period of "society in transition" in which we live today. Type A of behaviour is related to heart frequency and it's variability. In the hypertensive group, total influence of all independent variables on variability of the heart frequency during calm breathing was statistically important. The heart frequency is considerably negatively influenced by items which assess "tension", "expresion of emotions" and "total Bortner", and positively by "impetuosity". The type A behaviour is associated with increased heart frequency and it's variability in both groups.

An anger level was high within 65% of hypertonics. In the control group, 60% of examinees had high anger level, but difference was not statistically important. Some data from literature inform about connection between suppressed anger and high blood pressure. Haynes and associates and Goldberg didn't find constant connection between anger and blood pressure^{80,81}. Anger can be result of psychosocial stress which is associated with hypertension or it's aggravation^{82,83}.

In "Emotion profile", regarding the control group, hypertensive individuals had significantly lower emotional dimension "Rejecion", and higher emotional dimension "Exploration". These data

says that hypertensive individuals are more uncritical and unable to relax than individuals in the control group. According to the literature available to us, Plutchik's "Emotion profile" was rarely used in hypertensive patients. Čatipović-Veselica and associates did not find significant difference in average emotion profile between hypertensive and normotensive workers⁸⁴.

Our data shows that patients with hypertension have increased heart frequency and reduced variability of the heart frequency. Increased heart frequency and reduced variability of the heart frequency in patients with long-term hypertension can contribute to progression of cardial morbidity and mortality. The question is whether to use medicaments or non-medicament methods to improve variability of the heart frequency.

1. The method of measuring hundred sequential heart beats on EKG, during rest and durind deep breathing, showed to be sensitive and simple diagnostic measure of decreased function of the parasympathetic nervous system in essential arterial hypertension. Coefficient of variability calculated from standard deviation of RR interval is useful clinical indicator of cardial parasympathetic activity in essential hypertension.

2. Patients with essential arterial hypertension had significantly higher heart frequency at rest and during deep breathing then the control group. Variability of the heart frequency, at rest and during deep breathing, was considerably lower in hypertensive patients then in the control group, which goes in favour of increased sympatethic activity, and reduced vagal activity.

3. A body mass and a body mass index (BMI) were considerably higher in examinees with essential arterial hypertension. Systolic blood pressure, diastolic blood pressure, duration of the hypertension, have significant total influence on the heart frequency and it's variability. Variability of the heart frequency is negatively associated with duration of hypertension

4. Within examinees and controles prevailed "A" type behaviour. Items which assess "tension", "expression of emotions" and "total Bortner" had negative considerable influence on the heart frequency but "Impetuosity" had positive influence.

5. Eight basic emotions according to Plutchik influence the heart frequency and it's variability. Hypertensive patients were more uncritical and more unable to relax then the control group.

6. Our results show that patients with hypertension have increased heart frequency and reduced variability of the heart frequency, which are well known risk factors for increased cardial morbidity and mortality. This raises the question of therapeutic approach to the patient with essential hypertension, medicamentous treatment or some other methods like psychosocial to increase variability of the heart frequency.

REFERENCES

1. HUMERFELT SB, Acta Med Scand 401 (1963) 1. - 2. YOSEFY C, GINSBERG GM, DICKER D, VISKOPER JR, TULCHINSKY TH, LEIBOVITZ E, GAVISH D, Blood Press 12 (2003) 225. - 3. YOSEFY C, GINSBERG G, VISKOPER R, DICKER D, GAVISH D, Cost Eff Resour Alloc. 28 (2007) 5. - 4. KAUR P, RAO TV, SANKARASUBBAIYAN S, NARAYANAN AM, EZHIL R, RAO SR, GUPTE MD, J Assoc Physicians India 55 (2007) 771. - 5. CASTELLI WP, Am J Med 76 (1984) 4. - 6. DUPREZ DA, Am J Med 121 (2008) 179. - 7. BARQUERA S, DURAZO-ARVIZU RA, LUKE A, CAO G, COOPER RS, J Hum Hypertens 28 (2008) - 8. ALDERMAN MH, LAMPORT B, J Clin

Epidemiol 43 (1990) 195. - 9. WONG ND, LOPEZ VA, L'ITALIEN G, CHEN R, KLINE SE, FRANKLIN SS, Arch Intern Med 10 (2007) 2431. - 10. KI, CHANG HJ, CHO YS, YOUN TJ, CHUNG WY, CHAE IH, CHOI DJ, KIM CH, Hypertens Res. 31 (2008) 97. - 11. AMADO P, VASCONCELOS N, SANTOS I, ALMEIDA L, NAZARÉ J, CARMONA J, Rev Port Cardiol 18 (1999) 897. - 12. COLLINS K, GOUGH S, CLANCY M, Emerg Med J. 25 (2008) 196. - 13. COCA A, Clin Drug Investig. 28 (2008) 211. - 14. WHO REGIONAL OFFICE FOR EUROPE. 2001. Survey about hypertension control policies in CINDI countries (WHO, Copenhagen 1998). - 15. LITHELL H, HANSSON L, SKOOG I, ELMFELDT D, HOFMAN A, OLOFSSON B, TRENKWALDER P, ZANCHETTI A, J Hypertens 22 (2004) 1605. - 16. D'AGOSTINO RB SR, VASAN RS, PENCINA MJ, WOLF PA, COBAIN M, MASSARO JM, KANNEL WB, Circulation 12 (2008) 743. - 17. BABUS V, Epidemiologija. (Medicinska naklada, Zagreb, 1997). - 18. NATIONAL INSTITUTE OF HEALTH, The Fifth Report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure NIH Publication (1993) 1. - 19. BEREZAN DJ, DUNN KM, FALCK JR, DAVIDGE ST, J Cardiovasc Pharmacol 51 (2008) 327. - 20. FROHLICH ED, APSTEIN C, CHOBANIAN AV, DEVEREUX RB, DUSTAN HP, DZAU V, N Engl J Med 327 (1992) 998. - 21. SAVAGE D, Am J Hypertens 1 (1988) 12. - 22. HAINSWORTH R, Physiol Rev 71 (1991) 617. - 23. HAMLIN RL, SMITH CR, Am J Physiol 215 (1968) 560. - 24. LEVY MN, ZIESKE H, J Appl Physiol 27 (1969) 465. - 25. ZIPES DP, Genesis of cardiac arrhythmias: electrophysiological considerations. In: Braunwald E. Heart disease. Saunders, 1992: 588. - 26. COUMEL PH, HERMIDA IS, WENERBLOM B, LEENHARDT A, MAISON-BLANCHE P, CAUCHEMEZ B, Eur Heart J 12 (1991) 412. - 27. KARAPETIAN GK, ENGELS HJ, GRETEBECK RJ, Int J Sports Med. 29 (2008) 652. - 28. GILLUM RF, Am Heart J 116 (1988) 163. - 29. LANGEWITZ W, RUDEL H, SCHACHINGER H, Am Heart J 127 (1994) 122. - 30. CHAKKO S, MULINGTAPANG RF, HUIKURI HV, KESSLER KM, MASTERSON BJ, MYERBURG RJ, Am Heart J 126 (1993) 1364. - 31. MANCIA G, LUDBROOK J, FERRARI A, Circ Res 43 (1978) 170. - 32. ENGEL BT, Am J Cardiol 60 (1987) 3. - 33. ENGEL BT, Exp Gerontol. 28 (1993) 499. - 34. FRIEDMAN M, ROSENMAN RH, JAMA 169 (1959) 1286. - 35. ROSENMAN RH, The role of behavior patterns and neurogenic factors in the pathogenesis of coronary heart disease. In: Eliot Rs ed. Stress and Heart. New York: Futura Publishing Company. 1974:123. - 36. FONAROW GC, Rev Cardiovasc Med 8 (2007) 25. - 37. PLUTCHIK R, Psychol Rev 53 (1966) 2. - 38. CHAKRABARTI B, BULLMORE E, BARON-COHEN S, Soc Neurosci 1 (2006) 364. - 39. FRAPPORTI-ROGLIĆ M, Podaci o Indeksu profila emocija (IPE) (Naklada Slap, Jastrebarsko, 1994). - 40. CASSÉ-PERROT C, FAKRA E, JOUVE E, BLIN O, Encephale 33 (2007) 169. - 41. KEIL U, KUULASMA K, Int J Epidemiol 18 (1989) 46. - 42. WHO REGIONAL OFFICE FOR EUROPE. 2001. Survey about hypertension control policies in CINDI countries (WHO, Copenhagen 1998). - 43. LITHELL H, HANSSON L, SKOOG I, ELMFELDT D, HOFMAN A, OLOFSSON B, TRENKWALDER P, ZANCHETTI A, J Hypertens 22 (2004) 1605. - 44. INUKAI T, KOBAYASHI I, KOBASHY T, ISHII A, YAMAGUCHI T, YAMAGUCHI Y, IWASHITA A, OHSHIMA K, SHIAMURA Y, KOBAYASHI S, J Inter Med 228 (1990) 431. - 45. BORTNER RW, J Chron Dis 22 (1969) 87. - 46. THE BELGIAN-FRENCH POOLING PROJECT, Eur Heart J 5 (1984) 440. - 47. PLUTCHIK R, KELLERMAN H, Los Angeles: WPS (1974) 1. - 48. BAŠKOVAC-MILINKOVIĆ A, BELE-POTOČNIK Ž, HRUŠEVAR B, RAJSEK J. PIE, Profil indeksa emocija. (Zavod Slovenije za produktivnost dela, Ljubljana, 1979). - 49. HRNJICA S, Opšta psihologija sa psihologijom ličnosti (Naučna knjiga, Beograd, 1981). - 50. KAWACHI I, SPARROW D, SPIRO A, VOKONAS P, WEIS S, Circulation 94 (1996) 2090. - 51. LADER E, EGAN D, HUNSBERGER S, GARG R, CZAJKOWSKI S, MCSHERRY F, J Card Fail 9 (2003) 4. - 52. ROELAND J, GIBSON DG, Eur Heart J 1 (1980) 375. - 53. DEVREUX RB, REICHEK N, Circulation 55 (1997) 613. - 54. SAHN DJ, DEMARIA A, KISSLO J, WEJMAN A, Circulation 58 (1978) 1072. - 55. LEVINE RS, HENNEKENS CH, JESSE MJ, Br Med J 308 (1994) 298. - 56. CHRISTENSEN K, STØVRING H, MCGUE M, J Epidemiol Community Health 55 (2001) 583. - 57. MCNEILL G, TUYA C, SMITH WC, Int J Epidemiol 33 (2004) 995. - 58. STAMLER J, ROSE G, ELIOT P, MARMONT M, KESTELOOT H, STAMLER R, Hypertension 17 (1991) 9. - 59. STAMLER J, ELLIOTT P, APPEL L, CHAN Q, BUZZARD M, DENNIS B, DYER AR, ELMER P, GREENLAND P, JONES D, KESTELOOT H, KULLER L, LABARTHE D, LIU K, MOAG-STALBERG A, NICHAMAN M, OKAYAMA A, OKUDA N, ROBERTSON C, RODRIGUEZ B, STEVENS M, UESHIMA H, HORN LV, ZHOU B, J Hum Hypertens 17 (2003) 655. - 60. ZHOU BF, STAMLER J, DENNIS B, MOAG-STALBERG A, OKUDA N, ROBERTSON C, ZHAO L, CHAN Q, ELLIOTT P, J Hum Hypertens 17 (2003) 623. - 61. JONES PH, Drugs 59 (2000) 1127. - 62. STRANDBERG TE, FEELY J, SIGURDSSON EL, Clin Ther 26 (2004) 1821. - 63. LOMBARDI F, MALLIANI A, PAGANI M, CERUTTI S, Cardiovasc Res 32 (1996) 208. - 64. MOUROT L, BOUHADDI M, GANDELIN E, CAPPELLE S, NGUYEN NU, WOLF JP, ROUILLON JD, HUGHSON R, REGNARD J, Auton Neurosci 30 (2007) 27. - 65. GILLMAN MW, KANNEL WB, BELANGER A, D'AGOSTINO, Am Heart J 125 (1993) 1148. - 66. INOUE T, OSHIRO S, ISEKI K, TOZAWA M, TOUMA T, IKEMIYA Y, TAKISHITA S, Jpn Circ J 65 (2001) 969. - 67. INOUE T, ISEKI K, ISEKI K, KINJO K, OHYA Y, TAKISHITA S, Circ J 71 (2007) 1755. - 68. EWING DJ, NEILSON JMM, TRAVIS P, Br Heart J 52 (1984) 396. - 69. SUCHARITA S, BHARATHI AV, KURPAD AV, VAZ M, Physiol Meas 23 (2002) 347. - 70. KAGEYAMA S, MOCHIO S, TANIGUCHI M, Jikei Med J 28 (1981) 81. - 71. MADANMOHAN, PRAKASH ES, BHAVANANI AB, Indian J Physiol Pharmacol 49 (2005) 132. - 72. MALLIANI A, MONTANO N, Hypertension 39 (2002) 63. - 73. ALTER P, GRIMM W, VOLLRATH A, CZERNY F, MAISCH B, Am Heart J 151 (2006) 829. - 74. HOFFMANN J, GRIMM W, MENZ V, MAISCH B, Z Kardiol 89 (2000) 84. - 75. WICHTERLE D, SIMEK J, LA ROVERE MT, SCHWARTZ PJ, CAMM AJ, MALIK M, Circulation 110 (2004) 1183. - 76. ARONSON D, MITTLEMAN MA, BURGER AJ, Am J Cardiol 1 (2004) 59. - 77. LANZA GA, CIANFLONE D, REBUZZI AG, ANGELONI G, SESTITO A, CIRIELLO G, LA TORRE G, CREA F, MASERI A, Heart 92 (2006) 1055. - 78. BYRNE DG, ROSENMAN RH, Br J Med Psychol 59 (1986) 75. - 79. ZODPAY SP, RAUT SG, Indian J Med Sci 92 (1998) 348. - 80. DIMSDALE JE, PIERCE C, SCHOENFELD D, Psychosomatic Med 48 (1986) 430. - 81. OKKEN VS, NIEMEIJER MG, DIJKSTRA A, BAARS MW, SAID S, HOOGENBERG K, ORFGEN H, OTTEN S, CLEOPHAS TJ, Neth Heart J 16 (2008) 197. - 82. SPARRENBERGER F, CICHELO FT, ASCOLI AM, FONSECA FP, WEISS G, BERWANGER O, FUCHS SC, MOREIRA LB, FUCHS FD, J Hum Hypertens 23 (2009) 12. - 83. STEPTOE A, Hypertension 52 (2008) 220. - 84. ČATIPOVIĆ-VESELICA K, MERČEP A, VIZNER-LOVRIĆ I, BURIĆ D, Arh hig rada toksikol 42 (1991) 381.

K. Kožul

Institute of Public Health for the Osijek-baranja County, F. Krežme 1, 31000 Osijek, Croatia
e-mail: karlo.kozul@os.htnet.hr

TABLE 1

A CORRELATION BETWEEN R-R INTERVAL AND COEFFICIENT OF VARIABILITY (CV) OF R-R INTERVAL AND SOME HEALTH PARAMETERS AND LAB RESULTS IN HYPERTENSIVE PATIENTS

Health parameters	Beta-coefficient of R-R interval		Beta-coefficient of CV R-R interval	
	calm breathing	deep breathing	calm breathing	deep breathing
	Systolic heart pressure	0,063	0,082	-0,038
Diastolic heart pressure	0,009	-0,063	0,012	0,038
Duration of hypertension	-0,206	-0,218	-0,041	-0,291
Multiple regresion coefficient	0,569	0,558	0,989	0,975

TABLE 2

A CONNECTION BETWEEN R-R INTERVAL AND COEFFICIENT OF VARIABILITY (CV) OF R-R INTERVAL AND TYPE OF BEHAVIOUR

ACCORDING TO BORTNER IN THE CONTROL GROUP

Bortner scale (type of behaviour)	Beta coefficient of R-R interval		Beta coefficient of CV of R-R interval	
	calm breathing	deep breathing	calm breathing	deep breathing
Punctuality	0,058	0,074	0,145	0,113
Competitive spirit	0,151	0,041	-0,238	-0,194
Disobedience	-0,114	-0,024	0,147	0,153
Haste	-0,038	0,068	0,105	0,305
Impatience	0,079	0,065	0,003	-0,086
Insecurity	0,099	0,014	0,257	0,225
Doing several things simultaneously	-0,166	-0,179	-0,139	-0,189
Impetuosity	0,252	0,324	0,109	-0,085
Wish for acknowledgement	0,053	-0,072	0,153	-0,023
Promptitude	0,109	0,128	-0,050	-0,263
Tension	-0,453	-0,508	-0,249	0,147
Expression of emotions	-0,257	-0,314	-0,051	0,221
Several interest	0,182	0,031	-0,223	-0,222
Ambition	-0,086	-0,004	0,036	0,181
TOTAL	-0,222	-0,200	0,015	0,088
Multiple regression coefficient	0,474	0,561	0,363	0,415

TABLE 3
A CONNECTION BETWEEN R-R INTERVAL AND COEFFICIENT OF VARIABILITY (CV) OF R-R- INTERVAL AND BORTNER'S TYPE OF BEHAVIOUR IN THE GROUP OF HYPERTONICS

Bortner scale (type of behaviour)	Beta coefficient of R-R interval		Beta coefficient of CV R-R interval	
	calm breathing	deep breathing	calm breathing	deep breathing
Punctuality	-0,04	-0,07	0,14	0,14
Competitive spirit	-0,04	-0,08	0,21	0,12
Disobedience	0,04	0,09	-0,20	0,18
Haste	0,40	0,41	0,18	-0,01
Impatience	-0,19	-0,16	0,18	0,06
Insecurity	-0,06	-0,05	-0,02	-0,12
Doing several things simultaneously	0,03	0,05	-0,06	-0,15
Impetuosity	-0,36	-0,37	0,05	0,14
Wish for acknowledgement	-0,11	-0,05	-0,34	-0,14
Promptitude	-0,04	-0,03	-0,26	-0,02
Tension	0,10	0,07	0,18	0,06
Expression of emotions	-0,10	-0,12	0,04	-0,25
Several interest	-0,16	-0,11	-0,19	-0,25
Ambition	-0,13	-0,10	-0,111	-0,14
TOTAL	-0,222	-0,143	-0,046	-0,033
Multiple regression coefficient	0,452	0,434	0,509	0,460

TABLE 4
A CONNECTION BETWEEN R-R INTERVAL AND COEFFICIENT OF VARIABILITY (CV) OF THE R-R INTREVAL AND EIGHT BASIC EMOTIONS IN THE CONTROL GROUP

Plutchik's eight basic emotions	Beta coefficient of R-R interval		Coefficient of variability (CV) of the R-R interval	
	calm breathing	deep breathing	calm breathing	deep breathing
Incorporation	-0,106	-0,220	0,069	1,198
Orientation	-0,143	-0,051	-0,327	-0,218
Protection	-0,066	-0,189	-0,513	-1,077
Deprivation	0,021	0,080	-0,644	-0,815
Rejection	-0,546	-0,712	-0,622	-1,096
Exploration	0,193	0,057	0,293	0,499
Destruction	0,605	0,379	0,519	0,601
Reproduction	0,138	-0,017	-0,523	-1,509
Bias	-0,080	-0,148	-0,147	0,150
Multiple regression coefficient	0,384	0,484	0,286	0,429

TABLE 5
A CONNECTION OF R-R INTERVAL AND COEFFICIENT OF VARIABILITY (CV) OF R-R INTERVAL WITH EIGHT BASIC EMOTIONS IN THE

GROUP OF HYPERTONICS

Plutchik's eight basic emotions	Beta coefficient of R-R interval		Coefficient of variability (CV) of the R-R interval	
	calm breathing	deep breathing	calm breathing	deep breathing
Incorporation	-0,106	-0,220	0,069	1,198
Orientation	-0,143	-0,051	-0,327	-0,218
Protection	-0,066	-0,189	-0,513	-1,077
Deprivation	0,021	0,080	-0,644	-0,815
Rejection	-0,546	-0,712	-0,622	-1,096
Exploration	0,193	0,057	0,293	0,499
Destruction	0,605	0,379	0,519	0,601
Reproduction	0,138	-0,017	-0,523	-1,509
Bias	-0,080	-0,148	-0,147	0,150
Multiple regression coefficient	0,384	0,484	0,286	0,429