

Lethal and sublethal effects of carlina oxide on *Tetranychus urticae* (Acari: Tetranychidae) and *Neoseiulus californicus* (Acari: Phytoseiidae)

Roberto Rizzo,^a  Ernesto Ragusa,^b Giovanni Benelli,^c Gabriella Lo Verde,^b  Valeria Zeni,^c  Filippo Maggi,^d  Riccardo Petrelli,^d Eleonora Spinozzi,^d Marta Ferrati,^d Milko Sinacori^b and Haralabos Tsolakis^{b*} 

Abstract

BACKGROUND: *Tetranychus urticae* Koch, is a polyphagous and damaging pest, presenting several resistant populations worldwide. Among new and more environmentally friendly control tools, botanical pesticides represent a valuable alternative to synthetic ones within integrated pest management strategies. Accordingly, we investigated the lethal and sublethal effects of carlina oxide isolated from *Carlina acaulis* (Asteraceae) roots on *T. urticae* and its natural enemy, the predatory mite, *Neoseiulus californicus* (McGregor).

RESULTS: Carlina oxide (98.7% pure compound) was used for acaricidal tests on eggs, nymphs, and adult females of *T. urticae* (concentrations of 312.5, 625, 1250, 2500 and 5000 $\mu\text{L L}^{-1}$), and eggs and females of *N. californicus* (1250 and 5000 $\mu\text{L L}^{-1}$ on eggs and females, respectively). Behavioral two-choice tests were also conducted on phytoseiid females. Carlina oxide toxicity was higher on *T. urticae* females than nymphs (median lethal dose 1145 and 1825 $\mu\text{L L}^{-1}$, respectively), whereas egg mortality and mean hatching time were significantly affected by all tested concentrations. A decreasing daily oviposition rate for *T. urticae* was recorded with concentrations ranging from 625 to 5000 $\mu\text{L L}^{-1}$, whereas negative effects on the population growth rate were recorded only with the three higher concentrations (1250, 2500 and 5000 $\mu\text{L L}^{-1}$). No toxic effect on *N. californicus* females was found, but a strong repellent activity lasting for 48 h from application was recorded.

CONCLUSION: Carlina oxide reduced longevity and fecundity of *T. urticae* adults, but not of *N. californicus*. This selective property allows us to propose it as a novel active ingredient of ecofriendly acaricides for *T. urticae* management.

© 2023 The Authors. *Pest Management Science* published by John Wiley & Sons Ltd on behalf of Society of Chemical Industry.

Keywords: Asteraceae; *Carlina acaulis*; green acaricides; side effects; two-spotted spider mite; phytoseiid mites

1 INTRODUCTION

The two-spotted spider mite (TSSM) *Tetranychus urticae* Koch (Acariformes, Tetranychidae) is a global and economically important pest on various crops, both in the field and in the greenhouse.^{1,2} The phytoseiid mites *Phytoseiulus persimilis* Athias-Henriot and *Neoseiulus californicus* (McGregor) (Parasitiformes, Phytoseiidae) have been successfully used for decades in biological control programs against TSSM infestations.^{3–5} However, TSSM biocontrol is unsatisfactory on some crops because of the intrinsic characteristics of the host plant species (e.g., tomato), or environmental conditions (e.g., low relative humidity),^{6–8} forcing growers to still rely on chemical control. Synthetic acaricides have represented an appropriate control technique for phytophagous mites for decades.^{9,10} However, rapid evolution of resistance to acaricides in spider mites has also been recorded.^{11,12} In addition, acaricides also have detrimental effects on native phytoseiid populations inhabiting cultivated plants, causing major problems in integrated pest management (IPM) strategies,^{13–15} and leading researchers to focus their interest on botanical-based products.^{16–18}

Botanical pesticides were commonly used until World War II, but their role became marginal following the discovery of the synthetic pesticides.¹⁹ The former have no or less-negative effects on human health and the environment, as well as on natural enemies, and show low persistence.^{20,21} Among botanical products,

* Correspondence to: H. Tsolakis, Department of Agricultural, Food and Forest Sciences, University of Palermo, viale delle Scienze, Ed. 5, 90128, Palermo, Italy. E-mail: haralabos.tsolakis@unipa.it

a CREA - Research Centre for Plant Protection and Certification, Palermo, Italy

b Department of Agricultural, Food and Forest Sciences, University of Palermo, Palermo, Italy

c Department of Agriculture, Food and Environment, University of Pisa, Pisa, Italy

d Chemistry Interdisciplinary Project (ChIP) Research Center, School of Pharmacy, University of Camerino, Camerino, Italy

essential oils (EOs) and their bioactive constituents have been re-evaluated in past years as promising pesticides.^{22–24} EOs biocidal effects have been studied on many important arthropod pests such as thrips,^{25–31} ambrosia beetles,³² mites, ticks^{17,33–36}, and on weeds.^{37,38} Moreover, the high worldwide production of plant EOs for perfume and flavoring, makes them suitable from a commercial point of view.^{39,40}

Most studies on botanical pesticides focus on plant species belonging to the Apiaceae, Myrtaceae and Lamiaceae families. However, Asteraceae have also received considerable attention.^{41–44} The stemless carline thistle *Carlina acaulis* L., belongs to the latter family and is spread in xerothermic grasslands of Central and South Europe.⁴⁵ The plant was well known by ancient Greeks and Romans for its anthelmintic, antibiotic, dermatological and diuretic effects, and has been continuously included in the European Pharmacopoeia for more than two millennia.⁴⁶ Different attempts have recently been made to cultivate this plant on a large scale, with specific treatments to stimulate production of the active compounds in the plant.⁴⁷ The main constituent of the EO (>90%) is the polyacetylene carlina oxide [2-(3-phenylprop-1-ynyl) furan]^{48,49} and this has proven to be toxic on insect pests like *Lobesia botrana* (Denis & Schiffmüller),⁵⁰ *Ceratitis capitata* (Wiedemann)^{29,30} and *Bactrocera oleae* (Rossi)²⁸ and on nematodes such as *Meloidogyne incognita* (Kof. & White).⁵¹ Carlina oxide has potential as an ideal candidate ingredient in formulations for use in crop protection within the framework of sustainable agriculture aimed at reducing the risks from the use of conventional pesticides and to promote alternative strategies in IPM programs. *Carlina acaulis* and its EO are included in the list of botanicals to be used in food supplements shared with the health administrations of Belgium, France and Italy (BELFRIT) and this may facilitate biopesticide registration bypassing regulatory restrictions.⁵²

The scientific community has recently begun to focus on the potentially hazardous effects of natural products on non-target organisms such as predatory mites, aquatic organisms and earthworms.^{53–55} However, literature on the effects of botanical pesticides on non-target organism behavior is still poor.^{53,56} To the best of our knowledge, little is known on the behavioral responses of predatory mites to carlina oxide.

Within this framework, this study aimed to evaluate the effects of carlina oxide toward *T. urticae* eggs, nymphs and adults, also adopting a population-level approach.^{33,57} Moreover, because the study of side effects on natural enemies is essential when searching for biopesticides fitting with IPM principles, the toxicity of this polyacetylene was also evaluated on females and eggs of the phytoseiid predaceous mite *N. californicus*.

2 MATERIALS AND METHODS

2.1 Carlina oxide isolation

Carlina oxide was isolated as a yellow liquid from 1 kg of dry *C. acaulis* roots by hydrodistillation using 10 L of distilled water and was recovered through a Clevenger-type apparatus for 8 h (yield 0.78% dry weight). The plant material was bought from Minardi & Figli S.r.l. (Bagnacavallo, Ravenna, Italy; <https://www.minardierbe.it>; batch no C-210920250920, collected in 2020). The purity of the compound (98.7%, Fig. 1) was assessed by gas chromatography–mass spectrometry (GC–MS) analysis using a previously validated method,⁵⁸ and the chemical structure was checked by MS and NMR data using a standard previously obtained in the authors' laboratory.⁵⁹ Once obtained, carlina oxide was stored at –20 °C until biological assays.




Figure 1. Chemical structure of carlina oxide.

2.2 *T. urticae* and *N. californicus* rearing

Tetranychus urticae was collected on weeds and *Solanum melongena* L., near Palermo, Italy (38°1'41.49" N, 13°1'55.61" E), in June 2020, and kept in laboratory cultures on potted bean plants (*Phaseolus vulgaris* L.). *Neoseiulus californicus* was collected on strawberries infested by *T. urticae* at Partinico (Palermo) (38°4'22.21" N, 13°6'12.27" E) in December 2018 and reared on plexiglass arenas⁶⁰ supplying as food various stages of *T. urticae* and a mixture of pollens [*Carpobrotus edulis* (L.) N.E.Br., *Oxalis pes-caprae* L., *Typha latifolia* L.] as a food. Both rearings were maintained in a conditioned room [25 ± 1 °C, 70% ± 5% relative humidity (RH) and 16:8 h light/dark photoperiod]. The two rearings were regularly renewed or supplemented with field-collected specimens.

2.3 Toxicity of carlina oxide on *T. urticae*

Tests were carried out on eggs, nymphs and adults of *T. urticae*. The following concentrations of the carlina oxide were tested, i.e. 312.5, 625, 1250, 2500 and 5000 µL L⁻¹. Each experimental unit (EU) consisted of a bean leaf disk (3 cm in diameter) with the abaxial surface up, placed on wet cotton saturated daily with distilled water, in a Petri dish (diameter 100 mm, height 10 mm). For each EU, 8 mL of solution were sprayed using the Potter Precision Spray Tower,⁶¹ at 6.89 kPa of pressure. Carlina oxide was dissolved in acetone. Negative control tests were treated with acetone only.

Toxicity tests on *T. urticae* were carried out following the methodology described by Tsolakis and Ragusa.³³ To obtain a cohort of TSSM eggs, three females were placed on each EU, allowed to lay eggs for 24 h and then removed. Ten eggs per EU were left for treatments. Each test was replicated 30 times and lasted 7 days (when more than 95% of eggs hatched in the control). The mean hatching time was calculated at the end of the test period. The above procedure was adopted for obtaining coetaneous (max 24 h old) protonymphs. Six replicates were adopted for each toxicity test on five nymphs per EU; the test lasted 4 days.

To obtain a cohort of coetaneous females, 50 females were transferred with a fine brush (4/0) to the abaxial surface of five bean leaves (ten females per leaf), allowed to lay eggs for 24 h and then removed. Mites obtained from these eggs were reared until reaching adulthood; afterwards 60 young females (max 24 h old) were transferred to the EUs (one female per EU + one male originating from the colony) to ensure mating. After 48 h males were removed, and fertilized females were used for tests. Each test was replicated 60 times and lasted 4 days. The mortality of nymphs and adult females was recorded daily and the mean survival time and survival rate were calculated at the end of tests. Moreover, the number of eggs/female/day was recorded over the test period. To calculate the total toxic effect of carlina oxide on female mortality and oviposition rate, the following formula

proposed by Overmeer and van Zon⁶² was applied to data obtained from each concentration:

$$E = 100\% - (100\% - M) \times R$$

Where M is corrected mortality of females at the end of the test⁶³ and R is the effect on the reproduction (no. of eggs per treated female/no. of eggs per untreated female).

Four toxicity categories, as proposed by Hardman *et al.*⁶⁴ were applied for *T. urticae* considering the Abbott⁶³ corrected mortality: 1 = not toxic (<25% mortality), 2 = slightly toxic (25%–50% mortality), 3 = moderately toxic (51%–75% mortality), and 4 = very toxic (>75% mortality).

2.4 Effect of carlina oxide on *T. urticae* population growth rate

To measure the effect of carlina oxide on the populations growth of *T. urticae*, we calculated the instantaneous rate of increase (r_i).^{57,65} This index measures the population increase or decrease and is calculated adopting the following equation:

$$r_i = \frac{\ln\left(\frac{N_f}{N_0}\right)}{\Delta t}$$

where r_i is instantaneous rate of population increase, like that obtained with the intrinsic rate of increase (r_m),⁵⁷ N_f is final number of mites, N_0 is initial number of mites and Δt is time that the experiment lasted. Positive values of r_i show a growing population, $r_i = 0$ indicates a stable population, and negative values of r_i indicate a declining population directed toward extinction.⁵⁷ The r_i was calculated after 4 days.

2.5 Side effects of carlina oxide on *N. californicus*

To obtain young females of *N. californicus*, 100 eggs from the colony were transferred to a new arena and provided with an abundant mixture of pollens until adulthood was reached. Newly emerged females and males were transferred with a fine brush (4/0) to a new arena for 48 h to ensure mating (sex ratio females/males 3:1). Afterwards, one female predator was transferred on each EU for subsequent tests.

Four different tests were carried out on phytoseiid females: A, spraying directly on the EUs bearing each one female (30 replications); B, spraying the leaf disk, the female was placed on the leaf disk 4 min later when the leaf disk was dry (15 replications); C, spraying the female which was then placed on untreated leaf disk (30 replications); D spraying the leaf disk, the female was placed on leaf disk 48 h later (15 replications). In all the above tests a concentration of 5000 $\mu\text{L L}^{-1}$ was used, because only this caused a mortality rate of 95% on *T. urticae* females. Pollen grains were provided as food once the sprayed surface of leaf disk was dry. All tests lasted 3 days and were carried out at $25 \pm 1^\circ\text{C}$, $70\% \pm 5\%$ RH and 16:8 h light/dark photoperiod.

To obtain fresh eggs of *N. californicus* for toxicity tests, 50 females were placed in a new arena and allowed to lay eggs for 24 h. Afterwards, one egg was transferred with a fine brush (4/0) to each EU and then sprayed with carlina oxide at a concentration of 1250 $\mu\text{L L}^{-1}$, which was the dose causing 95% of mortality on *T. urticae* eggs. Each test was replicated 30 times and lasted 3 days.

The mortality of motile stages was daily recorded during the test period. We considered mortality to be natural or induced death

(dead mites found on leaf disks), whereas mites found drowned in the wet cotton wool surrounding the leaf disk were ascribed to a repellent effect. Egg mortality was ascertained when more than 95% of eggs hatched in the control (after 3 days).

The four toxicity categories proposed by the International Organization for Biological Control for natural enemies, like those applied for *T. urticae*, were adopted for *N. californicus* considering Abbott⁶³ corrected mortality: 1 = harmless (<30% mortality), 2 = slightly harmful (30%–79% mortality), 3 = moderately harmful (80%–99% mortality) and 4 = harmful (>99% mortality).⁶⁶

2.6 Y-tube olfactometer bioassays on *N. californicus* adults

2.6.1 Y-tube apparatus

To evaluate the potential attraction or repellency activity of carlina oxide on adult females of *N. californicus*, two-choice tests were performed in a Y-tube olfactometer as used by Canale *et al.*⁶⁷ and described in the Supplementary Information; the method of Fonseca *et al.*⁶⁸ with slight modifications was followed. Two different behavioral assays were carried out on *N. californicus* females: A, preliminary test on filter paper (50 replicates/dose); and B, spraying on a bean leaf disk and performing the test 15 min and 48 h after spraying (30 replicates/dose). The tested doses were 2500 and 5000 $\mu\text{L L}^{-1}$, the two highest doses tested on *T. urticae* females. All the bioassays were conducted under laboratory conditions ($25 \pm 1^\circ\text{C}$, $70\% \pm 5\%$ RH). Purified air was provided at 2 mL min^{-1} for both behavioral assays. Illumination was provided by a red LED light (12 W, 1050 lm) hanging vertically above the olfactometer unit (height 60 cm). In both behavioral assays, either the filter paper or the bean leaf were moistened with 10 μL of the product to be tested and the respective control. To avoid any bias, after testing five females the arena was flipped by 180° and cleaned as detailed in Carpita *et al.*,⁶⁹ the substances were then renewed.

2.6.2 Preliminary bioassay

Two doses (2500 and 5000 $\mu\text{L L}^{-1}$) of carlina oxide were formulated with Tween 80 (3%) (1:1) plus distilled water and applied to a filter paper (20 \times 20 mm, Whatman 1). Negative controls (the solution without carlina oxide) were also prepared. Females of *N. californicus* were introduced individually in the main arm of the olfactometer and each was allowed to choose between the two arms. A choice was recorded when the female reached the end of one of the two arms. After making a choice, the female was removed and a new female was introduced. Each observation lasted 5 min; females that did not move at all within 3 min were removed and scored as no choice. A total of 50 adult females of *N. californicus* were tested at the two carlina oxide concentrations.

2.6.3 Carlina oxide bioactivity over time

The repellent or attractive effect of carlina oxide was also evaluated over time in Y-tube bioassays, following the method described above. Fresh bean leaf disks (20 mm in diameter) were used to evaluate how the attractive or repellent activity of carlina oxide toward *N. californicus* females changed over time. Leaf disks were sprayed with 10 μL of carlina oxide solution and the respective negative control using an airbrush, and were air-dried for 1 h. The spraying apparatus consisted of a compressor (FD-186 Piston Type 125 W Air Compressor, Fengda, Zhejiang, China) set at a pressure of 0.17 bar and an airbrush (Badger Air Brush 200-9-GFX, Chicago, IL, USA) positioned perpendicularly at a height of 30 cm from the sample using an adjustable metal stand

(Frasconi et al., unpublished data). The bioactivity of both doses was evaluated at 0 h (t0) and 48 h (t1) after the spraying treatment; at the end of the first bioassay, leaf disks were kept in the fridge for 48 h. A total of 30 adult females of *N. californicus* were tested for carlina oxide concentrations.

2.7 Statistical analyses

Binary logistic regression (BLR) was adopted to compare data on the mortality of *T. urticae* females and nymphs as well as on the mortality and repellence of *N. californicus* females. Data on egg mortality, mean oviposition rate and mean hatching time of *T. urticae* eggs and *N. californicus* females were transformed using an arcsine-square-root equation prior to generalized linear modeling (GLM) analysis. Abbott's formula⁶³ was used to correct mortality data when control mortality was recorded. When significant differences were found, the means were separated using Tukey's HSD test ($P = 0.05$). For instantaneous rate of increase and total toxic effect data, the jackknife method was used to create pseudo values,⁷⁰ considering that sample distributions on the various tests were unknown. Goodness-of-fit tests to ascertain the normality of distribution, and analyses of variance followed by Student's *t*-tests, were performed on these values. Differences were considered significant when 95% of fiducial limits did not overlap. Lethal concentrations necessary for 10% (LC₁₀), 30% (LC₃₀), 50% (LC₅₀) and 90% (LC₉₀) mortality were calculated using the probit model implemented in the Minitab program, adopting a 95% confidence level. All analyses were computed using the Minitab 17.0 software (Minitab Inc., State College, PA, USA).

Concerning behavioral assays on *N. californicus*, a contingency analysis was carried out to analyze the biological activity of carlina oxide toward *N. californicus* adults. If Pearson's χ^2 test was significant, a residual analysis was also carried out to determine which category (choices) majorly contributed to rejecting the null

hypothesis. All statistical analyses on behavioral tests were done using RStudio software; $P = 0.05$ was set as the significance threshold.

3 RESULTS

3.1 Toxicity of carlina oxide on *T. urticae*

The BLR on survival rates of TSSM, showed significant differences among the various concentrations tested ($\chi^2 = 227.71$; $P < 0.001$) and the two stages ($\chi^2 = 14.63$; $P < 0.001$). No interactions between concentration and stage were noted ($\chi^2 = 0.72$; $P = 0.397$), indicating that each concentration had the same effect on females and nymphs (Table 1).

The highest concentration (5000 $\mu\text{L L}^{-1}$) killed all females within the first 2 days, whereas mortality with 2500 $\mu\text{L L}^{-1}$ reached similar values after 4 days. Less toxic effects were shown by the other three concentrations, with scalar mortality over time. Analyzing data regarding the two stages together, three main groups were found: (i) control, (ii) 625 and 1250 $\mu\text{L L}^{-1}$, and (iii) 2500 and 5000 $\mu\text{L L}^{-1}$, being a concentration of 312.5 $\mu\text{L L}^{-1}$ like both the first and second group ($P < 0.05$). All the dead motile stages were found on the leaf disk; no mites were found on the wet cotton wool surrounding the leaf disk, indicating the absence of a repellent effect of carlina oxide toward *T. urticae*. GLM analysis performed on mean survival time (with the carlina oxide concentration and survival rate at 24, 48, 72 and 96 h included as factors), showed significant differences in the survival trend among concentrations ($F_{15,1439} = 3.5$; $P < 0.001$); that is, TSSM death occurred at a different time regardless of the effect recorded for each concentration at the end of the test.

As regards the mean survival time of motile stages, we found significant differences among the tested carlina oxide concentrations ($F_{5,539} = 97.76$; $P < 0.001$) and the two stages ($F_{1,539} = 6.62$;

Table 1. Susceptibility of females and nymphs of *Tetranychus urticae* to different concentrations of carlina oxide

Ontogenetic stage	Carlina oxide concentration ($\mu\text{L L}^{-1}$)	Survival/day (%), mean \pm SE				Survival time (days) mean \pm SE	Overall mortality (%)	Fitted probability of mortality (95% CI)	Toxicity classes*
		Day 1	Day 2	Day 3	Day 4				
Female	Control	100.0 \pm 0.0	98.3 \pm 1.67	96.7 \pm 2.34	85.0 \pm 4.65	3.80 \pm 0.070 ^a	15.0 \pm 4.65	23.56 (18.2–29.9)	
	312.5	90.0 \pm 3.91	80.0 \pm 5.21	75.0 \pm 5.64	68.3 \pm 6.06	3.13 \pm 0.185 ^{abc}	31.7 \pm 6.06	31.16 (24.4–37.5)	1
	625	86.7 \pm 4.43	81.7 \pm 5.04	68.3 \pm 6.06	48.3 \pm 6.51	2.85 \pm 0.184 ^{bc}	51.7 \pm 6.51	39.96 (33.9–46.3)	2
	1250	83.3 \pm 4.85	75.0 \pm 5.64	56.7 \pm 6.45	48.3 \pm 6.51	2.63 \pm 0.200 ^c	51.7 \pm 6.51	58.97 (52.3–65.4)	2
	2500	61.7 \pm 6.33	26.7 \pm 5.76	11.7 \pm 4.18	10.0 \pm 3.90	1.10 \pm 0.159 ^d	90.0 \pm 3.91	87.02 (80.7–91.5)	3
	5000	38.3 \pm 6.33	0.0	0.0	0.0	0.38 \pm 0.063 ^d	100.0 \pm 0.0	99.32 (98.1–99.8)	4
Nymph	Control	100.0 \pm 0.0	96.7 \pm 3.33	90.0 \pm 4.47	90.0 \pm 4.47	3.78 \pm 0.133 ^{ab}	10.0 \pm 4.47	11.17 (7.2–16.9)	
	312.5	100.0 \pm 0.0	100.0 \pm 0.0	93.3 \pm 4.22	86.7 \pm 6.67	3.80 \pm 0.101 ^{ab}	13.3 \pm 6.67	15.59 (10.6–22.4)	1
	625	93.3 \pm 4.22	90.0 \pm 4.47	76.7 \pm 8.03	73.3 \pm 6.67	3.33 \pm 0.216 ^{abc}	26.7 \pm 6.67	21.36 (15.2–29.1)	1
	1250	80.0 \pm 5.16	73.3 \pm 9.89	66.7 \pm 13.3	66.7 \pm 13.3	2.90 \pm 0.301 ^{abc}	33.3 \pm 13.3	36.97 (28.4–46.4)	2
	2500	46.7 \pm 11.2	36.7 \pm 8.03	26.7 \pm 6.67	16.7 \pm 3.33	1.27 \pm 0.291 ^d	83.3 \pm 3.33	73.23 (62.4–81.9)	4
	5000	16.7 \pm 6.15	16.7 \pm 6.15	10.0 \pm 4.47	10.0 \pm 4.47	0.53 \pm 0.234 ^d	90.0 \pm 4.47	98.35 (95.6–99.4)	4

Different letters within the survival time column, show significant differences (analysis of variance followed by Tukey pairwise comparisons $P < 0.05$). Binary logistic regression (BLR) was performed on mortality data. $P_{(0)} = \exp(Y') / (1 + \exp(Y'))$ was used for probability of mortality after definition of Y' for females and nymphs.

*Calculated on corrected mortality.⁶³




Figure 2. Oviposition rate (mean ± SE) of *Tetranychus urticae* exposed to different concentrations of carlina oxide. Above each column, different letters indicate significant differences (generalized linear model followed by Tukey's HSD test, $P < 0.05$).




Figure 3. Total toxic effect (E) (mortality and reduction of fertility) of carlina oxide on *Tetranychus urticae* females. Different letters above columns indicate significant differences (one-way analysis of variance followed by Tukey's HSD test, $P < 0.05$).

$P < 0.001$), although no interaction of concentrations × stage was found ($F_{5,539} = 0.88$; $P = 0.497$). Tukey's HSD test showed no significant effect of the lowest concentration ($312.5 \mu\text{L L}^{-1}$) on females and of the lower three concentrations (312.5 , 625 and $1250 \mu\text{L L}^{-1}$) on nymphs, compared with control. However, the two higher concentrations (2500 and $5000 \mu\text{L L}^{-1}$) significantly reduced the mean survival time of both motile stages (Table 1). The highest concentration of carlina oxide ($5000 \mu\text{L L}^{-1}$) showed very toxic effects (class 4), after 4 days, toward females, but was moderately toxic (class 3) toward *T. urticae* juveniles. Concentrations of 2500 and $1250 \mu\text{L L}^{-1}$ showed the same trend for females and nymphs (classes 3 and 2, respectively). By contrast, the two lowest concentrations (625 and $312.5 \mu\text{L L}^{-1}$) showed no toxic effect for juveniles but were slightly toxic for *T. urticae* females.

The different concentrations of carlina oxide also showed a secondary effect on the oviposition performance of treated TSSM females (Fig. 2). In fact, the tested concentrations significantly reduced the mite daily oviposition rate ($F_{5,872} = 105.63$; $P < 0.001$) and this was obviously reflected on the total toxic effect (E) of the various concentrations (Fig. 3). All concentrations showed significant negative effects toward TSSM eggs ($F_{5,179} = 455.0$; $P < 0.001$) (Table 2). Slightly toxic effects were found with $312.5 \mu\text{L L}^{-1}$ (22% mortality), whereas the remaining concentrations induced mortality rates of 83.7% to 100% (Table 2). Also, the mean hatching time was significantly affected

by carlina oxide in a concentration-dependent manner ($F_{4,609} = 57.83$; $P < 0.001$) (Table 2).

Carlina oxide LC_{50} was calculated by probit analysis for all *T. urticae* stages. Female and nymph mortalities fitted well with both the normal and Weibull distributions, whereas egg mortality did not fit a linear model. We adopted the Weibull distribution to define percentiles for the two motile stages and the normal distribution for eggs. The LC_{50} values obtained for *T. urticae* females and nymphs were 1145.1 and $1825.0 \mu\text{L L}^{-1}$, respectively (Table 3), showing that females were more susceptible than nymphs ($z = -2.53$, $P = 0.012$). As regards eggs, the LC detailed in Table 3 should be considered indicative because the goodness-of-fit test (Pearson) showed a lack of fit to the linear model of the probit analysis ($\chi^2 = 84.0$, $P < 0.001$).

3.2 Effect of carlina oxide on *T. urticae* population growth rate

The different carlina oxide concentrations significantly influenced the instantaneous rate of population growth ($F_{5,359} = 752.69$; $P < 0.001$). In particular, the growth rate calculated for the lowest concentration ($312 \mu\text{L L}^{-1}$) did not differ from the control ($r_i = 0.847$ and 0.955 , respectively). A positive growth index was also obtained with the $625 \mu\text{L L}^{-1}$ concentration ($r_i = 0.343$), even if it was statistically different from the lowest concentration.

Table 2. Susceptibility of *Tetranychus urticae* eggs to different concentrations of carlina oxide

Carlina oxide concentration ($\mu\text{L L}^{-1}$)	Cumulative percentage of hatched eggs (mean ± SE)				Hatching time (days) (mean ± SE)	Mortality (%)
	Elapsed days after egg laying					
	4	5	6	7		
Control	93.7 ± 1.55	97.7 ± 0.92	98.3 ± 0.84	100.0 ± 0.0	4.10 ± 0.026 ^a	0.0 ^a
312.5	55.7 ± 4.69	71.3 ± 3.77	76.7 ± 3.44	78.0 ± 2.89	4.39 ± 0.044 ^b	22.0 ± 2.89 ^b
625	2.7 ± 1.43	13.3 ± 2.81	16.3 ± 3.13	16.3 ± 3.13	5.02 ± 0.085 ^c	83.7 ± 3.13 ^c
1250	0.0	2.0 ± 0.88	4.3 ± 1.14	4.3 ± 1.14	5.60 ± 0.131 ^d	95.7 ± 1.14 ^d
2500	0.0	1.0 ± 0.56	2.3 ± 0.92	2.3 ± 0.92	5.57 ± 0.202 ^{cd}	97.7 ± 0.92 ^{de}
5000	0.0	0.0	0.0	0.0	0.0 ^e	100.0 ± 0.0 ^e

Different letters within a column show significant differences (one-way analysis of variance followed by Tukey pairwise comparisons, $P < 0.05$).

Table 3. Lethal concentrations (LC) of carlina oxide against different ontogenetic stages of *Tetranychus urticae*

Ontogenetic stage	LC ₁₀ $\mu\text{L L}^{-1}$ (95% CI)	LC ₃₀ $\mu\text{L L}^{-1}$ (95% CI)	LC ₅₀ $\mu\text{L L}^{-1}$ (95% CI)	LC ₉₀ $\mu\text{L L}^{-1}$ (95% CI)	LC ₉₅ $\mu\text{L L}^{-1}$ (95% CI)	Intercept \pm SE	Slope \pm SE	χ^2 (d.f.)
Female	181.5 (116.4–254.0)	560.7 (415.5–708.4)	1145.1 (918.1–1395.1)	5137.4 (3880.2–7516.7)	7434.5 (5330.9–11 893.5)	3.2447 \pm 0.045	6.21436 \pm 0.601	7.85 (7) $P = 0.346$ ns*
Nymph	256.1 (162.2–366.8)	852.3 (604.5–1152.1)	1825.0 (1333.6–2520.9)	9042.3 (5848.0–16 332.0)	13 409.4 (8195.0–26 771.3)	3.4594 \pm 0.075	3.7138 \pm 0.203	84.03 (3) $P < 0.001$
Egg	196.2 (170.8–220.1)	313.8 (286.1–339.6)	434.3 (405.5–462.6)	961.3 (887.2–1055.6)	1204.2 (1093.5–1351.0)	−9.79632 \pm 0.560	−9.79632 \pm 0.560	84.03 (3) $P < 0.001$

*Not significant.




Figure 4. Instantaneous rate of increase values estimated for *Tetranychus urticae* mites exposed for different concentrations carlina oxide. Different letters within columns denote significant differences among tests (generalized linear model followed by Tukey's HSD test, $P < 0.05$).

By contrast, negative values were obtained with the other three concentrations ($r_i = -0.202, -0.775$ and -0.991 for the 1250, 2500 and 5000 $\mu\text{L L}^{-1}$, respectively), showing a decreasing trend in population toward extinction (Fig. 4).

3.3 Side effects of carlina oxide on *N. californicus*

Side effects of carlina oxide on *N. californicus* females are reported in Table 4. Mortality in the four tests was low and not statistically different from the control ($\chi^2 = 0.00$; $P = 0.993$), whereas a different repellent effect was recorded among tests ($\chi^2 = 12.63$; $P < 0.001$). The oviposition rate of *N. californicus* in test D (females placed 48 h after the treatment) was the same as that recorded in the control, whereas it was significantly lower in the other tests ($F_{4,355} = 53.57$; $P < 0.001$).

The toxic effect (mortality) on *N. californicus* females occurred entirely in the first day, and the four tests were not significantly different from control showing a harmless effect (class 1). However, a strong repellent effect was recorded both when females were directly sprayed and remained on the sprayed leaf disk, and when unsprayed females were placed on freshly sprayed leaf disk (tests A and B, Table 4). In these tests, females trying to escape the treated leaf disk, drowned on the wet cotton wool surrounding the leaf disk. A significant repellent effect was also recorded when treated females were put on untreated leaf disks (test C, Table 4). It should be mentioned that the repellent effect of carlina oxide lasted for about 2 days. In fact, when unsprayed females were put on treated leaf disks 2 days after spraying (test D, Table 4), only a slight repellent effect was recorded within the first day ($P < 0.05$) and the oviposition rate did not differ from the control.

Carlina oxide caused no side effects on phytoseiid eggs (100% hatching) ($F_{1,166} = 0.57$; $P = 0.453$) (Table 5). However, the highest hatching percentages were recorded on the second day in the control (93.4%), and the third day in treated eggs (70.0%), indicating a significant slowdown in egg hatching ($F_{2,166} = 57.4$; $P < 0.001$) (Table 5).

3.4 Y-tube olfactometer bioassays on *N. californicus* adults

3.4.1 Preliminary bioassays

The contingency analysis highlighted a significant difference among the mite choices made at different carlina oxide concentrations ($\chi^2 = 46.192$, d.f. = 2, $P < 0.001$) (Fig. 5).

The residual analysis indicates a correlation between the escape of *N. californicus* females and the highest dose, which indicates a

Table 4. Side effects of carlina oxide (5000 $\mu\text{L L}^{-1}$) on *Neoseiulus californicus* females in the four different tests and in the control

Test	No.	Daily mortality (%) (mean \pm SE)			Toxicity classes*	Daily repellence (%) (mean \pm SE)			Eggs/female/day	Total negative effect (%)
		1	2	3		1	2	3		
A. Spraying on a female placed on the leaf disk	30	10.0 \pm 5.57	—	—	1	90.0 \pm 5.57 ^a	—	—	0.00 ^a	100.0 ^a
B. Spraying on the leaf disk; the female was placed on leaf disk 4 min later	15	6.7 \pm 6.67 ^a	0.0	—	1	80.0 \pm 10.7 ^a	13.3 \pm 9.09 ^a	—	0.02 ^a	96.7 ^a
C. Spraying on a female which was then moved on an untreated leaf disk	30	6.7 \pm 4.63 ^a	0.0	0.0	1	23.3 \pm 7.85 ^b	30.0 \pm 8.51 ^a	0.0	0.09 ^a	86.8 ^b
D. Spraying on the leaf disk; the female was placed on leaf disk after 48 h	15	0.0 ^a	0.0	0.0	1	13.3 \pm 9.09 ^b	0.0 ^b	0.0	0.62 ^b	1.6 ^c
Control	30	6.7 \pm 4.63 ^a	0.0	0.0	1	0.0 ^c	0.0 ^b	0.0	0.63 ^b	

Different letters denote significant differences among tests for oviposition rate and total negative effect (one-way analysis of variance followed by Tukey pairwise comparisons for $P < 0.05$).

*Calculated on corrected mortality.⁶³

repellent activity of the compound at 5000 $\mu\text{L L}^{-1}$. Indeed, 58% of predatory mites responded by escaping immediately after the release.

By contrast, there was a positive correlation between the choice of the control arm and the lowest dose (2500 $\mu\text{L L}^{-1}$).

3.4.2 Carlina oxide bioactivity over time

Investigating how the biological activity of carlina oxide changed in relation to a different substrate (the fresh bean leaf) and over time, the previously obtained results were partially corroborated. As in preliminary bioassays, a significant difference was found between the highest tested dose and the choice made by *N. californicus*, immediately after the leaf was sprayed (t0) ($\chi^2 = 24.649$, d.f. = 2, $P < 0.0001$). Residual analysis highlighted only an association between mite response and treatment (5000 $\mu\text{L L}^{-1}$), which translates to a greater preference for escape (non-random) than for the control or the treated arm.

At 2500 $\mu\text{L L}^{-1}$ of carlina oxide, the repellent action found in the preliminary results may be mitigated by the volatile compounds released by the leaf (Fig. 6). However, analyzing data after 48 h (t1), no significant differences were found between the dose and the choice made by the mite ($\chi^2 = 0.7213$, d.f. = 2,

$P = 0.6972$), meaning that carlina oxide no longer exerted a repellent effect on the predatory mite (Fig. 7).

Y-Tube results showed a significant repellent activity of the highest dose of carlina oxide at the time of application on both fresh bean leaf and filter paper. However, the repellent activity was reduced or even disappeared after 48 h.

4 DISCUSSION

Research for eco-compatible methods to control insect or mite infestations on crops are desirable in modern agriculture to achieve four main goals: (i) provide viable alternatives to synthetic




Figure 5. Contingency analysis on the dose of carlina oxide sprayed on filter paper and *Neoseiulus californicus* female choices in Y-tube olfactometer assays (based on a Pearson chi-squared test of independence, P -value reported in the plot). The area of each tile is proportional to the count of choices made by the predatory mite. The shading of the cells refers to the sign and magnitude of the respective Pearson residuals. The number of observations per cell is presented and each value is color-coded according to the dose ($P < 0.05$).

Table 5. Side effects of carlina oxide on eggs of *Neoseiulus californicus*

Treatment	Daily percentage of hatched eggs (mean \pm SE)			Hatching (%) (mean \pm SE)
	1	2	3	
1250 $\mu\text{L L}^{-1}$	13.3 \pm 6.31 ^a	16.67 \pm 6.92 ^a	70.0 \pm 4.66 ^a	100.0
Control	3.3 \pm 3.33 ^b	93.4 \pm 4.63 ^b	3.3 \pm 3.33 ^b	100.0

Within a column, different letters indicate significant differences among values (generalized linear model followed by Tukey pairwise comparisons, $P < 0.05$).




Figure 6. Contingency analysis on the dose of carlina oxide just sprayed (t0) on fresh bean leaves and *Neoseiulus californicus* female choices in Y-tube olfactometer assays (based on a Pearson chi-squared test of independence, P -value reported in the plot). The area of each tile is proportional to the count of choices made by the predatory mite. The shading of the cells refers to the sign and magnitude of the respective Pearson residuals. The number of observations per cell is presented and each value is color-coded according to the dose ($P < 0.05$).




Figure 7. Contingency analysis on the dose of carlina oxide sprayed 48 h before (t1) on fresh bean leaves and *Neoseiulus californicus* female choices in Y-tube olfactometer assays (based on a Pearson chi-squared test of independence, P -value reported in the plot). The area of each tile is proportional to the count of choices made by the predatory mite. The shading of the cells refers to the sign and magnitude of the respective Pearson residuals. The number of observations per cell is presented and each value is color-coded according to the dose ($P < 0.05$).

chemicals, (ii) reduce environmental pollution, (iii) reduce chemical residues in the agri-food chain, and (iv) tackle the thorny problem of resistant strains induced by the continuous use of synthetic pesticides. Moreover, the use of acaricides that are selective for natural enemies is particularly useful in IPM strategies.^{6,71}

Several studies have investigated the acaricidal activity, repellence and life-time parameters of different EOs and their bioactive constituents against *T. urticae*^{23,72–76} and, to a lesser extent, the side effects on predatory mites.^{33,73,75,77} To the best of our knowledge, this work is the first to study the toxic effects of carlina oxide on *T. urticae*, as well its side effects on the important biocontrol phytoseiid *N. californicus*.

Carlina oxide showed greater toxicity against females than nymphs of *T. urticae* ($LC_{50} = 1145.1$ and $1825 \mu\text{L L}^{-1}$ for females and nymphs, respectively), but the effects on mean survival time were the same for both stages, indicating a similar action against all mobile stages.

Carlina oxide can be considered the chemical main responsible for the plant insecticidal and acaricidal activity previously reported for *C. acaulis*.^{29,58,59} The triple bond of the propynyl chain in the molecule produces radicals generating oxidative damage.⁷⁸ This can be boosted under ultraviolet light.³⁰ Carlina oxide may also interact with the insect γ -aminobutyric acid (GABA) receptor.⁷⁸ Further studies on its mechanism of action are needed.

The toxic effects of other Asteraceae species have also been reported on *T. urticae* females. Chiasson *et al.*¹⁷ reported the lethal effects of *Artemisia absinthium* L. and *Tanacetum vulgare* L. EOs on tetranychid females, adopting 16-fold higher concentrations ($80\,000 \mu\text{L L}^{-1}$) in comparison with the highest ($5000 \mu\text{L L}^{-1}$) used in our experiments. In addition, carlina oxide showed 26-fold more toxicity compared with *Lippia gracilis* Schauer (Verbenaceae) EO, 34-fold more toxicity than azadirachtin, and 5-fold more toxicity than the synthetic acaricide fenpyroximate.⁷⁹

In addition to toxicity, a negative effect on the daily oviposition rate by all but the lowest concentration was also noted. However, toxic effects recorded on treated eggs from doses as low as

$625 \mu\text{L L}^{-1}$ indicate ovicidal activity of carlina oxide, which was confirmed by probit analysis estimating lethal concentrations. The lack of fit to a linear model was due to the high mortality ($>80\%$) observed by four of the five concentrations of carlina oxide (625 – $5000 \mu\text{L L}^{-1}$). These data indicate that small concentration increments between 312.5 and $625 \mu\text{L L}^{-1}$ result in a large increment in eggs mortality. Three days after treatment, *T. urticae* eggs were empty, with a dry residue inside the chorion, indicating arrest of the embryo development. Over time, the negative impact of carlina oxide on *T. urticae* females was reflected in the population growth rate. The negative values recorded with the three highest doses showed a combined effect of toxicity and decreased female fecundity, as demonstrated by the total effect of carlina oxide on females (Fig. 3). In contrast to various EOs and their major bioactives,^{33,80,81} no repellent effects on *T. urticae* have been observed for carlina oxide, indicating the absence of disturbing olfactory cues for all mobile stages. Otherwise, carlina oxide elicited a strong repellent effect on *N. californicus*, but no toxic effect was observed against phytoseiid females. The different behavioral responses of the two mite species exposed to the same compound dose could be linked to their chemoreceptor types. Recently, Su *et al.*⁸² reported that predatory mites, like *N. cucumeris*, rely mostly on gustatory receptors and ionotropic receptors for chemosensation. Ionotropic receptors represent the largest class of chemoreceptors in Acari and are particularly abundant in phytoseiid mites, with approximately 60 homologs, ten times those in spider mites.⁸² Thus, the abundance of chemoreceptor genes in predatory mites may imply a different chemosensory ability.⁸² The repellent effect, as also confirmed by our behavioral tests, vanishes after 48 h, allowing the predatory mite to again colonize the treated area.

Carlina oxide was shown to be harmless to *N. californicus* eggs, but all concentrations of the compound caused a significant delay in predator egg hatching. This has also been observed on tetranychid eggs. The harmless effects on predator eggs in comparison with the high toxicity toward the *T. urticae* eggs could be due to

the different exposure time as also seen for *P. persimilis* eggs treated with caraway EO.³³ In fact, *N. californicus* eggs hatch within 2 days, whereas tetranychid eggs hatch after 4–5 days; this longer period allows the active ingredient to penetrate the chorion and block embryonic development. The significative delay in eggs hatching recorded for *N. californicus* eggs, strengthens this hypothesis.

5 CONCLUSION

Carlina oxide was shown to be a new candidate ingredient for the development of sustainable products for controlling *T. urticae* infestations. Because cultivation of *C. acaulis* has been scarcely developed so far, the supply chain of this compound relies only on spontaneous plant populations. Thus, to boost the scalability of carlina oxide-based insecticidal formulations, several cropping systems should be carried out, especially in marginal areas, in addition to possible sustainable synthetic routes for the production of this polyacetylene⁸³ by agrochemical companies. At the same time, new advanced extraction techniques should be pursued to increase the yield of the active ingredient, as well as novel formulations to boost its bioactivity over time.^{52,84} Our findings show that carlina oxide reduces the longevity and fecundity of *T. urticae* adults. However, to predict the fate of a treated population over several generations, it is necessary to consider other important aspects, especially the development of resistance to these botanical products. Obviously, a reiterated use of the same compound increases the possibility of resistance development in *T. urticae*. Also, semi-field and field studies aiming to evaluate the efficacy of carlina oxide are needed for practical implementation under field and greenhouse conditions. Furthermore, the low negative impact on *N. californicus* makes this polyacetylene a good candidate in IPM programs, despite its repellent effects that, however, last for a very short period.

AUTHOR CONTRIBUTIONS

R Rizzo: conceptualization, methodology, investigation, validation, visualization, writing – original draft preparation, reviewing and editing. E Ragusa: investigation, data curation, reviewing and editing. G Benelli: conceptualization, methodology, resources, validation, formal analysis, supervision, writing – original draft preparation, reviewing and editing. G Lo Verde: conceptualization, methodology, validation, formal analysis, visualization, writing – original draft preparation, reviewing and editing. V Zeni: investigation, formal analysis, data curation, writing – original draft preparation, reviewing and editing. F Maggi: resources, writing – original draft preparation, reviewing and editing. R Petrelli: investigation, reviewing and editing. E Spinozzi: investigation, reviewing and editing. M Ferrati: investigation, reviewing and editing. M Sinacori: investigation, reviewing and editing. H Tsolakis: supervision, conceptualization, methodology, validation, formal analysis, visualization, writing – original draft preparation, reviewing and editing.

ACKNOWLEDGEMENTS

The authors are deeply indebted to Professor Alberto Lombardo for his suggestions on statistical analyses and to the two anonymous reviewers for their constructive comments. The study was partially funded by the University of Palermo (Fondo di Finanziamento per la Ricerca FFR 2018/2021, D13, by Gabriella Lo Verde and Haralabos Tsolakis).

CONFLICT OF INTEREST

The authors have no relevant financial or non-financial interests to disclose.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

- Knapp M, Sarr I, Gilioli G and Baumgärtner J, Population models for threshold-based control of *Tetranychus urticae* in small-scale Kenyan tomato fields and for evaluating weather and host plant species effects. *Exp Appl Acarol* **39**:195–212 (2006). <https://doi.org/10.1007/s10493-006-9018-1>.
- Pascual-Ruiz S, Aguilar-Fenollosa E, Ibáñez-Gual V, Hurtado-Ruiz MA, Martínez-Ferrer MT and Jacas JA, Economic threshold for *Tetranychus urticae* (Acari: Tetranychidae) in clementine mandarins *Citrus clementina*. *Exp Appl Acarol* **62**:337–362 (2013).
- Greco NM, Liljesthröm GG and Sánchez NE, Spatial distribution and coincidence of *Neoseiulus californicus* and *Tetranychus urticae* (Acari: Phytoseiidae, Tetranychidae) on strawberry. *Exp Appl Acarol* **23**:567–580 (1999).
- Easterbrook MA, Fitzgerald JD and Solomon MG, Biological control of strawberry tarsonemid mite *Phytonemus pallidus* and two-spotted spider mite *Tetranychus urticae* on strawberry in the UK using species of *Neoseiulus* (*Amblyseius*) (Acari: Phytoseiidae). *Exp Appl Acarol* **25**:25–36 (2001).
- Abad-Moyano R, Pina T, Pérez-Panadés J, Carbonell EA and Urbaneja A, Efficacy of *Neoseiulus californicus* and *Phytoseiulus persimilis* in suppression of *Tetranychus urticae* in young clementine plants. *Exp Appl Acarol* **50**:317–328 (2010).
- Van Lenteren JC and Woets J, Biological and integrated pest control in glasshouses. *Annu Rev Entomol* **33**:239–269 (1988).
- Nihoul P, Effect of temperature and relative humidity on successful control of *Tetranychus urticae* Koch by *Phytoseiulus persimilis* Athias-Henriot (Acari: Tetranychidae Phytoseiidae) in tomato crops under glasshouse conditions. *Int Symp Crop Prot* **57**:949–957 (1992).
- Cédola CV, Sánchez NE and Liljesthröm GG, Effect of tomato leaf hairiness on functional and numerical response of *Neoseiulus californicus* (Acari: Phytoseiidae). *Exp Appl Acarol* **25**:819–831 (2001).
- Wood E, Latli B and Casida JE, Fenazaquin acaricide specific binding sites in NADH: ubiquinone oxidoreductase and apparently the ATP synthase stalk. *Pestic Biochem Phys* **54**:135–145 (1996).
- Nauen R, Stumpf N, Elbert A, Zebitz CPW and Kraus W, Acaricide toxicity and resistance in larvae of different strains of *Tetranychus urticae* and *Panonychus ulmi* (Acari: Tetranychidae). *Pest Manage Sci* **57**:253–261 (2001).
- Van Leeuwen T, Vontas J, Tsagakarakou A, Dermauw W and Tirry L, Acaricide resistance mechanisms in the two-spotted spider mite *Tetranychus urticae* and other important Acari: a review. *Insect Biochem Mol Biol* **40**:563–572 (2010).
- Marcic D, Acaricides in modern management of plant-feeding mites. *J Pest Sci* **85**:395–408 (2012).
- Grafton Cardwell EE and Hoy MA, Comparative toxicity of avermectin B1 to the predator *Metaseiulus occidentalis* (Nesbitt) (Acari: Phytoseiidae) and the spider mites *Tetranychus urticae* Koch and *Panonychus ulmi* (Koch) (Acari: Tetranychidae). *J Econ Entomol* **76**:1216–1220 (1983).
- Ibrahim YB and Yee TS, Influence of sublethal exposure to abamectin on the biological performance of *Neoseiulus longispinosus* (Acari: Phytoseiidae). *J Econ Entomol* **93**:1085–1089 (2000).
- Hamedi N, Fathipour Y, Saber M and Garjan AS, Sublethal effects of two common acaricides on the consumption of *Tetranychus urticae* (Prostigmata: Tetranychidae) by *Phytoseius plumifer* (Mesostigmata: Phytoseiidae). *Syst Appl Acarol* **14**:197–205 (2009).
- Schmutterer H and Ascher KRS, Natural pesticides from the neem tree (*Azadirachta indica* A. Juss) and other tropical plants, in *Proceedings of the 3rd International Neem Conference; 1986 July 10–15; Nairobi, Kenya. Bonn, Germany*, ed. by Schmutterer H and Ascher KRS. Germany Agency for Technical Cooperation Eschbom, Germany (1987).

- 17 Chiasson H, Bostanian NJ and Vincent C, Acaricidal properties of a *Chenopodium*-based botanical. *J Econ Entomol* **97**:1373–1377 (2004). <https://doi.org/10.1093/jee/97.4.1373>.
- 18 Pavela R and Benelli G, Essential oils as ecofriendly biopesticides? Challenges and constraints. *Trends Plant Sci* **21**:1000–1007 (2016).
- 19 Jacobson M, Botanical pesticides. Past, Present, and Future, in *Insecticides of Plant Origin*, Vol. **387**, ed. by Arnason JT, Philogène BJR and Morand P. American Chemical Society, Washington DC, USA, pp. 1–10 (1989). American Chemical Society Symposium Series.
- 20 Regnault-Roger C, Vincent C and Arnason JT, Essential oils in insect control: low-risk products in a high-stakes world. *Annu Rev Entomol* **57**:405–424 (2012).
- 21 Hernández-Moreno D, Soffers AE, Falke HE, Rietjens IM and Murk AJ, Consumer and farmer safety evaluation of application of botanical pesticides in black pepper crop protection. *Food Chem Toxicol* **56**:483–490 (2013).
- 22 Isman MB and Grieneisen ML, Botanical insecticide research: many publications, limited useful data. *Trends Plant Sci* **19**:140–145 (2014).
- 23 Benelli G, Pavela R, Canale A, Nicoletti M, Petrelli R, Cappellacci L et al., Isofuranodiene and germacrone from *Smyrniolum olusatrum* essential oil as acaricides and oviposition inhibitors against *Tetranychus urticae*: impact of chemical stabilization of isofuranodiene by interaction with silver triflate. *J Pest Sci* **90**:693–699 (2017).
- 24 Kavallieratos NG, Boukouvala MC, Ntalaka CT, Skourti A, Nika EP, Maggi F et al., Efficacy of 12 commercial essential oils as wheat protectants against stored-product beetles, and their acetylcholinesterase inhibitory activity. *Entomol Gen* **41**:385–414 (2021).
- 25 Lengai GM, Muthomi JW and Mbega ER, Phytochemical activity and role of botanical pesticides in pest management for sustainable agricultural crop production. *Sci Afr* **7**:e00239 (2020).
- 26 Ghabbari M, Guarino S, Caleca V, Saiano F, Sinacori M, Baser N et al., Behavior-modifying and insecticidal effects of plant extracts on adults of *Ceratitis capitata*. *J Pest Sci* **91**:907–917 (2018).
- 27 Rizzo R, Lo Verde G, Sinacori M, Maggi F, Cappellacci L, Petrelli R et al., Developing green insecticides to manage olive fruit flies? Ingestion toxicity of four essential oils in protein baits on *Bactrocera oleae* (Rossi). *Ind Crops Prod* **143**:111884 (2020).
- 28 Rizzo R, Pistillo M, Germinara GS, Lo Verde G, Sinacori M, Maggi F et al., Bioactivity of *Carlina acaulis* essential oil and its Main component towards the olive fruit Fly, *Bactrocera oleae*: ingestion toxicity, electrophysiological and behavioral insights. *Insects* **12**:880 (2021).
- 29 Benelli G, Rizzo R, Zeni V, Govigli A, Samková A, Sinacori M et al., *Carlina acaulis* and *Trachyspermum ammi* essential oils formulated in protein baits are highly toxic and reduce aggressiveness in the medfly, *Ceratitis capitata*. *Ind Crops Prod* **161**:113191 (2021).
- 30 Benelli G, Ceccarelli C, Zeni V, Rizzo R, Lo Verde G, Sinacori M et al., Lethal and behavioural effects of a green insecticide against an invasive polyphagous fruit fly pest and its safety to mammals. *Chemosphere* **287**:132089 (2022).
- 31 Zeni V, Benelli G, Campolo O, Giunti G, Palmeri V, Maggi F et al., Toxics or lures? Biological and behavioral effects of plant essential oils on Tetranychidae fruit flies. *Molecules* **26**:5898 (2021).
- 32 Gugliuzzo A, Francardi V, Simoni S, Roversi PF, Ferrati M, Spinuzzi E et al., Role of plant essential oil nanoemulsions on host colonization by the invasive ambrosia beetle *Xylosandrus compactus*. *Ind Crops Prod* **195**:116437 (2023).
- 33 Tsolakis H and Ragusa S, Effects of a mixture of vegetable and essential oils and fatty acid potassium salts on *Tetranychus urticae* and *Phytoseiulus persimilis*. *Ecotoxicol Environ Saf* **70**:276–282 (2008).
- 34 Pavela R, Stepanycheva E, Shchenikova A, Chermenskaya T and Petrova M, Essential oils as prospective fumigants against *Tetranychus urticae* Koch. *Ind Crops Prod* **94**:755–761 (2016).
- 35 Camilo CJ, Alves Nonato CF, Galvão-Rodrigues FF, Costa WD, Clemente GG, Sobreira M et al., Acaricidal activity of essential oils. *Trends Phytochem Res* **1**:83–198 (2017).
- 36 de Santana MF, Câmara C, Monteiro VB, de Melo JR and de Moraes MM, Bioactivity of essential oils for the management of *Tetranychus urticae* Koch and selectivity on its natural enemy *Neoseiulus californicus* (McGregor): a promising combination for agroecological systems. *Acarologia* **61**:564–576 (2021).
- 37 Benvenuti S, Cioni PL, Flamini G and Pardossi AJWR, Weeds for weed control: Asteraceae essential oils as natural herbicides. *Weed Res* **57**:342–353 (2017).
- 38 Pouresmaeil M, Nojaded MS, Movafeghi A and Maggi F, Exploring the bio-control efficacy of *Artemisia fragrans* essential oil on the perennial weed *Convolvulus arvensis*: inhibitory effects on the photosynthetic machinery and induction of oxidative stress. *Ind Crops Prod* **155**:112785 (2020).
- 39 Isman MB, Miresmaili S and Machial C, Commercial opportunities for pesticides based on plant essential oils in agriculture, industry and consumer products. *Phytochem Rev* **10**:197–204 (2011).
- 40 Lubbe A and Verpoorte R, Cultivation of medicinal and aromatic plants for specialty industrial materials. *Ind Crops Prod* **34**:785–801 (2011).
- 41 Umpiérrez ML, Lagrega ME, Cabrera R, Grille G and Rossini C, Essential oils from Asteraceae as potential biocontrol tools for tomato pests and diseases. *Phytochem Rev* **11**:339–350 (2012).
- 42 Czerniewicz P, Chrzanoski G, Sprawka I and Sytykiewicz H, Aphicidal activity of selected Asteraceae essential oils and their effect on enzyme activities of the green peach aphid, *Myzus persicae* (Sulzer). *Pestic Biochem Phys* **145**:84–92 (2018).
- 43 Spinuzzi E, Maggi F, Bonacucina G, Pavela R, Boukouvala MC, Kavallieratos NG et al., Apiaceae essential oils and their constituents as insecticides against mosquitoes. A review. *Ind Crops Prod* **171**:113892 (2021).
- 44 Kavallieratos NG, Nika EP, Skourti A, Perinelli DR, Spinuzzi E, Bonacucina G et al., Apiaceae essential oil nanoemulsions as effective wheat protectants against five arthropod pests. *Ind Crops Prod* **186**:115001 (2022).
- 45 Iamónico D and Carlina L, in *Flora d'Italia*, Vol. **3**, 2nd edn, ed. by Pignatti S, Guarino R and La Rosa M. Edizioni Agricole di New Business Media, Bologna, Italy, pp. 1025–1031 (2018).
- 46 Strzemski M, Plachno BJ, Mazurek B, Kozłowska W, Sowa I, Lustofin K et al., Morphological, anatomical, and phytochemical studies of *Carlina acaulis* L. cypselas. *Int J Mol Sci* **21**:9230 (2020).
- 47 Strzemski M, Dzida K, Dresler S, Sowa I, Kurzepa J, Szymczak G et al., Nitrogen fertilisation decreases the yield of bioactive compounds in *Carlina acaulis* L. grown in the field. *Ind Crops Prod* **170**:113698 (2021).
- 48 Chalchat JC, Djordjevic S and Gorunovic M, Composition of the essential oil from the root of *Carlina acaulis* L. Asteraceae. *J Essent Oil Res* **8**:577–578 (1996).
- 49 Stojanović-Radić Z, Čomić L, Radulović N, Blagojević P, Mihajilov-Krstević T and Rajković J, Commercial *Carlinae* radix herbal drug: botanical identity, chemical composition and antimicrobial properties. *Pharm Biol* **50**:933–940 (2012).
- 50 Benelli G, Pavoni L, Zeni V, Ricciardi R, Cosci F, Cacopardo G et al., Developing a highly stable *Carlina acaulis* essential oil nanoemulsion for managing *Lobesia botrana*. *Nanomaterials* **10**:1867 (2020).
- 51 Ntalli N, Zochios G, Nikolaou P, Winkiel M, Petrelli R, Bonacucina G et al., *Carlina acaulis* essential oil nanoemulsion for managing *Meloidogyne incognita*. *Ind Crops Prod* **193**:116180 (2023).
- 52 Pavela R, Pavoni L, Bonacucina G, Cespi M, Cappellacci L, Petrelli R et al., Encapsulation of *Carlina acaulis* essential oil and carlina oxide to develop long-lasting mosquito larvicides: microemulsions versus nanoemulsions. *J Pest Sci* **94**:899–915 (2021).
- 53 Ferraz CA, Pastorinho MP, Palmeira-de-Oliveira A and Sousa ACA, Ecotoxicity of plant extracts and essential oils: a review. *Environ Pollut* **292**:118319 (2022).
- 54 dos Santos MC, Teodoro AV, Santos Menezes M, Pinto-Zevallos DM, de Fatima A-BM, Cruz Oliveira EM et al., Bioactivity of essential oil from *Lippia gracilis* Schauer against two major coconut pest mites and toxicity to a non-target predator. *Crop Prot* **125**:104913 (2019).
- 55 Giunti G, Benelli G, Palmeri V, Laudani F, Ricupero M, Ricciardi R et al., Non-target effects of essential oil-based pesticides for crop protection: impact on natural enemies, pollinators, and soil invertebrates. *Biol Control* **176**:105071 (2022).
- 56 Toledo PDF, Ferreira TP, Bastos IMAS, Rezende SM, Viteri Jumbo LO, Didonet J et al., Essential oil from *Negramina* (*Siparuna guianensis*) plants controls aphids without impairing survival and predatory abilities of non-target ladybeetles. *Environ Pollut* **255**:113153 (2019).
- 57 Walthall WK and Stark JD, Comparison of two population level ecotoxicological endpoints: the intrinsic (rm) and instantaneous (ri) rates of increase. *Environ Toxicol Chem* **16**:1068–1073 (1997).
- 58 Rosato A, Barbarossa A, Mustafa AM, Bonacucina G, Perinelli DR, Petrelli R et al., Comprehensive evaluation of the antibacterial and antifungal activities of *Carlina acaulis* L. essential oil and its nanoemulsion. *Antibiotics* **10**:1451 (2021).
- 59 Benelli G, Pavela R, Petrelli R, Nzekoué FK, Cappellacci L, Lupidi G et al., *Carlina* oxide from *Carlina acaulis* root essential oil acts as a potent mosquito larvicide. *Ind Crops Prod* **137**:356–366 (2019).

- 60 Swirski E, Amitai S and Dorzia N, Laboratory studies on the feeding habits, post-embryonic survival and oviposition of the predaceous mites *Amblyseius chilensis* Dosse and *Amblyseius hibisci* Chant (Acarina: Phytoseiidae) on various kinds of food substances. *Entomophaga* **15**:93–106 (1970).
- 61 Potter C, An improved laboratory apparatus for applying direct sprays and surface films, with data on the electrostatic charge on atomized spray fluids. *Ann Appl Biol* **39**:1–28 (1952).
- 62 Overmeer WPJ and Van Zon AQ, A standardized method for testing the side effects of pesticides on the predaceous mite, *Amblyseius potentillae* (Acarina: Phytoseiidae). *Entomophaga* **27**:357–363 (1982).
- 63 Abbott WS, A method of computing the effectiveness of an insecticide. *J Econ Entomol* **18**:265–267 (1925).
- 64 Hardman JM, Franklin JL, Moreau DL and Bostanian NJ, An index for selective toxicity of miticides to phytophagous mites and their predators based on orchard trials. *Pest Manage Sci* **59**:1321–1332 (2003).
- 65 Hall DJ, An experimental approach to the dynamics of a natural population of *Daphnia galeata mendotae*. *Ecology* **45**:94–112 (1964).
- 66 Sterk G, Hassan SA, Baillod M, Bakker F, Bigler F, Blümel S *et al.*, Results of the seventh joint pesticide testing programme carried out by the IOBC/WPRS-working group “pesticides and beneficial organisms”. *BioControl* **44**:99–117 (1999).
- 67 Canale A, Benelli G, Germinara GS, Fusini G, Romano D, Rapalini F *et al.*, Behavioural and electrophysiological responses to overlooked female pheromone components in the olive fruit fly, *Bactrocera oleae* (Diptera: Tephritidae). *Chemoecology* **25**:147–157 (2015).
- 68 Fonseca MM, Pallini A, Marques PH, Lima E and Janssen A, Compatibility of two predator species for biological control of the two-spotted spider mite. *Exp Appl Acarol* **80**:409–422 (2020).
- 69 Carpita A, Canale A, Raffaelli A, Saba A, Benelli G and Raspi A, (Z)-9-tricosene identified in rectal gland extracts of *Bactrocera oleae* males: first evidence of a male-produced female attractant in olive fruit fly. *Naturwissenschaften* **99**:77–81 (2012).
- 70 Meyer JS, Ingersoll CG, McDonald LL and Boyce MS, Estimating uncertainty in population growth rates: jackknife vs. bootstrap techniques. *Ecology* **67**:1156–1166 (1986).
- 71 Hoy MA and Ouyang YL, Selectivity of the acaricides clofentezine and hexythiazox to the predator *Metaseiulus occidentalis* (Acarina: Phytoseiidae). *J Econ Entomol* **79**:1377–1380 (1986).
- 72 Chiasson H, Bélanger A, Bostanian N, Vincent C and Poliquin A, Acaricidal properties of *Artemisia absinthium* and *Tanacetum vulgare* (Asteraceae) essential oils obtained by three methods of extraction. *J Econ Entomol* **94**:167–171 (2001).
- 73 Choi WI, Lee SG, Park HM and Ahn YJ, Toxicity of plant essential oils to *Tetranychus urticae* (Acarina: Tetranychidae) and *Phytoseiulus persimilis* (Acarina: Phytoseiidae). *J Econ Entomol* **97**:553–558 (2004).
- 74 Miresmailli S, Bradbury R and Isman MB, Comparative toxicity of *Rosmarinus officinalis* L. essential oil and blends of its major constituents against *Tetranychus urticae* Koch (Acarina: Tetranychidae) on two different host plants. *Pest Manage Sci* **62**:366–371 (2006).
- 75 Han J, Choi BR, Lee SG, Il Kim S and Ahn YJ, Toxicity of plant essential oils to acaricide-susceptible and -resistant *Tetranychus urticae* (Acarina: Tetranychidae) and *Neoseiulus californicus* (Acarina: Phytoseiidae). *J Econ Entomol* **103**:1293–1298 (2010).
- 76 da Camara CA, Akhtar Y, Isman MB, Seffrin RC and Born FS, Repellent activity of essential oils from two species of *Citrus* against *Tetranychus urticae* in the laboratory and greenhouse. *Crop Prot* **74**:110–115 (2015).
- 77 Araújo MJC, da Câmara CAG, Born FDS and de Moraes MM, Acaricidal activity of binary blends of essential oils and selected constituents against *Tetranychus urticae* in laboratory/greenhouse experiments and the impact on *Neoseiulus californicus*. *Exp Appl Acarol* **80**:423–444 (2020).
- 78 Negri R, Polyacetylenes from terrestrial plants and fungi: recent phytochemical and biological advances. *Fitoterapia* **106**:92–109 (2015).
- 79 de Souza BF, da Camara CAG, de Melo JPR and de Moraes MM, Acaricidal property of the essential oil from *Lippia gracilis* against *Tetranychus urticae* and a natural enemy, *Neoseiulus californicus*, under greenhouse conditions. *Exp Appl Acarol* **75**:491–502 (2018).
- 80 Motazedian N, Ravan S and Bandani AR, Toxicity and repellency effects of three essential oils against *Tetranychus urticae* Koch (Acarina: Tetranychidae). *J Agric Sci Technol* **14**:275–284 (2012).
- 81 Mozaffari F, Abbasipour H, Garjan AS, Saboori A and Mahmoudvand M, Toxicity and oviposition deterrence and repellency of *Mentha pulegium* (Lamiaceae) essential oils against *Tetranychus urticae* Koch (Tetranychidae). *J Essent Oil Bear Plants* **16**:575–581 (2013).
- 82 Su Y, Zhang B and Xu X, Chemosensory systems in predatory mites: from ecology to genome. *Syst Appl Acarol* **26**:852–865 (2021).
- 83 Spinozzi E, Ferrati M, Baldassarri C, Maggi F, Pavela R, Benelli G *et al.*, Synthesis of Carlina oxide analogues and evaluation of their insecticidal efficacy and cytotoxicity. *J Nat Prod* **86**:1307–1316 (2023).
- 84 Pavoni L, Pavela R, Cespi M, Bonacucina G, Maggi F, Zeni V *et al.*, Green micro- and nanoemulsions for managing parasites, vectors and pests. *Nanomaterials* **9**:1285 (2019).