

An overview of the Leucospidae (Hymenoptera, Chalcidoidea) of the Arabian Peninsula with description of a new species

Syed Kamran Ahmad¹, Syeda Uzma Usman², Farmanur Rahman Khan³, Hossein Lotfalizadeh⁴, Hassan A. Dawah⁵, Parvez Qamar Rizvi¹, Prince Tarique Anwar⁶

1 Department of Plant Protection, Faculty of Agricultural Sciences, Aligarh Muslim University, Aligarh – 202002, India

2 Department of Zoology, Mohammad Ali Jauhar University, Rampur – 244901, Uttar Pradesh, India

3 Department of Biology, Deanship of Educational Services, Qassim University, Buraidah – 51452, Al-Qassim, Saudi Arabia

4 Department of Plant Protection, East-Azerbaijan Agricultural and Natural Resources Research Center, Agricultural Research, Education and Extension Organization (AREEO), Tabriz, Iran

5 Centre for Environmental Research and Studies, Jazan University, P.O. Box 2095, Jazan, Saudi Arabia

6 Department of Zoology, School of Sciences, Aligarh College of Education, Chherat, Aligarh – 202122, India

Corresponding authors: Syeda Uzma Usman (insect.11lab@gmail.com); Prince Tarique Anwar (ta.friday@gmail.com)

Abstract

An overview of the family Leucospidae (Hymenoptera, Chalcidoidea) is provided for the leucospid fauna of the Arabian Peninsula. Two genera containing four species are identified based on morphometrics and colour patterns. One species, *Leucospis ayezae* Usman, Anwar & Ahmad, **sp. nov.**, is described. *Leucospis elegans* Klug had been previously recorded from Arabia Felix (= Yemen) and is recorded here for the first time from Saudi Arabia. The status of *Leucospis* aff. *namibica* from Yemen has been clarified, and this species is placed here in the genus *Micrapion* Kriechbaumer as *M. clavaforme* Stefan. An updated key and a map showing the distribution of the family Leucospidae in the Arabian Peninsula is provided. The occurrence and color morphs of all leucospid species that have been recorded so far from the region are briefly discussed.

Key words: Biodiversity, ectoparasitoids, new species, taxonomy

Introduction

Members of the family Leucospidae (Hymenoptera, Chalcidoidea) are large chalcid wasps (6–15 mm) and develop as ectoparasitoids on aculeate wasps or bees (Lima and Dias 2018). They are mostly dark brown, red, or yellow, with a patterned, orange or white body, metafemur enlarged with teeth, and strongly curved metatibia. Females typically have a recurved ovipositor which lies along the dorsal side of the metasoma. Leucospids are cosmopolitan in their distribution but rarely encountered, and there are 144 described species worldwide which belong to four genera (Noyes 2019).

The family is mostly represented by the genus *Leucospis* Fabricius, which accounts for more than 86% of the total number of species. Bouček (1974) provided a comprehensive taxonomic revision of the Leucospidae and provided separate keys to American, African, and Asiatic-Australian *Leucospis* species. More recently, Ye et al. (2017) recognized and provided a key to 12 *Leucospis*



Academic editor: Andreas Köhler

Received: 6 October 2023

Accepted: 10 December 2023

Published: 16 January 2024

ZooBank: <https://zoobank.org/8D05BBE1-922C-4E76-8256-5B02AF30D857>

Citation: Ahmad SK, Usman SU, Khan FR, Lotfalizadeh H, Dawah HA, Rizvi PQ, Anwar PT (2024) An overview of the Leucospidae (Hymenoptera, Chalcidoidea) of the Arabian Peninsula with description of a new species. ZooKeys 1189: 185–202. <https://doi.org/10.3897/zookeys.1189.113635>

Copyright: © Syed Kamran Ahmad et al. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Table 1. List of sampling sites with coordinates, altitude, and sampling methods for Leucospidae collected from Southwest of Saudi Arabia.

Locality	Coordinates	Altitude (m)	Method
Abha, Hay Al-Menhel, vegetable farm	18°12'N, 42°29'E	2214	MT
Abha, Hay Al-Nusub (Abha Farm Centre) vegetable farm	18°13'N, 42°30'E	2226	MT, SN
Jazan, Farasan Island, Aziz Yousef Village	16°40'N, 42°50'E	3	MT, SN
Najran, Al-Shurfa	17°31'N, 44°15'E	1342	MT

of card-mounted specimens were taken using a Nikon SMZ 1000 stereozoom binocular microscope. Figs 3B, 5B, 8C, D were taken using a video camera and Synaptics Automontage software to produce a montage image of the species. Photographs of the slide-mounted parts were taken with a Leica DFC295 digital camera attached to a Leica DM 2500 compound microscope with automontage facility. The final figures were prepared using Adobe Photoshop v. 7.0.

Measurements were made with the use of an ocular micrometer attached to the eyepiece of the microscope and were later converted into micrometers (μm). All the determined and type materials were deposited at the Insect Collections Department of Zoology, Aligarh Muslim University, Aligarh, Uttar Pradesh, India.

The terms mentioned in the text follow Bouček (1974) and Lima and Dias (2018).

List of abbreviations used in the text: **AOL**, anterior-ocular length; **MOD**, median ocellar diameter; **OCL**, ocular–occipit length; **OOL**, ocello–ocular line; **POL**, posterior ocellar line; **psa**, parascrobal area; **F**, antennal flagellomere; **PMV**, postmarginal vein of fore wing; **STV**, stigmal vein; **GT**, gastral or metasomal tergite; **MT**, Malaise trap; **SN**, Sweep net.

The following acronym is used for the depository:

- BMNH** Natural History Museum [formerly British Museum (Natural History)], Department of Entomology, London, UK;
- KSMA** King Saud University Museum of Arthropods, Plant Protection Department, College of Food and Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia;
- MNHN** Muséum National d'Histoire Naturelle, Paris, France;
- NMWC** The National Museum of Wales, Cardiff, UK;
- ZMHU** Zoological Museum, Humboldt University, Berlin, Germany;
- ZDAMU** Department of Zoology, Aligarh Muslim University, Aligarh, India.

Results

Four species, including one new species, are among the materials examined. These belong to one of two genera, either *Leucospis* or *Micrapion*. The family Leucopsidae has seven species in total known from the Arabian Peninsula, and all seven species are keyed below.

An updated key to females of Leucospidae from the Arabian Peninsula

Modified from Schmid-Egger 2010 and Gadallah et al. 2018.

- 1 Clypeus curved convexly at posterior margin and without a median tooth; mandibles thin, setose, and notched at apex; gaster distinctly clavate,

- basally narrow (Fig. 9A, C); GT4 with hind margin always produced backward and sharply angulate (Fig. 9A, C); GT6 fused to epipygium (gaster with coarse punctures; GT1 without band and with shiny interspaces; apical band of GT5 0.5× as broad as length of sheaths; metafemur relatively slender) ***Micrapion clavaforme* Steffan**
- Clypeus bilobed at posterior margin and often with a median tooth; mandibles robust, without setae; lower tooth stronger; gaster less clavate (Figs 3C, 4A, 5B, 7B, 8B); GT4 posteriorly straight; GT6 distinctly separated from epipygium **2 (genus *Leucospis* Fabricius)**
- 2 Pronotum with three distinct transverse carinae (Figs 3A, 5A, 7A, 8A) **3**
- Pronotum with at most two transverse, less-developed carinae (Fig. 4A) ... **6**
- 3 Ovipositor short, not reaching anterior margin of GT5 (Fig. 7A) ***Leucospis africana* Cameron**
- Ovipositor long, reaching at least posterior margin of GT4 or beyond (Figs 3A, 4A, 5B, 8B) **4**
- 4 Discal carina on pronotum weak and straight (Fig. 8A); metafemur slender with nine ventral teeth, basal tooth angular and pointed (Fig. 8A); ovipositor distinctly reaching beyond (hind fifth) posterior margin of GT1 (Fig. 8B) ***Leucospis insularis* Kirby**
- Discal carina on pronotum strong and angulate; metafemur oval with eight or nine ventral teeth, basal tooth triangular and robust; ovipositor hardly reaching posterior margin of GT1 **5**
- 5 Metafemur with nine ventral teeth (Fig. 3B); ovipositor hardly reaching posterior margin of GT4 (Fig. 3C) ***Leucospis ayezae* Usman, Anwar & Ahmad sp. nov.**
- Metafemur with eight ventral teeth (Fig. 5A); ovipositor clearly reaching posterior margin of GT1 (Fig. 5B) ***Leucospis elegans* Klug**
- 6 Pronotum red or orange, except black at base of mesopleuron; middle teeth of metafemur distinctly longer than basal triangular tooth ***Leucospis vanharteni* Schmid-Egger**
- Pronotum dark brown, except with a transverse yellow strip posteriorly between preapical and marginal carinae, continuing to lateral panel of pronotum as an oblique marking above ventral depression of panel (Fig. 4A); basal tooth of metafemur longer and more robust than any of the following teeth ***Leucospis arabica* Gadallah & Soliman**

Taxonomy

New species

***Leucospis ayezae* Usman, Anwar & Ahmad, sp. nov.**

<https://zoobank.org/02A1622A-412E-423C-B50C-028EE20D31EF>

Figs 2, 3

Type material. Holotype: SAUDI ARABIA • ♀; Asir, Abha, Hay Al-Nusub; 18°13'N, 42°30'E; 2226 m alt.; 24.vii.2013; H.A. Dawah leg.; ZDAMU Reg. No. HYM.CH.873, body, dissected head with one antenna and one dissected hind leg on card; one pair of fore wing and antenna on slide under two coverslips, slide HYM.06.




Figure 2. *Leucospis ayezae* Usman, Anwar & Ahmad sp. nov. holotype, female **A** head, frontal view **B** antenna **C** wings.




Figure 3. *Leucospis ayezae* Usman, Anwar & Ahmad sp. nov. holotype, female, habitus **A** lateral view **B** dorsal view.

Diagnosis. The new species is similar to *L. insularis* in having a yellow band on the pronotum and scutellum and distinct discal, preapical, and marginal carinae, but the new species differs from *L. insularis* as follows: discal carina on pronotum strong and angulate (discal carina on pronotum weak and straight in *L. insularis*); metafemora oval with eight ventral teeth, basal tooth triangular and robust (metafemora slender with nine ventral teeth, basal tooth angular and pointed in *L. insularis*); pubescence on sides of propodeum and metatibia relatively short and less dense (pubescence on sides of propodeum and metatibia long and more dense in *L. insularis*) ovipositor hardly reaching posterior margin of GT4 (ovipositor distinctly reaching beyond posterior margin of GT1 in *L. insularis*).

Description. Colour (Figs 2A, 3). Head dark brown; maxillary and labial palps yellowish brown; antenna dark brown except scape with posterior margin yellow. Mesosoma dark brown except a yellow transverse strip in front of discal carina, not continuing to sides of pronotum, and a narrow, transverse yellow strip on scutellum just above apex. Gaster largely reddish to dark brown, with transverse yellow strips medially on GT4 and apically on GT5. Basal two-thirds of ovipositor reddish brown; the rest dark brown. Pro- and mesofemur brown, with yellow tips where joining tibia; pro- and mesotibia reddish brown, with their margins yellow; hind legs dark brown except apex of coxa in ventral view and margins of femur yellow; all tarsi yellow. Fore wing below PMV and in apical half strongly infuscate, the rest hyaline.

Head (Fig. 2A). Head in frontal view 1.2× as broad as high and as wide as posterior margin of pronotum; less densely punctate, medially at psa smooth, setae on face and eyes silvery, erect, and less dense; POL 2× OOL and 3× MOD; scrobe 1.5× as wide as parascrobal area, transversely carinate; occipital carina distinctly visible between and beyond posterior ocelli; malar space 0.35× eye height and as long as F7; flagellum with erect, black setae; F1 widened apically, as long as broad and shortest of all funicular segments individually; F3–F5 subequal in length; F7 longest; clava 1.7× as long as broad, distinctly longer than F7 and F8 combined (Fig. 2B); mandible tridentate.

Mesosoma (Fig. 3A, C). Mesosoma densely punctate, punctures setigerous, with dense, long, pale setae; setae denser on propodeal callus; discal, preapical, and marginal carinae well developed, raised, and angulate; posterior margin of scutellum with punctures in a line with margins and with carina; propodeum medially with a complete carina. Hind leg with coxa punctate, carinate and subserrate posterodorsally; metafemur oval, 1.9× as long as broad, punctate, and setose, with eight ventral teeth; basal tooth robust, 3–5 longer than rest and subequal (Fig. 3B); metatibia with spine subequal to spur. Fore wing 3.2× as long as broad, with dense, black setation towards apical margin; STV with bifurcate uncus; uncus longer than apical process of stigmal vein (Fig. 2C). Hind wing 4.4× as long as broad (Fig. 2C).

Metasoma (Fig. 3A, C). Gaster moderately punctate, with dense, pale setae; density of setae more at epipygium. GT1 wider than long, interiorly with triangular process attached to petiole and, medially with a raised carina, narrower than GT4 in dorsal view; GT4 with posterior margin entire; ovipositor sheaths long, nearly reaching anterior margin of GT3.

Measurements (holotype, mm): head width:length:height, 1.4:0.7:1.2; AOL, 0.14; MOD, 0.12; OCL, 0.03; OOL, 0.19; POL, 0.38; sh [scrobe height], 0.51; sw [scrobe weight], 0.4; psa, 0.33; ceh [compound eye height], 0.82; mls [malar space], 0.28; antennal segments length:width – radicle, 0.13:0.16; scape, 0.64:0.2; pedicel, 0.24:0.17; F1, 0.16:0.16; F2, 0.22:0.2; F3, 0.25:0.24; F4, 0.25:0.25; F5, 0.25:0.27; F6, 0.25:0.28; F7, 0.28:0.22; F8, 0.24:0.32; clava, 0.56:0.33; pronotum, 0.56; mesoscutum, 0.72; scutellum, 0.61; dorsellum, 0.16; propodeum, 0.24; fore wing length:width, 4.4:1.35; hind wing length:width, 3.1:7; metacoxa, 0.96:0.85; metafemur, 1.24:0.64; metatibia, 1.12; metatarsus, 1.12; petiole, 0.16:0.37; gaster, 2.8; GT1, 0.9; GT3, 0.1; GT4, 0.4; GT5, 0.7; GT6, 0.2; ovipositor, 1.4; hypopygeum, 0.8.

Male. Unknown.

Host. Unknown.

Distribution. Saudi Arabia: Asir.

Etymology. The species name after Ayeza Tarique, daughter of the authors SUU and PTA.

Other species

***Leucospis arabica* Gadallah & Soliman, 2018**

Fig. 4

Leucospis arabica Gadallah & Soliman in Gadallah et al. 2018: 2079, female, male. Holotype, female (KSMA), Saudi Arabia (Jazan, Farasan Islands), not examined.

Materials examined. 8♀, 9♂. SAUDI ARABIA • Jazan, Farasan Island, Aziz Yousef Village; 16°40'N, 42°50'E; 3 m alt.; 6♀, 9♂ (each on cards; 2 females, 1 male with one pair of wings on slide under 1 coverslip, slide No. HYM.02, 03, 11), 15.v.2017; S.K. Ahmad leg.; 2♀ (on cards); 13.v.2017; H.A. Dawah leg.; ZDAMU.

Remarks. The examined specimens were collected from the type locality and differ from the holotype in size. The females were 5–15 mm long and males 2–10 mm long. In both sexes, the size of the yellow patch on the metafemur varies minute to broad.

Host. Unknown.

Distribution. Saudi Arabia: Jazan (Farasan Islands) and Egypt (Sinai Peninsula).

***Leucospis elegans* Klug, 1834**

Figs 5, 6

Leucospis elegans Klug, 1834: 26. Holotype, female (ZMHU), Yemen, not examined.

Materials examined. 2♀, 1♂. SAUDI ARABIA • Jazan, Farasan Island, Aziz Yousef Village; 16°40'N, 42°50'E; 3 m alt.; 2♀ (on cards, one pair of fore wing of one female specimen on slide under 1 coverslip, slide No. HYM.04; one pair of fore wing and antenna of other female specimen on slide under 2 coverslips, slide No. HYM.05), 15.v.2017; S.K. Ahmad leg.; 1♂ (on card, one pair of fore wing and antenna on slide under 2 coverslips, slide No. HYM.10), 1.ii.2015; H.A. Dawah leg.; ZDAMU.




Figure 4. *Leucospis arabica* Gadallah & Soliman, habitus A female, lateral view B male, lateral view.




Figure 5. *Leucospis elegans* Klug, female, habitus **A** lateral view **B** dorsal view.




Figure 6. *Leucospis elegans* Klug, males, habitus, lateral view **A** reddish brown-yellow morph **B** dark brown-pale yellow morph.

Remarks. This is the first record of *L. elegans* from Saudi Arabia. However, Bouček (1974) included it in the fauna of Saudi Arabia but referred to Arabia Felix, which is a former name for Yemen. He briefly provided a diagnosis of *L. elegans* and described the male for the first time.

Host. Unknown.

Distribution. Afrotropical, Palaearctic, Oriental (Klug 1834; Bouček 1959, 1974; Narendran 1986; Schmid-Egger 2010; Madl and Schwarz 2014; Gadallah et al. 2018). Yemen (as Arabia Felix; Bouček 1974). Saudi Arabia (new record).

***Leucospis africana* Cameron, 1907**

Fig. 7

Leucospis africana Cameron, 1907: 204. Lectotype, female (BMNH), designated by Bouček 1974: 104, South Africa (Cape Province), examined (illustrations only).

Remarks. Bouček (1974) recorded *L. africana* from several African countries and provided a brief diagnosis of females and described the males. He further recorded its host for the first time. Gadallah et al. (2018) recorded males from Saudi Arabia and provided a detailed diagnosis of males and a key to identify it from other species of Saudi Arabia. Here, we figure the lectotype for the first time.

Host. *Serapista denticulata* (Smith) (Hymenoptera, Megachilidae) (Bouček 1974); *Megachile spinarum* Cockerell (Hymenoptera, Megachilidae) (Gess 1981).

Distribution. Afrotropical: Burundi, Central African Republic, Democratic Republic of Congo, Eritrea, Ethiopia, Ghana, Ivory Coast, Kenya, Lesotho, Malawi, Mozambique, Nigeria, Rhodesia, South Africa, Tanzania, Uganda, Zambia, Zimbabwe (Cameron 1907; Bouček 1974; Noyes 2019); Saudi Arabia (Gadallah et al. 2018).

***Leucospis insularis* Kirby, 1900**

Fig. 8

Leucospis insularis Kirby, 1900: 13. Holotype, female (BMNH), Yemen (Socotra Island), examined (illustrations only).

Remarks. *Leucospis insularis* is only known from the type locality Socotra Islands (Yemen). Bouček (1974) included *L. insularis* in a key to African *Leucospis*. Schmid-Egger (2010) included it in his key to Arabian species. Here, we figure the holotype for the first time.

Host. Unknown.

Distribution. Afrotropical: Socotra Islands (Yemen) (Kirby 1900).

***Micrapion clavaforme* Steffan, 1948**

Fig. 9

Micrapion clavaforme Steffan, 1948: 85, female. Lectotype, female (MNHN), designated by Bouček 1974: 220, Gabon (Ogowe), not examined.

Material examined. 3♀, 2♂. SAUDI ARABIA • Asir, Abha, Hay Al-Menhel; 18°12'N, 42°29'E; 2214 m alt.; 2♀ (one on card; one on card with one pair of fore wing on slide under 1 coverslip, slide No. HYM.01), 20.xii.2014; H.A. Dawah leg.; Najran • Al-Shurfa, Saleh Maqbol Farm, 17°31'N, 44°15'E; 1342 m alt.; 1♀ (on card, one




Figure 7. *Leucospis africana* Cameron, lectotype, female (photographs courtesy of Natalie Dale-Skey Papilloud, BMNH) **A** lateral view **B** dorsal view **C** labels.



Figure 8. *Leucospis insularis* Kirby, holotype, female (photographs courtesy of Natalie Dale-Skey Papilloud, BMNH) **A** lateral view **B** dorsal view **C** labels.




Figure 9. *Micrapion clavaforme* Steffan, habitus, lateral view **A, B** dark brown-pale yellow morph **A** female **B** male **C, D** reddish brown-yellow morph **C** female **D** male.

pair of fore wing on slide under 1 coverslip, slide No. HYM.09), 17.ix.2014; H.A. Dawah leg.; Asir, Abha, Hay Al-Nusub, 18°13'N, 42°30'E; 2226 m alt.; 2♂ (one on card; one on card with one pair of fore wing on slide under 1 coverslip, slide No. HYM.07), 3.vi.2015; H.A. Dawah leg.; ZDAMU.

Remarks. Females and males were collected in the present study from two sites in Saudi Arabia. They agree fairly well with the original description of *M. clavaforme* and the diagnoses by Bouček (1974) and Gadallah et al. (2018). In both sexes there are two colour morphs, one brown with ivory stripes and another reddish brown with yellow stripes. All specimens, however, exhibit almost no variation in stripe patterns and wing infuscation.

Schmid-Egger (2010) tentatively identified two *Leucospis* specimens from Yemen as *L. aff. namibica*. On close examination of his figure (Schmid-Egger 2010: 321, pl. 3) there is no doubt that these specimens are not a *Leucospis* species but *Micrapion* Kriechbaumer instead. Here, these specimens are re-identified as *M. clavaforme*.

Host. Solitary bees: *Ceratina* Latreille (Bouček 1974).

Distribution. Afrotropical: (Steffan 1948; Bouček 1974). Saudi Arabia (Al Bahah, Asir, Najran) (Gadallah et al. 2018); Yemen (Schmid-Egger 2010 as *Leucospis* aff. *namibica*).

Acknowledgements

We thank Dr Mohammad Hayat and Dr Shahid Bin Zeya (ZDAMU) for encouragement and for providing space to carry out our research. Natalie Dale-Skey Papilloud (BMNH), kindly provided photographs of type materials of *Leucospis africana* and *L. insularis*. The National Museum Wales is gratefully acknowledged for their painstaking efforts in imaging Figs 3B, 5B, and 8C. D. Farmanur R. Khan thanks the Deanship of Scientific Research, Qassim University, for funding the publication of this study.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by Qassim University.

Author contributions

SK Ahmad and HA Dawah carried out sampling identified the family Leucospidae, conceived the idea of the research and, helped in preparing the map work. SU Usman and PT Anwar identified the genus and species and, prepared the manuscript. H Lotfalizadeh provided various information and literature on the Leucospidae and, critically revised the MS. FR Khan helped in preparation of the manuscript and provided several information about the Arabian fauna of chalcid wasps. The entire work was carried out under the mentorship of PQ Rizvi.

Author ORCIDs

Syed Kamran Ahmad  <https://orcid.org/0000-0002-6211-2345>

Syeda Uzma Usman  <https://orcid.org/0000-0003-2578-2515>

Farmanur Rahman Khan  <https://orcid.org/0000-0001-8906-709X>

Hossein Lotfalizadeh  <https://orcid.org/0000-0002-7927-819X>

Hassan A. Dawah  <https://orcid.org/0000-0001-5642-7247>

Parvez Qamar Rizvi  <https://orcid.org/0000-0002-2965-1806>

Prince Tarique Anwar  <https://orcid.org/0000-0002-1682-0033>

Data availability

All of the data that support the findings of this study are available in the main text.

References

Anwar PT, Zeya SB, Veenakumari K (2020) Fairyfly genus *Camptoptera* Foerster (Hymenoptera: Chalcidoidea: Mymaridae) in India and Sri Lanka, with descriptions of

- eleven new species. In: Paulus HF (Ed.) *Zoologica* (Vol. 165). Schweizerbart Science Publishers, Frankfurt, 89 pp.
- Bouček Z (1959) A revised key to the West-Palaeartic species of *Leucospis* (Hym., Chalc.), with some new synonymy. *Acta Entomologica Musei Nationalis Pragae* 33(562): 435–444.
- Bouček Z (1974) A revision of the Leucospidae (Hymenoptera: Chalcidoidea) of the world. *Bulletin of the British Museum (Natural History). Entomology* 23: 1–241. <https://doi.org/10.5962/p.140909>
- Cameron P (1907) Descriptions of species of parasitic Hymenoptera, chiefly in the collections of the South African Museum, Cape Town. (Second paper). *Annals of the South African Museum* 5: 203–225.
- Gadallah NS, Soliman AM, Abu El-Ghiet UM, Elsheikh TY, Al Dhafer HM (2018) The family Leucospidae (Hymenoptera: Chalcidoidea) from the South of Saudi Arabia, with the first report of the genus *Micrapion* and description of *Leucospis arabica* sp. nov. *Journal of Natural History* 52(31–32): 2071–2096. <https://doi.org/10.1080/00222933.2018.1510557>
- Gess FW (1981) Some aspects of an ethological study of the aculeate wasps and the bees of a karroid area in the vicinity of Grahamstown, South Africa. *Annals of the Cape Provincial Museums. Natural History* 14(1): 1–80.
- Hesami S, Akrami MA, Baur H (2005) *Leucospis dorsigera* Fabricius (Hymenoptera, Leucospidae) as a hyperparasitoid of Cerambycidae (Coleoptera) through Xoridae (Hymenoptera: Ichneumonidae) in Iran. *Journal of Hymenoptera Research* 14(1): 66–68.
- Kareem AA, Lotfalizadeh H, Aljaafari RK (2020) First record of *Leucospis bifasciata* Klug (Hymenoptera: Leucospidae) in Iraq. *Journal of Insect Biodiversity and Systematics* 6(3): 223–228. <https://doi.org/10.52547/jibs.6.3.223>
- Kirby WF (1900) [The expedition to Sokotra.] XII. Description of new Hymenoptera. *Bulletin of the Liverpool Museums* 3: 13–24.
- Klug FF (1834) *Symbolae physicae, seu icons et descriptiones insectorum, quae in itinere per Africam borealem et Asiam occidentalem Frederici Guilelmi Hemprich et Christiani Godofredi Ehrenberg, Studio, novae aut illustratae redierunt. Pars Zoologica. Insecta*, 4 Berlin: G. Reimer, 31–40.
- Lima AR, Dias PG (2018) The New World species of *Leucospis* Fabricius, 1775 (Hymenoptera, Chalcidoidea, Leucospidae): an update of Bouček's revision with description of two new species from Brazil. *Zootaxa* 4441(1): 001–045. <https://doi.org/10.11646/zootaxa.4441.1.1>
- Lotfalizadeh H, Fakhrzadeh N (2012) A short review of the family Leucospidae (Hym.: Chalcidoidea) in Iran. *Biharean Biologist* 6(1): 51–54.
- Madl M, Schwarz M (2014) Notes on Palaeartic species of the family Leucospidae (Hymenoptera, Chalcidoidea), with new records from North Africa and Middle East. *Linzer Biologische Beiträge* 46(2): 1569–1580.
- Narendran TC (1986) Family Leucospidae. In: Subba Rao BR, Hayat M (Eds) *The Chalcidoidea (Insecta; Hymenoptera) of India and the adjacent countries. Oriental Insects* 20: 43–45. <https://doi.org/10.1080/00305316.1986.10433717>
- Noyes JS (1982) Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). *Journal of Natural History* 16(3): 315–334. <https://doi.org/10.1080/00222938200770261>
- Noyes JS (2019) Universal Chalcidoidea Database. <https://www.nhm.ac.uk/our-science/data/chalcidoids>

- Schmid-Egger C (2010) Order Hymenoptera, family Leucospidae. *Arthropod Fauna of the UAE* 3: 319–324.
- Steffan JR (1948) Le genre *Micrapion* Kriechb. (Hym. Chalcididae). Description de trois espèces nouvelles. *Bulletin de la Société Entomologique de France* 53(5): 81–88. <https://doi.org/10.3406/bsef.1948.16036>
- Ye X-H, Achterberg CV, Yue Q, Xu Z-F (2017) Review of the Chinese Leucospidae (Hymenoptera: Chalcidoidea). *ZooKeys* 651: 107–157. <https://doi.org/10.3897/zookeys.651.11235>