

The effectiveness of the use of the drug "Furinaid" in complex therapy in the treatment of idiopathic cystitis in small pets

Tatyana Alekseeva¹, Olga Krotova^{1,2*}, Konstantin Savenkov³, Dayana Ubushaeva⁴, Anna Nimeeva⁴, and Hongr Chuev⁴

¹Don State Agrarian University, Krivoshlykov str., 24, 346493 Rostov region, P. Persianovsky, Russia

²Don State Technical University, 1, Gagarin Sq., 344003 Rostov-on-Don, Russia

³St. Petersburg State Agrarian University, Peterburgskoe shosse, 2, 196601 Pushkin, St. Petersburg, Russia

⁴Kalmyk State University named after B.B. Gorogovikov, Pushkin St., 11, 358000 Elista, Russia

Abstract. The disease of the lower urinary tract in cats is clinically significant pathology affecting about 7% of cats. Idiopathic feline cystitis (ICC) is diagnosed in more than half of cases of lower urinary tract disease in cats under the age of ten years. The cat HCC is a diagnosis of exclusion, in other words, urolithiasis and bacterial infections of the urinary tract should be excluded from the diagnosis. There are several reasons leading to the occurrence of diseases of the lower urinary system of cats. However, the idiopathic form of this disease (idiopathic cat cystitis) is one of the most common today. Stress, as it is now known, plays a very important role in the initiation and/or aggravation of the course of HCC. Worldwide, idiopathic cystitis is the most common cause of diseases of the lower urinary tract of cats, both males and females. The development of effective methods for the treatment of idiopathic cystitis in cats is one of the promising areas of veterinary science and practice. The article presents the results of studies obtained in assessing the effectiveness of the use of the drug "Furinaid" in complex therapy in the treatment of idiopathic cystitis in cats. Based on the obtained research results, it was found that 100% of the animals recovered in all the test groups, which indicates that the animals were properly treated, but the scheme using the drug "Furinaid" was therapeutically more effective. Therefore, it can be argued that in pharmacotherapy of idiopathic cystitis in cats, the use of drugs containing glucosaminoglycans is of great importance.

1 Introduction

Idiopathic cystitis is an inflammatory disease of the bladder in cats, the cause of which remains unknown ("idiopathic" means "without a known cause"). This is a fairly common condition and can occur in cats of any age and breed. Although the exact mechanisms of the development of idiopathic cystitis are unknown, it is believed that a combination of factors

* Corresponding author: alb9652@yandex.ru

such as changes in urine composition, stress, poor nutrition, decreased immunity and genetic predisposition occur [1-2].

This disease is one of the most common diseases of NMP in cats. The widespread spread of the disease, the considerable frequency of recurrence, indicate that the issues of prevention and treatment of HCC remain open and extremely necessary [3-5].

The discussion on the problem of the prevalence and etiology of idiopathic cat cystitis is characterized by significant differences in views. The causes of interstitial or idiopathic cystitis have not yet been established. It is believed that one of the provoking factors is stress (physical, emotional), namely, a change in the hypothalamic-pituitary-adrenal axis. Stress and anxiety can act as triggers in the process of the appearance and maintenance of clinical manifestations of idiopathic cystitis [6-8].

It is assumed that as a result of violation of the integrity of the glycosaminoglycan (GAG) barrier of the bladder, which, in turn, passes into the transitional epithelium, the nerve endings of the transitional epithelium of the bladder are exposed. When signals are transmitted to the overlying centers, an axon reflex develops with the final release of histamine and provocation of an inflammatory process in the urethra, which can cause the development of symptoms typical of cystitis. When making a differential diagnosis, it is necessary to exclude pathologies such as urolithiasis, bacterial infections of the urinary tract, bladder tumors and urinary incontinence. Successful diagnosis depends on the possibility of a comprehensive and careful study of clinical signs that may affect various organ systems, and the study of behavioral changes. [9-11].

At the moment, there are no consistently reliable diagnostic markers that make it possible to diagnose idiopathic cystitis and differentiate various clinical and pathological phenotypes of the disease.

Correction of BCC consists of two main stages: relief of stress and reduction of urine density by increasing the volume of fluid consumed. Additional methods, depending on the degree of damage to the bladder wall and the severity of cystitis symptoms, may include the use of anti-inflammatory, painkillers and antispasmodic drugs [12-14].

The drug "Furinaid" (manufacturer TRM, Ireland) was specially developed for cats to restore the protective layer of the urinary tract and treat idiopathic cystitis of various etiologies. The active substance of the drug — N-acetylglucosamine — is a structural unit of glycosaminoglycans that form a protective film on the mucous membrane of the bladder. In this form, the substance is well absorbed through the gastrointestinal tract, attaches to the damaged epithelium and increases the tightness of the bladder and the resistance of the mucous membrane to harmful effects and inflammation. Thus, the main cause of cystitis of any etiology is eliminated — a lack of glycosaminoglycans [15-17].

In connection with the above, the purpose of our work is to study the effectiveness of the use of the drug "Furinaid" in complex therapy in the treatment of idiopathic cystitis in cats.

2 Materials and methods

The research was carried out by the veterinary clinic "White Fang" (Novocherkassk) from September 2022 to June 2023 18 cats with clinical signs of cystitis of various breeds and genders aged 1-6 years, with a live weight of 3.5 - 6.0 kg were selected for research in the test period on the principle of analogues. Two groups were formed of them.

Diagnosis of idiopathic cystitis was carried out comprehensively, taking into account anamnestic data, clinical signs, as well as laboratory tests of blood and urine of sick animals. To exclude neoplasms, abnormal development of the urinary system organs, urolithiasis, ultrasound and X-ray examination of the bladder was performed. Bacterial cystitis was excluded during bacteriological examination of urine of animals of both groups.

During the collection of anamnesis, the number of manifestations of cystitis and the presence/absence of stress factors were taken into account. Clinical manifestations were determined before the treatment and 10 days after its completion. The main clinical signs of idiopathic cystitis in both groups are: frequent urination, tension, vocalization during urination, hematuria, urination outside the tray, increased groin care and the presence of signs of neurotic behavior.

The animals were on outpatient treatment. The animals of both groups were transferred to dietary nutrition for the period of the experiment. A prerequisite for the owners was strict compliance with the conditions of watering for cats, frequent change of clean water and unlimited access to water. Also, for all animals, it was recommended to create a favorable environment that helps minimize the impact of stress triggers.

Patients of the first and second experimental groups were transferred to the therapeutic food of Farmina Vet Life Struvite Management (dietary nutrition for cats developed to manage relapses of urolithiasis and for the treatment of idiopathic cystitis).

Scheme I. Animals of the first experimental group were treated with the analgesic and anti-inflammatory drug Meloxicvet 0.2% solution for injection at a dose of 0.05–0.1 mg / kg of body weight 1 time / day course - 3 days; oral suspension of Stop-cystitis for cats 2 times a day, 2 ml per 5 kg of MT for 10 days; the drug Relaxes 1 table. per kg of MT 1 time per day orally together with food for 1 month, the drug for pheromotherapy "Felivey" (a fraction of pheromones of the facial glands in a cat) in the form of a spray was applied once a day to cat habitat objects (couches, scratching posts, trays, upholstered furniture and others).

Scheme II. Animals of the second experimental group were treated with the analgesic and anti-inflammatory drug Meloxicvet 0.2% solution for injection at a dose of 0.05–0.1 mg / kg of body weight 1 time / day in / m course - 3 days; oral suspension of Stop-cystitis for cats 2 times a day, 2 ml per 5 kg MT for 10 days; the drug for pheromotherapy "Felivey" in the form of a spray was applied once a day to the objects of the habitat of cats; the drug Furinaid: during the first 14 days, the dosage for cats is 2.5 ml of the drug orally per animal 1 time per day, the next 14 days - 1.25 ml of the drug 1 time per day; the drug Relaxes 1 table. per kg of MT 1 time per day orally with food for 1 month.

Clinical studies were carried out using examination and palpation, and anamnesis data were taken into account. Attention was paid to appetite, thirst, urination, body temperature, pulse, respiratory rate, condition of mucous membranes and hair, degree of dehydration, body position in space, fatness, presence/absence of anxiety in animals.

During palpation of the abdominal cavity, special attention was paid to palpation of the kidneys to determine their size, shape, surface condition (smooth, bumpy), soreness. Palpation of the bladder was performed in order to determine the degree of its filling, tone, soreness.

The urine test was performed on the 1st, 15th, and 30th day of the study. The clinical analysis of urine included an assessment of the physical and chemical characteristics of urine, an analysis of mineral and organic substances contained in urine by microscopy of urine sediment

by an approximate method using test strips Combiscrin 11 ASUS. Urine sampling was carried out by natural urination and cystocentesis for bacteriological examination with the position of the animal on its side. Urine density was determined using a hydrometer (urometer) with a scale range from 1.001 to 1.050.

Hematological studies were performed on an automatic hematological analyzer URIT-3020 VetPlus at the time of treatment at the clinic, on the 15th and 30th days of curation. To do this, blood was taken from cats from vena saphena. At the same time, the total number of erythrocytes, leukocytes, hematocrit and hemoglobin levels were determined. The

biochemical composition of the blood serum was examined on an analyzer "Statfax1904" on the 1st, 15th, 30th day of the study. The levels of creatinine, urea, total protein, total calcium, inorganic phosphorus, potassium and sodium were determined. The study was carried out in blood serum by taking blood from cats from vena saphena in an amount of 3 ml, stood for an hour to form a clot, then centrifuged.

Ultrasonographic studies were performed at the time of treatment at the clinic. The fur on the abdomen was shaved in the usual way, the skin was treated with medical alcohol, after that gel for ultrasound diagnostics of the company "Geltek" was used. The animal for the study of the genitourinary system was fixed in a dorsal position or lying on its side. Ultrasound examinations were performed on the Mindray UMT-150 device.

The test animals were monitored every day, up to the moment of recovery. The onset of recovery was judged by the change in the general condition of the animal, the absence of clinical signs and the results of hematological studies.

3 Results

A retrospective analysis of the frequency of manifestations of pathologies of the urinary system organs from the total number of diseases of cats admitted to the veterinary clinic "White Fang" (Novocherkassk) from September 2022 to June 2023 showed that the most common diseases of the urinary system in cats are urolithiasis, cystitis (including idiopathic cystitis), nephritis, chronic renal failure (CRF) and acute renal failure (ARF). Polycystic kidney disease (PBP) and nephropathies are less frequently registered. These diseases accounted for 25.2% of the total number of sick cats. Of the diseases of the urinary tract in cats admitted to the veterinary clinic "White Fang" (Novocherkassk), urolithiasis and idiopathic cystitis were most often detected (Table 1).

Table 1. Percentage ratio of various pathologies of the urinary system of cats for 2022-2023 who were admitted to the veterinary clinic "White Fang" (Novocherkassk)

Name of the disease	Number of cases, %
Urolithiasis	30,2
Cystitis (including idiopathic)	19,6
Acute renal failure	14,4
Chronic renal failure	14,8
Nephropathies	2,9
Jade	18,1

According to the data of our retrospective analysis, diseases of the urinary organs were recorded in cats throughout the calendar year. Urolithiasis was detected in cats throughout the year, but the peak incidence occurred in autumn. The high incidence of cystitis occurred in the autumn and winter periods. Idiopathic cystitis is less dependent on seasonality. There is a slight increase in the disease in the autumn and spring periods. Acute and chronic renal failure was detected mainly in summer and autumn. Polycystic kidney disease was diagnosed mainly in winter, and nephritis and nephropathy in autumn.

An analysis of the prevalence of idiopathic cystitis in domestic cats, depending on gender and physiological condition, showed that cats are more likely to be susceptible to idiopathic cystitis, regardless of the activity of the functioning of the reproductive system (29.9% unsterilized cats and 31.3% sterilized, respectively). In cats, minor differences in the spread of the disease were recorded depending on the physiological state: in non-castrated cats - 20.1%, in castrated cats - 18.7% of the total number of sick animals. Idiopathic cystitis, as a rule, most often affects cats. Cats with this disease, in comparison with other animals, behave

quite nervously and react excessively violently to the environment. Animals with free access to walks may also be susceptible to this condition, especially with a large population density of cats in the surrounding area (Figure 1).

The age of sick animals varies in very large age ranges from 6 months to 12 years, but the greatest percentage of morbidity falls on the age from 1 year to 3 years (53.9%), from 3 to 7 years -26.1% and from 7 to 12 years - 16.2% of the total number of cases. Thus, idiopathic cystitis is less common in the age group under one year, and most often idiopathic cystitis affects cats in young and middle age (Fig.2).




Fig. 1. The incidence of idiopathic cystitis, depending on gender




Fig. 2. Incidence of idiopathic cystitis of cats of various age groups.

It was also found that animals living in apartments and not having access to the street, in comparison with those who are on a samovygula, are more likely to suffer from idiopathic cystitis – 61.21%, respectively, of the total number of cases.

anamnesis was collected in cats who were admitted to the reception with suspected idiopathic cystitis for diagnosis, a general clinical examination was performed, body temperature was measured, hematological and urine tests were performed, ultrasonographic studies, if necessary, X-ray studies. Since idiopathic cat cystitis is a diagnosis of exclusion, it is necessary to exclude other diseases from the group of ZNMP (diseases of the lower urinary tract). To confirm the diagnosis, in all cases, a bacteriological analysis of urine was necessarily performed.

During the collection of anamnesis, the number of manifestations of cystitis and the presence/absence of stress factors were taken into account. An analysis of the anamnesis data showed that all the sick animals were exposed to a certain influence of various stress factors. The most common triggers were conflicts with other cats, the appearance of children or new pets in the family, change of residence. In 60% of cases, two or more stress factors were detected simultaneously in the examined animals. Stress factors can provoke the manifestation of clinical signs of idiopathic cystitis by activating the efferent part of the sympathetic nervous system, stimulating the dorsal root ganglia. The latter react to such an effect by inducing the release in peripheral tissues of neuropeptides and mediators responsible for the development of an inflammatory reaction and the emergence of a feeling of pain.

Ultrasound examination revealed signs of inflammation of the bladder in all animals, without the presence of concretions, which was also confirmed by X-ray examination.

Table 2. Clinical monitoring of cats with idiopathic cystitis

A symptom of the disease	Frequency of occurrence, %
Pollakiuria	59,7%
Hematuria	45%
Ishuria	30,3%
Violation of appetite and thirst	18,3%
The presence of signs of neurotic behavior	34,7%
Localization during urination	39,2%
Stranguria	6,4%
Absence of symptoms	5,5%
Increased groin care	44,6%
Signs of aggression	13,7%

As can be seen from Table 2, the main symptoms treated by cat owners with a subsequent confirmed diagnosis of idiopathic cystitis are pollakiuria, hematuria, ischuria, the presence of signs of neurotic behavior, vocalization during urination, increased groin care.

The analysis of therapeutic efficacy showed that in animals in group I, the clinical manifestations of the disease under discussion were manifested longer and more intensively than in animals of group II. Thus, the absence of the main clinical manifestations of idiopathic cystitis was noted in animals of the II nd group on the sixth day. Signs of undesirable behavior of cats were not noted. It is noteworthy that the animals of the second experimental group showed no signs of aggression. In animals of the 1st experimental group, the disappearance of the main clinical signs of cystitis was noted on the 8th day, some animals with signs of neurotic disorders in the history of the disease retained signs of aggression even after normalization of indicators. Complete stabilization of physiological parameters and behavioral reactions in group 2 animals was noted by the 20th day of treatment, in group 1 by the 30th day, which indicates the effectiveness of including N-acetylglucosamine (Furinaid drug) in the therapeutic regimen.

Studies have shown in the first day significant deviations from the physiological norm in terms of protein content in urine, the presence of a flat, transitional epithelium, the presence of hematuria, pronounced leukocyturia.

Table 3. Changes in clinical urine analysis during correction of idiopathic cystitis in cats

Indicator	Day	FP	Animal groups	
			I	II
pH	1	5,0-7,0	7,00±0,50	6,50±0,50
	15		6,50±0,10	6,00±0,50

	30		6,50±0,10	6,30±0,15
Density, g/cm ³	1	1,020–	1,050± 0,005	1,050±0,005
	15	1,040	1,035± 0,005	1,030±0,002
	30		1,025± 0,002	1,030±0,005
Protein, g/l	1	0 – 0,1	0,60 ±0,10	0,60 ±0,10
	15		0,40 ±0,10	0,15 ±0,10
	30		0,10 ±0,10	0±0,10
Transitional epithelium, pcs/pp	1	0	12,00 ± 1,00	12,00 ± 1,00
	15		5,00 ± 0,05	3,00 ± 0,05
	30		1,00 ± 0,05	0 ± 0,05
Squamous epithelium, pcs/pp.	1	0-0,5	15,00±0,50	16,00±0,05
	15		10,00±0,50	5,00±0,05
	30		3,00±0,05	1,00±0,05
Renal epithelium, pcs/pp.	1	0	2,00 ± 0,50	2,00 ± 0,05
	15		1,00 ± 0,10	0 ± 0,05
	30		0 ± 0,05	0 ± 0,05
Erythrocytes, pcs /p.zr.	1	0-2	250,00± 0,50	250,00±0,50
	15		150,00± 0,05	0±0,10
	30		0±0,05	0±0,005
Leukocytes, pcs /p.zr.	1	0-2	25,00 ± 0,50	33,00 ± 0,50
	15		15,00 ± 0,10	9,00 ± 0,10
	30		3,00 ± 0,10	2,00 ± 0,05
Hyaline cylinders, pcs/p.zr.	1	0-1	1 ± 0,05	1 ± 0,05
	15		0 ± 0,05	0 ± 0,05
	30		0± 0,05	0± 0,05
Inorganic precipitation, pcs/pp	1	0	ots	ots
	15		ots	ots
	30		ots	ots
Bacteria	1	single	ots	ots
	15		ots	ots
	30		ots	ots

As can be seen from the table below, on the first day of observation, hyperproteinuria, or nonselective proteinuria (protein content in urine 0.60 ± 0.10 g/l) was diagnosed in patients of the experimental group. On the 30th day of treatment, the protein level in the urine in both groups fell within the reference interval.

Mild or moderate leukocyturia in cats does not always indicate an infectious disease of the urinary system. During the study of urine samples on day 1, pronounced leukocyturia was detected in all groups. By day 30, the content of leukocytes in the urine in group I only slightly exceeded the physiological norm (3.00 ± 0.10), in group II it did not go beyond the reference values and amounted to 2.00 ± 0.10 .

On the first day, a large number of cells of the flat (12.00 ± 1.00 and 15.00 ± 0.05) transitional epithelium were detected ($15,00 \pm 0,50$; $12,00 \pm 1,00$). On the 30th day, the amount of epithelium in urine samples did not exceed physiological parameters in both groups. When analyzing the content of the renal epithelium, it was revealed that on the 15th day in group I it was present in a small amount (1.00 ± 0.05), while in group II there was no renal epithelium. By the 30th day of treatment, renal epithelium was not detected in any of the groups. Thus, it can be concluded that during treatment, uneven improvements in urine samples were observed in patients of all the study groups. In the second group, the improvements were

more significant and were traced already from the 15th day, and by the 30th day, the main indicators returned to the physiological norm.

Biochemical studies of blood serum in cats were carried out in order to control the work of the filtration apparatus of the kidneys, as well as to detect secondary changes as a result of acute urinary retention (Table 4). the main attention was paid to the work of the kidneys.

Table 4. Biochemical blood parameters in idiopathic cystitis of cats.

Indicator	Day	FP	Animal groups	
			I	II
urea mmol/l	1	7,1- 15	13,2±0,4	15,3 ±0,5
	15		12,4±0,6	13,4±0,3
	30		12,3±0,4	13,0±0,3
creatinine mmol/l	1	44-160	168,8± 5,4	172,8±5,1
	15		163,1± 3,9	159,1±4,9
	30		165,3± 3,5	145,2±3,7
phosphorus mmol/l	1	1.3-2.1	1,9±0,15	2,3±0,2
	15		1,5±0,10	1,9±0,05
	30		1,5±0,10	1,6±0,05
Total calcium, mmol/l	1	2.1-2.5	2,6±0,20	2,2±0,20
	15		2,2±0,20	2,1±0,10
	30		2,1±0,10	2,4±0,05
sodium mmol/l	1	145-150	138,2± 4,40	135,2±5,10
	15		141,5± 3,90	144,1±3,80
	30		148,2± 3,30	148,4±3,90
potassium mmol/l	1	4.2-5.5	3,7±0,30	3,9±0,30
	15		4,1±0,20	4,5±0,10
	30		4,5±0,10	4,7±0,10
Total protein g/l	1	58-76	78,2± 2,30	85,5±2,20
	15		75,6± 1,70	71,4±2,00
	30		68,5± 1,50	65,5±1,40

An increase in the level of nitrogenous metabolites in the blood can be interpreted as a decrease in the detoxification function of the kidneys as a result of the development of OBP, the result of dehydration / hypovolemia due to impaired appetite and thirst. At the time of admission, a sufficiently high content of total protein in the blood serum of animals of all groups was observed, the values were 78.2 ± 2.30 in the I-th, 85.5 ± 2.20 in the II-th groups, respectively. Hypokalemia and hyponatremia were observed in the study groups on the first day. The potassium level was 3.7 ± 0.30 mmol/l in the I-th, 3.9 ± 0.30 in the II-th. The sodium level is 138.2 ± 4.40 in the I-th, 135.2 ± 5.10 mmol/l in the II-th. An increase in intracellular sodium content and a decrease in intracellular potassium content is observed when the transmembrane fluid flow is disrupted.

The results of the analysis of venous blood samples on the 15th day of treatment showed that in the first group the creatinine level decreased, but still remained above the threshold values (163.1 ± 3.9), in the second group this indicator returned to normal and was 159.1 ± 4.9 mmol/L. Thus, it was found that in the second group, a more pronounced decrease in blood creatinine concentration was detected – by 7.96% versus 3.4% in group I. In addition, the level of total protein and calcium decreased more markedly in the second group, which also indicates the stabilization of the body's metabolic processes and normalization of kidney

function. The level of total calcium returned to normal in group I (2.2 ± 0.20). The phenomenon of hypokalemia and hyponatremia became less pronounced by the 15th day, but it was still traced. At the same time, potassium values were slightly reduced in group I (4.1 ± 0.20) and returned to normal in group II (4.5 ± 0.10).

It should be noted that by the 30th day of observation, all the studied values in both experimental groups were within the reference values, however, a more positive dynamics of normalization of indicators and a more stable result was noted in animals of the II experimental group. This indicates that when correcting idiopathic cystitis, the use of drugs that contribute to the restoration of the protective layer of the urinary tract (the drug Furinaid) is of great importance. In addition, adequate anti-stress therapy is not the last place in the treatment.

Table 5. Changes in clinical blood analysis in cats with idiopathic cystitis

Indicator	Day	FP	Animal groups	
			I	II
Hemoglobin (Hb), g/l	1	90 – 150	$120,3 \pm 4,1$	$125,3 \pm 4,8$
	15		$129,3 \pm 3,9$	$130,2 \pm 3,9$
	30		$135,7 \pm 5,3$	$142,3 \pm 5,1$
Hematocrit (HCT), %	1	30 – 47	$52,2 \pm 1,5$	$50,3 \pm 3,5$
	15		$45,1 \pm 1,9$	$45,2 \pm 3,2$
	30		$41,0 \pm 2,2$	$42,5 \pm 2,6$
Erythrocytes (RBC), $10^{12}/l$	1	5,6–10,7	$6,5 \pm 2,5$	$5,5 \pm 1,9$
	15		$7,2 \pm 3,1$	$7,3 \pm 2,5$
	30		$8,4 \pm 2,9$	$8,9 \pm 3,2$
White blood cells (WBC), $10^9/l$	1	5,5–18,5	$20,4 \pm 3,6$	$23,0 \pm 3,5$
	15		$18,3 \pm 2,7$	$16,2 \pm 3,2$
	30		$15,0 \pm 2,7$	$14,5 \pm 2,7$
Erythrocyte sedimentation rate (ESR), mm/h	1	1-13	$32,3 \pm 3,1$	$40,0 \pm 2,8$
	15		$23,1 \pm 2,8$	$24,1 \pm 2,1$
	30		$12,0 \pm 1,5$	$11,2 \pm 1,2$

In the course of our study, from the first day to the last, we did not observe a violation of the production of red blood cells. Further, during the entire study period, the level of red blood cells in all groups was within the physiological boundaries. The level of hemoglobin throughout the study was also within the reference values.

When analyzing the number of leukocytes, it was found that on the 1st day, leukocytosis was registered in cats of all groups, which indicated that severe inflammatory reaction. By day 15, the number of leukocytes returned to normal in the study groups and amounted to 18.3 ± 2.7 , 16.2 ± 3.2 , $\times 10^9/l$ in groups I and II, respectively. By day 30, more significant improvements were observed in the study groups. The presence of an inflammatory process in the body is also indicated by an increase in the erythrocyte sedimentation rate (ESR). Thus, in all groups up to 30 days, a pronounced increase in ESR was recorded, which returned to normal in the studied groups.

4 Conclusions

As a result of the conducted studies, it was found that a promising direction in the treatment of inflammatory diseases of the bladder is the use of drugs containing glucosaminoglycans. The use of the drug Furinaid (N-acetyl glucosamine) in the therapeutic regimen for the treatment of idiopathic cystitis had a positive effect on the stabilization of physiological parameters and contributed to the protection of the bladder mucosa from aggressive urine components, preventing the development of

the inflammatory process. We recommend using the drug "Furinaid" as part of complex therapy in the treatment of HCC according to the following scheme: during the first 14 days, 2.5 ml of the drug orally per animal 1 time per day, the next 14 days - 1.25 ml of the drug 1 time per day.

References

1. A.O. Michel, T.A. Donovan, B. Roediger, Q. Lee, C.J. Jolly, S. Monette, *Veterinary Pathology*, **59(1)**, 120-126 (2022) doi: 10.1177/03009858211045439
2. S.Chaiyasak, C. Piewbang, J. Yostawonkul, S. Boonrungsiman, T. Kasantikul, A. Rungsipipat, S. Techangamsuwan, *Veterinary Pathology*, **59(1)**, 127-131 (2022) doi: 10.1177/03009858211045441
3. M. Katayama, Y. Uemura, *Veterinary Sciences*, **8(12)**, 328 (2021) doi: 10.3390/vetsci8120328
4. M. Davenport, A. Muntean, N. Hadzic, *Journal of Clinical Medicine*, **10(23)**, 5675 (2021) doi: 10.3390/jcm10235675
5. S.R. Compton, *Comparative medicine*, **71(5)**, 333-341 (2021) doi: 10.30802/AALAS-CM-21-000007
6. D.A. Meekins, N.N. Gaudreault, J.A. Richt, *Viruses*, **13(10)**, 1993 (2021) doi: 10.3390/v13101993
7. H. Nan, D.J. Yoo, S.H. Kang, *Journal of Separation Science*, **36(2)**, 350-355 (2013) doi: 10.1002/jssc.201200721
8. D.S. Chatzopoulos, L.V. Athanasiou, V.Spyrou, G.C. Fthenakis, C.Billinis, *Journal of the Hellenic Veterinary Medical Society*, **64(2)**, 145-160 (2013) doi: 10.12681/jhvms.15489
9. D. Jas, V. Frances-Duvert, D. Vernes, P.-M. Guigal, H. Poulet, *Veterinary Microbiology*, **177(1-2)**, 123-131 (2015) doi: 10.1016/j.vetmic.2015.03.009
10. C. Pressanti, M.-C. Cadiergues, *Pratique Medicale et Chirurgicale de l'Animal de Compagnie*, **48(4)**, 105-112 (2013) doi: 10.1016/j.anicom.2013.10.001
11. H.-Y. Jeoung, S.-I. Lim, B.-H. Shin, J.-A. Lim, J.-Y. Song, D.-S. Song, B.-K. Kang, H.-J. Moon, D.-J. An, *Veterinary Microbiology*, **165(3-4)**, 281-286 (2013) doi: 10.1016/j.vetmic.2013.03.021
12. D.J. Wiener, M.M. Welle, F.C. Origgi, *Veterinary Dermatology*, **24(5)**, 543-e130 (2013) doi: 10.1111/vde.12063
13. S.J. McDonnel, M.L. Liepnieks, B.G. Murphy, *Antiviral Research*, **108(1)**, 74-78 (2014) DOI: 10.1016/j.antiviral.2014.05.014
14. A.L. Litster, *Veterinary Journal*, **201(2)**, 184-188 (2014) doi:10.1016/j.tvjl.2014.02.030
15. P. McDonagh, P.A. Sheehy, A. Fawcett, J.M. Norris, *Veterinary Microbiology*, **176(3-4)**, 370-377 (2015) DOI: 10.1016/j.vetmic.2015.02.007
16. O. Krotova, T. Alekseeva, K. Savenkov, S. Sklyarov, M. Savenkova, XV International Scientific Conference "Interagromash 2022". Collection of materials of the 15th International Scientific Conference. Global Precision Ag Innovation 2022. Rostov-on-Don, 897-905 (2023) doi:10.1007/978-3-031-21219-2_101
17. O. Krotova, N. Bashkatova, K.Savenkov, K.Sklyarov, A. Manjikova, IOP Conference Series: Earth and Environmental Science. Cep. "Fundamental and Applied Scientific Research in the Development of Agriculture in the Far East, AFE 2021 - Papers", 022016 (2021) doi: 10.1088/1755-1315/937/2/022016