

Frequency And Determinants Of Chronic Lower Back Pain Among Patients Presenting To Primary Care Clinics In Pakistan: A Cross-Sectional Analysis

Rabbiya Amir¹, Fatima Jehangir², Misbah Muhammad³, Nida J. Ahsan⁴, Tariq Adnan⁵

Abstract

Objective: To ascertain the determinants of chronic lower back pain among patients presenting in primary care clinics.

Methods: 300 participants from 18 to 75 years of age were enrolled from Primary Care Clinics, Primary Health Center, Sikanderabad and Family Medicine Health Center, Clifton, for a Cross-sectional study that was done from December 1, 2020, to June 30, 2021. Consultant family physicians took informed consent and then asked relevant history questions and performed a relevant physical examination, such as a straight leg raise on the patients.

Results: The frequency of chronic lower back pain came out to be 16.7% (50). The median age was 31.99+15.7 years. For those who had chronic back pain, the majority were regularly taking Cholecalciferol supplements (20.8% p-value 0.05). The effect of depression increases the chances (0.688) of chronic lower back pain. (p-value 0.006)

Conclusion: In our study, we concluded that lower back pain, especially chronic in duration, is very rampant in the community with male predominance. The majority reported a dull type of pain. Depressed people are more vulnerable to developing chronic back pain.

Keywords: LBP (lower back pain), magnitude, restricted mobility, S.I. joint (Sacroiliac joint), community.

^{1,3,4} Specialist Family Physician, Ziauddin University, Sukkur; ² Associate Professor, Ziauddin Medical College; ⁵ Associate Professor Community Health Sciences.

Correspondence: Dr. Fatima Jehangir, Associate Professor, Associate Professor, Ziauddin Medical College, Sukkur. **Email:** fatimajehangir@zu.edu.pk

Cite this Article: Amir R, Jehangir F, Muhammad M, Ahsan NJ, Adnan T. Frequency And Determinants Of Chronic Lower Back Pain Among Patients Presenting To Primary Care Clinics In Pakistan: A Cross-Sectional Analysis. JRMC. 2023 Dec. 30;27(4). <https://doi.org/10.37939/jrmc.v27i4.2493>.

Received January 19, 2023; accepted October 09, 2023; published online December 30, 2023

1. Introduction

Low back pain is a pain, stiffness, or muscle tension located between the costal margin and the inferior gluteal folds. It can be present with or without sciatica.^[1] It is classified as acute back pain when it lasts for less than six weeks, subacute when it occurs for six weeks to three months, and chronic when it lasts for more than three months.^[2]

Low back pain is a recurring cause of a visit to a primary healthcare doctor in developed countries. It is a common musculoskeletal disorder worldwide and one of the leading causes of years lived with disability. LBP occurs in about 60–80% of people at some point in their lives and can begin in childhood.^{[3][4][5][6][7]} It has been determined that up to 9% of the world's population has LBP at any given time.^[8] Some studies have reported that around 60–90% of adults will experience LBP at some point in their lifetime.^{[9][10][11]} Another study showed that it was the most common type of pain reported by patients, with 25% of U.S. adults reporting LBP in the prior three months.^[12]

Low back pain has multiple risk factors, including physical and psychosocial risk factors. Incorrect sitting posture and frequently carrying and lifting

heavy loads cause pain in the lower back.^[13] Ageing is also a well-known risk factor for LBP, as degenerative changes in the disc and the spine are among the significant causes of LBP.^[14] Also, prolonged sitting time is associated with LBP intensity.^[15]

Lifestyle, short sleep hours, and lack of physical exercise also increase the risk of LBP. Unfortunately, the level of evidence for most psychosocial factors is limited, and the prevalence and burden of LBP continue to grow with time.

We decided to conduct our study on chronic LBP because there is a shortage of data on chronic LBP from Pakistan. There have been prior studies on the prevalence of low back pain in nurses and bankers of Pakistan. We want to conduct a study in a primary healthcare setting to determine the prevalence and determinants of lower back pain in Pakistan. We want to find out the relationship between backache with age and the BMI of the patients. We would also like to know the association of low back pain with lifestyle factors and what mental impact it has. We also want to find out the protective factors such as sun exposure and vitamin D supplements. We believe this study will help primary health practitioners prevent recurrent and chronic LBP.

2. Materials & Methods

A cross-sectional study was conducted in the primary health care centre of Ziauddin University from December 1, 2020, to June 30, 2021. Three hundred patients coming to the primary health care centre, aged 18 years to 75 years, were enrolled in the study. Exclusion criteria included pregnant and lactating women, patients with malignancy, congenital structural deformities, subjects with kyphosis and scoliosis and back pain related to menstrual periods. After taking the informed and voluntary written consent to participate in the study, consultant family physicians and authors of the study took the history of the subjects and performed a relevant physical examination.

History questions included questions related to lower back pain, such as onset, course, duration, quality, exacerbating, relieving factors, and pain severity. Patients were requested to rate their pain on a scale of 1 to 10, where 1 was the mildest pain, and 10 meant the worst possible pain. Patients were also asked about pain triggers, history, family history, and socioeconomic questions. The patients were also screened for depression using Patient Health Questionnaire-2 (PHQ-2) and Patient Health Questionnaire-9 (PHQ-9). The Patient Health Questionnaire (PHQ) is a self-directing form of the PRIME-MD diagnostic instrument for common mental disorders. The PHQ-2 is a 2 question-screening tool used to screen for depression in individuals. If the score is three or greater, there is a higher likelihood of depression.^[16] The patients who screened positive were further evaluated using the PHQ-9. The PHQ-9 is the depression module, which uses a scale to score each of the 9 DSM criteria as "0" which means not at all to "3" which stands for nearly every day. Major Depression is then diagnosed if five or more of the nine depressive symptom criteria have been present, with depressed mood or anhedonia being one symptom and the symptoms are present more than half the days in the past fourteen days.^[17] The 7-item anxiety scale (GAD-7) was used to screen for GAD.^[18] The patient's BMI was calculated with the help of their height and weight measured at the clinic. A physical exam was then performed, including inspection, palpation, range of motion of the back, and special tests such as straight leg raise and the Flexion, Abduction and External Rotation (FABER) test. The examination was performed while the patient was standing, lying, and sitting. The patient's gait, heel, and toe walking were also observed.

After asking the questions and performing the examination, the information was documented for each participant by the consultant family physicians.

In a study conducted in 2018 in Lahore, Pakistan, 167 subjects were recruited to see the prevalence of lower back pain. We almost doubled the sample size in our study.^[19] Data entry was done using SPSS version 20. Shapiro Wilk test for normality of numerical variables seen. For non-parametric variables, the Median and interquartile range were analyzed by Mann-Whitney's test. The BMI was gauged by dividing the weight in kilograms by the square of height in meters. The frequency and percentage were computed for categorical variables such as type, location, nature of pain, pain score, pain duration, aggravating and relieving factors, and presence of depression. The association of chronic back pain with variables was analyzed using chi-square, and binary logistic regression was also computed with a 95% Confidence interval. Multivariate logistic regression was analyzed for all significant variables. A P-value of 0.05 was considered significant.

3. Results

The frequency of chronic lower back pain came out to be 16.7% (50). The mean age was 31.99 ± 15.7 years. The mean BMI was $23.23 \pm 4.9 \text{ kg/m}^2$. The mean Pain Score as per the Visual analogue Score (VAS) came out to be 2.8 ± 2.9 in our study. Table 1 demonstrates the demographics and association of chronic back pain, various risk factors and sequelae. Table 2 demonstrates Odd's ratio of various factors with chronic back pain with a 95% Confidence Interval. Graph 1 demonstrates the proportion of subjects having varying natures of pain. The majority of subjects, 53.3% (160), had lumbar pain, followed by 28% (84) having sacral joint pain. Out of that, 26.2% had right sacroiliitis, while 6.7% had left sacroiliitis. A statistically significant (p-value 0.006) increase effect of depression (0.688) was seen with chronic back pain. When lifestyle factors were evaluated, most participants (20%) did prolonged sitting, followed by 6.7% who did a lot of bending. 6% lifted heavyweight, and 4.7% had to stand for long hours.

When we looked at the association of risk factors with chronic back pain, we analyzed that those who had chronic back pain majority were regularly taking Cholecalciferol supplements (20.8% p-value 0.05).

Table-1 Comparison of Risk Factors among Healthy and Chronic Lower Back Pain Afflicted Patients

Numerical variables			With Chronic Back Pain Median (IQR)	Without Chronic Back Pain Median (IQR)	p-value
Age in years			23(12)	24(19)	0.336
BMI kg/m ²			22.23(3.92)	22.65(5.97)	0.512
Pain Score			0.00(8)	3(6)	0.00
Categorical Variables		N(%)	With ch. back pain n(%)	Without ch. back pain n(%)	p-value
Gender	Males	134(44.7)	28(20.9)	106(79.1)	0.00
	Females	166(55.3)	22(13.3)	144(86.7)	
Marital status	single	184(61.3)	35(19.1)	148(80.9)	0.541
	Married	116(38.7)	15(12.9)	101(87.1)	
Education	illiterate	26 (8.7)	1(3.8)	25(96.2)	0.2
	Primary	20(6.7)	4(20)	16(80)	
	Secondary	16(5.3)	1(6.2)	15(93.8)	
	College	24(8)	2(8.3)	22(91.7)	
	university	214(71.3)	42(19.6)	172(80.4)	
Sleep on a matted bed		246(82)	44(17.9)	202(82.1)	0.021
	Floor	22(7.3)	5(22.7)	17(77.3)	
	Charpai	32(10.7)	1(3.1)	31(96.9)	
back trauma	present	58(19.3)	6(10.3)	52(89.7)	0.000
	absent	200(66.7)	43(21.5)	157(78.5)	
Screen time(Hrs)	<2	170(56.7)	28(16.5)	142(83.5)	0.310
	2.1-6	96(32)	18(18.8)	78(81.2)	
	6	32(10.7)	4(12.5)	28(87.5)	
	>6				
Sun Exposure m 1hr	<15m	90(30)	13(14.4)	77(85.6)	0.844
	15-30	90(30)	18(20)	72(80)	
	30m-1hr	54(18)	9(16.7)	45(83.3)	
	1hr	16(5.3)	2(12.5)	14(87.5)	
	1hr-2hr	32(10.7)	4(12.5)	28(87.5)	

	2hr-3hr >3hrs	18(6)	4(22.2)	14(77.8)	
Vit D3	Don't take	202(67.3)	5(20.8)	19(79.2)	0.05
	Regular intake	24(8)	5(20.8)	19(79.2)	
	Irregular intake	72(24)	8(11.1)	64(88.9)	
MDD	present	58(19.3)	5(8.6)	53(91.4)	0.006
	absent	174(58)	27(15.5)	147(84.5)	
Anxiety	present	184(61.3)	34(18.5)	150(81.5)	0.074
	absent	116(38.7)	16(13.8)	100(86.2)	

Table-2 Association of variables with chronic back pain using binary regression.

Variables		Crude OR	95% CI		p-value	Adjusted OR	95% CI		p-value
			Lower	Upper			Lower	Upper	
VIT D supplements	Regular intake	1	-	-	-	-	-	-	-
	Irregular intake	0.47	0.139	1.624	0.235	0.548	0.151	1.986	0.360
	Don't take	0.84	0.295	2.400	0.747	0.982	0.333	2.895	0.973
Gender	Male	1.729	0.937	3.189	0.08	2.22	1.14	4.323	0.19
	Female	1	-	-	-	-	-	-	-
Age		0.985	0.964	1.007	0.183	0.978	0.954	1.002	0.073
BMI		0.97	0.904	1.04	0.396	-	-	-	-
Marital status	single	1	-	-	-	-	-	-	-
	married	0.158	0.821	3.046	0.17	1.19	0.45	3.15	0.724
Education	illiterate	0.164	0.22	1.244	0.08	1.253	0.104	15.08	0.859
	primary	1.024	0.325	3.222	0.968	7.493	1.074	52.27	0.042
	secondary	0.273	0.035	2.125	0.215	0.614	0.065	5.77	0.670
	College	0.372	0.084	1.646	0.193	0.956	0.175	5.228	0.959
	university	1	-	-	-	-	-	-	-
Major Depressive Disorder	Yes	0.514	0.188	1.403	0.194	0.678	0.237	1.940	0.469
	No	1	-	-	-	-	-	-	-

Sun exposure	<15 mins	1	-	-	-	-	-	-	-
	15-30 mins	1.481	0.677	3.238	0.325	-	-	-	-
	30mins-1hr	1.185	0.469	2.991	0.720	-	-	-	-
	1-2hrs	0.846	0.172	4.166	0.837	-	-	-	-
	2-3hrs	0.846	0.255	2.813	0.785	-	-	-	-
	>3hrs	1.692	0.481	5.948	0.412	-	-	-	-
Anxiety	Absent	1				-	-	-	-
	Present	1.417	0.743	2.702	0.290		--	-	-
Back Trauma	Present	0.421	0.170	1.047	0.063	0.504	0.195	1.308	0.159
	Absent	1	-	-	-	-	-	-	-
Screen time	<2hrs	1	-	-	-	-	-	-	-
	2-6hrs	1.466	0.471	4.559	0.509	-	-	-	-
	>6hrs	1.871	0.579	6.045	0.295	-	-	-	-
Sleep on	Mattressed bed	1	-	-	-	-	-	-	-
	Floor	7.273	0.963	54.9	0.054	5.27	0.603	46.03	0.133
	Charpai	9.748	1.042	91.229	0.046	6.34	0.590	68.3	0.127

5. Discussion

This prospective cross-sectional study was carried out to determine the magnitude and associations of lower back pain in our community in Pakistan. Three hundred patients from primary health care centres between 18 to 75 years of age were enrolled in the study.

In this current study, the frequency of chronic lower back pain was 16.7%(50) which was quite high. Out of 300 participants, 244 participants had lower back pain. Out of these 244 participants, 50 participants had chronic lower back pain.


In the present study, the mean age for low back pain was found to be 31.99+15.7. In contrast, a study conducted between November 2007 and December 2008 showed that patients with lower backache have a mean age of 52.8 years (standard deviation [S.D.] = 15.0).^[20] This difference might be because of our small

sample size and the candidates who took part in our study were comparatively young.

In this research, the mean BMI of the participants was found to be 23.23+4.9 kg/m². The mean BMI fell in the normal weight range for the participants in our study. A previous study conducted in Norway showed that obese women and men had an approximately 20% increased risk of chronic low back pain.^[21] Another study conducted in Norway showed that an increased prevalence of low back pain was significantly associated with high body mass index in both men and women.^[22] In our current study, the association of BMI with low back pain was not significant.

One study showed that the prevalence of chronic lower back pain generally declined with greater levels of education and increasing income.^[12]

Whereas our study showed that the prevalence of lower back pain increased with the level of education. 19.6 % of the individuals with chronic lower back pain were either in university or had graduated from university. This could be related to sitting for prolonged periods while studying or working in the office.



Graph-1 Proportion of subjects having varying nature of Pain

Our study shows that lower back pain, especially chronic in duration, is rampant in our community with male predominance. Another study showed that the median overall prevalence of low back pain was higher among females than males across all age groups, contradicting our findings.^[23]

The current study showed that patients with chronic lower back pain were not depressed (15.5% p-value 0.00). A Jerusalem Longitudinal Study consisting of 277 subjects aged 70 years at baseline and 77 years at follow-up showed that chronic back pain was associated with the female gender, and depression was associated with age 70 years in these patients.^[24] A prospective cross-sectional study was conducted in 2010 at the Department of Neurosurgery at Liaquat National Hospital in Karachi, Pakistan, that showed an abnormal level of anxiety and depression. In chronic low back pain patients, 77 (55%) participants had anxiety, and 68 (48.57%) patients had depression.^[25] Another study carried out in the U.S. showed that individuals with chronic low backache had an association with depression.^[26] Our findings were different from the rest of the studies, which could be due to the low sample size and reluctance towards talking about mood disorders in our society.

5. Conclusion

Our study concluded that lower back pain, especially chronic in duration, is very rampant in our community with male predominance. The majority report a dull type of pain, and it does not lead to mood disorders such as anxiety; however, depression increases the likelihood of developing chronic back pain.

CONFLICTS OF INTEREST- None

Financial support: None to report.

Potential competing interests: None to report

Contributions:

R.A, F.J - Conception of study

M.M, N.A - Experimentation/Study Conduction

F.J, M.M, N.A - Analysis/Interpretation/Discussion

F.J, - Manuscript Writing

R.A, F.J, M.M, N.A, T.A - Critical Review

R.A, F.J, M.M, N.A, T.A - Facilitation and Material analysis

References

1. Kebede T, Bedane A, Admassie D, Zenebe G. Patterns of lumbar myelographic findings in patients with LBP a 5 years retrospective study at Yehuleshet Higher Clinic, Addis Ababa, Ethiopia. *Ethiop Med J.* 2010 Jul;48(3):229-36. PMID: 21073084.
2. Goertz M, Thorson D, Bonsell J, Bonte B, Campbell R, Haake B, Johnson K, Kramer C, Mueller B, Peterson S, Setterlund L, Timming R. Institute for Clinical Systems Improvement. *Adult Acute and Subacute Low Back Pain.* Updated November 2012.
3. Manek NJ, MacGregor AJ. Epidemiology of back disorders: prevalence, risk factors, and prognosis. *Curr Opin Rheumatol.* 2005 Mar;17(2):134-40. doi: 10.1097/01.bor.0000154215.08986.06. PMID: 15711224.
4. Martin BI, Deyo RA, Mirza SK, et al. Expenditures and Health Status Among Adults With Back and Neck Problems. *JAMA.* 2008;299(6):656-664. doi:10.1001/jama.299.6.656
5. Nagi SZ, Riley LE, Newby LG. A social epidemiology of back pain in a general population. *J Chronic Dis* 1973;26:769-79.
6. Rejsbord LS, Greenland S. Factors associated with self-reported back-pain prevalence: a population-based study. *Journal of Chronic Diseases.* 1985 ;38(8):691-702. DOI: 10.1016/0021-9681(85)90023-2. PMID: 3160720.
7. Burton AK, Clarke RD, McClune TD, Tillotson KM. The natural history of low back pain in adolescents. *Spine* 1996;21:2323-8.
8. Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, Williams G, Smith E, Vos T, Barendregt J, Murray C, Burstein R, Buchbinder R. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis.* 2014 Jun;73(6):968-74. doi: 10.1136/annrheumdis-2013-204428. Epub 2014 March 24. PMID: 24665116.

9. Cassidy JD, Carroll LJ, Côté P. The Saskatchewan health and back pain survey. The prevalence of low back pain and related disability in Saskatchewan adults. *Spine (Phila Pa 1976)*. 1998 Sep 1;23(17):1860-6; discussion 1867. doi: 10.1097/00007632-199809010-00012. PMID: 9762743.
10. Walker BF, Muller R, Grant WD. Low back pain in Australian adults: the economic burden. *Asia Pac J Public Health*. 2003;15(2):79-87. doi: 10.1177/101053950301500202. PMID: 15038680.
11. Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*. 1999 August 14;354(9178):581-5. doi: 10.1016/S0140-6736(99)01312-4. PMID: 10470716.
12. Deyo RA, Mirza SK, Martin BI. Back pain prevalence and visit rates: estimates from U.S. national surveys, 2002. *Spine (Phila Pa 1976)*. 2006 Nov 1;31(23):2724-7. doi: 10.1097/01.brs.0000244618.06877.cd. PMID: 17077742.
13. Claus M, Kimbel R, Spahn D, et al. prevalence and influencing factors of chronic back pain among staff at special schools with multiple and severely handicapped children in Germany: results of a cross-sectional study. *BMC Musculoskeletal Disorders*. 2014 Feb;15:55. DOI: 10.1186/1471-2474-15-55.
14. Luoma K, Riihimäki H, Luukkonen R, Raininko R, Viikari-Juntura E, Lamminen A. Low back pain in relation to lumbar disc degeneration. *Spine (Phila Pa 1976)*. 2000 Feb 15;25(4):487-92. doi: 10.1097/00007632-200002150-00016. PMID: 10707396.
15. Gupta N, Christiansen CS, Hallman DM, Korshøj M, Carneiro IG, Holtermann A. Is Objectively Measured Sitting Time Associated with Low Back Pain? A Cross-Sectional Investigation in the NOMAD study. *PLoS ONE*. 2015; 10(3): e0121159. <https://doi.org/10.1371/journal.pone.0121159>
16. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care*. 2003 Nov;41(11):1284-92. doi: 10.1097/01.MLR.0000093487.78664.3C. PMID: 14583691.
17. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001 Sep;16(9):606-13. doi: 10.1046/j.1525-1497.2001.016009606.x. PMID: 11556941; PMCID: PMC1495268.
18. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006 May 22;166(10):1092-7. doi: 10.1001/archinte.166.10.1092. PMID: 16717171.
19. Tauqeer, S., F. Amjad, A. Ahmad, and S. A. Gillani. "PREVALENCE OF LOW BACK PAIN AMONG BANKERS OF LAHORE, PAKISTAN". *KHYBER MEDICAL UNIVERSITY JOURNAL*, vol. 10, no. 2, June 2018, pp. 101-4, doi:10.35845/kmu.j.2018.17948.
20. Michael JD, Jessica MK, Thomas S. What Is the Source of Chronic Low Back Pain and Does Age Play a Role? *Pain Medicine*. 2011 Feb;12(2):224-233. <https://doi.org/10.1111/j.1526-4637.2010.01045.x>
21. Nilsen TI, Holtermann A, Mork PJ. Physical exercise, body mass index, and risk of chronic pain in the low back and neck/shoulders: longitudinal data from the Nord-Trøndelag Health Study. *Am J Epidemiol*. 2011 August 1;174(3):267-73. doi: 10.1093/aje/kwr087. Epub 2011 June 1. PMID: 21633119.
22. Heuch I, Hagen K, Heuch I, Nygaard Ø, Zwart JA. The impact of body mass index on the prevalence of low back pain: the HUNT study. *Spine (Phila Pa 1976)*. 2010 April 1;35(7):764-8. doi: 10.1097/BRS.0b013e3181ba1531. PMID: 20228714.
23. Hoy, D., Bain, C., Williams, G., March, L., Brooks, P., Blyth, F., et al. A systematic review of the global prevalence of low back pain. *Arthritis & Rheumatism*. 2012; 64: 2028-2037. <https://doi.org/10.1002/art.34347>
24. Jacobs, Jeremy M. MB, BS; Hammerman-Rozenberg, Robert MD; Cohen, Aaron MD; Stessman, Jochanan MD Chronic Back Pain Among the Elderly: Prevalence, Associations, and Predictors, *Spine: April 1, 2006 - Volume 31 - Issue 7 - p E203-E207* doi: 10.1097/01.brs.0000206367.57918.3c
25. Sagheer MA, Khan MF, Sharif S. Association between chronic low back pain, anxiety and depression in patients at a tertiary care centre. *J Pak Med Assoc*. 2013 Jun;63(6):688-90. PMID: 23901665
26. Shmagel A, Foley R, Ibrahim H. Epidemiology of Chronic Low Back Pain in U.S. Adults: Data From the 2009-2010 National Health and Nutrition Examination Survey. *Arthritis Care Res (Hoboken)*. 2016 Nov;68(11):1688-1694. doi: 10.1002/acr.22890. PMID: 26991822; PMCID: PMC5027174.