

氏名	熊井 健
所属	人間健康科学研究科 人間健康科学専攻
学位の種類	博士（理学療法学）
学位記番号	健博 第253号
学位授与の日付	令和5年3月25日
課程・論文の別	学位規則第4条第1項該当
学位論文題名	Brain and muscle activation patterns during postural control affect static postural control (姿勢制御時の脳と筋の活動パターンは静的姿勢制御能力に影響する)
論文審査委員	主査 教授 池田 由美 委員 准教授 金子 文成 委員 准教授 儀間 裕貴

【論文の内容の要旨】

Background: Previous studies have reported existence of coordinated brain and muscle activity patterns that affect postural control. However, what differences in these activity patterns affect postural control are still unclear. The purpose of this study was to clarify brain and muscle activity pattern affecting postural control.

Research question: Does the difference in brain and muscle activity patterns during postural control affect postural control ability?

Method: Nineteen healthy men (mean age: 24.8 ± 4.1 years, height: 171.8 ± 5.5 cm, and weight: 63.5 ± 12.5 kg) performed a postural control task on a balance board, and their brain and muscle activities and body sway during the task were measured using functional near-infrared spectroscopy, surface electromyography, and three-dimensional accelerometry. Hierarchical cluster analysis was conducted to extract subgroups based on brain and muscle activities and postural control, and correlation analysis was performed to investigate the relationship between brain activity, muscle activity, and postural control.

Results: Two subgroups were found. Subgroup 1 ($n = 9$) showed higher brain activity in the supplementary motor area ($p = 0.04$), primary motor cortex ($p = 0.04$) and stable postural control in the mediolateral ($p < 0.01$) planes, and subgroup 2 ($n = 10$) showed higher muscle activity in the tibialis anterior ($p < 0.01$), a higher shank muscles co-contraction ($p = 0.02$) and unstable postural control. Furthermore, the

supplementary motor area activity is negatively correlated with body sway of mediolateral plane ($r = -0.51$, $p = 0.02$), and tibialis anterior activity is positively correlated with body sway on the mediolateral plane ($r = 0.62$, $p = 0.004$).

Significance: Higher brain activity in motor-related areas, lower activity in the lower limb muscles and lower co-contraction of shank muscles were observed in stable postural control. These characteristics of stable postural control shown in this study will facilitate the planning of new rehabilitation methods for improving postural control ability.