

Document downloaded from the institutional repository of the University of Alcalá: <https://ebuah.uah.es/dspace/>

This is a postprint version of the following published document:

De la Cueva-Alique, I. et al. (2019) 'Study of the anticancer properties of optically active titanocene oximate compounds', *Journal of organometallic chemistry*, 881, pp. 150–158.

Available at <https://doi.org/10.1016/j.jorgchem.2018.12.011>

© 2018 Elsevier

(Article begins on next page)



This work is licensed under a

Creative Commons Attribution-NonCommercial-NoDerivatives
4.0 International License.

1 Study of the Anticancer Properties of Optically
2 Active Titanocene Oximate Compounds

3 *Isabel de la Cueva-Alique,[‡] Sara Sierra,[‡] Adrián Pérez-Redondo,[‡] Isabel Marzo,[⊥]*
4 *Lourdes Gude,[‡] Tomás Cuenca,[‡] Eva Royo^{‡*}*

5

6 *Dedicated to the memory of Prof. Dr. Pascual Royo, who loved aquo titanium chemistry*

7

8 [‡] Departamento de Química Orgánica y Química Inorgánica, Instituto de Investigación
9 en Química Andrés del Río (IQAR), Universidad de Alcalá, 28805 Alcalá de Henares,
10 Madrid, Spain

11 [⊥] Departamento de Bioquímica y Biología Molecular y Celular, Universidad de Zaragoza,
12 50009 Zaragoza, Spain

13 **Keywords**

14 titanium, chiral, enantiomer, DNA, cytotoxic

15 **Corresponding Author**

16 *E-mail: eva.royo@uah.es. Phone Nr. (+34) 918854629; Fax Nr. (+34) 918854683

17 ABSTRACT

1 New water soluble and optically active cyclopentadienyl titanium derivatives $[(\eta^5-$
2 $C_5H_5)_2Ti\{(1R,4S)\text{-}\kappa ON,(R)NH\}Cl]$ (R = Bn (Benzyl) **1a'**, 2-pic (2-picolylamine) **1b'**)
3 have been synthesized. The novel compounds along with those previously described $[(\eta^5-$
4 $C_5H_5)_2Ti\{(1S,4R)\text{-}\kappa ON,(R)NH\}Cl]$ (R = Bn **1a**, 2-pic **1b**) were evaluated by polarimetry,
5 ultra-violet and circular dichroism spectroscopy. The structure of **1b** was determined by
6 single crystal X-ray crystallography and showed a unique terminal monohapto Ti-O
7 disposition of the oximato ligand. All enantiomers have been tested against several cancer
8 cell lines *in vitro*: prostate PC-3 and DU-145, lung A-549, pancreas MiaPaca-2, colorectal
9 HCT-116, leukemia Jurkat and cervical HeLa. In addition, **1a**, **1b** and **1b'** were tested
10 against non-tumorigenic prostate RWPE-1 cell line. After 24 h of incubation, **1b** and **1b'**
11 were moderately active against Jurkat and A-549 cells. The anti-proliferative effect of
12 titanium compounds on prostate PC-3, DU-145 and RWPE-1 cell lines was also assessed
13 after 72 h of drug exposure. The cytotoxic profile of the enantiomers was similar,
14 exception made for the PC-3 cells, with *S,R*-isomers exhibiting cytotoxicities 2 to 3 times
15 higher than *R,S*-compounds. Under these conditions, derivative **1b** showed calculated
16 IC_{50} values better than those of Tacke's Titanocene-Y (bis-[(*p*-
17 methoxybenzyl)cyclopentadienyl]titanium(IV) dichloride) on both the prostate PC-3 and
18 DU-145 cells. **1a** and **1b** cytotoxic behaviour shows certain selectiveness, with activities
19 2-4 times lower on normal prostate RWPE-1 than on cancer PC-3 cells. Furthermore, **1b**
20 produces higher cytotoxicity on prostate PC-3, DU-145 and RWPE-1 cells than the
21 additive dose of titanocene dichloride and pro-ligand **b**·HCl. Additionally, compound-
22 DNA interactions have been investigated by equilibrium dialysis, Fluorescence
23 Resonance Energy Transfer (FRET) melting assays and viscometric titrations, which
24 suggest that these metal complexes and/or their hydrolysis products bind DNA either in
25 the minor groove or externally.

1 1. Introduction

2 Since the successful introduction of cisplatin (*cis*-[PtCl₂(NH₃)₂]) as an anticancer drug,
3 much effort has been devoted to investigation of the anticancer activity of other
4 coordination and/or organometallic transition metal compounds [1-7]. The titanium
5 derivatives titanocene dichloride ([(η^5 -C₅H₅)₂TiCl₂], TDC) [8,9] and budotitane (*cis*-
6 diethoxybis(1-phenylbutane-1,3-dionato)titanium(IV)) [10,11] were the first metal
7 compounds to enter clinical trials after platinum complexes. Although these derivatives
8 showed promising properties in preliminary studies, they failed advanced clinical trials
9 due to low antitumor efficacy *in vivo*, rapid hydrolysis and limited solubility in biological
10 media [12-19]. Since then, a plethora of modified titanium based compounds have been
11 synthesized and studied as potential antitumor agents [17-29].


12 The effect of stereochemistry on biological activity is of great importance in medicinal
13 chemistry, as many of the biological targets are chiral [30,31]. The anticancer properties
14 of chiral metal derivatives have been largely studied [32-46], but the role of the
15 stereochemistry in the biological activity of non-platinum based compounds has been less
16 investigated [22,47-61]. Effect of the absolute configuration on the anticancer efficiency
17 of titanium compounds was firstly explored by Tshuva in 2010 [50]. The enantiomers of
18 C₂-symmetrical Ti(IV) compounds with chiral diamine bis(phenolato) ligands showed
19 different antitumor activities by factors of 2-4 on human colorectal (HT-29) and ovarian
20 (OVCAR-1) carcinoma cells [50,51,56,60]. According to these results, the authors
21 proposed that stereochemistry should be considered in the design, modification, and
22 improvement of active compounds [60]. The same year, Baird published a family of
23 enantiomerically pure titanocene derivatives bearing chiral alkylammonium groups, but
24 a relationship between the anticancer activity and chirality could not be established due
25 to the low cytotoxicity showed on the cancer cell lines evaluated [62]. Enantiomer-

1 dependent activity was found in chiral substituted titanocene compounds by Cini et al
2 [22,58], with the (*S,S*) enantiomer of Cp^R₂TiCl₂ (Cp^R = η⁵-C₅H₄CH(CH₂CH₃)C₆H₅OMe)
3 being twice as active as the (*R,R*) isomer towards pancreatic, breast and colon cancer cell
4 lines, after 24 h of treatment. Interestingly, lack of enantiomer recognition was observed
5 at 72 h when screening the compounds in MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-
6 diphenyltetrazolium bromide) assays.

7 Within this context, enantiomerically pure, naturally occurring terpenes are useful
8 building blocks for asymmetric synthesis [63,64]. They are inexpensive and
9 commercially available reagents in optically pure form, and easily tailored by
10 stereoselective functionalization [65]. On the other hand, oxime groups are presented as
11 excellent chemical modifiers, with a wide versatility of coordination modes going from
12 mono κNO, κON, to dihapto κ²N,O; either with *side on* or bridging coordination, which
13 could offer an increased stability of the final compounds when bonded to Ti(IV) acid
14 centres [66-68].


15 We have recently reported a new family of enantiopure cyclopentadienyl titanium(IV)
16 compounds with amino-oximato ligands derived from *R*-limonene, of formula [(η⁵-
17 C₅H₅)₂Ti{(1*S*,4*R*)-κON,(*R*)NH}Cl] (R = Bn **1a**, 2-pic **1b**) (Fig. 1), with relevant
18 antitumor properties. Our compounds show significant effects on cytotoxicity, cell
19 adhesion to collagen and migration of androgen-independent prostate cancer cells while
20 they do not seem to exhibit strong interactions with plasmid DNA by electrophoretic
21 mobility shift assays. Compounds **1a** or **1b** suffered hydrolysis in water or phosphate
22 buffered saline (PBS) solutions. However, the additive doses of TDC and **a**·HCl or **b**·HCl
23 produced lower antiproliferative effects on prostate cancer PC3 cells than those observed
24 after treatment with oximato titanocenes **1a** or **1b**, respectively. This fact led us to the

1 conclusion that the active operating titanium species was positively influenced by the
 2 presence of the oximato ligand [69].



3
 4 **Fig. 1** Optically active titanocene compounds containing ligands derived from *R*-
 5 limonene

6 Encouraged by these previous results, we decided to explore the reactions of TDC with
 7 the already described amino-oxime chiral compounds (1*R*,4*S*)-{NH(*R*),NOH} (R = Bn
 8 **a'**, 2-pic **b'**), see Fig. 2) [65,70,71], derived from *S*-limonene.



9
 10
 11 **Fig. 2** Synthesis of optically active titanocene oximato compounds

12 We report here on the synthesis and characterization of corresponding cyclopentadienyl
 13 Ti(IV) enantiomers $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}\{(1*R*,4*S*)\text{-}\kappa\text{ON},(\text{R})\text{NH}\}\text{Cl}]$ (R = Bn **1a'**, 2-pic **1b'**).

1 Their hydrolytic behaviour has been studied by ^1H NMR, Ultraviolet-visible (UV-Vis)
2 spectroscopy and circular dichroism (DC). These novel compounds along with those
3 previously described have been evaluated against several cancer cell lines *in vitro*:
4 prostate PC-3 and DU-145, lung A-549, pancreas MIA PaCa-2, colorectal HCT-116,
5 leukemia Jurkat and cervical HeLa. In addition, the compounds were tested against the
6 non-tumorigenic human prostate RWPE-1 cell line. DNA interactions of the metal
7 derivatives and/or their hydrolysis products have been further investigated by FRET
8 melting assays, equilibrium dialysis and viscometric titrations experiments.

9

10 **2. Experimental Section**

11 2.1. Chemicals and synthesis

12 Manipulations involving the synthesis of titanium compounds **1a**, **1a'**, **1b** and **1b'** and
13 Titanocene-Y (bis-[(*p*-methoxybenzyl)cyclopentadienyl]titanium(IV) dichloride) were
14 performed at an argon/vacuum manifold using standard Schlenk techniques or in a
15 MBraun MOD System glove-box. Solvents were dried by known procedures and used
16 freshly distilled. Titanocene-Y [72], (1*S*,4*R*)-{NH(R),NOH}, (R = Bn **a** [70], 2-pic **b**);
17 (1*R*,4*S*)-{NH(R),NOH} (R = Bn **a'**; 2-pic **b'**); corresponding adducts (1*S*,4*R*)-
18 {NH(R)·HCl,NOH}, (R = Bn **a**·HCl, 2-pic **b**·HCl); (1*R*,4*S*)-{NH(R)·HCl,NOH} (R =
19 Bn **a'**·HCl, 2-pic **b'**·HCl) [63,73] and metal compounds $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}\{(1*S*,4*R*)-$
20 $\kappa\text{ON},(\text{R})\text{NH}\}\text{Cl}]$ (R = Bn **1a**, 2-pic **1b**) [69] were prepared according to previous reports.
21 *R*- or *S*-limonene and isopentyl nitrite were reacted following the standard method
22 described by Carman et al in 1977 [73]. *R*-limonene, *S*-limonene, TDC and cisplatin were
23 purchased from Sigma-Aldrich. Commercially available reagents were used without
24 further purification. Nuclear Magnetic Resonance (NMR) spectra were recorded on a
25 Bruker 400 Ultrashield. ^1H and ^{13}C chemical shifts are reported relative to

1 tetramethylsilane. ^{15}N chemical shifts are reported relative to liquid ammonia (25 °C).
2 Coupling constants J are given in Hertz. Elemental analysis was performed on a LECO
3 CHNS 932 Analyzer at the Universidad de Alcalá or, alternatively, at the Universidad
4 Autónoma de Madrid. Fourier Transform Infrared (FT IR) spectra were recorded on IR
5 FT Perkin Elmer (Spectrum 2000) spectrophotometer on KBr pellets. The pH was
6 measured in a HANNA HI208 pHmeter in distilled water solutions. Circular Dichroism
7 (CD) spectra were recorded on a J-715 CD spectropolarimeter (Jasco, UK) at ambient
8 temperature (297 K). The spectra were determined at a concentration of 0.5 mM in water
9 using a quartz cuvette of 0.5 cm path length, scan speed of $20 \text{ nm}\cdot\text{min}^{-1}$, 0.1 nm band
10 width, 0.5 nm data pitch and 0.5 s of response time. Optical rotations of all the compounds
11 solutions were recorded on a Perkin Elmer 341 polarimeter, using the sodium D line (589
12 nm) at ambient temperature (297 K) in a quartz cell of 1 dm path length. Specific optical
13 rotation values were calculated according to the equation $[\alpha]_{\text{D}}^{25} = 100 \cdot \alpha_{\text{obs}} / l \cdot c$ [74].
14 Analytical balance and volumetric pipettes (2.0 mL) were used to prepare CHCl_3
15 solutions of the compounds at concentrations within a range of $7.50\text{-}7.80 \text{ g}\cdot\text{dL}^{-1}$. UV-Vis
16 spectra were measured at room temperature on water solutions of the compounds with a
17 Perkin Elmer Lambda 35 spectrophotometer.

18 2.1.1. (*1R,4S*)-{NH(2-pic),NOH} (**b'**). An analogous procedure to that described before
19 for the synthesis of **b** [63] was used, starting from *S*-limonene [70,71,73]. $[\alpha]_{\text{D}}^{23}$ ($\text{deg}\cdot\text{dm}^{-1}\cdot\text{cm}^3\cdot\text{g}^{-1}$) -126 ± 1.3 (**b'** at $c = 0.7839 \text{ g}\cdot\text{dL}^{-1}$, $\alpha_{\text{obs}} = -0.957 \text{ deg}$); $+127 \pm 1.3$ (**b** at $c =$
20 $0.7604 \text{ g}\cdot\text{dL}^{-1}$, $\alpha_{\text{obs}} = +0.954 \text{ deg}$). All analytical and spectroscopic data are identical to
21 those observed for **b**. Anal. Calcd. for $\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}$: C, 70.30; H, 8.48; N, 15.37; Found:
22 C, 70.13; H, 8.07; N, 15.20. FT IR (KBr, $\lambda_{\text{max}}/\text{cm}^{-1}$): 3086-3314 (br, $\nu_{\text{OH/NH}}$), 1650,
23 1598 ($\nu_{\text{C=N}}$). UV-Vis (0.1 mM in $\text{H}_2\text{O}:\text{DMSO}$ 99:1): λ_{max} (ϵ): 261 (316), 340 (10). ^1H
24 NMR (plus two dimensional correlation spectroscopy (COSY), 400.1 MHz, 293 K,
25

1 chloroform-*d*₁): δ 9.80 (=NOH), 8.49, 7.60, 7.28, 7.11 (m, each 1H, NC₅H₄), 4.75 (br,
2 2H, =CH₂), 3.87, 3.61 (both d, each 1H, ³J_{HH} = 6, -CH₂-C₅H₄N), 3.28 (d, 1H, ²J_{HH} = 12, -
3 CH₂³), 2.60 (br, 1H, NH), 2.09 (m, 1H, -CH⁴), 2.03 (dd, 1H, ²J_{HH} = 12, ³J_{HH} = 3, -CH₂³),
4 2.00, 1.69 (m, each 1H, -CH₂⁶ + -CH₂⁵), 1.85 (m, 1H, CH₂⁶), 1.75 (s, 3H, CH₃-C=), 1.65
5 (m, 1H, CH₂⁵), 1.32 (s, 3H, -CH₃-C_q-N). ¹³C NMR (plus Attached Proton Test (APT),
6 plus gradient Heteronuclear Single Quantum Coherence (gHSQC), plus Heteronuclear
7 Multiple Bond Correlation (HMBC), 100.6 MHz, 293 K, chloroform-*d*₁): δ 162.4
8 (C_q=NOH, C_q is quaternary carbon), 161.1 (C_{ipso}-C₅H₄N), 148.9 (C=CH₂), 149.2, 136.8,
9 122.7, 122.1 (C₅H₄N), 109.6 (=CH₂), 56.9 (C_q-NH), 48.1 (CH₂-C₅H₄N), 45.0 (CH⁴), 40.5
10 (-CH₂⁶), 26.4 (-CH₂⁵), 25.6 (-CH₂³), 23.5 (-CH₃-CNH), 21.0 (CH₃-C=). ¹⁵N NMR
11 (gHMBC, 40.5 MHz, 293 K, chloroform-*d*₁): δ 346.7 (C=N-), 305.3 (C₅H₄N), 51.8 (-
12 NHpic).

13 2.1.2. [(η^5 -C₅H₅)₂Ti{(1*R*,4*S*)- κ ON,(Bn)NH}Cl] (**1a'**). An analogous procedure to that
14 described for [(η^5 -C₅H₅)₂Ti{(1*S*,4*R*)- κ ON,(Bn)NH}Cl] [69] was followed, starting from
15 TDC (0.20 g, 0.80 mmol), (1*R*,4*S*)-{NH(Bn),NOH} (0.22 g, 0.80 mmol) and NEt₃ (0.11
16 mL, 0.80 mmol). Compound **1a'** was obtained as a yellow-orange solid. Yield: 0.32 g
17 (88%). [α]_D²³ (deg·dm⁻¹·cm³·g⁻¹) -88.9 \pm 1.2 (**1a'** at c = 0.7602 g·dL⁻¹, α_{obs} = -0.676 deg),
18 +89.2 \pm 1.2 (**1a** at c = 0.7497 g·dL⁻¹, α_{obs} = +0.681 deg). Analytical and spectroscopic
19 data of the compound are identical to those already reported [69]. Solubility in H₂O at 24
20 °C (mM): 6.6 \pm 0.2. Value of pH ([2.0 mM]) in H₂O at 24 °C: 5.54. Anal. Calcd for
21 C₂₇H₃₃ClN₂O₂Ti: C, 66.88; H, 6.86; N, 5.78; Found: C, 66.80; H, 6.90; N, 5.76. FT IR
22 (KBr, λ_{max} /cm⁻¹): 3370 (m, NH), 1646, 1601 (both m, C=N). ¹H NMR (plus HSQC,
23 plus HMBC, plus COSY, 400.1 MHz, 293 K, chloroform-*d*₁): δ 7.32 (m, 5H, -C₆H₅),
24 6.39, 6.39 (both s, each 5H, C₅H₅), 4.76, 4.74 (both s, each 1H, =CH₂), 3.76, 3.55 (both

1 m, each 1H, -CH₂Ph), 2.92 (m, 1H, -CH₂³), 2.05 (m, 1H, -CH-C=), 1.90 (m, 1H, -CH₂⁶),
2 1.72 (m, 1H, -CH₂³), 1.68 (m, 1H, -CH₂⁵), 1.59 (m, 1H, -CH₂⁶), 1.56 (m, 1H, -CH₂⁵), 1.25
3 (br, 1H, NH), 1.47, 1.25 (both s, each 3H, NC-CH₃ + CH₃C=). ¹³C NMR (plus APT, plus
4 gHSQC, plus HMBC, 100.6 MHz, 293 K, chloroform-*d*₁): δ 159.2 (Cq=N), 149.3 (=Cq-
5 Me), 141.6 (C_{ipso}Ph), 128.7, 128.7, 127.2 (C₆H₅), 117.1, 117.1 (C₅H₅), 109.4 (=CH₂), 57.1
6 (Cq-NH), 47.3 (-CH₂Ph), 45.6 (-CH⁴), 41.2 (-CH₂⁶), 27.8 (-CH₂³), 26.2 (-CH₂⁵), 23.9,
7 21.3 (CH₃-CNH + CH₃-C=). ¹⁵N NMR (gHMBC, 40.5 MHz, 293 K, chloroform-*d*₁): δ
8 398.9 (C=N), 60.0 (NHBn).

9 2.1.3. [(η⁵-C₅H₅)₂Ti{(1*R*,4*S*)-κON,(2-pic)NH}Cl] (**1b'**). An analogous procedure to
10 that described for [(η⁵-C₅H₅)₂Ti{(1*S*,4*R*)-κON,(2-pic)NH}Cl] [69] was followed, starting
11 from TDC (0.30 g, 1.20 mmol), (1*R*,4*S*)-{NH(2-pic),NOH} (0.33 g, 1.20 mmol) and NEt₃
12 (0.11 mL, 1.20 mmol). Compound **1b'** was obtained as a yellow-orange solid. Yield: 0.35
13 g (60%). [α]²³_D (deg·dm⁻¹·cm³·g⁻¹) -75.7 ± 1.2 (**1b'** at c = 0.7534 g·dL⁻¹, α_{obs} = -0.570
14 deg), +74.2 ± 1.2 (**1b** at c = 0.7772 g·dL⁻¹, α_{obs} = +0.570 deg). Solubility in H₂O at 24 °C
15 (mM): 15.7 ± 1.7. Value of pH ([2.0 mM]) in H₂O at 24 °C: 5.22. Anal. Calcd for
16 C₂₆H₃₂ClN₃OTi: C, 64.27; H, 6.64; N, 8.65; Found: C, 64.62; H, 7.25; N, 8.54. FT IR
17 (KBr, λ_{max}/cm⁻¹): ν̄ 3304 (m, NH), 1640, 1591, 1569 (all s, C=N). ¹H NMR (plus HSQC,
18 plus HMBC, plus COSY, 400.1 MHz, 293 K, chloroform-*d*₁): δ 8.50, 7.60, 7.30, 7.12 (all
19 m, each 1H, -NC₅H₄), 6.38, 6.38 (both s, each 5H, C₅H₅), 4.77, 4.74 (both s, each 1H,
20 =CH₂), 3.91, 3.70 (both m, each 1H, CH₂-C₅H₄N), 2.84 (m, 1H, -CH₂³), 2.07 (m, 1H, -
21 CH-C=), 1.98 (m, 2H, overlapped -CH₂⁶⁺³), 1.78 (s, 3H, CH₃C=), 1.64 (m, 1H, -CH₂⁶),
22 1.62 (m, 1H, -CH₂⁵), 1.60 (m, 1H, -CH₂⁵), 1.48 (br, 4H, NC-CH₃ + NH). ¹³C NMR (plus
23 APT, plus gHSQC, plus HMBC, 100.6 MHz, 293 K, chloroform-*d*₁): δ 157.6 (Cq=N),
24 148.1 (=Cq-Me), 160.2 (C_{ipso}C₅H₄N), 149.3, 136.7, 122.9, 122.9 (C₅H₄N), 117.1, 117.1

1 (C₅H₅), 109.6 (=CH₂), 48.5 (-CH₂-C₅H₄N), 45.3 (-CH⁴), 41.1 (-CH₂⁶), 27.7 (-CH₂³), 26.2
2 (-CH₂⁵), 23.9, 21.3 (CH₃-CNH + CH₃-C=). ¹⁵N NMR (gHMBC, 40.5 MHz, 293 K,
3 chloroform-*d*₁): δ 402.1 (C=N), 312.5 (C₅H₄N), 52.6 (NHpic).

4 2.1.5. ¹H NMR experiments at physiological pH. Phosphate buffered saline solution
5 (PBS) was prepared according to Cold Spring Harbor Protocols
6 (<http://cshprotocols.cshlp.org/content/2006/1/pdb.rec8247>) using NaCl, KCl, Na₂HPO₄
7 and K₂HPO₄ in D₂O. Adjustment of pD (pD = pH* + 0.4, where pH* = pHmeter reading
8 in D₂O) was carried out using a solution of DCl (0.01M) or NaOD (0.01M) in D₂O, with
9 the help of a HANNA HI208 pHmeter. Titanium compounds were then dissolved in 2000
10 μL of the freshly prepared PBS, final pD measured (7.30-7.38) and time-dependent ¹H
11 NMR spectra of 500 μL aliquots of final solutions were carried out at 25 °C.

12 2.2. Single-crystal X-ray structure determination

13 Yellow crystals of pure enantiomer **1b** were grown from a hexane-toluene solution. The
14 crystals were removed from the vial and covered with a layer of a viscous
15 perfluoropolyether. A suitable crystal was selected with the aid of a microscope,
16 mounted on a cryo-loop, and placed in the low-temperature nitrogen stream of the
17 diffractometer. The intensity data sets were collected at 200 K on a Bruker-Nonius
18 Kappa CCD diffractometer equipped with an Oxford Cryostream 700 unit. The
19 molybdenum radiation (λ = 0.71073) was used in both cases, graphite monochromated,
20 and enhanced with an MIRACOL collimator.

21 The structure was solved, using WINGX package [75], by intrinsic phasing methods
22 (SHELXT) [76], and refined by least-squares against F² (SHELXL-2014/7) [77]. Crystals
23 of **1b** were refined as a two-component inversion twin, and also had two independent
24 molecules in the asymmetric unit with no significant differences. All non-hydrogen atoms

1 were anisotropically refined. Positions of the amine hydrogen atoms, H(2) and H(21),
2 were located in the difference Fourier map. H(2) was refined isotropically, while U_{iso} for
3 H(21) was fixed with a value of 0.05. The rest of the hydrogen atoms were positioned and
4 refined by using a riding model. *Crystal data for 1b*: (C₂₆H₃₂ClN₃OTi), FW = 485.89,
5 Monoclinic, space group $P2_1$, crystal dimensions (mm³) 0.30 x 0.27 x 0.27, $a = 10.470(1)$,
6 $b = 11.631(1)$, $\beta = 91.53(1)$, $c = 19.856(3)$ Å, $V = 2417.2(5)$ Å³, $Z = 4$, $\rho_{\text{calcd}} = 1.335$ g
7 cm⁻³, $\mu = 0.488$ mm⁻¹, $F(000) = 1024$, θ range = 3.08 to 27.50 deg, no. of rflns collected
8 = 42638, no. of indep rflns / $R_{\text{int}} = 10939 / 0.074$, no. of data / restraints / params = 10939
9 / 1 / 589, $R1 / wR2 (I > 2\sigma(I)) = 0.068 / 0.141$, $R1 / wR2$ (all data) = 0.089 / 0.151, GOF
10 (on F^2) = 1.167, Absolute structure parameter = 0.04(5). Final difference Fourier maps
11 did not show peaks higher than 0.695 nor deeper than -0.329 eÅ⁻³. CCDC-1572920
12 contains the supplementary crystallographic data for this paper. These data can be
13 obtained free of charge from The Cambridge Crystallographic Data Centre via
14 www.ccdc.cam.ac.uk/structures.

15 2.3. Cell culture, cytotoxicity assays and cell death analysis

16 2.3.1. Cell culture

17 The prostate androgen-unresponsive cancer cell line PC-3 was obtained from the
18 American Type Culture Collection (Manassas, VA) and may be related to recurrent
19 prostate cancers that have achieved androgen independence. All culture media were
20 supplemented with 1% penicillin/streptomycin/amphoterycin B (Life Technologies,
21 Barcelona, Spain). The culture was performed in a humidified 5% CO₂ environment at
22 37 °C. After the cells reached 70–80% confluence, they were washed with PBS, detached
23 with 0.25% trypsin/0.2% ethylenediaminetetraacetic acid (EDTA) and seeded at 30,000–
24 40,000 cells·cm⁻². The culture medium was changed every 3 days. A549 (lung carcinoma)
25 cells were maintained in high glucose DMEM (Dulbecco's Modified Eagle's Medium)

1 and RWPE-1 (non-tumorigenic prostate) cells in DMEM/F12 (Dulbecco's Modified
2 Eagle Medium: Nutrient Mixture F-12), supplemented with 5% fetal bovine serum (FBS),
3 $200 \text{ U}\cdot\text{mL}^{-1}$ penicillin, $100 \text{ }\mu\text{g}\cdot\text{mL}^{-1}$ streptomycin and 2 mM L-glutamine. DU-145
4 (prostate carcinoma), MIA PaCa-2 (pancreas carcinoma), HCT-116 (colorectal
5 carcinoma), HeLa (cervical cancer) and Jurkat (leukemic cancer) cells were maintained
6 in RPMI (Roswell Park Memorial Institute) 1640 medium supplemented with 5% FBS,
7 $200 \text{ U}\cdot\text{mL}^{-1}$ penicillin, $100 \text{ }\mu\text{g}\cdot\text{mL}^{-1}$ streptomycin and 2 mM L-glutamine. Cultures were
8 maintained in a humidified atmosphere of 95% air:5% CO_2 at 37 °C. Adherent cells were
9 allowed to attach for 24 h prior to addition of compounds.

10 2.3.2. MTT Toxicity Assays

11 For toxicity assays, cells (5×10^4 for Jurkat cells and 10^4 for adherent cell lines) were
12 seeded in flat-bottom 96-well plates (100 μL /well) in complete medium. Adherent cells
13 were allowed to attach for 24 h prior to addition of cisplatin or tested compounds. Stock
14 solutions of Titanocene-Y, TDC and ammonium-oxime pro-ligands were freshly
15 prepared in 1% of dimethyl sulfoxide (DMSO) in water, while cisplatin and oximato
16 titanium compounds were dissolved in culture medium. The stock solutions were then
17 diluted in complete medium and used for sequential dilutions to desired concentrations.
18 The final concentration of DMSO in the cell culture medium did not exceed 0.1%. Control
19 groups with and without DMSO (0.1%) were included in the assays. Compounds were
20 then added at different concentrations in quadruplicate. Cells were incubated with
21 compounds for 24 h or 72 h, and then cell proliferation was determined by a modification
22 of the MTT-reduction method. Briefly, 10 μL /well of [3-(4,5-dimethylthiazol-2-yl)-2,5-
23 diphenyltetrazolium bromide] (MTT) ($5 \text{ mg}\cdot\text{mL}^{-1}$ in PBS) was added, and plates were
24 incubated for 1–3 h at 37 °C. Finally, formazan crystals were dissolved by adding 100
25 μL /well *i*PrOH (0.05 M HCl) and gently shaking. The optical density was measured at

1 550 nm using a 96-well multi-scanner auto-reader Enzyme-Linked Immuno Sorbent
2 Assay (ELISA).

3 2.4. DNA interaction studies

4 2.4.1. Equilibrium Dialysis

5 Duplex DNA from calf thymus (CT DNA), (Deoxyribonucleic acid, Activated, Type
6 XV) was directly purchased from Sigma Aldrich and used as provided. Duplex-forming
7 oligonucleotides ds17-1 (5'-CCA GTT CGT AGT AAC CC-3') and ds17-2 (5'-GGG TTA
8 CTA CGA ACT GG-3') were acquired High Performance Liquid Chromatography
9 (HPLC) -purified and desalted from Integrated DNA Technologies (IDT). Dialysis
10 membranes (Spectra/Por® molecular porous membrane tubing MWCO: 3.5–5.0 kDa; 6.4
11 mm diameter) were purchased from Spectrum Laboratories Inc. Aqueous solutions of
12 surfactant sodium dodecyl sulphate (SDS) (10%) were purchased from Sigma Aldrich.
13 The buffer employed in this experiment was 10 mM phosphate buffer
14 $\text{NaH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4$, pH = 7.2, with either 10 mM or 100 mM NaCl. The solutions of
15 DNA were prepared in the working phosphate buffer at 75 μM monomeric unit (mum.)
16 concentrations, in base pairs. For the preparation of the short oligonucleotide solution, an
17 annealing step was needed, with heating at 90 °C for 10 min and then gradually cooling
18 to 25 °C during 3 h. The solutions were left at 4 °C overnight.

19 Dialysis bags, previously washed with milli-Q water, were filled with 75 μM (m.u.) of
20 DNA duplex (200 μL each bag) and placed in a beaker containing 225 mL of ca. 2 μM
21 solution of the tested compound. The beaker was covered with parafilm and aluminium
22 foil and allowed to equilibrate during 24 h at room temperature. Experiments were run,
23 at least, in triplicate. Once the dialysis process had been completed, the solutions from
24 each dialysis bag were transferred to Eppendorf tubes. The content of each bag was then
25 mixed with an aqueous detergent solution (10%) to reach a 1% concentration (v/v) of

1 SDS. The concentrations of free compound in the dialysate solution and compound in the
2 dialysis bags were determined by absorbance measurements using the extinction
3 coefficients of the metal complexes (determined in the presence and absence of the
4 detergent) and apparent association constants were calculated [78].

5 2.4.2. DNA FRET melting assay

6 The DNA melting assay was performed on a quantitative PCR kit ABI PRISM® 7000
7 Sequence Detection System (Applied Biosystems) in a 96-well plate format (96-Well
8 Optical MicroAmp® Reaction Plate, Applied Biosystems, Life Technologies
9 Corporation). The oligonucleotide sequence employed in this experiment, F10T (5'-
10 FAM-AGC TAT TA /sp18/ TA TA GCT ATA-TAMRA-3') was produced, HPLC-
11 purified and desalted by IDT. FAM is 6-carboxyfluorescein and TAMRA is
12 carboxytetramethylrhodamine. The buffer system used in this experiment was: 10 mM
13 sodium cacodylate, 100 mM LiCl, (pH = 7.3). First, the duplex-forming oligonucleotide
14 was dissolved in water (grade BPC) and a 50 µM stock solution was prepared, which was
15 then diluted to 0.5 µM. Then, the diluted DNA solution was mixed with the working
16 buffer (2x) and water Biotechnology Performance Certified (BPC) grade. The DNA
17 solution was heated at 90 °C for 10 min, cooled down slowly for 3 h and left at 4 °C
18 overnight. Compounds to be tested were dissolved in water and approximately 1 mM
19 stock solutions were prepared. The exact concentrations were checked by UV-Vis. Stock
20 solutions were then diluted with buffer to obtain 50 µM solutions of each compound. In
21 a 96-well microplate, DNA solutions were mixed with solutions of tested compound and
22 buffer to reach a total volume of 50 µL with a F10T concentration of 0.2 µM and a
23 compound concentration ranging between 1 and 10 µM.

24 The experimental protocol consisted of an incubation for 5 min at 24 °C, followed by
25 a temperature ramp with heating rate 1 °C/min. Fluorescence values corresponding to the

1 fluorophore FAM at wavelength of 516 nm (after excitation at 492 nm) were collected at
2 each degree of temperature. Afterwards, the fluorescence data were normalized, plotted
3 against temperature (°C) at each compound concentration, and T_m values were
4 determined.

5 2.4.3. Viscometric titrations

6 Duplex DNA from CT (Deoxyribonucleic acid, Activated, Type XV) was purchased
7 from Sigma Aldrich and used as provided. The buffer employed in this experiment was
8 10 mM phosphate buffer $\text{NaH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4$, pH = 7.2. The viscosity measurements
9 were performed in a Visco System AVS 470 at 25.00 ± 0.01 °C, using a microUbbelohde
10 (K = 0.01) capillary viscometer. 6 mL of DNA solution (0.4 mM in nucleotides) in
11 phosphate buffer were equilibrated for 20 min at 25.00 °C and then 20 flow times were
12 registered. Small aliquots (30–50 μL) of solutions of metal complexes (1.6–2.3 mM) were
13 added to the same DNA solution. Before each flow time registration, the solutions were
14 equilibrated for 20 min to 25.00 °C and then 20 flow times were measured. With the
15 averaged time of the different flow time measurements and the viscometer constant, the
16 viscosities (μ) for each point were calculated. The viscosity results were plotted as
17 $(\mu/\mu_0)^{1/3}$, where μ_0 represents the DNA solution viscosity in the absence of the ligand,
18 versus (r), representing the ratio [ligand]/[DNA].

19 2.5. Data analysis

20 Results were subjected to computer-assisted statistical analysis using One-Way
21 Analysis of Variance ANOVA, Bonferroni's post-test, and Student's t-test. Data are
22 shown as the means of individual experiments and presented as the mean \pm SD (Standard
23 deviation). Differences of $P < 0.05$ were considered to be significantly different from the
24 controls.


3. Results and Discussion

3.1. Synthesis and characterization of metal compounds

Synthesis of the novel Ti(IV) compounds was carried out analogously to that of previously described enantiomers **1a** and **1b** [69]. Treatment of TDC and amino-oxime derivatives **a'** or **b'** in the presence of NEt_3 allows isolation of novel chiral-at-ligand titanium compounds **1a'** or **1b'**, respectively (Fig. 2), which are formed together with $\text{Et}_3\text{N}\cdot\text{HCl}$.

Analytical and spectroscopic data of the novel compounds **1a'** and **1b'** are identical to those reported before for **1a** and **1b**, respectively (see ref [69], Experimental Section and Online Resource, Fig. S3-S9).

Calculated data of specific optical rotation in chloroform solution for the ligands and novel metal derivatives ($[\alpha]^{23}_{\text{D}}$ ($\text{deg}\cdot\text{dm}^{-1}\cdot\text{dL}\cdot\text{g}^{-1}$) = -127 ± 1.3 **a'**, $+130 \pm 1.3$ **a**, -126 ± 1.3 **b'**, $+127 \pm 1.3$ **b**, -88.9 ± 1.2 **1a'**, $+89.2 \pm 1.2$ **1a**, -75.7 ± 1.2 **1b'**, $+74.2 \pm 1.2$ **1b**) evidence the enantiomeric relationship of the stereoisomers. Furthermore, absolute configuration of the compound **1b** has been confirmed through X-ray structure determination (Fig. 3, and Online Resource Table S1, S2 and Fig. S16).



1

2 **Fig. 3** ORTEP drawing of compound **1b** with 50% probability ellipsoids. Hydrogen
 3 bonded to carbon atoms have been omitted for clarity. Representative bond lengths (Å)
 4 and angles (deg): Ti(1)-Ct(1) 2.073; Ti(1)-Ct(2) 2.065; Ti(1)-Cl(1) 2.380(2); Ti(1)-O(1)
 5 1.899(4); Ti(1)···N(1) 2.866(5); N(1)-O(1) 1.403(6); N(1)-C(21) 1.273(8); Cl(1)-Ti(1)-
 6 O(1) 92.4(2); Ti(1)-O(1)-N(1) 119.6(3); O(1)-N(1)-C(21) 114.1(5); Ct(1)-Ti(1)-Ct(2)
 7 130.3; (Ct(1) is the centroid of the C(11)-C(15) ring, Ct(2) is the centroid of the C(16)-
 8 C(20) ring)

9

10 The X-ray crystal structure determination of **1b** shows the presence of two independent
 11 molecules in the asymmetric unit, with the same absolute configuration of the two chiral
 12 centers; an ORTEP diagram of one of them is presented in Fig. 3. The crystallographic
 13 study confirms a monohapto coordination of the oximate unit to the titanium atom. The
 14 compound shows a pseudotetrahedral environment around the metal centre, with Ti-O
 15 bond distances and O-N-C angles slightly shorter and closer (Online Resource Fig. S16),
 16 respectively, than those found in analogous biscyclopentadienyl oximate titanium(IV)


1 derivatives [66,68] or alcoxoximate titanium(IV) compounds [79-82] with a dihapto
2 $\kappa^2\text{NO}$ coordination of the oximate unit to the titanium centre.

3 To the best of our knowledge, this is the first example found of an oximate titanium
4 derivative with a terminal monohapto Ti-ON= coordination, where this coordination
5 mode is probably caused by the large steric requirements of the functionalized
6 cyclohexane residue. This terminal coordination may account for the hydrolysis suffered
7 for the compounds in aqueous media. In contrast, dihapto titanocene oximate compounds
8 $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{H}_2\text{O})(\kappa^2\text{O}=\text{NR})]^+$ (R = CMe₂; C₆H₁₀), reported by Thewalt et al [66], were
9 described as surprisingly stable against air and water.

10 The reactions in water or PBS solutions of **1a** or **1b** were elucidated in a previous report
11 and afford soluble ammonium-oxime pro-ligands (1*S*,4*R*)-{NH(R)·HCl,NOH} (R = Bn
12 **a**·HCl or 2-pic **b**·HCl, respectively), together with aqua-oxo or -hydroxo
13 biscyclopentadienyltitanium(IV) species [69,83,84] which are detected at least during the
14 first three hours after dilution. The same behavior as that described before has now been
15 observed for novel compounds **1a'** and **1b'** when their solutions in water-*d*₂ or PBS were
16 studied by ¹H NMR spectroscopy (see Online Resource, Fig. S10).

17 We decided to further investigate the existence of an amino-oxime ligand containing
18 Ti(IV) species, which could account for the observed stereoisomer-dependent cytotoxic
19 behaviour of the compounds on the prostate cancer PC-3 cell line. Since UV-Vis
20 spectroscopy is considered a more sensitive technique than NMR, we recorded time-
21 dependent UV-Vis spectra for compounds **1a'** and **1b'** in PBS solution. Right after
22 dilution, UV-Vis spectrum of **1a'** (Online Resource Fig. S13) and **1b'** (Fig. 4) shows two
23 very broad absorption bands centered at 240 and 325, and 246 and 322 nm, respectively,
24 ascribed to overlapping of LMCT bands due to cyclopentadienyltitanium aquo cations
25 and the absorption bands corresponding to proligands **a'**·HCl and **b'**·HCl. After 24 h,

1 only the absorption bands assigned to $\mathbf{a}'\cdot\text{HCl}$ or $\mathbf{b}'\cdot\text{HCl}$, at 250 and 332, and 260 and
2 332 nm, respectively, are detected. Similar UV-Vis spectra are obtained after 72 h.
3 Analogous results were obtained when the compounds are diluted in pure water.



4
5 **Fig. 4** Comparison of time-dependent UV-Vis spectra of $\mathbf{1b}'$ with $\mathbf{b}'\cdot\text{HCl}$ and TDC
6 spectra in PBS solution

7 CD spectra were also recorded for each pair of enantiomers. However, the spectra of
8 derivatives $\mathbf{1a}$, $\mathbf{1b}$ and $\mathbf{1a}'$, $\mathbf{1b}'$ are identical to those obtained for ammonium-oxime
9 compounds $\mathbf{a}\cdot\text{HCl}$, $\mathbf{b}\cdot\text{HCl}$, $\mathbf{a}'\cdot\text{HCl}$, $\mathbf{b}'\cdot\text{HCl}$ (see Online Resource Fig. S14, S15), 15 min
10 after dilution or after 72 h, leading to the assumption that those are the only detectable
11 optically active, soluble in water products of the hydrolysis of titanium oximate
12 compounds.

13 3.2. *In vitro* cell studies

3.2.1. Anti-proliferative studies

Chiral compounds **1a** and **1b** have already shown their promising anticancer properties on the human prostate and renal cancer cell lines PC-3 and Caki-1. Both titanocenes, especially **1b**, are considerable less toxic to the non-tumorigenic human embryonic kidney cell line HEK-293T than to Caki-1 renal cells (7-15-fold less toxic) [69].

In order to compare and evaluate the versatility of the different enantiomers, the cytotoxic activity of pro-ligands **a·HCl**, **a'·HCl**, **b·HCl**, **b'·HCl** and metal compounds TDC, Tacke's Titanocene-Y [72,85,86], **1a**, **1a'**, **1b** and **1b'** was now assessed after 24 h of incubation time on a wide variety of human cancer cell lines, i.e. prostate PC-3 and DU-145, lung A-549, pancreas MIA PaCa-2, colorectal HCT-116, leukemia Jurkat and cervical HeLa. The *in vitro* effect of the compounds on cytotoxicity was firstly evaluated by monitoring their ability to inhibit cell growth using the MTT assay.

Under these conditions, pro-ligands **a·HCl**, **a'·HCl**, **b·HCl**, **b'·HCl** and metal compounds **1a** and **1a'**, TDC, and Titanocene-Y are poorly cytotoxic in all tested cell lines ($IC_{50} > 150 \mu\text{M}$ under these experimental conditions). Enantiomers **1b** and **1b'** are also not effective, after 24 h of exposure, in prostate PC-3, pancreatic MIA PaCa-2 or colon HCT-116 human carcinoma cell lines, but show inhibitory activities of 40-50% and 20-25% at concentrations of 50 μM against human lung carcinoma A-549 (Online Resource Fig. S17) and leukemia Jurkat-T cell lines respectively. Cell morphology evaluation of A-549 cells indicated that titanium derivatives **1b** and **1b'** did not induced apoptotic cell death, since no apoptotic cells, characterized by condensed nuclei and membrane blebbing, were detected. Cisplatin was included in the experiment as a positive control of apoptosis.

Since compounds **1a** and **1b** had shown to be efficiently cytotoxic on the PC-3 cell line after 72 h of incubation with the cells [69], we decided to assess the anti-proliferative

1 effect of titanium compounds on prostate PC-3 and DU-145 cell lines as the IC₅₀ value
 2 after 72 h of drug exposure. The results are summarized in Table 1.

3 **Table 1.** IC₅₀ values (μM) of cisplatin, Titanocene-Y and enantiomers **1a**, **1a'**, **1b** and
 4 **1b'** in prostate cancer PC-3, DU-145 and non-tumorigenic RWPE-1 cell lines,^a (n.m. not
 5 measured)

Compound	PC-3	DU-145	RWPE-1
1a	> 150 (24 h) 48.7 ± 3.2 (72 h)	> 150 (72 h)	> 200 (72 h)
1a'	> 150 (24 h) > 150 (72 h)	> 150 (72 h)	n.m.
1b	> 150 (24 h) 14.5 ± 3.1 (72 h)	27.1 ± 1.1 (72 h)	30.8 ± 0.57 (72 h)
1b'	> 150 (24 h) 49.9 ± 7.0 (72 h)	23.9 ± 8.6 (72 h)	43.8 ± 7.2 (72 h)
1b + 1b'	37.5 ± 5.1 (72 h)	n.m.	n.m.
Titanocene-Y	> 200 (24 h) 58.1 ± 11.2 (72 h)	> 150 (72 h)	42.9 ± 0.73 (72 h)
cisplatin	104.2 ± 8.1 (24 h) 14.5 ± 2.5 (72 h)	3.7 ± 0.6 (72 h)	19.9 ± 1.1 (72 h)

6 ^a Each value represents the mean ± S.D. (n = 3)

7

8 Under these conditions, the enantiomer **1b** shows IC₅₀ values on the prostate PC-3 and
 9 DU-145 cell lines 2-5 times lower than Tacke's compound, Titanocene-Y. The
 10 cytotoxicity on PC-3 cells of the titanium enantiomers **1a** and **1b**, with the absolute
 11 configuration *S,R*-, is higher than that of the *R,S*-stereoisomers by a factor of ca. 2-3,
 12 while the racemic mixture of **1b** and **1b'** afford IC₅₀ values average between the two
 13 enantiomers. In contrast, no enantiomer recognition is observed on the prostate DU-145
 14 cells for derivatives **1b**, **1b'**, while **1a**, **1a'** resulted not to be efficient in this non-hormone
 15 dependent cancer cell line.

16 Titanocene-Y has already shown an encouraging activity in PC-3 tumour-bearing mice
 17 [85]. Other titanium compounds which have proved their *in vitro* antitumor activity in

1 prostate cancer cell lines under similar time exposure conditions are Schiff-base titanium
2 (IV) derivatives [87] (IC_{50} values within the range 5-18 μM , in PC-3) or heterometallic
3 titanocene-gold compounds (IC_{50} values ranged from 27-40 μM in PC-3 [88,89], and
4 11.8-27.6 μM in DU-145 [24,90]).

5 In order to analyse the cytotoxic selectiveness to healthy cells, the isomers **1a**, **1b** and
6 **1b'** were also tested in the non-tumorigenic human prostate (RWPE-1) cells. Regarding
7 selectivity, **1a** and **1b** are less toxic to the non-tumorigenic RWPE-1 than to the cancer
8 PC-3 cells (from 2 to 4 times less toxic), while **1b'** shows a similar behaviour relative to
9 the cancer DU-145 cells.

10 Titanium compound **1b** was selected for a further study *in vitro*. We evaluated a
11 combination of TDC and pro-ligand **b·HCl** on the cellular viability after 72 h of exposure
12 to the drug. As already described in the PC-3 cell line,[69] the additive dose of both
13 starting materials also produced lower anti-proliferative effects than those observed after
14 treatment with only **1b** (Table 2) in the prostate DU-145 and RWPE-1 cell lines. These
15 results are consistent with the involvement of metal oxime containing species in the
16 cytotoxicity mechanism. While water soluble hydrolysis species detected in our studies
17 are the same as those formed from a mixture of TDC and amino-oxime proligand, the
18 existence of polinuclear, ligand influenced species formed in a colloidal phase of
19 hydrolysis cannot be ruled out.

20

21

22

23

24 **Table 2.** Comparison of IC_{50} values (μM) of **1b**, **b·HCl**, TDC and TDC+**b·HCl** in
25 prostate cancer PC-3, DU-145 and non-tumorigenic RWPE-1 cell lines^a (after 72 h of
26 exposure to the drug).

Compound	PC-3	DU-145	RWPE-1
1b	14.5 ± 3.1	27.1 ± 1.1	30.8 ± 0.57
b·HCl	> 100	106.1 ± 10.4	140.5 ± 23.0
TDC+b·HCl	39.5 ± 2.1	54.9 ± 13.5	> 150
TDC	> 150	> 150	> 150

^a Each value represents the mean ± S.D. (n = 3)

3.3. DNA binding

To date, various distinct mechanisms have been proposed for titanium-based therapeutics. DNA binding is still thought to be one important potential mode of action for titanocene compounds, although interactions with DNA have been found to be generally very weak at physiological pH conditions [16,17]. The study of DNA interactions for these particular metal complexes does often represent an experimental challenge, since the compounds can easily hydrolyse in water solutions. That said, investigation in this area may be used to shed some light about the nature of the interactions that may partially account for the biological activity observed in physiologically relevant aqueous environments, albeit the results obtained should be interpreted cautiously. Our previous results showed that titanocenes **1a** or **1b** did not exhibit strong interactions with plasmid DNA by electrophoretic mobility shift assays, but the absence of a shift in the electrophoretic bands did not allow us to rule out DNA binding. Having established the interesting antitumor properties of metal compounds **1a**, **1a'**, **1b** and **1b'**, our aim with the study presented now was to further analyse and compare the kind of potential interactions of the enantiomers with DNA, by using other techniques to complement previous studies.

1 Dialysis experiments, based on the fundamental thermodynamic principle of
2 equilibrium dialysis [78,91], were performed to determine apparent binding constants
3 between DNA and the metal compounds, following the protocol described by Chaires
4 [78] with some modifications. As the DNA targets, we selected CT DNA and a short
5 oligonucleotide duplex of known sequence (ds17, 17 bp).

6 Unfortunately, under the conditions employed, large dispersion data sets were obtained,
7 which prevented the precise determination of association constants between the
8 titanium(IV) compounds and DNA. This is likely to be a consequence of the hydrolysis
9 of these complexes in aqueous media. However, even if the results should be interpreted
10 with caution, a significant increase in compound concentration was invariably observed
11 in the dialysis bags of replicate experiments, suggesting effective DNA binding by metal
12 complexes **1a**, **1a'**, **1b**, **1b'** and/or their hydrolysis products.

13 With the purpose of determining the effect that these compounds may exert on the DNA
14 denaturing temperature, T_m , we used a variable-temperature (FRET-melting) assay. This
15 experiment requires little DNA consumption, allows the assessment of a wide range of
16 compound concentrations, can be adapted to a high-throughput fashion, and it has been
17 extensively used to determine the degree of thermal stabilization of different DNA
18 structures in the presence of potential ligands [92]. Thus, FRET experiments were used
19 to establish whether metal complexes **1a**, **1a'**, **1b** and **1b'** were able to thermally stabilize
20 duplex DNA structures.


21 In these experiments, a 10-bp oligonucleotide (F10T) labelled with two fluorophores,
22 FAM at its 5' end and TAMRA at the 3' end, was selected [93]. If the metal complex binds
23 to DNA affecting the stability of the helix, changes in the value of DNA T_m should be
24 expected. Stabilization of duplex DNA usually results in increased values of T_m .

1 Compounds **1a**, **1a'**, **1b** and **1b'** were analysed for their ability to affect duplex DNA
2 melting within the 1-10 μM concentration range. However, under these conditions, the
3 titanium(IV) derivatives were not able to produce a significant change in the DNA
4 melting temperature. Furthermore, none of the enantiomers of the precursor ligand,
5 **a**·HCl, **a'**·HCl, showed DNA stabilization (see Online Resource Fig. S18). These results
6 suggest that the compounds may interact with DNA in an external, mainly electrostatic
7 fashion or through partial recognition of the DNA grooves.

8 Finally, DNA viscometric titrations were carried out because viscosity measurements
9 can provide a simple way to discriminate between the different binding modes of potential
10 DNA ligands (such as intercalation *versus* groove or external binding) [94]. According to
11 the theory of Cohen and Eisenberg [95], from gradual titration of DNA solutions with the
12 compounds of interest, linear plots of the cubed root of the relative DNA viscosity
13 $(\eta/\eta_0)^{1/3}$ versus the molar ratio of bound ligand to DNA nucleotide (r) can be obtained.
14 The slope values in these plots correlate well with the DNA-ligand binding modes.
15 Groove binding compounds normally display a slope close to 0.0, whereas classical
16 mono-intercalants result in a slope close to 1.0 [94,95].

17 The tested compounds showed a linear $(\eta/\eta_0)^{1/3}$ versus r correlation in the typical r
18 range used in these experiments (Fig. 5). Complexes **1a**, **1a'**, **1b** and **1b'**, irrespective of
19 the amino-bound ligand and the stereochemistry of the metal complex, gave rise to slope
20 values practically equal to zero.

21



1

2 **Fig. 5** Viscometric titrations of CT DNA and metal complexes **1a**, **1a'**, **1b** and **1b'** at 25
 3 °C (10 mM sodium phosphate buffer, pH 7.2)

4

5 These results are in good agreement with the FRET DNA melting assays and point
 6 towards an external or groove interaction of the titanium metal complexes and/or their
 7 hydrolysis products that does not result in overall changes of contour length or thermal
 8 stabilization of the DNA double helix structure.

9

10 **4. Conclusions**

11 Optically active amino-oxime ligands derived from natural products are useful and
 12 inexpensive starting materials for the design of enantiopure titanocene compounds. In
 13 contrast with the resistance to hydrolysis of other $\kappa^2\text{N,O}$ oximato-Ti biscyclopentadiene
 14 compounds described before, our systems suffer hydrolysis in water at physiological
 15 conditions, most likely due to the monohapto κON coordination mode of the highly

1 sterically demanding limonene residue of the oximato ligand. Regarding their cytotoxic
2 behaviour, the oxime-containing Ti(IV) compound **1b** has shown potent anticancer
3 activities against both prostate cancer PC-3 and DU-145 cell lines after 72 h of incubation
4 time. The cytotoxicity of the enantiomers **1a**, **1a'** and **1b'**, **1b'** towards all the cancer cell
5 lines tested showed no significant differences, exception made for the PC-3 cells. In
6 addition, isomers **1a** and **1b** showed certain selectivity in their toxicity against prostate
7 cancer PC-3 versus non-tumorigenic RWPE-1 cells. Furthermore, compound **1b** shows
8 higher activity than the additive dose of TDC and proligand **b·HCl** on the prostate PC-3,
9 DU-145 and RWPE-1 cell lines. These results point towards the existence of an influence
10 of the oximato-Ti unit on the hydrolysis process and/or the cytotoxicity mechanism.
11 Compound-DNA interactions have been investigated by equilibrium dialysis, FRET
12 melting assays and viscometric titrations. The experimental results suggest that these
13 metal complexes and/or their hydrolysis products can bind DNA either in the minor
14 groove or externally, irrespective of the ligand stereochemistry.

1

2 **Acknowledgments**

3 Financial support from Ministerio de Economía y Competitividad (MICINN CTQ2014-
4 58270-R), Comunidad Autónoma de Madrid (CAM, I3 Program) and the Universidad de
5 Alcalá (UAH, Projects CCG2016/EXP-044, CCG2016/EXP-028 and UAH-AE-2017-2)
6 is acknowledged. I.C.A. and S.S. are grateful to UAH for their FPI-UAH fellowships.

7 **Online Resource.** Supplementary data associated with this article can be found in the
8 online version, at <http://>. These data include: Representative NMR, UV-Vis and CD
9 spectra of compounds **a**, **a'**, **b**, **b'**, **1a**, **1a'**, **1b**, **1b'**. Selected biological data. Selected
10 crystallographic data and bond lengths and angles for X-ray molecular structures of **1b**.

11

- 1 [1] S. Medici, M. Peana, V. M. Nurchi, J. I. Lachowicz, G. Crisponi and M. A.
2 Zoroddu, *Coord. Chem. Rev.* 284 (2015) 329-350. DOI:10.1016/j.ccr.2014.08.002.
- 3 [2] G. Palermo, A. Magistrato, T. Riedel, T. von Erlach, C. A. Davey, P. J. Dyson and
4 U. Rothlisberger, *ChemMedChem* 11 (2016) 1199-1210. DOI:10.1002/cmdc.201500478.
- 5 [3] B. S. Murray, M. V. Babak, C. G. Hartinger and P. J. Dyson, *Coord. Chem. Rev.*
6 306 (2016) 86-114. DOI:10.1016/j.ccr.2015.06.014.
- 7 [4] T. C. Johnstone, K. Suntharalingam and S. J. Lippard, *Chem. Rev.* 116 (2016)
8 3436-3486. DOI:10.1021/acs.chemrev.5b00597.
- 9 [5] A. Casini and M. Contel, *J. Inorg. Biochem.* 165 (2016) 54-55.
10 DOI:10.1016/j.jinorgbio.2016.11.029.
- 11 [6] P. Y. Zhang and P. J. Sadler, *J. Organomet. Chem.* 839 (2017) 5-14.
12 DOI:10.1016/j.jorganchem.2017.03.038.
- 13 [7] E. Alessio, *Eur. J. Inorg. Chem.* (2017) 1549-1560. DOI:10.1002/ejic.201600986.
- 14 [8] H. Kopf and P. Kopfmaier, *Angew. Chem.-Int. Edit. Engl.* 18 (1979) 477-478.
15 DOI:10.1002/anie.197904771.
- 16 [9] P. Kopfmaier, S. Grabowski and H. Kopf, *Eur. J. Med. Chem.* 19 (1984) 347-352.
- 17 [10] H. J. Keller, B. Keppler and D. Schmahl, *Arzneimittelforschung* 32-2 (1982) 806-
18 807.
- 19 [11] B. K. Keppler and D. Schmahl, *Arzneimittelforschung* 36-2 (1986) 1822-1828.
- 20 [12] P. Kopfmaier, B. Hesse, R. Voigtlander and H. Kopf, *J. Cancer Res. Clin. Oncol.*
21 97 (1980) 31-39. DOI:10.1007/bf00411276.

- 1 [13] B. K. Keppler, C. Friesen, H. G. Moritz, H. Vongerichten and E. Vogel, *Struct.*
2 *Bond.* 78 (1991) 97-127.
- 3 [14] B. K. Keppler, *Metal Complexes in Cancer Chemotherapy*, Wiley VCH,
4 Weinheim, (1993).
- 5 [15] E. Y. Tshuva and D. Peri, *Coord. Chem. Rev.* 253 (2009) 2098-2115.
6 DOI:10.1016/j.ccr.2008.11.015.
- 7 [16] K. M. Buettner and A. M. Valentine, *Chem. Rev.* 112 (2012) 1863-1881.
8 DOI:10.1021/cr1002886.
- 9 [17] S. A. Loza-Rosas, M. Saxena, Y. Delgado, K. Gaur, M. Pandrala and A. D.
10 Tinoco, *Metallomics* 9 (2017) 346-356. DOI:10.1039/c6mt00223d.
- 11 [18] H. Skoupilova, R. Hrstka and M. Bartosik, *Med. Chem.* 13 (2017) 334-344.
12 DOI:10.2174/1573406412666161228113650.
- 13 [19] M. Cini, T. D. Bradshaw and S. Woodward, *Chem. Soc. Rev.* 46 (2017) 1040-
14 1051. DOI:10.1039/c6cs00860g.
- 15 [20] J. Schur, C. M. Manna, A. Deally, R. W. Koster, M. Tacke, E. Y. Tshuva and I.
16 Ott, *Chem. Commun.* 49 (2013) 4785-4787. DOI:10.1039/c3cc38604j.
- 17 [21] M. Grutzke, T. K. Zhao, T. A. Immel and T. Huhn, *Inorg. Chem.* 54 (2015) 6697-
18 6706. DOI:10.1021/acs.inorgchem.5b00690.
- 19 [22] M. Cini, H. Williams, M. W. Fay, M. S. Searle, S. Woodward and T. D. Bradshaw,
20 *Metallomics* 8 (2016) 286-297. DOI:10.1039/c5mt00297d.

- 1 [23] R. M. Lord, J. J. Mannion, B. D. Crossley, A. J. Hebden, M. W. McMullon, J.
2 Fisher, R. M. Phillips and P. C. McGowan, *ChemistrySelect* 1 (2016) 6598-6605.
3 DOI:10.1002/slct.201601290.
- 4 [24] Y. F. Mui, J. Fernandez-Gallardo, B. T. Elie, A. Gubran, I. Maluenda, M. Sanau,
5 O. Navarro and M. Contel, *Organometallics* 35 (2016) 1218-1227.
6 DOI:10.1021/acs.organomet.6b00051.
- 7 [25] S. Meker, O. Braitbard, M. D. Hall, J. Hochman and E. Y. Tshuva, *Chem.-Eur. J.*
8 22 (2016) 9986-9995. DOI:10.1002/chem.201601389.
- 9 [26] Y. Ellahioui, S. Prashar and S. Gomez-Ruiz, *Inorganics* 5 (2017) 4-27.
10 DOI:10.3390/inorganics5010004.
- 11 [27] K. E. Jones, K. L. Batchler, C. Zalouk and A. M. Valentine, *Inorg. Chem.* 56
12 (2017) 1264-1272. DOI:10.1021/acs.inorgchem.6b02399.
- 13 [28] S. A. Loza-Rosas, A. M. Vazquez-Salgado, K. I. Rivero, L. J. Negron, Y.
14 Delgado, J. A. Benjamin-Rivera, A. L. Vazquez-Maldonado, T. B. Parks, C. Munet-
15 Colon and A. D. Tinoco, *Inorg. Chem.* 56 (2017) 7788-7802.
16 DOI:10.1021/acs.inorgchem.7b00542.
- 17 [29] N. Ganot and E. Y. Tshuva, *RSC Adv.* 8 (2018) 5822-5827.
18 DOI:10.1039/c8ra00229k.
- 19 [30] E. Francotte and W. Lindner, *Chirality in Drug Research*, Wiley VCH, Weinheim,
20 (2006). DOI:10.1002/cmdc.200700060.
- 21 [31] M. J. Romero and P. J. Sadler in *Chirality in Organometallic Anticancer*
22 *Complexes*, Bioorganometallic Chemistry Applications in Drug Discovery, Biocatalysis

1 and Imaging, Eds.: G. Jaouen and M. Salmain, Wiley CH-VCH Verlag GmbH, (2015)
2 pp. 85-115. DOI:10.1002/9783527673438.ch03.

3 [32] F. Arnesano, A. Pannunzio, M. Coluccia and G. Natile, *Coord. Chem. Rev.* 284
4 (2015) 286-297. DOI:10.1016/j.ccr.2014.07.016.

5 [33] S. Y. Bi, A. D. Wang, C. F. Bi, Y. H. Fan, Y. Xiao, S. B. Liu and Q. Wang, *Inorg.*
6 *Chem. Commun.* 15 (2012) 167-171. DOI:10.1016/j.inoche.2011.10.016.

7 [34] S. Blanck, Y. Geisselbrecht, K. Kraling, S. Middel, T. Mietke, K. Harms, L. O.
8 Essen and E. Meggers, *Dalton Trans.* 41 (2012) 9337-9348. DOI:10.1039/c2dt30940h.

9 [35] D. Csokas, B. I. Karolyi, S. Bosze, I. Szabo, G. Bati, L. Drahos and A. Csampai,
10 *J. Organomet. Chem.* 750 (2014) 41-48. DOI:10.1016/j.jorganchem.2013.10.057.

11 [36] A. Dobrova, S. Platzer, F. Bacher, M. N. M. Milunovic, A. Dobrov, G. Spengler,
12 E. A. Enyedy, G. Novitchi and V. B. Arion, *Dalton Trans.* 45 (2016) 13427-13439.
13 DOI:10.1039/c6dt02784a.

14 [37] M. Frik, J. Fernandez-Gallardo, O. Gonzalo, V. Mangas-Sanjuan, M. Gonzalez-
15 Alvarez, A. S. del Valle, C. H. Hu, I. Gonzalez-Alvarez, M. Bermejo, I. Marzo and M.
16 Contel, *J. Med. Chem.* 58 (2015) 5825-5841. DOI:10.1021/acs.jmedchem.5b00427.

17 [38] Y. Fu, A. Habtemariam, A. Basri, D. Braddick, G. J. Clarkson and P. J. Sadler,
18 *Dalton Trans.* 40 (2011) 10553-10562. DOI:10.1039/c1dt10937e.

19 [39] Y. Fu, M. J. Romero, A. Habtemariam, M. E. Snowden, L. J. Song, G. J. Clarkson,
20 B. Qamar, A. M. Pizarro, P. R. Unwin and P. J. Sadler, *Chem. Sci.* 3 (2012) 2485-2494.
21 DOI:10.1039/c2sc20220d.

- 1 [40] W. Ginzinger, G. Muhlgassner, V. B. Arion, M. A. Jakupec, A. Roller, M.
2 Galanski, M. Reithofer, W. Berger and B. K. Keppler, *J. Med. Chem.* 55 (2012) 3398-
3 3413. DOI:10.1021/jm3000906.
- 4 [41] H. Glasner and E. Y. Tshuva, *Inorg. Chem.* 53 (2014) 3170-3176.
5 DOI:10.1021/ic500001j.
- 6 [42] A. Kurzwernhart, W. Kandioller, C. Bartel, S. Bachler, R. Trondl, G.
7 Muhlgassner, M. A. Jakupec, V. B. Arion, D. Marko, B. K. Keppler and C. G. Hartinger,
8 *Chem. Commun.* 48 (2012) 4839-4841. DOI:10.1039/c2cc31040f.
- 9 [43] M. G. Mendoza-Ferri, C. G. Hartinger, R. E. Eichinger, N. Stolyarova, K. Severin,
10 M. A. Jakupec, A. A. Nazarov and B. K. Keppler, *Organometallics* 27 (2008) 2405-2407.
11 DOI:10.1021/om800207t.
- 12 [44] S. Newcombe, M. Bobin, A. Shrikhande, C. Gallop, Y. Pace, H. Yong, R. Gates,
13 S. Chaudhuri, M. Roe, E. Hoffmann and E. M. E. Viseux, *Org. Biomol. Chem.* 11 (2013)
14 3255-3260. DOI:10.1039/c3ob27460h.
- 15 [45] S. Tabassum, A. Asim, R. A. Khan, F. Arjmand, D. Rajakumar, P. Balaji and M.
16 A. Akbarsha, *RSC Adv.* 5 (2015) 47439-47450. DOI:10.1039/c5ra07333b.
- 17 [46] S. F. Xi, L. Y. Bao, J. G. Lin, Q. Z. Liu, L. Qiu, F. L. Zhang, Y. X. Wang, Z. D.
18 Ding, K. Li and Z. G. Gu, *Chem. Commun.* 52 (2016) 10261-10264.
19 DOI:10.1039/c6cc05743h.
- 20 [47] K. S. M. Smalley, R. Contractor, N. K. Haass, A. N. Kulp, G. E. Atilla-Gokcumen,
21 D. S. Williams, H. Bregman, K. T. Flaherty, M. S. Soengas, E. Meggers and M. Herlyn,
22 *Cancer Res.* 67 (2007) 209-217. DOI:10.1158/0008-5472.can-06-1538.

- 1 [48] J. Maksimoska, L. Feng, K. Harms, C. L. Yi, J. Kissil, R. Marmorstein and E.
2 Meggers, *J. Am. Chem. Soc.* *130* (2008) 15764-15765. DOI:10.1021/ja805555a.
- 3 [49] S. A. Abramkin, U. Jungwirth, S. M. Valiahdi, C. Dworak, L. Habala, K. Meelich,
4 W. Berger, M. A. Jakupec, C. G. Hartinger, A. A. Nazarov, M. Galanski and B. K.
5 Keppler, *J. Med. Chem.* *53* (2010) 7356-7364. DOI:10.1021/jm100953c.
- 6 [50] C. M. Manna and E. Y. Tshuva, *Dalton Trans.* *39* (2010) 1182-1184.
7 DOI:10.1039/b920786b.
- 8 [51] C. M. Manna, G. Armony and E. Y. Tshuva, *Chem.-Eur. J.* *17* (2011) 14094-
9 14103. DOI:10.1002/chem.201102017.
- 10 [52] S. Blanck, J. Maksimoska, J. Baumeister, K. Harms, R. Marmorstein and E.
11 Meggers, *Angew. Chem.-Int. Edit.* *51* (2012) 5244-5246. DOI:10.1002/anie.201108865.
- 12 [53] Y. Fu, R. Soni, M. J. Romero, A. M. Pizarro, L. Salassa, G. J. Clarkson, J. M.
13 Hearn, A. Habtemariam, M. Wills and P. J. Sadler, *Chem.-Eur. J.* *19* (2013) 15199-15209.
14 DOI:10.1002/chem.201302183.
- 15 [54] K. J. Kilpin, S. M. Cammack, C. M. Clavel and P. J. Dyson, *Dalton Trans.* *42*
16 (2013) 2008-2014. DOI:10.1039/c2dt32333h.
- 17 [55] E. Menendez-Pedregal, J. Diez, A. Manteca, J. Sanchez, A. C. Bento, R. Garcia-
18 Navas, F. Mollinedo, M. P. Gamasa and E. Lastra, *Dalton Trans.* *42* (2013) 13955-13967.
19 DOI:10.1039/c3dt51160j.
- 20 [56] M. Miller and E. Y. Tshuva, *Eur. J. Inorg. Chem.* *2014* (2014) 1485-1491.
21 DOI:10.1002/ejic.201301463.

- 1 [57] Z. F. Chen, Q. P. Qin, J. L. Qin, J. Zhou, Y. L. Li, N. Li, Y. C. Liu and H. Liang,
2 *J. Med. Chem.* 58 (2015) 4771-4789. DOI:10.1021/acs.jmedchem.5b00444.
- 3 [58] M. Cini, T. D. Bradshaw, S. Woodward and W. Lewis, *Angew. Chem.-Int. Ed.* 54
4 (2015) 14179-14182. DOI:DOI: 10.1002/anie.201508034.
- 5 [59] X. Q. Zhou, Q. Sun, L. Jiang, S. T. Li, W. Gu, J. L. Tian, X. Liu and S. P. Yan,
6 *Dalton Trans.* 44 (2015) 9516-9527. DOI:10.1039/c5dt00931f.
- 7 [60] C. M. Manna, G. Armony and E. Y. Tshuva, *Inorg. Chem.* 50 (2011) 10284-
8 10291. DOI:10.1021/ic201340m.
- 9 [61] I. de la Cueva-Alique, S. Sierra, L. Munoz-Moreno, A. Perez-Redondo, A. M.
10 Bajo, I. Marzo, L. Gude, T. Cuenca and E. Royo, *J. Inorg. Biochem.* 183 (2018) 32-42.
11 DOI:10.1016/j.jinorgbio.2018.02.018.
- 12 [62] G. D. Potter, M. C. Baird and S. P. C. Cole, *Inorg. Chim. Acta* 364 (2010) 16-22.
13 DOI:10.1016/j.ica.2010.05.020.
- 14 [63] M. S. Ibn El Alami, M. A. El Amrani, A. Dahdouh, P. Roussel, I. Suisse and A.
15 Mortreux, *Chirality* 24 (2012) 675-682. DOI:10.1002/chir.22073.
- 16 [64] S. V. Larionov, *Russ. J. Coord. Chem.* 38 (2012) 1-23; and references therein.
17 DOI:10.1134/s1070328412010058.
- 18 [65] D. J. Brecknell, R. M. Carman, B. Singaram and J. Verghese, *Aust. J. Chem.* 30
19 (1977) 195-203. DOI:10.1071/ch9770195.
- 20 [66] U. Thewalt and R. Friedrich, *Z.Naturforsch.(B)* 46 (1991) 475-482.
21 DOI:10.1515/znb-1991-0409.

- 1 [67] O. P. Pandey, S. K. Sengupta and C. M. Tripathi, *Molecules* 10 (2005) 653-658.
2 DOI:10.3390/10060653.
- 3 [68] M. Carvalho, A. M. Galvao, J. Kredatusova, J. Merna, P. F. Pinheiro and M. M.
4 Salema, *Inorg. Chim. Acta* 383 (2012) 244-249. DOI:10.1016/j.ica.2011.11.019.
- 5 [69] I. de la Cueva-Alique, L. Munoz-Moreno, Y. Benabdelouahab, B. T. Elie, M. A.
6 El Amrani, M. E. G. Mosquera, M. Contel, A. M. Bajo, T. Cuenca and E. Royo, *J. Inorg.*
7 *Biochem.* 156 (2016) 22-34. DOI:10.1016/j.jinorgbio.2015.12.002.
- 8 [70] A. V. Tkachev, A. V. Rukavishnikov, A. M. Chibiryaev, A. Y. Denisov, Y. V.
9 Gatilov and I. Y. Bagryanskaya, *Aust. J. Chem.* 45 (1992) 1077-1086.
10 DOI:10.1071/CH9921077.
- 11 [71] M. Fernandez-Millan, M. Temprado, J. Cano, T. Cuenca and M. E. G. Mosquera,
12 *Dalton Trans.* 45 (2016) 10514-10518. DOI:10.1039/c6dt02116f.
- 13 [72] N. J. Sweeney, O. Mendoza, H. Muller-Bunz, C. Pampillon, F. J. K. Rehmann, K.
14 Strohfeltdt and M. Tacke, *J. Organomet. Chem.* 690 (2005) 4537-4544.
15 DOI:10.1016/j.jorganchem.2005.06.039.
- 16 [73] R. M. Carman, P. C. Mathew, G. N. Saraswathi, B. Singaram and J. Verghese,
17 *Aust. J. Chem.* 30 (1977) 1323-1335. DOI:10.1071/CH9771323.
- 18 [74] G. E. Tranter in *Spectroscopic Analysis: Polarimetry and Optical Rotatory*
19 *Dispersion, Vol. 8*, Comprehensive Chirality, Eds.: E. M. Carreira and H. Yamamoto,
20 Elsevier, (2012) pp. 411-421. DOI:10.1016/B978-0-08-095167-6.00843-0
- 21 [75] L. J. Farrugia, *J. Appl. Crystallogr.* 45 (2012) 849-854.
22 DOI:10.1107/s0021889812029111.

- 1 [76] G. M. Sheldrick, *Acta Crystallogr. Sect. A* 71 (2015) 3-8.
2 DOI:10.1107/s2053273314026370.
- 3 [77] G. M. Sheldrick, *Acta Crystallogr. Sect. C-Struct. Chem.* 71 (2015) 3-8.
4 DOI:10.1107/s2053229614024218.
- 5 [78] J. B. Chaires in *Structural selectivity of drug-nucleic acid interactions probed by*
6 *competition dialysis, Vol. 253, DNA Binders and Related Subjects*, Eds.: M. J. Waring
7 and J. B. Chaires, Springer-Verlag Berlin, Berlin, (2005) pp. 33-53.
8 DOI:10.1007/b100441.
- 9 [79] M. G. Davidson, A. L. Johnson, M. D. Jones, M. D. Lunn and M. F. Mahon,
10 *Polyhedron* 26 (2007) 975-980. DOI:10.1016/j.poly.2006.09.055.
- 11 [80] S. O. Baumann, M. Bendova, H. Fric, M. Puchberger, C. Visinescu and U.
12 Schubert, *Eur. J. Inorg. Chem.* (2009) 3333-3340. DOI:10.1002/ejic.200900381.
- 13 [81] S. O. Baumann, M. Bendova, M. Puchberger and U. Schubert, *Eur. J. Inorg.*
14 *Chem.* (2011) 573-580. DOI:10.1002/ejic.201000881.
- 15 [82] A. Chaudhary, V. Dhayal, M. Nagar, R. Bohra, S. M. Mobin and P. Mathur,
16 *Polyhedron* 30 (2011) 821-831. DOI:10.1016/j.poly.2010.12.025.
- 17 [83] J. H. Toney and T. J. Marks, *J. Am. Chem. Soc.* 107 (1985) 947-953.
18 DOI:10.1021/ja00290a033.
- 19 [84] A. Erxleben, J. Claffey and M. Tacke, *J. Inorg. Biochem.* 104 (2010) 390-396.
20 DOI:10.1016/j.jinorgbio.2009.11.010.

- 1 [85] C. M. Dowling, J. Claffey, S. Cuffe, I. Fichtner, C. Pampillon, N. J. Sweeney, K.
2 Strohfeltdt, R. W. G. Watson and M. Tacke, *Lett. Drug Des. Discov.* 5 (2008) 141-144.
3 DOI:10.2174/157018008783928463.
- 4 [86] R. A. Hilger, D. Alex, A. Deally, B. Gleeson and M. Tacke, *Lett. Drug Des.*
5 *Discov.* 8 (2011) 904-910.
- 6 [87] A. Obeid, A. El-Shekeil, S. Al-Aghbari and J. Al-Shabi, *J. Coord. Chem.* 65
7 (2012) 2762-2770. DOI:10.1080/00958972.2012.703780.
- 8 [88] J. Fernandez-Gallardo, B. T. Elie, T. Sadhukha, S. Prabha, M. Sanau, S. A.
9 Rotenberg, J. W. Ramos and M. Contel, *Chem. Sci.* 6 (2015) 5269-5283.
10 DOI:10.1039/c5sc01753j.
- 11 [89] J. Fernandez-Gallardo, B. T. Elie, F. J. Sulzmaier, M. Sanau, J. W. Ramos and M.
12 Contel, *Organometallics* 33 (2014) 6669-6681. DOI:10.1021/om500965k.
- 13 [90] J. F. Gonzalez-Pantoja, M. Stern, A. A. Jarzecki, E. Royo, E. Robles-Escajeda, A.
14 Varela-Ramirez, R. J. Aguilera and M. Contel, 50 (2011) 11099-11110.
15 DOI:10.1021/ic201647h.
- 16 [91] W. Muller and D. M. Crothers, *Eur. J. Biochem.* 54 (1975) 267-277.
17 DOI:10.1111/j.1432-1033.1975.tb04137.x.
- 18 [92] D. Renciuik, J. Zhou, L. Beaurepaire, A. Guedin, A. Bourdoncle and J. L. Mergny,
19 *Methods* 57 (2012) 122-128. DOI:10.1016/j.ymeth.2012.03.020.
- 20 [93] R. KIELTYKA, P. Englebienne, J. Fakhoury, C. Autexier, N. Moitessier and H. F.
21 Sleiman, *J. Am. Chem. Soc.* 130 (2008) 10040-10041. DOI:10.1021/ja8014023.

1 [94] D. Suh and J. B. Chaires, *Bioorg. Med. Chem.* 3 (1995) 723-728.

2 DOI:10.1016/0968-0896(95)00053-j.

3 [95] G. Cohen and H. Eisenberg, *Biopolymers* 8 (1969) 45-55.

4 DOI:10.1002/bip.1969.360080105.

5

6

7