

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

School of Medicine Publications and Presentations

School of Medicine

1-2023

Quality and Reliability of Spanish-language YouTube Videos on Breast Cancer

Fernando C. Diaz

Michael LaPelusa

Hansapani Rodrigo

The University of Texas Rio Grande Valley

Ismael Polo Perez

The University of Texas Rio Grande Valley, Ismael.poloperez@utrgv.edu

Dailis S. Corria Cedeno

See next page for additional authors

Follow this and additional works at: https://scholarworks.utrgv.edu/som_pub



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Diaz, F., LaPelusa, M., Rodrigo, H., Perez, I., Cedeno, D., & Kwang, H. (2023). Quality and Reliability of Spanish-language YouTube Videos on Breast Cancer. *Oncology & Haematology*, 19, 2. <https://doi.org/10.17925/OHR.2023.19.2.2>

This Article is brought to you for free and open access by the School of Medicine at ScholarWorks @ UTRGV. It has been accepted for inclusion in School of Medicine Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

Authors

Fernando C. Diaz, Michael LaPelusa, Hansapani Rodrigo, Ismael Polo Perez, Dailis S. Corria Cedeno, and Henry Kwang

Quality and Reliability of Spanish-language YouTube Videos on Breast Cancer

Fernando C Diaz,¹ Michael B LaPelusa,² Hansapani Rodrigo,³ Ismael Polo Perez,⁴ Dailis B Corria Cedeno¹ and Henry T Kwang⁴

1. Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; 2. Division of Cancer Medicine, MD Anderson Cancer Center, Houston, TX, USA; 3. School of Mathematical and Statistical Sciences, University of Texas Rio Grande Valley, Edinburg, TX, USA; 4. Department of Internal Medicine, University of Texas Rio Grande Valley School of Medicine, Harlingen, TX, USA

DOI: <https://doi.org/10.17925/OHR.2023.19.2.2>

As the number of Spanish-speaking patients with cancer in the USA grows, it is vital that reliable online breast cancer information is available in Spanish. The main objective of our study was to determine the quality and reliability of online videos in Spanish regarding breast cancer. We conducted a YouTube search using the phrase "cancer de mama" (breast cancer) and graded their quality and reliability using the Journal of the American Medical Association (JAMA) Benchmark Criteria, Global Quality Score (GQS), and the Modified DISCERN Questionnaire. Videos were characterized by descriptive variables. Videos were primarily from Spain (36.4%) and Mexico (17.9%). 'professional' videos had the highest JAMA mean rank (99.53), the highest GQS mean rank (102.86) and the second-highest DISCERN (101.10) mean rank. 'Personal' videos, by contrast, had the lowest mean rank values for GQS (72.28) and DISCERN (68.83) scores. A significant majority of online breast cancer videos in Spanish were of poor quality and reliability. 'Professional' videos are of higher quality and reliability than videos of other typologies; however, they have a limited range of content. In addition, a small percentage of videos were uploaded from the USA.

Keywords

Breast cancer, education, medical education, patient advocacy, YouTube, Spanish

Disclosures: Fernando C Diaz, Michael B LaPelusa, Hansapani Rodrigo, Ismael Polo Perez, Dailis B Corria Cedeno and Henry T Kwang have no financial or non-financial relationships or activities to declare in relation to this article.

Review process: Double-blind peer review.

Compliance with ethics: This study was performed in accordance with the Helsinki Declaration of 1964, and its later amendments. Ethics approval was not required as this study was granted exemption from institutional IRB oversight as it did not meet HHS or FDA definition for human subjects research. This article does not contain any identifying information about participants, therefore consent was not sought from participants for publication.

Data availability: The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Authorship: The named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval for the version to be published.

Access: This article is freely accessible at touchONCOLOGY.com © Touch Medical Media 2023.

Received: 30 May 2023

Accepted: 14 August 2023

Citation: *touchREVIEWS in Oncology & Haematology*. 2023;19(2):Online ahead of journal publication

Corresponding author: Fernando C Diaz, Hematology-Oncology Fellow, University of North Carolina at Chapel Hill, 101 Manning Dr., Chapel Hill, 27707, NC, USA. E: fernando.diaz.md@gmail.com

Support: No funding was received in the publication of this article.

Breast cancer is the leading cause of cancer death in Hispanic women in the USA. According to the American Cancer Society, as many as one in nine Hispanic women will develop invasive breast cancer during their lifetime and one in 49 Hispanic women will die from invasive breast cancer.¹ Female breast cancer is also the most common and leading cause of cancer death in Latin American and Caribbean countries, with an estimated 114,900 new diagnoses and 37,000 deaths each year.^{2,3}

Patients who receive a cancer diagnosis are often faced with difficult decisions regarding their treatment and prognosis and thus they rely on multiple sources of information to make an informed decision. The internet is an increasingly important source of health-related information for patients in the USA and worldwide. According to a National Health Interview Study survey from 32,139 adults, approximately 43% reported searching the internet for health information.⁴ Fox and colleagues found that 83% of Hispanic adults living in the USA obtained health information from media sources, with 79% of those acting on the health-related information obtained from these sources.⁵

YouTube, a global repository of videos and social networking platforms, is a popular way for people to access health-related information, with over a billion hours viewed every day.⁶ According to a study conducted by the Pew Research Center, 72% of adults online use some form of video-sharing platform, of whom 63% use YouTube as a source of health-related information.⁷ A variety of online outlets – including advertisers, government agencies, news agencies, patient health portals, patients and health professionals – produce health-related videos and upload them to YouTube. However, several studies have questioned the quality and reliability of health-related information on YouTube.⁸⁻¹⁰ A study published in 2018 evaluating breast self-examination videos on YouTube found 54 out of 87 videos misleading.¹¹

Objectives

While previous studies have evaluated the quality and reliability of online information in English, no studies have assessed the quality and reliability of YouTube videos in Spanish containing health-related information pertaining to breast cancer.¹¹⁻¹⁶ For Spanish-speaking-only patients, who face language barriers in receiving health information, high-quality, reliable online health-related information that is available in Spanish is essential.^{17,18} Our main objective was to determine the quality and reliability of online videos in Spanish regarding breast cancer.

Table 1: Description of video typologies

Typology	Description
Commercial	Videos that buy, sell or offer a service for a fee
Government/news agency	Videos created and/or managed by an official government body
Health portal	Videos created by an organization that offers health information on a variety of health topics
Personal	Videos created by individuals, patients and families of patients
Professional	Video created by health professionals, experts and professional organizations

Methods

We conducted a descriptive analysis of Youtube videos using the search terms "*cancer de mama*" (translation: "breast cancer") to investigate what patients would find when they are seeking information via YouTube. We chose these words over other options because this is the most broadly used set of words to describe breast cancer in Spanish. Search history, cookies and caches were cleared from the personal computers of the investigators to avoid any possible influence from prior browsing history. We created a new YouTube account in Harlingen, Texas. The first 200 video results from uniform resource locators of the search engine results page were transferred onto a playlist to avoid daily variances in search results using the default search filters. All inactive videos, duplicates, non-Spanish duplicates, or those with audio only, or without accompanying audio were excluded. Out of the initial 200 videos, a total of 173 were included in the study for further analysis. Twenty-seven videos (15%) were excluded because they had no audio (n=5), no relevant information (n=8), were in a non-Spanish language (n=10), in multiple parts (n=3) or were duplicates (n=1).

All videos were viewed in their entirety, with information collected from each video, including the number of 'likes', 'dislikes', total number of views, country of origin, video length and video upload date. Videos were categorized into one of five typology groups according to the source of upload: 'commercial' (for-profit companies), 'government/news agencies', 'health portal' (health information website), 'personal' (patient or family), and 'Professional' (university, hospital, physician groups or non-profit physician) (Table 1). The content of each video was assessed for the following: signs and symptoms, screening and detection, risk factors, diagnosis, research, treatment, reconstruction, survivorship/advocacy, and male breast cancer.

Video quality and reliability were assessed using three validated tools. The Journal of the American Medical Association (JAMA) Benchmark Criteria utilizes four core standards to evaluate online health information and identify the essential components of transparency and reliability: authorship, attribution of references, currency and disclosure of any potential conflict of interest. A point is given for the presence of each criterion, with a minimum score of 0 and a maximum score of 4.⁴ The Global Quality Score (GQS) is a subjective tool graded on a 5-point Likert scale, with a score of 1 representing poor quality and not being beneficial for patients, and a score of 5 representing excellent quality and very useful for patients.¹⁹ The modified 5-point DISCERN score is a questionnaire used to evaluate the quality of information with a maximum potential score of 5 points indicating high quality and a minimum of 0 points indicating poor quality (Table 2). In summary, the JAMA score evaluates information reliability, while the GQS and DISCERN scores assess information quality. Combining the scores provides a broader portrait of the quality and reliability of the video, as a whole. Scores were generated by two internal medicine resident physicians who were fluent in English and Spanish (IL and DC) independently at different times, who also selected the videos

from the original search. The corresponding Cohen's kappa coefficients for GQS and DISCERN scores were 0.923 and 0.960, respectively. A third physician (FD) resolved any disagreements on scores.

The ordinal scales were used to measure the video quality and reliability scores. Since these measurements are in an ordinal scale, we summarize them using median and interquartile ranges (IQR). All numerical measures are summarized as median \pm IQR and all qualitative variables were summarized as counts (percentages), as the best summary measures for categorical variables are percentages. The qualitative variables were recorded as percentages (year video uploaded, length of video, country of origin and typology). The differences in the quality and reliability measures were evaluated by typology using the Kruskal-Wallis (KW) non-parametric test.²⁰ This is mainly due to having ordinal type measures (quality and reliability scores) for small sample sizes within certain typology categories, and hence not being able to satisfy the normality assumption to use the one-way analysis of variance.²¹ It should be noted that, in the KW test, we deal with either the median (when the data distribution has the same shapes) or mean rank (when the data distribution do not have similar shapes). In our case, the shape of the data distribution for JAMA, GQS and DISCERN scores, within typologies are not the same and hence we report the mean ranks. Bonferroni adjusted pairwise comparisons with Dunn test were used to identify the significant differences in the three scores, between each

Table 2: Evaluation tools for reliability and quality

JAMA score (1 point per question if present; maximum 4 points)
1. Authorship
2. Attribution of references
3. Currency
4. Disclosure
Global Quality Score (choose one)
1. Poor quality, poor flow, most information missing, not helpful for patients
2. Generally poor, some information given but of limited use to patients
3. Moderate quality, some important information is adequately discussed
4. Good quality good flow, most relevant information is covered, useful for patients
5. Excellent quality and excellent flow, very useful for patients
DISCERN score (1 point per question if answered yes; maximum 5 points)
1. Are the explanations given in the video clear and understandable?
2. Are useful reference sources given? (publication cited, from valid studies)
3. Is the information in the video balanced and neutral?
4. Are additional sources of information given from which the viewer can benefit?
5. Does the video evaluate areas that are controversial or uncertain?

JAMA = Journal of the American Medical Association.

Table 3: Video characteristics

Year video uploaded	Total N=173 N (%)
<2010	7 (4%)
2010	2 (1.2%)
2011	17 (9.8%)
2012	8 (4.6%)
2013	10 (5.8%)
2014	17 (9.8%)
2015	16 (9.2%)
2016	22 (12.7%)
2017	26 (15%)
2018	28 (16.2%)
2019	18 (10.4%)
2020	2 (1.2%)
Length of video	
Median	5.28 mins
Interquartile range	9.30
0–5 minutes	83 (48%)
5–10 minutes	33 (19%)
10–20 minutes	34 (20%)
20–30 minutes	15 (9%)
>30 min	6 (3%)
Number of views	
Median	23,539
Number of likes	
Median	159
Range	0–47,000
Interquartile range	468
Number of dislikes	
Median	10
Range	0–721
Interquartile range	38
Country of origin	
Argentina	10 (5.8%)
Brazil	1 (0.6%)
Chile	1 (0.6%)
Colombia	8 (4.6%)
Costa Rica	4 (2.3%)
Cuba	1 (0.6%)
Dominican Republic	2 (1.2%)
Guatemala	1 (0.6%)
Mexico	31 (17.9%)
Panama	1 (0.6%)
Paraguay	1 (0.6%)
Peru	12 (6.9%)
Spain	63 (36.4%)
USA	18 (10.4%)
Venezuela	4 (2.3%)
Unknown	15 (8.7%)
Typology	
Commercial	13 (7.5%)
Government/news agency	43 (24.9%)
Health portal	14 (8.1%)
Personal	63 (36.4%)
Professional	40 (23.1%)

mins = minutes; N = number.

pair of typology.²² To detect a significant association among qualitative variables within each typology group, either the Pearson's chi-square or Fisher's exact tests were utilized.

Linear regressions were used to evaluate the impact of variables (number of likes, dislikes, duration of videos and number of views) on the three quality and reliability scores after adjusting for the typology and the contents of the videos. If the assumptions of linear regression models were not satisfied, a corresponding generalized additive model (GAM) was used to evaluate the impact of each variable. GAM provide a general framework for extending a standard linear model by allowing non-linear functions of each of the variables. All tests were two-tailed, and a 0.05 significance level was used to assess the statistical significance. Data analysis was performed using R (V.4.1.2).

Results

Approximately 4% of videos were uploaded before 2010, 31% uploaded 2010–2014, and 63.5% uploaded 2015–2019. The median length of the videos was approximately 5 minutes, with an IQR of 9.30. Approximately 48% of videos lasted less than 5 minutes, 19% were between 5 and 10 minutes, 20% were between 10 and 20 minutes, and 12% lasted longer than 20 minutes. The videos in our sample had a median number of views of 23,539. The median number of 'likes' per video was 159, with a range of 0–47,000 (IQR: 468). The median number of 'dislikes' was 10, with a range of 0–721 (IQR: 38). Most videos were published in Spain (36.4%), followed by Mexico (17.9%) and the USA (10.4%). Of the remaining videos, 26.6% were uploaded from 12 other Latin American countries and, in 8.7%, the country of origin could not be identified (*Table 3*).


With regards to video typology, 63 videos (36.4%) were uploaded by patients or their family or friends ('personal'), 43 videos (24.9%) by a government or news agency ('government/news agency'), 40 videos (23.1%) by medical professionals ('professional'), 14 videos (8.1%) by a stand-alone health portal ('health portal'), and 13 videos (7.5%) by commercial companies ('commercial') (*Table 3*).

We next evaluated the distribution of video content based on their typology. Of the 'professional' videos, 50% focused on treatment modalities, while very few referenced breast reconstruction, survivorship and advocacy, or breast cancer in men. Videos that discussed signs and symptoms of breast cancer were mainly uploaded by 'commercial' typologies (54%), videos discussing screening and detection were uploaded primarily by 'government/news agency' (44%), and videos discussing risk factors were uploaded mostly by standalone 'health portals' (21%) (*Figure 1*).

The details of video characteristics were stratified by the source of upload as shown in *Table 4*. 'Personal' videos had the longest duration on average (8 minutes) while 'health portal' videos had the shortest average (2.8 minutes). No significant differences were found in the median time since upload among the different typology types (p-value: 0.053).

The overall median JAMA score of all videos was 2, with a range of 0 to 3 (IQR: 0.00), in which 82.7% scored between 2 and 4 possible points. Only 3.5% of videos achieved 3 points, and no videos reached the maximum score of 4 points. The overall median GQS score was 2.00, with a range of 1 to 4 (IQR: 1.00). Approximately 21% of videos scored 1 point, 37% scored 2 points, 38% scored 3 points, 4% achieved 4 points, and no videos obtained all 5 possible points. The overall median DISCERN score was 2 with a range of 0 to 5 points (IQR: 2.00). Approximately 8% scored 0 points, 15% scored 1 point, 39% scored 2 points, 14.5% scored

Figure 1: Content of videos organized by typology (%)



3 points, 2.9% scored 4 points and only one video (0.6%) scored all 5 points (Table 5).

The correlation among the three reliability scores and the length of videos, likes, dislikes and views were calculated overall, and by typology (Figure 2). Overall, the length of the videos had shown a significant positive correlation with the GQS ($r=0.28$) and DISCERN ($r=0.23$) scores.

Similar positive correlations for GQS and DISCERN scores were observed for length of videos within all typology types except for commercial videos ($r=-0.56$). Note that the JAMA score correlation was not calculated, as the scores were the same for all commercial videos.

The mean rank and median scores by the source of upload (typology) were analysed and are shown in Table 6. Notably, for the JAMA score,

Table 4: Characteristics organized by typology*

	Commercial	Government/news agency	Health portal	Personal	Professional	p-value
N (%)	13 (7.5)	43 (24.9)	14 (8.1)	63 (36.4)	40 (23.1)	
Video length (mins)	3.17 ± 1.0	7.21 ± 10.8	2.83 ± 4.2	8.00 ± 12.1	4.93 ± 8.1	0.003
Time since upload (days)	1,936.0 ± 2,087.0	1,291.0 ± 1,110.0	2,026.5 ± 2,399.0	986.0 ± 1,390.0	1,299.5 ± 2,078.0	0.053
Views	39,947.0 ± 173,472.0	22,808.0 ± 111,208.0	26,943.0 ± 86,607.0	36,489.0 ± 165,906.0	8,975.5 ± 25,680.0	0.011
Likes	272.0 ± 536.0	100.5 ± 339.0	56.5 ± 187.0	383.5 ± 1036.0	53.0 ± 179.0	0.000
Dislikes	16.0 ± 45.0	10.0 ± 29.0	10.0 ± 20.0	17.5 ± 97.0	3.0 ± 11.0	0.000

*Values based on Kruskal–Wallis test and presented as median ± interquartile range. mins = minutes; N = number.


Table 5: Summary characteristics for quality and reliability scores

	JAMA	GQS	DISCERN
Median ± IQR	2.0 ± 0.0	2.0 ± 1.0	2.0 ± 2.0
Range	0–3	1–4	0–5
Points	Score N=173 (%)		
0	2 (1.2%)	N/A	49 (28.3%)
1	22 (12.7%)	36 (20.8%)	26 (15.0%)
2	143 (82.7%)	64 (37.0%)	67 (38.7%)
3	6 (3.5%)	66 (38.2%)	25 (14.5%)
4	0 (0.0%)	7 (4.0%)	5 (2.9%)
5	N/A	0 (0.0%)	1 (0.6%)

GQS = Global Quality Score; IQR = interquartile range; JAMA = Journal of the American Medical Association; N/A = not applicable.

the mean rank for 'professional' videos was highest among typologies at 99.5 (standard deviation [SD]: 24.3) compared with 73.6 (SD: 37.3) for 'personal' and 71.0 (SD: 50.3), for 'health portal' the lowest. Concerning the GQS score, the mean rank for 'professional' was at 102.9 (SD: 40.7), 98.5 (SD: 44.9) for 'health portal' (second highest), and 72.3 (SD: 50.8) for 'personal', the lowest. Concerning DISCERN score, 'health portal' had the highest mean rank at 110.1 (SD: 48.9) followed by 101.1 (SD: 46.5) for 'professional' while 'personal' was lowest at 68.9 (SD: 48.2). Table 6 depicts the significant mean rank differences for JAMA, GQS, and DISCERN scores as identified by pairwise comparisons. We observe that the mean ranks of three quality and reliability scores between Professional and Personal to be significantly different with p-values: JAMA:0.001; GQS:0.013; and DISCERN: 0.009, respectively. Additionally, we found that the mean rank for JAMA between 'government/news agency' and 'personal' (p-value: 0.002) and the mean rank for DISCERN

Figure 2: Correlation matrices by quality and reliability tools and video characteristics



The correlation matrices among the variables of length of videos, likes, dislikes, views, Global Quality Score (GQS) and Journal of the American Medical Association (JAMA) Benchmark Criteria, and the Modified DISCERN Questionnaire scores for: (a) full set of videos; (b) commercial; (c) government/news agency; (d) health portal; (e) personal videos; and (f) = professional. Overall, length of the videos had shown a significant positive correlation with the GQS ($r=0.28$) and DISCERN ($r=0.23$) scores. Similar positive correlations for GQS and DISCERN scores were observed for length of videos within all typology types except for commercial videos ($r=-0.56$). Note that the JAMA score correlation was not calculated, as the scores were the same for all commercial videos.

between 'health portal' and 'personal' (p-value: 0.002) was significantly different.

As per GAM analyses, video length was the only variable that had shown a significant impact (p-value: 0.0002) on the DISCERN score, after adjusting for typology and the contents of the videos. None of the variables were associated with either the JAMA or GQS scores (Tables 7 and 8).

Discussion

Hispanic people are the largest minority group in the USA, and nearly one-third identify as foreign-born.²³ According to estimates from the United States Census Bureau, approximately 58 million Americans, or nearly 19% of the total population, identified as Hispanic or Latino in 2016, and this is expected to double over the next four decades.²⁴ Additionally, data from the North American Central Cancer Registries (NACCR) showed that Hispanic females were less likely than non-Hispanic whites to be diagnosed with localized stage breast cancer

Table 6: Summary characteristics for quality and reliability scores by typology*

	Commercial	Government/news agency	Health portal	Personal	Professional	Kruskal-Wallis p-value
JAMA (0-4 points)						
Median ± IQR	2.0 ± 0.0	2.00 ± 0.00	2.00 ± 1.00	2.00 ± 1.00	2.00 ± 0.00	-
Mean rank ± SD	96.0 ± 0.0	97.6 ± 20.6	71.0 ± 50.4	73.6 ± 37.3	99.5 ± 24.3	0.000
GQS (1-5 points)						
Median ± IQR	2.0 ± 1.0	2.0 ± 1.0	3.0 ± 1.0	2.0 ± 2.0	3.0 ± 1.0	-
Mean rank	89.6 ± 38.5	89.3 ± 45.6	98.5 ± 44.9	72.3 ± 50.8	102.9 ± 40.7	0.020
DISCERN (0-5 points)						
Median ± IQR	2.0 ± 2.0	2.0 ± 1.0	2.0 ± 2.0	1.0 ± 2.0	2.0 ± 2.0	-
Mean rank	79.5 ± 44.1	95.2 ± 41.2	110.1 ± 48.9	68.9 ± 48.2	101.1 ± 46.5	0.0002

*The assumption for one-way analysis of variance was not satisfied, hence the Kruskal-Wallis (KW) test was used. The mean ranks were used in the KW test, as the distribution of the data within each typology was not similar within each quality and reliability score measurement. GQS = Global Quality Score; IQR = interquartile range; JAMA = Journal of the American Medical Association; SD = standard deviation.

Table 7: Generalized additive models summaries for JAMA and DISCERN scores

GAM model for JAMA score							
ANOVA for parameter effects		DF	Sum Sq	Mean Sq	F value	Pr (>F)	Significance
Likes		1	0.028	0.028	0.173	0.6780	
Dislikes		1	0.000	0.000	0.000	0.9880	
Video length		1	0.342	0.342	2.147	0.1449	
Views		1	0.073	0.073	0.458	0.4996	
Typology		4	5.019	1.255	7.887	0.0000	***
Content type	D1	1	0.712	0.712	4.475	0.0360	*
	D2	1	1.267	1.267	7.965	0.0054	**
	D3	1	0.088	0.088	0.555	0.4574	
	D4	1	0.826	0.826	5.194	0.0241	*
	D5	1	0.000	0.000	0.000	0.9964	
	D6	1	0.265	0.265	1.667	0.1986	
	D7	1	0.133	0.133	0.836	0.3620	
	D8	1	0.088	0.088	0.554	0.4579	
	D9	1	0.755	0.755	4.744	0.0310	*
	D10	1	0.025	0.025	0.155	0.6941	
	D11	1	0.021	0.021	0.133	0.7160	
Residuals		151	24.020	0.159			
GAM for DISCERN score							
ANOVA for parameter effects		DF	Sum Sq	Mean Sq	F value	Pr(>F)	Significance
Likes		1	0.339	0.339	0.332	0.5651	
Dislikes		1	3.175	3.175	3.112	0.0798	
Video length		1	14.520	14.520	14.233	0.0002	***
Views		1	1.388	1.388	1.361	0.2453	
Typology		4	24.418	6.105	5.984	0.0002	***
Content type	D1	1	8.104	8.105	7.944	0.0055	**
	D2	1	8.693	8.693	8.521	0.0040	**
	D3	1	1.706	1.706	1.672	0.1979	
	D4	1	8.194	8.194	8.032	0.0052	**
	D5	1	2.682	2.682	2.629	0.1070	
	D6	1	0.356	0.356	0.349	0.5555	
	D7	1	0.086	0.086	0.084	0.7724	
	D8	1	0.030	0.030	0.030	0.8638	
	D9	1	4.795	4.795	4.700	0.0317	*
	D10	1	0.210	0.210	0.206	0.6507	
	D11	1	0.006	0.006	0.006	0.9385	
Residuals		151	154.045	1.020			

Significance codes: *** <0.001; ** <0.01; * <0.05

ANOVA = analysis of variance; GAM = generalized additive model; JAMA = Journal of the American Medical Association.

(57% versus 65%, respectively).²⁵ Another study also found that Hispanic women were less likely than non-Hispanic white women to present with stage I disease.²⁶ This difference has been hypothesized to be related to the fact that Hispanic people, on average, belong to communities of lower socioeconomic status, which tend to have inadequate access to care.²⁷ Individuals in lower socioeconomic groups are also less

Table 8: Linear regression model summary for Global Quality Score

Parameter effects		Estimate	Std. error	T value	Pr(> t)	Significance
Intercept		1.76	0.196	8.999	0.0000	***
Likes		0.00	0.000	0.470	0.6389	
Dislikes		0.00	0.001	1.371	0.1724	
Video length		0.01	0.006	1.795	0.0747	.
Views		0.00	0.000	0.660	0.5101	
Typology2		0.28	0.223	1.253	0.2121	
Typology3		0.06	0.258	0.217	0.8284	
Typology4		0.39	0.208	1.896	0.0599	.
Typology5		0.22	0.216	1.041	0.2997	
Content type	D1	0.46	0.121	3.780	0.0002	***
	D2	0.44	0.136	3.215	0.0016	**
	D3	0.09	0.161	0.534	0.5938	
	D4	0.42	0.160	2.660	0.0086	**
	D5	0.23	0.205	1.114	0.2671	
	D6	0.04	0.176	0.251	0.8024	
	D7	0.17	0.123	1.369	0.1729	
	D8	0.27	0.264	1.038	0.3009	
	D9	0.85	0.270	3.135	0.0021	**
	D10	0.19	0.114	1.668	0.0974	.
	D11	0.08	0.264	0.303	0.7622	
R ² : 0.4796						
Adjusted R ² : 0.4141						

Significance codes: *** <0.001; ** <0.01; * <0.05

likely to be insured. One study, using data from the National Cancer Database, found that uninsured and Medicaid-insured patients had higher proportions of distant stage cancer at presentation than those with private insurance, or Medicare plus supplemental insurance.²⁸ Data from the NACCR also show that the incidence of breast cancer among Hispanic women increased from 2006 to 2015, at an average rate of 0.4% per year. Other studies have also suggested that Hispanic women are less likely than non-Hispanic white women to receive appropriate and timely breast cancer treatment and are more likely to be diagnosed at an advanced stage.^{29,30} One multi-institutional analysis of breast cancer found, in addition to confirming the aforementioned differences in stage at presentation compared with non-Hispanic populations, that Hispanic women were more likely to present with triple-negative breast cancer (an aggressive subtype of breast cancer) and were less likely to receive breast-conserving surgery than non-Hispanic women.³¹

Interestingly, foreign-born Hispanic women experience lower rates of breast cancer compared with USA-born matched controls. This is thought to be related to reproductive factors such as younger age at first birth, higher parity, higher rates of breast cancer, and less use of hormone therapy for contraception and menopause, which confer a reduced risk of breast cancer.³² However, lower screening rates may also explain the decreased prevalence in Hispanic women in the USA compared with non-Hispanic white women in the USA and the reduced prevalence in foreign-born Hispanic women compared with USA-born Hispanic women.^{33,34}

Screening for breast cancer occurs less frequently in Hispanic women than in non-Hispanic white women.³⁵ Data from the National Health Interview Survey show that the prevalence of screening mammography

was 61% in Hispanic women, compared with 65% in non-Hispanic white women.

With over 2 billion users in over 100 countries, YouTube has become an increasingly important source of health information for patients and physicians.⁶ As the platform grows in popularity, so does the number of videos published on medical topics. Previous studies have assessed the quality of videos concerning breast self-examination, mammography and mastectomy in English, using the same tools we utilized.^{11,15,16} Other studies have also assessed the quality and reliability of information found in videos on various medical topics, from lumbar spinal fusion to food allergies.³⁶ As a company, YouTube employs community guidelines enforced by human reviewers and machine learning technology, in an attempt to “make YouTube a safer community” while still giving creators the freedom to share a broad range of experiences and perspectives.³⁷ YouTube’s community guidelines outline which type of content is kept off the platform and covers a broad range of topics, including spam and deceptive practices, sensitive content, violent or dangerous content, regulated goods, and misinformation.³⁷ Notably, YouTube’s community guidelines are fluid and continually changing in the midst of recent political trends and misinformation in the setting of the COVID-19 pandemic.

To our knowledge, this is the first study that analyses the quality, reliability and content of online videos relating to breast cancer exclusively in Spanish. On average, the videos examined in this study were viewed more than 128,000 times, demonstrating the popularity of the videos among the Spanish-speaking community. However, the unregulated nature of videos posted on this platform raises concerns about the quality and reliability of material available to the patient population. Our study demonstrates low JAMA, GQS and DISCERN scores for videos uploaded in Spanish. Notably, no video achieved the maximum possible scores for JAMA or GQS. When the scores of the videos were stratified according to typology, we found differences amongst upload sources to be statistically significant, with those uploaded by professional organizations having the highest mean rank scores. Professional videos had the highest mean rank for JAMA score and GQS, while the highest mean rank for DISCERN scores was recorded for 'health portal' videos. When stratifying the content of videos by typology, we found that roughly half of the videos uploaded by a 'professional' typology discussed treatment modalities and one-quarter of the videos spoke about the signs and symptoms found in individuals with breast cancer. The remaining content categories, including screening, risk factors, diagnosis, advocacy, breast cancer in men, disease pathophysiology, and types and stages, were addressed in only a small percentage of videos. However, this topic was found in higher frequency in other video typologies of lower quality. We also found that only approximately 10% of videos were uploaded from a source in the USA, with Spain and Mexico generating most of the videos.

Several limitations of our study exist. Firstly, our search strategy (“*cancer de mama*”) may not capture what patients search for on YouTube for

information. Secondly, we only examined 200 YouTube videos, despite thousands of videos resulting when our search strategy was employed. This likely excluded mostly videos classified as 'personal'. Other limitations of our study include the lack of data collection pertaining to the average number of views per video and the specific source of content provided. Lastly, two reviewers reviewed the videos. While there was only a minimal discrepancy in scoring between the two viewers, including additional viewers may have added to the validity of the scores given to each video.

Conclusion

Our study aimed to evaluate the quality and reliability of breast cancer videos produced in Spanish and found on YouTube, the most popular social media platform, with over 2 billion active users. We found that the overall quality and reliability of videos in Spanish were poor, and those of higher quality did not reliably cover some of the subjects. This is particularly important to federal agencies such as the National Institutes of Health and the National Cancer Institute, and academic institutions, where efforts have been made to recruit Hispanic patients with cancer and reduce disparities. The importance of social media platforms in the Hispanic community should not go unrecognized. Providing high-quality and reliable health information through videos targeting Hispanic people can be a helpful modality in strengthening the recruitment of these individuals to educate them on their illness so that they can ultimately make more informed decisions with their physician. Considering the growing Hispanic demographic in the USA and worldwide, hospitals and specialists should consider producing educational videos to increase the number of high-quality videos found online and dilute those promoting scientifically unsubstantiated information. A mechanism to achieve this may be via a peer-review process, which is employed in the publishing of patient information by Cancer.Net, the American Society of Clinical Oncology’s patient information website.³⁸

The lack of high-quality videos covering these topics from a reputable source is concerning. We believe false or misleading information in all subjects, especially those not broadly covered, contributes to public misperception and may result in ill-informed and potentially harmful decision-making. The importance of accurate and reliable information becomes especially relevant in socioeconomically disadvantaged groups with inadequate health literacy and who may rely heavily on video media for their medical information. For videos published by non-professional sources, the HONcode is a certification system that exists and could be utilized by YouTube. The HONcode is a non-governmental organization with a code of ethics that guides website managers on how best to disseminate reliable and high-quality health information and provides yearly assessments for websites seeking its certification.³⁹

Moving forward, major academic institutions and national organizations should conduct a comprehensive needs assessment to ensure accurate information is thoroughly presented for patients and their families. □

1. Miller KD, Ortiz AP, Pinheiro PS, et al. Cancer statistics for the US Hispanic/Latino population, 2021. *CA Cancer J Clin*. 2021;71:466–87. DOI: 10.3322/caac.21695.
2. de Lemos LLP, Carvalho de Souza M, Pena Moreira D, et al. Stage at diagnosis and stage-specific survival of breast cancer in Latin America and the Caribbean: A systematic review and meta-analysis. *PLoS One*. 2019;14:e0224012. DOI: 10.1371/journal.pone.0224012.
3. Justo N, Wilking N, Jönsson B, et al. A review of breast cancer care and outcomes in Latin America. *Oncologist*. 2013;18:248–56. DOI: 10.1634/theoncologist.2012-0373.

4. Silberg WM, Lundberg GD, Musacchio RA. Assessing, controlling, and assuring the quality of medical information on the internet: Caveant lector et viewor—let the reader and viewer beware. *JAMA*. 1997;277:1244–5.
5. Fox S, Rainie L. The online health care revolution: How the web helps Americans take better care of themselves. 1999. Available at: <https://api.semanticscholar.org/CorpusID:69027258> (Date last accessed: 13 October 2023).
6. Google. YouTube About. 2020. Available at: <https://youtube.com/about/press/> (Date last accessed: 17 May 2020).
7. Anderson M. 5 facts about online video, for YouTube’s 10th birthday. 2015. Available at: www.pewresearch.org/fact-tank/2015/02/12/5-facts-about-online-video-for-youtubes-10th-birthday/ (Date last accessed: 13 October 2023).

8. Briones R, Nan X, Madden K, Waks L. When vaccines go viral: An analysis of HPV vaccine coverage on YouTube. *Health Commun*. 2012;27:478–85. DOI: 10.1080/10410236.2011.610258.
9. Nguyen BT, Allen AJ. Social media and the intrauterine device: A YouTube content analysis. *BMJ Sex Reprod Health*. 2018;44:28–32. DOI: 10.1136/bmjshr-2017-101799.
10. Singh AG, Singh S, Singh PR. YouTube for information on rheumatoid arthritis—A wakeup call *J Rheumatol*. 2012;39:899–903. DOI: 10.3899/jrheum.111114.

11. Clarke Hillyer G, Basch CH, Guerro S, et al. YouTube videos as a source of information about mastectomy. *Breast J*. 2019;25:349–50. DOI: 10.1111/tbj.13222.
12. Arif N, Ghezzi P. Quality of online information on breast cancer treatment options. *Breast*. 2018;37:6–12. DOI: 10.1016/j.breast.2017.10.004.
13. Basch CH, Hillyer GC, MacDonald ZL, et al. Characteristics of YouTube(TM) videos related to mammography. *J Canc Educ*. 2015;30:699–703. DOI: 10.1007/s13187-014-0769-9.
14. Bruce JG, Tucholka JL, Steffens NM, Neuman HB. Quality of online information to support patient decision-making in breast cancer surgery. *J Surg Oncol*. 2015;112:575–80. DOI: 10.1002/jso.24046.
15. Esen E, Aslan M, Sonbahar BÇ, Kerimoğlu RS. YouTube English videos as a source of information on breast self-examination. *Breast Cancer Res Treat*. 2019;173:629–35. DOI: 10.1007/s10549-018-5044-z.
16. Tan MLH, Kok K, Ganesh V, Thomas SS. Patient information on breast reconstruction in the era of the world wide web. A snapshot analysis of information available on YouTube.com. *Breast*. 2014;23:33–7. DOI: 10.1016/j.breast.2013.10.003.
17. News From the Centers for Disease Control and Prevention. Access to health-care and preventive services among Hispanics and non-hispanics—United States, 2001–2002. *JAMA*. 2004;292:2331. DOI: 10.1001/jama.292.19.2331.
18. Mayberry RM, Milli F, Ofili E. Racial and ethnic differences in access to medical care. *Med Care Res Rev*. 2000;57 Suppl 1:108–45. DOI: 10.1177/1077558700057001S06.
19. Oremule B, Patel A, Orekoya O, et al. Quality and reliability of YouTube videos as a source of patient information on rhinoplasty. *JAMA Otolaryngol Head Neck Surg*. 2019;145:282–3. DOI: 10.1001/jamaoto.2018.3723.
20. Kruskal WH, Wallis WA. Use of ranks in one-criterion variance analysis. *J Am Stat Assoc*. 1952;47:583–621. DOI: 10.1080/01621459.1952.10483441.
21. McDonald JH. *Handbook of Biological Statistics*. University of Delaware: Sparky House Publishing, 2009.
22. Dinno A. Nonparametric pairwise multiple comparisons in independent groups using Dunn's test. *Stata J*. 2015;15:292–300. DOI: 10.1177/1536867X1501500117.
23. Ward Net al. Frequently requested statistics on immigrants and immigration in the United States. Available at: <https://reliefweb.int/report/united-states-america/frequently-requested-statistics-immigrants-and-immigration-united-states#:~:text=White%20the%20immigrant%20population%20has,of%2014.8%20percent%20in%201890> (Date last accessed: 6 August 2023).
24. Funk C, Lopez MH. A brief statistical portrait of U.S. Hispanics. Available at: www.pewresearch.org/science/2022/06/14/a-brief-statistical-portrait-of-u-s-hispanics/ (Date last accessed: 13 October 2023).
25. Miller KDet al. Cancer statistics for Hispanics/Latinos 2021. *Cancer J Clin*. 2021;71:466–87. DOI: 10.3322/caac.21695.
26. Iqbal J, Ginsburg O, Rochon PA, et al. Differences in breast cancer stage at diagnosis and cancer-specific survival by race and ethnicity in the United States. *JAMA*. 2015;313:165–73. DOI: 10.1001/jama.2014.17322.
27. Fang CY, Tseng M. Ethnic density and cancer: A review of the evidence. *Cancer*. 2018;124:1877–903. DOI: 10.1002/cncr.31177.
28. Ward EM, Fedewa SA, Cokkinides V, Virgo K. The association of insurance and stage at diagnosis among patients aged 55 to 74 years in the National cancer database. *Cancer J*. 2010;16:614–21. DOI: 10.1097/PP0.0b013e3181ff2aec.
29. Chen L, Li CI. Racial disparities in breast cancer diagnosis and treatment by hormone receptor and Her2 status. *Cancer Epidemiol Biomarkers Prev*. 2015;24:1666–72. DOI: 10.1158/1055-9965.EPI-15-0293.
30. Sineshaw HM, Gaudet M, Ward EM, et al. Association of race/ethnicity, socioeconomic status, and breast cancer subtypes in the National cancer data base (2010–2011). *Breast Cancer Res Treat*. 2014;145:753–63. DOI: 10.1007/s10549-014-2976-9.
31. Nahleh Z, Otoukesh S, Mirshahidi HR, et al. Disparities in breast cancer: A multi-institutional comparative analysis focusing on American hispanics. *Cancer Med*. 2018;7:2710–7. DOI: 10.1002/cam4.1509.
32. John EM, Phipps AJ, Davis A, Koo J. Migration history, acculturation, and breast cancer risk in Hispanic women. *Cancer Epidemiol Biomarkers Prev*. 2005;14:2905–13. DOI: 10.1158/1055-9965.EPI-05-0483.
33. Gangnon RE, Sprague BL, Stout NK, et al. The contribution of mammography screening to breast cancer incidence trends in the United States: An updated age–period–cohort model. *Cancer Epidemiol Biomarkers Prev*. 2015;24:905–12. DOI: 10.1158/1055-9965.EPI-14-1286.
34. Sauer AG, Siegel RL, Jemal A, Fedewa SA. Updated review of prevalence of major risk factors and use of screening tests for cancer in the United States. *Cancer Epidemiol Biomarkers Prev*. 2017;26:1192–208. DOI: 10.1158/1055-9965.EPI-17-0219.
35. National Center for Health, US. *Health, United States, in Health, United States, 2016: With Chartbook on Long-Term Trends in Health*. Hyattsville (MD): National Center for Health Statistics (US), 2017.
36. Ahuja K, Aggarwal P, Sareen JR, et al. Comprehensiveness and reliability of YouTube as an information portal for lumbar spinal fusion: A systematic review of Video content. *Int J Spine Surg*. 2021;15:179–85. DOI: 10.14444/8023.
37. YouTube. Community guidelines. 2022. Available at: www.youtube.com/howyoutubeworks/policies/community-guidelines/ (Date last accessed: 4 April 2022).
38. Oncology, A.S.o.C. Cancer.Net. 2022. Available at: www.cancer.net (Date last accessed: 5 April 2022).
39. Conduct, T.H.o.t.N.F.C.o. HONcode. 2019. Available at: <https://myhon.ch/HONcode/Conduct.html?HONConduct695800> (Date last accessed: 4 April 2022).