



This work has been submitted to **NECTAR**, the **Northampton Electronic Collection of Theses and Research**.

#### Article

**Title:** The macroecology of animal versus wind pollination: ecological factors are more important than historical climate stability

**Creators:** Rech, A. R., Dalsgaard, B., Sandel, B., Sonne, J., Svenning, J.-C., Holmes, N. and Ollerton, J.

**DOI:** [10.1080/17550874.2016.1207722](https://doi.org/10.1080/17550874.2016.1207722)

**Example citation:** Rech, A. R., Dalsgaard, B., Sandel, B., Sonne, J., Svenning, J.-C., Holmes, N. and Ollerton, J. (2016) The macroecology of animal versus wind pollination: ecological factors are more important than historical climate stability. *Plant Ecology & Diversity*. **9**(3), pp. 253-262. 1755-0874.

It is advisable to refer to the [publisher's version](#) if you intend to cite from this work.

**Version:** Accepted version

**Official URL:** <http://dx.doi.org/10.1080/17550874.2016.1207722>

<http://nectar.northampton.ac.uk/8928/>



1 NOTE: This is the manuscript version of the paper. Please refer to the final published version for  
2 the definitive text and findings.

3

4

5 Original Article

6 **The macroecology of animal versus wind pollination: The macroecology of animal versus**  
7 **wind pollination: ecological factors are more important than historical climate stability**

8 André Rodrigo Rech<sup>1</sup>, Bo Dalsgaard<sup>2</sup>, Brody Sandel<sup>3</sup>, Jesper Sonne<sup>2</sup>, Jens-Christian Svenning<sup>3</sup>,  
9 Naomi Holmes<sup>4</sup>, Jeff Ollerton<sup>4\*</sup>

10

11 <sup>1</sup>*Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, Minas Gerais,*  
12 *Brazil;* <sup>2</sup>*Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark,*  
13 *University of Copenhagen, Copenhagen, Denmark;* <sup>3</sup>*Section for Ecoinformatics and Biodiversity,*  
14 *Department of Bioscience, Aarhus University, Aarhus, Denmark;* <sup>4</sup>*Landscape and Biodiversity*  
15 *Research Group, School of Science and Technology, University of Northampton, Northampton, UK.*

16 Short running-title: The macroecology of pollination

17 \* Corresponding author. Email: Jeff.Ollerton@northampton.ac.uk

18

1 **Abstract**

2 **Background:** The relative frequency of wind and animal pollinated plants is non-randomly  
3 distributed across the globe and numerous hypotheses have been raised for the greater occurrence of  
4 wind pollination in some habitats and towards higher latitudes. To date, however, there has been no  
5 comprehensive global investigation of these hypotheses.

6 **Aims:** Investigating a range of hypotheses for the role of biotic and abiotic factors as determinants  
7 of the global variation in animal versus wind pollination.

8 **Methods:** We analysed 67 plant communities ranging from 70° North to 34° South. For these we  
9 determined habitat type, species richness, insularity, topographic heterogeneity, current climate and  
10 Late-Quaternary climate change. The predictive effects of these factors on the proportion of wind-  
11 and animal-pollinated plants were tested using correlations, ordinary least squares (OLS) and  
12 logistic regression analyses with information-theoretic model selection.

13 **Results:** The proportion of animal-pollinated plant species was positively associated with plant  
14 species richness and current temperature. Furthermore, in forest, animal pollination was positively  
15 related to precipitation. Historical climate was only weakly and idiosyncratically correlated with  
16 animal pollination.

17 **Conclusion:** Results were consistent with the hypothesised reduced chance for wind-transported  
18 pollen reaching conspecific flowers in species-rich communities, fewer constraints on nectar  
19 production in warm and wet habitats, and reduced relative effectiveness of wind dispersal in humid  
20 areas. There was little evidence of a legacy of historical climate change affecting these patterns.

21 **Keywords:** abiotic, biotic, community ecology, forest, mutualism, open vegetation, pollen  
22 dispersal, precipitation, species richness, temperature.

23

24

## 1 **Introduction**

2 Pollination is a fundamental ecological process that is crucial for the functioning of most terrestrial  
3 ecosystems (Kearns et al. 1998; Ollerton et al., 2011; Lever et al., 2014). Plants achieve pollen  
4 transfer among conspecifics using biotic (animal) and abiotic (wind and, rarely, water) pollen  
5 vectors. Wind pollination occurs in about 20% of angiosperm families and most groups of  
6 gymnosperms (Ackerman 2000), an estimated 12.5% of all angiosperm species are wind pollinated  
7 (Ollerton et al. 2011). Wind pollination is a derived condition that has evolved independently more  
8 than 60 times within otherwise animal-pollinated clades (Ackerman 2000; Friedman and Barrett  
9 2009; Hu et al. 2012), with occasional reversions to biotic pollination (e.g. Wragg and Johnson  
10 2011).

11 Wind pollination is currently understood to be a reproductive strategy that evolves when  
12 pollinators become unpredictable or unavailable (Barrett 1996; Culley et al. 2002; Friedman and  
13 Barrett 2009). Supporting this idea, the proportion of wind-pollinated plants is higher in the  
14 temperate zone (Culley et al. 2002; Ollerton et al. 2011), where climate is less stable seasonally  
15 (Cramer et al. 1999) and over longer glacial–interglacial time scales (Sandel et al. 2011). In  
16 temperate regions usually < 80% of the plant species within a community are animal pollinated,  
17 whereas animal pollination is more common in the tropics where on average  $\geq 90$  % of all co-  
18 occurring plants are pollinated by animals (Whitehead 1969; Regal 1982; Ollerton et al. 2011).  
19 Although this latitudinal pattern is already known, no study has quantitatively assessed the possible  
20 factors related to the global variation of community-level differences in wind *versus* animal  
21 pollination (Barrett 1996; Schemske et al. 2009; Ollerton et al. 2011).

22 The efficiency of wind and animal pollination may depend on the biotic and abiotic  
23 environment where plants occur. A number of testable hypotheses for geographic variation in wind  
24 *versus* animal pollination have been put forward (Whitehead 1969; Culley et al. 2002). For instance,

1 extensive, dense and species-rich forests with high canopy and low wind speeds are supposed to be  
2 unfavourable places for wind pollination because the chance of wind transporting pollen to a  
3 conspecific should be rather low (Whitehead 1969; Regal 1982; Ollerton et al. 2006, 2011; Culley  
4 et al. 2002). Climate may also affect the efficiency of wind pollination. High humidity should make  
5 the cytoplasm of the pollen grains to collapse or become heavier and cause them to clump, reducing  
6 pollen dispersal distance (Whitehead 1969; Niklas 1985). Additionally, high temperature and  
7 precipitation may increase productivity and support more nectar production for animal pollination.

8         Thus, both mean annual temperature and precipitation should correlate positively with the  
9 proportion of animal pollination. On the other hand, high temperature and precipitation seasonality  
10 may promote wind pollination as pollinator abundances will fluctuate through the year, becoming  
11 less reliable pollen vectors (Regal 1982), unless pollination is in synchrony with these seasonal  
12 fluctuations in pollinator abundances, thereby causing no measurable effect on pollination mode  
13 (McKinney et al. 2012). Finally, in topographically flat environments such as coastal plains wind  
14 pollination could be favoured due to more constant and fast winds. Likewise wind pollination could  
15 be more frequent on islands because wind-pollinated plants are independent of pollinator  
16 colonisation to maintain reproduction (Kühn et al. 2006; Crawford et al. 2011).

17         Paleoclimate has been linked to contemporary ecological patterns, such as species  
18 distributions and endemism (Svenning and Skov 2007; Cárdenas et al. 2011; Sandel et al. 2011;  
19 Kissling et al. 2012), and the structure of interactions within communities of plants and pollinators  
20 (Dalsgaard et al., 2011, 2013; Groom et al., 2014). Past climatic instability may have broken up  
21 mutualistic associations between plant species and their pollinating animals (Dalsgaard et al., 2011,  
22 2013, McKinney *et al.*, 2012) that may have favoured wind pollination and could be related to the  
23 contemporary composition of plant communities.

1           Here, we use a global dataset to describe (1) latitudinal patterns in pollination mode, and (2)  
2 the relationships between putative contemporary and historical factors, and the proportion of animal  
3 *versus* wind pollinated species in plant communities around the world. Specifically, following on  
4 from the discussion of abiotic and biotic factors that other authors have presented, we test the  
5 following hypotheses:

6           1. Wind pollination is less frequent in closed forest habitats compared to more open habitats,  
7 such as grasslands.

8           2. Animal pollination is more frequent in areas with higher mean annual temperature and  
9 precipitation.

10          3. Conversely, wind pollination is more frequent in regions with lower temperature and  
11 greater seasonality of precipitation.

12          4. Animal pollination is more frequent in communities with high plant species richness.

13          5. Wind pollination is more frequent in topographically flat environments.

14          6. Animal pollination is less frequent on islands.

15          7. Wind pollination is more frequent in those areas of the world that have experienced  
16 greater climatic instability during the Quaternary.

17

## 18 **Materials and methods**

### 19 *Animal versus wind pollination datasets*

20 The dataset was taken from 82 published and unpublished community-based assessments of the  
21 proportion of wind and animal pollinated species. These studies are censuses of all native  
22 angiosperm species in flower in a given community at a specific time. Gymnosperms were not  
23 considered by most of the published studies, hence we excluded gymnosperms from all  
24 communities. The latitudes ranged from 34° S to 70° N (Figure 1). When the same geographic

1 coordinates were attributed to communities within the same vegetation type by the original authors,  
2 we pooled them together and used the mean of pollination mode proportion as a descriptor of the  
3 proportion of animal pollination for those coordinates. Doing this, we reduced the dataset to 67 data  
4 points for the spatial analysis (Appendix Table S1).

5           In some of the published community studies no species list was available, just data on  
6 the proportions of wind and animal pollinated species observed, preventing us from attributing  
7 genera or families to the data analysed by the original authors. For those studies having a species  
8 list, pollination mode at the species level was highly correlated with higher taxa, i.e. genera and  
9 families (Table 1). This does not mean that species within the same genus have always the same  
10 pollination mode. It implies that proportions are constant across taxonomic levels and therefore any  
11 level may be used in the analysis. For the published studies we followed the information provided  
12 by the original authors as to the pollination mode (wind or animal) of a given species. For our own  
13 field observations, flowers were assessed in terms of presence or absence of: mass pollen release  
14 when an inflorescence was shaken, reduced corolla, and feathery stigmas (wind pollination); or  
15 large and colourful flowers with scent or nectar, and the presence and behaviour of potential flower  
16 visitors (animal pollination) (Figure 2; see also Table 1 in Friedman and Barrett 2009). We paid  
17 particular attention to species with small, dull flowers, e.g. some tropical trees, where only the  
18 presence of scent, nectar, and/or insect visitation distinguished biotic from abiotic pollination.

19           The proportion of animal-pollinated plant species could be underestimated because some  
20 plant species that possess wind pollination traits can also be visited and sometimes pollinated by  
21 animals (Figueredo and Sazima 2000). There might also be cases of obligatory self-pollinating and  
22 non-sexually reproducing species coded as wind or animal pollinated, though this should be a  
23 relatively minor and not spatially structured bias and, hence, should not affect our results (Ollerton  
24 et al. 2011). Similarly to Ollerton et al. (2011), we took a community-level approach as we were

1 interested in the ecological question of how large a proportion of plant species in terrestrial  
2 communities are animal or wind pollinated, and the possible factors related to these proportions.  
3 However, first, we measured the phylogenetic signal for pollination mode by means of Phylogenetic  
4 Eigenvector Regression (PVR) considering the species lists where plants were identified up to the  
5 species level (Diniz-Filho et al. 2012). Species lists were available for 56 plant communities,  
6 including 1689 genera. Phylogenetic information was extracted at genus level from Phylomatic  
7 (R20120829 stored tree, Webb and Donoghue 2005) and the tree generated by using Phylocom  
8 (Webb et al. 2008). PVR was calculated by using a principal coordinate analysis (PCoA) of the  
9 phylogenetic distance matrix and selecting the 50 first axes. Phylogenetic signal was obtained by  
10 the logistic regression coefficient ( $R^2$ ) between the selected phylogenetic eigenvectors and the  
11 binomial pollination mode (Diniz-Filho et al. 2012). Phylogenetic signal significance was tested by  
12 measuring PVR on 1000 randomisations of the pollination mode in the phylogeny. The analysis was  
13 carried out by using the package PVR for R (Santos et al. 2013). Since phylogenetic signal of  
14 pollination mode was very low (PVR for 50 eigenvectors = 0.0411;  $P = 0.47$ ;  $DF = 50$ ; Deviance =  
15 50.2) no taxonomic correction was required. The low phylogenetic signal probably came out of  
16 many independent origins of wind pollination and, due to many genera having one pollination mode  
17 also showing a minority of species possessing the other pollination mode.

18

### 19 *Correlates of pollination mode*

20 For each site, we extracted variables of contemporary and historical climate conditions  
21 hypothesised to affect the degree of wind *versus* animal pollination. Current climate descriptors  
22 included mean annual temperature (MAT), mean annual precipitation (MAP), temperature  
23 seasonality (MAT seasonality) and precipitation seasonality (MAP seasonality) extracted at 2.5 arc-  
24 minute resolution (approximately 16 km<sup>2</sup>) from the WorldClim dataset ([www.worldclim.org](http://www.worldclim.org);



1 Hijmans et al. 2005). As different climate change metrics may capture different effects of climate  
2 change (Garcia et al. 2014), we considered two measures of historical climate stability: (1) the  
3 velocity of mean annual temperature (MAT velocity) and that of mean annual precipitation (MAP  
4 velocity) between the Last Glacial Maximum (LGM) and the present; and (2) the anomaly of mean  
5 annual temperature (MAT anomaly) and mean annual precipitation (MAP anomaly) between the  
6 LGM and the present (current minus LGM precipitation/temperature – i.e. positive values indicate  
7 wetter or warmer contemporary conditions, respectively). Velocities reflect the speed of climate  
8 change irrespective of the direction, integrating macroclimatic changes and local spatial topoclimate  
9 gradients (Loarie et al. 2009; Sandel et al. 2011), whereas anomalies reflect the direction of climate  
10 change. Paleoclimate estimates were obtained from the CCSM3 model (Collins et al. 2006; Otto-  
11 Bliesner et al. 2006), and statistically downscaled to 2.5 arc-minute resolution to match the current  
12 climate data (Hijmans et al. 2005).

13 In addition, we determined potentially important non-climate descriptors of local  
14 conditions at each site, including whether the site was on the mainland (coded as 0) or on an island  
15 (1), topographic heterogeneity, regional proportion of tree cover, plant species richness, and  
16 whether the vegetation was forest (0; n = 16) or open vegetation type (1; grassland, scrubland, and  
17 other low and open vegetation types, n = 51). Topographic heterogeneity was calculated as the  
18 range of elevation values observed in a  $0.2^\circ \times 0.2^\circ$  window (ca. 20 km  $\times$  20 km) centered on the  
19 sampling location, using the WorldClim 1-km DEM (Farr et al. 2007). The regional proportion of  
20 tree cover was calculated as the average tree cover observed within a window of 64 km  $\times$  64 km  
21 around each site, based on the Vegetation Continuous Fields product applied to MODIS data  
22 (Hansen et al. 2003).

23

24 *Statistical analyses*

1           For all analyses, plant species richness, MAT anomaly, precipitation and temperature  
2 velocities were  $\log_{10}$ -transformed; contemporary mean annual precipitation and topography were  
3 square-root transformed; the proportion of regional tree cover was arcsine square-root transformed.  
4 The proportion of animal pollinated plant species in each community was arcsine square-root  
5 transformed for ordinary least squares (OLS) regression, but was left untransformed for logistic  
6 regression. All other variables were left untransformed.

7           First, the latitudinal pattern in pollination mode was tested with a quadratic regression  
8 between latitude and the proportion of animal pollination. Second, for all vegetation types, and  
9 separately for open vegetation types and forest, we tested the univariate associations between all  
10 predictor variables and between the proportion of animal pollinated plant species and all predictor  
11 variables (Appendix Table S2). This identified predictor variables that were strongly inter-  
12 correlated. For all further analyses, we excluded temperature seasonality as it was strongly  
13 correlated with mean annual temperature ( $r = -0.73$ ,  $P < 0.001$ ,  $n = 67$ ), and we excluded the  
14 proportion of regional tree cover as it was unavailable for three island datasets and strongly  
15 correlated with contemporary precipitation ( $r = 0.73$ ,  $P < 0.001$ ,  $n = 64$ ; see Appendix Table S2).  
16 Due to the strong positive relationship between precipitation and the proportion of regional tree  
17 cover, we adjusted our above hypothesis of a unidirectional positive effect of precipitation on  
18 animal pollination. For forest, we still expected a positive correlation between precipitation and the  
19 proportion of animal pollination. For open vegetation types, on the other hand, we expected a less  
20 strong (or even zero) effect of precipitation due to the possible opposite effects of increasing  
21 precipitation (favouring animal pollination) and associated increasing fragmentation of open  
22 vegetation communities surrounded by increasing amounts of forests as precipitation and regional  
23 tree cover coincide (increased fragmentation favouring wind pollination).

1           To test the study hypotheses, we used ordinary least squares (OLS) and logistic  
2 regression, conducting information-theoretic model selection, as outlined by Diniz-Filho et al.  
3 (2008). In doing so, we tested the effect of both measures of historical climate stability, i.e.  
4 temperature/precipitation velocity and anomaly; the anomaly models generally performed the best  
5 (highest  $R^2$  and lowest AIC), so we focus the discussion on these models (but we also report models  
6 including velocity). First, we fitted models with all combinations of the explanatory variables:  
7 forest versus open vegetation, insularity, plant species richness, contemporary precipitation,  
8 contemporary temperature, precipitation seasonality, topography and historical climate stability as  
9 measured by temperature and precipitation anomaly. Furthermore, due to the possible different  
10 effect of precipitation in open and forest vegetation, we included an interaction term between  
11 precipitation and forest versus open vegetation. We then identified minimum adequate models  
12 (MAMs) among these as any model with  $\Delta AIC_c < 2$  ( $n = 67$ , 1023 alternative models; Table 2).  
13 When using temperature and precipitation velocity as a measure of climate stability, we excluded  
14 topography as it is strongly correlated with velocities (Appendix S2), i.e. when using velocity as our  
15 historical climate variable we compared 511 models. We were unable to estimate velocities for one  
16 island data point, therefore in models using velocities there is one data point fewer.

17           Standardised regression coefficients are reported for both a multi-model average  
18 regression model based on weighted  $w_i$  and the MAMs (Diniz-Filho et al. 2008). We tested whether  
19 significant ( $P < 0.05$ ) positive spatial autocorrelation remained in OLS model residuals, tested using  
20 10 distance classes and applying a permutation test with 10,000 iterations. As no spatial  
21 autocorrelation was found in our anomaly models that performed the best, we did not use spatial  
22 models. All correlations and OLS regression analyses were conducted using Spatial Analysis in  
23 Macroecology 4.0 (Rangel et al. 2010). Logistic model selection analyses were carried out in R (R  
24 Development Core Team, 2014) using the 'MuMIn' package (Barton 2015).

1

## 2 **Results**

3 The proportion of animal-pollinated plants in a community decreased with latitude (Figure 1). This  
4 pattern appeared to be largely symmetrical across the Equator, though there was a geographical bias  
5 to the distribution of the studies, and more data from Africa and Asia, in particular, are needed to  
6 confirm the observed pattern.

7           The models to assess the relative importance of the factors tested as determinants of  
8 the proportion of animal pollination showed that overall the explanatory power was considerably  
9 higher for current climate variables (ca.  $32\% < R^2 < 36\%$ ) than for species richness (ca.  $4\% < R^2 <$   
10  $5\%$ ) and historical climate variables (ca.  $0\% < R^2 < 2\%$ ; Table 2 and Appendix S3). The proportion  
11 of animal pollination in both simple univariate correlations and in regression models was strongly  
12 and positively associated with current temperature and species richness (Figure 3; Table 2;  
13 Appendix S3). Furthermore, the interaction term between precipitation and vegetation type was  
14 included in the best-fit regression model (Table 2; Appendix S3), indicating that precipitation  
15 correlated strongly with the proportion of animal pollination in forest (Figure 3; Appendix S2).  
16 There were only weak associations with topography and temperature anomaly and velocity (Table  
17 2, Appendix S2). In addition to explaining little variation in pollination mode, topography and  
18 historical temperature anomaly/velocity associated positively with animal pollination in regression  
19 models (Table 2), but were unrelated in simple univariate correlations (Table S1). In logistic, but  
20 not in OLS, regression islands were found to have a lower proportion of animal pollination.  
21 Seasonality, precipitation anomaly, and precipitation velocity were not statistically significant in  
22 simple univariate correlations nor were they included in any MAMs (Table 2 and Appendix S3).

23

## 24 **Discussion**

1 The proportion of plant species using animal versus wind pollination in a given community is non-  
2 random with respect to latitude, animal pollination being especially dominant in the tropics (Figure  
3 1), confirming previous studies, but with a much larger and geographically widespread data set. We  
4 found that differences in the proportion of animal pollinated species were associated mainly with  
5 current climate (Figure 3). Notably, temperature was overall positively related to the prevalence of  
6 animal pollination (partly supporting hypothesis 2), and in forests animal pollination was more  
7 frequent in areas with higher precipitation (supporting both hypothesis 1 and, in part, hypotheses 2  
8 and 3). However, the ‘more wind pollination with greater precipitation seasonality’ aspect of  
9 hypothesis 3 was not supported, perhaps due to the different nature of seasonality in the lowland  
10 tropics (wet, dry) compared with the temperate or high altitude zones (cold and warm). In addition  
11 to current climate, we found that communities with greater local plant species richness have a  
12 higher proportion of animal pollination (supporting hypothesis 4). On the other hand, topography  
13 had little effect on frequency of pollination mode, not supporting hypothesis 5. Kühn et al. (2006)  
14 showed that wind pollination in Germany is more frequent in the flat landscapes of north relative to  
15 other parts of the country, although it increases also toward the southern Alpine uplands. This  
16 suggests that the effect of topography is scale- and context-dependent, which may explain its weak  
17 effect in our global analysis. Insularity also had only a minor influence on pollination-mode, with  
18 islands tending to have lower proportions of animal-pollination, but only when also taking climate  
19 into account (Table 2), which only partly supports hypothesis 6. In contrast to the strong  
20 associations with current factors (Table 2; Appendix S2), historical climate was associated only  
21 weakly and idiosyncratically with animal pollination, varying independently when using regression  
22 models or simple univariate correlations (Table 2 and Appendix S2), and therefore not supporting  
23 hypothesis 7. Overall, this suggests that the relative proportions of different pollination modes are  
24 more linked to current climate and ecological factors than to historical legacies.

1           Although these results support the greater influence of contemporary ecological  
2 factors over historical ones, it is not straightforward to determine the exact mechanism responsible  
3 for this pattern. For example, current precipitation was strongly related to the percentage of regional  
4 tree cover, hence, animal pollination is associated with both high precipitation and forest habitat.  
5 Wet forests clearly offer poor conditions for pollen dispersal, but it is difficult to disentangle  
6 whether this is because of mechanical restrictions imposed by dense vegetation or because air  
7 humidity decreases pollen dispersability, or both (Whitehead 1969; Niklas 1985). In open areas,  
8 especially in the tropics, the distribution of plant individuals tends to be sparse and floral resources  
9 less abundant. For example, an estimate of nectar energy available per hectare in the Atlantic rain  
10 forest in Brazil showed that forests produced twice as much floral energy as the nearby open  
11 ‘restinga’ (coastal scrub) vegetation (Fonseca 2013, Fonseca et al. 2015). Moreover, the distribution  
12 of an Andean plant species was associated with a progressive disruption of its animal pollination  
13 mutualism accompanied by the gradual decrease in precipitation (Chalcoff et al. 2012). This  
14 illustrates the potential role of vegetation structure and precipitation in determining the functioning  
15 of animal-dependent mutualisms, but disentangling the roles of correlated factors is a major  
16 challenge for macroecology.

17           Interestingly, current climate may play a similar role for seed dispersal as there is also  
18 a pattern of higher importance of animal dispersers in rain forests, whereas in dry forests wind and  
19 self-dispersal are more common (Howe and Smallwood 1982; Jordano 2000 and references  
20 therein). For instance, current precipitation patterns are largely correlated with proportion of  
21 endozoochory across the Atlantic forest of Brazil, animals being more important dispersers in the  
22 wetter areas (Almeida-Netto et al. 2008), although large-scale studies are not available to confirm  
23 this hypothesis. One tentative joint explanation for the lower frequency of animal-pollinated and  
24 animal-dispersed plants at drier sites could be the higher metabolic costs of producing nectar and

1 fleshy fruit, as was originally proposed only for fruit (Wilson et al. 1989; Almeida-Neto et al.  
2 2008). Associated with humidity, high temperature may also increase nectar production and  
3 therefore influence animal pollination, although, as temperature and temperature seasonality were  
4 found to be negatively correlated, we cannot rule out the possibility that seasonality promotes wind  
5 pollination (Regal 1982).

6           Another factor positively associated with animal pollination was plant species  
7 richness. Other trophic interactions between plants and animals respond to the bottom-up effect of  
8 plant species richness (Kissling et al. 2007; Scherber et al. 2010); for instance, in the Neotropics the  
9 diversity of pollinating birds and bats is associated with their food plant species richness (Fleming  
10 2005; Kissling et al. 2007). Meanwhile, as plant species richness increases, the density of  
11 conspecifics per unit area tends to decrease (Comita et al. 2010), which may also reduce the  
12 efficiency of pollen dispersion by wind (Whitehead 1969; Regal 1982). Plant species richness may  
13 also play an important role in sustaining pollinator communities (Ebeling et al. 2008; Dorado and  
14 Vázquez 2014) via two main mechanisms: (1) more plant species per unit area may ensure a more  
15 predictable and diverse food supply through complementarity (Waser and Real 1979; Rathcke 1983;  
16 Blüthgen and Klein 2011; Yang et al. 2013); and (2) more plant species also increases species  
17 redundancy within functional groups ('biodiversity insurance effect'), reducing the extinction risk  
18 of functionally specialised interactions (Bartomeus et al. 2013; Fründ et al. 2013). For instance,  
19 manipulative experiments have shown that high functional diversity of plants enhances the number  
20 of pollinators, and the functional diversity of pollinators increases individual plant fitness and  
21 community persistence over time (Fontaine et al. 2006; Albrecht et al. 2012).

22

## 23 **Conclusions**

1 We tested a range of previously proposed hypotheses and found that a subset of contemporary  
2 climate and ecological processes are the factors more strongly related to the global pattern of animal  
3 than to wind pollination at the community level. The deduced limited influence of historical climate  
4 instability on the contemporary prevalence of animal and wind pollination is in contrast with  
5 previous findings of high importance of historical factors on the structuring of plant-pollinator  
6 interaction networks (Dalsgaard et al., 2011, 2013; Martín González et al. 2015) and the distribution  
7 of animal and plant life on earth (Svenning and Skov 2007; Cárdenas et al. 2011; Sandel et al. 2011;  
8 Kissling et al. 2012), but is in accordance with studies of the organisation of plant-frugivore  
9 interaction networks (Schleuning et al. 2014). In future studies, it would be interesting to examine if  
10 other measures of historical processes, e.g. geological differences or historical measures of plant  
11 migration rates, influence plant reproduction. The large scale drivers of plant reproductive ecology  
12 will be better understood with similar studies examining also the role of both historical and  
13 contemporary factors on other processes, such as plant reproduction via autogamy and seed  
14 dispersal. In addition, to improve the certainties of the conclusions presented here will require  
15 additional sampling effort in Africa, Asia, Russia and oceanic islands, where so few data have been  
16 collected on plant reproductive ecology at the community level.

17

## 18 **Acknowledgements**

19 Special thanks to Dr. Andrew Moldenke, who shared with us his great personal community  
20 assessments dataset, and to all the researchers who helped us during field work, pollination mode  
21 confirmation, community assessments and/or with extra data to perform this study. ARR was  
22 supported by FAPESP (2009/54491-0); CAPES (PDSE 0197/12-2); CNPq (140912/2010-0);  
23 Unicamp (FAEPEX 519292) and Santander Universities. BD and JS acknowledge the Danish  
24 National Research Foundation for funding the Center for Macroecology, Evolution and Climate



1 (grant-number D NRF96). JCS was supported by the European Research Council (ERC-2012-StG-  
2 310886-HISTFUNC). JO was supported by FAPESP; The Royal Society; The Leverhulme Trust;  
3 The British Ecological Society; Church and Co. PLC; EVOLINK; The Percy Sladen Memorial  
4 Fund; The Royal Entomological Society and the University of Northampton. NH was supported by  
5 the British Exploring Society.

6

7

## 8 **References**

9

10 Ackerman JD. 2000. Abiotic pollen and pollination, ecological, functional, and evolutionary  
11 perspectives. *Plant Systematics and Evolution* 222: 167-185.

12 Albrech M, Schmid B, Hautier Y, Müller CB. 2012. Diverse pollinator communities enhance plant  
13 reproductive success. *Proceedings of the Royal Society B* 279: 4845-4852. Almeida-Neto

14 M, Campassi F, Galetti M, Jordano P, Oliveira-Filho A. 2008. Vertebrate dispersal  
15 syndromes along the Atlantic forest: broad-scale patterns and macroecological correlates.

16 *Global Ecology and Biogeography* 17: 503–513.

17 Barton K. 2015. MuMIn: Multi-Model Inference. R package version 1.13.4. Available at:  
18 <http://cran.r-project.org/package=MuMIn>.

19 Barrett SCH. 1996. The reproductive biology and genetics of island plants. *Philosophical*  
20 *Transactions of the Royal Society Ser. B.* 351: 725–733.

21 Bartomeus I, Park MG, Gibbs J, Danforth BN, Lakso AN, Winfree R. 2013. Biodiversity ensures  
22 plant–pollinator phenological synchrony against climate change. *Ecology Letters* 11: 1331–  
23 1338.

- 1 Blüthgen N, Klein AM. 2011. Functional complementarity and specialisation: Why biodiversity is  
2 important in plant-pollinator interactions. *Basic and Applied Ecology* 12: 282–291.
- 3 Cárdenas ML, Gosling WD, Sherlock SC, Poole I, Pennington RT, Mothes P. 2011. The response  
4 of vegetation on the Andean flank in western Amazonia to pleistocene climate change.  
5 *Science* 331: 1055-1058.
- 6 Chalcoff VR, Aizen MA, Ezcurra C. 2012. Erosion of a pollination mutualism along an  
7 environmental gradient in a south Andean treelet, *Embothrium coccineum* (Proteaceae).  
8 *Oikos* 121: 471-480.
- 9 Collins WD, Bitz CM, Blackman ML et al. 2006. The Community Climate System Model version 3  
10 (CCSM3). *Journal of Climate* 19: 2122-2143.
- 11 Comita LS, Muller-Landau HC, Aguilar S, Hubbell SP. 2010. Asymmetric density dependence  
12 shapes species abundances in a tropical tree community. *Science* 329: 330-332.
- 13 Cramer W, Kicklighter DW, Bondeau A, Iii BM, Churkina G, Nemry B, Ruimy A, Schloss AL.  
14 1999. Comparing global models of terrestrial net primary productivity (NPP): overview and  
15 key results. *Global Change Biology* 5: 1–15.
- 16 Crawford DJ, Anderson GJ, Bernardello G. 2011. The reproductive biology of Island plants. *The*  
17 *Biology of Island Floras* (ed. by D. Bramwell and J. Caujapé-Castells), pp. 11-36,  
18 Cambridge University Press.
- 19 Culley TM, Weller SG, Sakai AK. 2002. The evolution of wind pollination in angiosperms. *Trends*  
20 *in Ecology and Evolution* 17: 361-369.
- 21 Dalsgaard B, Magård E, Fjeldså J. et al. 2011. Specialization in plant-hummingbird networks is  
22 associated with species richness, contemporary precipitation and quaternary climate-change  
23 velocity. *PLoS One* 6: e25891.

- 1 Dalsgaard B, Trøjelsgaard K, González AMM. et al. 2013. Historical climate-change influences  
2 modularity and nestedness of pollination networks. *Ecography* 36: 1331-1340.
- 3 Diniz-Filho JAF, Bini LM, Rangel TF, Morales-Castilla I, Olalla-Tárraga MÁ, Rodríguez MÁ,  
4 Hawkins BA. 2012. On the selection of phylogenetic eigenvectors for ecological analyses.  
5 *Ecography* 35: 239.
- 6 Diniz-Filho JAF, Rangel TFLVB, Bini LM. 2008. Model selection and information theory in  
7 geographical ecology. *Global Ecology and Biogeography* 17: 479-488.
- 8 Dorado J, Vázquez DP. 2014. The diversity–stability relationship in floral production. *Oikos* 123:  
9 1137–1143.
- 10 Ebeling A, Klein A-M, Schumacher J, Weisser WW, Tschardt T. 2008. How does plant richness  
11 affect pollinator richness and temporal stability of flower visits? *Oikos* 117: 1808–1815.
- 12 Farr TG, Rosen PA, Caro E. et al. 2007. The shuttle radar topography mission. *Reviews of*  
13 *Geophysics* 45: 1-33.
- 14 Figueiredo RA, Sazima M. 2000. Pollination Biology of Piperaceae species in Southeastern Brazil.  
15 *Annals of Botany* 85: 455-460.
- 16 Fleming TH. 2005. The relationship between species richness of vertebrate mutualists and their  
17 food plants in tropical and subtropical communities differs among hemispheres. *Oikos* 111:  
18 556-562.
- 19 Fonseca LCN, Vizentin-Bugoni J, Rech AR, 2015. Hummingbird-plant interactions and temporal  
20 nectar availability in a restinga from Southeast Brazil. *Anais da Academia Brasileira de*  
21 *Ciências* 87: 2163-2175.
- 22 Fontaine C, Dajoz I, Meriguet J, Loreau M. 2006. Functional Diversity of Plant–Pollinator  
23 Interaction Webs Enhances the Persistence of Plant Communities. *PLoS Biol* 4: e1.

- 1 Friedman J, Barrett SCH. 2009. Wind of change, new insights on the ecology and evolution of  
2 pollination and mating in wind-pollinated plants. *Annals of Botany* 103: 1515-1527.
- 3 Fründ J, Dormann CF, Holzschuh A, Tschardtke T. 2013. Bee diversity effects on pollination  
4 depend on functional complementarity and niche shifts. *Ecology* 94: 2042–2054.
- 5 Garcia RA, Cabeza M, Rahbek C, Araújo MB. 2014. Multiple dimensions of climate change and  
6 their implications for biodiversity. *Science* 334: 6183.
- 7 Groom SVC, Stevens MI, Schwarz MP. 2014. Parallel responses of bees to Pleistocene climate  
8 change in three isolated archipelagos of the southwestern Pacific. *Proceedings of the Royal  
9 Society B* 281: 20133293.
- 10 Hansen MC, DeFries RS, Townshend JRG, Carroll M, Dimiceli C, Sohlberg R A. 2003. Global  
11 Percent Tree Cover at a Spatial Resolution of 500 Meters: First Results of the MODIS  
12 Vegetation Continuous Fields Algorithm. *Earth Interactions* 7: 1–15.
- 13 Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated  
14 climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.
- 15 Howe HF, Smallwood J 1982. Ecology of seed dispersal. *Annual Review of Ecology and  
16 Systematics* 13: 201–228.
- 17 Hu S, Dilcher DL, Taylor DW. 2012. Pollen evidence for the pollination biology of early flowering  
18 plants. *Evolution of plant-pollinator relationships* (ed. by S. Patiny), pp. 165-236.  
19 Cambridge, Cambridge University Press.
- 20 Jordano P. 2000. Fruits and frugivory. *Seeds: the ecology of regeneration in plant communities*(ed.  
21 by M. Fenner), pp. 125–166. CAB International, Wallingford, UK.
- 22 Kearns CA, Inouye, DW, Waser NM. 1998. Endangered mutualisms: the conservation of plant–  
23 pollinator interactions. *Annual Review Ecology Systematics* 29: 83–112.

- 1 Kissling WD, Eiserhardt WL, Baker WJ, Borchsenius F, Couvreur TLP, Balslev H, Svenning JC.  
2 2012. Cenozoic imprints on the phylogenetic structure of palm species assemblages  
3 worldwide. *Proceeding of the National Academy of Science* 109: 7379-7384.
- 4 Kissling WD, Rahbek C, Böhning-Gaese K. 2007. Food plant diversity as broad-scale determinant  
5 of avian frugivore richness. *Proceedings of the Royal Society B* 274: 799-808.
- 6 Kühn I, Bierman SM, Durka W, Klotz S. 2006. Relating geographical variation in pollination types  
7 to environmental and spatial factors using novel statistical methods. *New Phytologist* 172:  
8 127-139.
- 9 Lever JJ, van Nes EH, Scheffer M, Bascompte J. 2014. The sudden collapse of pollinator  
10 communities. *Ecology Letters* 17: 350-359.
- 11 Loarie SR, Dufy PB, Hamilton H, Asner GP, Field CB, Ackerly DD. 2009. The velocity of climate  
12 change. *Nature* 462: 1052-1055.
- 13 Martín González AM, Dalsgaard B, Nogués-Bravo D, Graham CH, Schleuning M, Maruyama PK,  
14 Abrahamczyk S, Alarcón R, Araujo AC, Araújo FP, Mendes de Azevedo S Jr, Baquero AC,  
15 Cotton PA, Ingwersen TT, Kohler G, Lara C, Guedes Las-Casas FM, Machado AO, Machado  
16 CG, Maglianesi MA, McGuire JA, Moura AC, Oliveira GM, Oliveira PE, Ornelas JF, Rodrigues  
17 LdC, Rosero-Lasprilla L, Rui AM, Sazima M, Timmermann A, Varasin IG, Vizentin-Bugoni J,  
18 Wang Z, Watts S, Rahbek C, Martinez ND. 2015. The macroecology of phylogenetically  
19 structured hummingbird-plant networks. *Global Ecology and Biogeography* 24: 1212-1224.
- 20 McKinney AM, CaraDonna PJ, Inouye DW, Barr B, Bertelsen CD, Waser NM. 2012.  
21 Asynchronous changes in phenology of migrating broad-tailed hummingbirds and their  
22 early-season nectar resources. *Ecology* 93:1987-1993.
- 23 Niklas KJ. 1985. The aerodynamics of wind pollination. *Botanical Review* 51: 328-386.

- 1 Ollerton J, Johnson SD, Hingston AB. 2006. Geographical variation in diversity and specificity of  
2 pollination systems. Pp. 283-308. *In*, Waser, N.M. and Ollerton, J. (eds.) Plant-pollinator  
3 interactions, from specialization to generalization. University of Chicago Press, Chicago,  
4 USA.
- 5 Ollerton J, Winfree R, Tarrant S. 2011. How many flowering plants are pollinated by animals?  
6 *Oikos* 120: 321-326.
- 7 Otto-Bliesner BL, Brady EC, Clauzet G, Tomas R, Levis S, Kothavala Z. 2006. Last Glacial  
8 Maximum and Holocene Climate in CCSM3. *Journal of Climate* 19: 2526-2544.
- 9 R Development Core Team. 2014. R, A language and environment for statistical computing. R  
10 Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>.
- 11 Rangel TFLVB, Diniz-Filho JAF, Bini LM. 2010. SAM, a comprehensive application for Spatial  
12 Analysis in Macroecology. *Ecography* 33: 46-50.
- 13 Rathcke B. 1983. Competition and facilitation among plants for pollination. *In*, Real, L. *Pollination*  
14 *Biology*. Academic Press. pp. 305-325.
- 15 Regal PJ. 1982. Pollination by wind and animals: ecology of geographic patterns. *Annual Review*  
16 *of Ecology and Systematics* 13: 497-524.
- 17 Sandel B, Arge L, Dalsgaard B, Davies RG, Gaston KJ, Sutherland WJ, Svenning JC. 2011. The  
18 influence of late Quaternary climate-change velocity on species endemism. *Science* 334:  
19 660-664.
- 20 Santos T, Diniz-Filho JA, Rangel T, Bini LM. 2013. PVR: Computes phylogenetic eigenvectors  
21 regression (PVR) and phylogenetic signal-representation curve (PSR) (with null and  
22 Brownian expectations). R package version 0.2.1. <http://CRAN.R-project.org/package=PVR>

- 1 Schemske DW, Mittelbach GG, Cornell HV, Sobel JM, Roy K. 2009. Is there a latitudinal gradient  
2 in the importance of biotic interactions? *Annual Review of Ecology, Evolution, and*  
3 *Systematics* 40: 245-269.
- 4 Scherber C, Eisenhauer N, Weisser WW. et al. 2010. Bottom-up effects of plant diversity on  
5 multitrophic interactions in a biodiversity experiment. *Nature* 468: 553-556.
- 6 Schleuning M, Ingmann L, Strauß R. et al. 2014. Ecological, historical and evolutionary  
7 determinants of modularity in weighted seed-dispersal networks. *Ecology Letters* 17: 454-  
8 463.
- 9 Svenning CJ, Skov F. 2007. Ice age legacies in the geographical distribution of tree species richness  
10 in Europe. *Global Ecology and Biogeography* 16: 234-245.
- 11 Waser NM, Real LA. 1979. Effective mutualism between sequentially flowering plant species.  
12 *Nature* 281: 670-672.
- 13 Webb CO, Ackerly DD, Kembel SW. 2008. Phylocom: software for the analysis of phylogenetic  
14 community structure and trait evolution. *Bioinformatics* 24: 2098-2100.
- 15 Webb CO, Donoghue MJ. 2005. Phylomatic: tree assembly for applied phylogenetics. *Molecular*  
16 *Ecology Notes* 5: 181-183.
- 17 Whitehead DR. 1969. Wind-pollination in the angiosperms, evolutionary and environmental  
18 considerations. *Evolution* 23: 28-35.
- 19 Willson MF, Irvine AK, Walsh NG. 1989. Vertebrate dispersal syndromes in some Australian and  
20 New Zealand plant communities, with geographic comparisons. *Biotropica* 21: 133-147.
- 21 Wragg PD, Johnson SD. 2011. Transition from wind pollination to insect pollination in sedges,  
22 experimental evidence and functional traits. *New Phytologist* 191: 1128-1140.

1 Yang CF, Wang QF, Guo YH. 2013. Pollination in a patchily distributed lousewort is facilitated by  
2 presence of a co-flowering plant due to enhancement of quantity and quality of pollinator  
3 visits. *Annals of Botany* 112: 1751-1758.

4

5 **Supporting information**

6 Appendix Table S1. Site descriptions

7 Appendix Table S2. Correlations between predictor variables

8 Appendix Table S3. Contemporary and historical determinants of the pollination mode



### Table and Figures

Table 1. Correlations among the proportion of units in different taxonomic categories regarding pollination mode (animal *versus* wind pollination).

| Taxa                 | Pearson correlation | t-value | Degrees of freedom | P-value |
|----------------------|---------------------|---------|--------------------|---------|
| Species and Genera   | 0.971               | 31.2131 | 59                 | < 0.001 |
| Species and Families | 0.776               | 9.4789  | 59                 | < 0.001 |
| Genera and Families  | 0.836               | 11.7435 | 59                 | < 0.001 |

Table 2. Contemporary and historical correlates (precipitation and temperature anomalies) of the proportion of animal-pollinated plant species in plant communities worldwide (n=67). The standardised regression coefficients are reported both for ordinary least square (OLS) and logistic regression, and reported for both an averaged model based on weighted  $w_i$  and minimum adequate models (MAMs), as in Diniz-Filho et al. (2008). For all MAMs based on OLS, we give  $AIC_c$ , Condition Number, Moran's I, and coefficients of determination ( $R^2$ ). Finally, " $R^2_{\text{species richness}}$ ", " $R^2_{\text{topography}}$ ", " $R^2_{\text{current climate}}$ " and " $R^2_{\text{historical climate}}$ " reflect the unique variation explained by species richness, topography, current climate and historical climate, respectively. Note that historical climate stability is represented by temperature and precipitation anomaly between 21,000 years ago and the present (current minus Last Glacial Maximum - LGM precipitation/temperature, i.e. positive values reflect areas having been drier or colder at LGM than at present). See Appendix Table 2 for similar calculations when using temperature and precipitation velocities as historical climate stability measures.

|                        | OLS          |          |                  | Logistic     |          |                  |
|------------------------|--------------|----------|------------------|--------------|----------|------------------|
|                        | $\Sigma w_i$ | Averaged | MAM <sup>†</sup> | $\Sigma w_i$ | Averaged | MAM <sup>¶</sup> |
| Insularity             | 0.22         | +0.02    |                  | 0.96         | -0.33    | -0.33            |
| Topography             | 0.84         | +0.28    | +0.28            | 1.00         | +0.45    | +0.42            |
| Plant species richness | 0.83         | +0.23    | +0.22            | 0.93         | +0.24    | +0.25            |

|   |      |       |                         |      |       |       |
|---|------|-------|-------------------------|------|-------|-------|
| Open vegetation vs. forest                      | 0.42 | -0.22 |                         | 1.00 | +0.27 | +0.29 |
| MAP x Open vegetation vs forest                 | 0.64 | +0.37 | +0.18                   | 1.00 | +1.18 | +1.23 |
| MAP (Mean Annual Precipitation)                 | 0.25 | -0.05 |                         | 1.00 | +0.13 | +0.18 |
| MAT (Mean Annual Temperature)                   | 1.00 | +0.69 | +0.74                   | 1.00 | +1.42 | +1.34 |
| MAP seasonality                                 | 0.31 | -0.11 |                         | 0.48 | -0.17 |       |
| MAT anomaly                                     | 0.70 | +0.25 | +0.25                   | 1.00 | +0.84 | +0.85 |
| MAP anomaly                                     | 0.24 | +0.00 |                         | 0.51 | +0.18 |       |
| AIC (Akaike Information Criteria <sub>c</sub> ) |      |       | -73.29                  |      |       | 668.9 |
| Moran's Index                                   |      |       | $\leq 0.08^{\text{NS}}$ |      |       |       |
| Condition Number                                |      |       | 2.6                     |      |       |       |
| R <sup>2</sup>                                  |      |       | 0.49                    |      |       |       |
| R <sup>2</sup> <sub>species richness</sub>      |      |       | 0.04                    |      |       |       |

|                                   |      |
|-----------------------------------|------|
| $R^2_{\text{topography}}$         | 0.01 |
| $R^2_{\text{current climate}}$    | 0.36 |
| $R^2_{\text{historical climate}}$ | 0.02 |

---

\*\* ,  $P < 0.01$ ; \* ,  $P < 0.05$ ; <sup>NS</sup>, non-significant. † three models were equally fit (i.e.  $\Delta AIC_c \leq 2$ ) containing the following variables, 1) Open vegetation vs forest, plant species richness, MAT, MAP x Open vegetation versus forest, topography, MAT anomaly; 2) plant species richness, MAT, topography, MAT anomaly; 3) Open vegetation vs forest, plant species richness, MAT, topography, MAT anomaly. † three other models were equally fit (i.e.  $\Delta AIC_c \leq 2$ ) containing the following variables, 1) Open vegetation vs forest, insularity, plant species richness, MAT, MAP, MAP x Open vegetation versus forest, MAP seasonality, topography, MAT anomaly, MAP anomaly ; 2) Open vegetation vs forest, insularity, plant species richness, MAT, MAP, MAP x Open vegetation versus forest, MAP seasonality, topography, MAT anomaly; 3) Open vegetation vs forest, insularity, plant species richness, MAT, MAP, MAP x Open vegetation versus forest, topography, MAT anomaly, MAP anomaly.

**Figure 1.** Geographic patterns of animal pollination in 67 communities surveyed across the world. Fitted curves are second order polynomials;  $r^2 = 0.28$ ,  $F_{2, 47}$ , 13.85,  $P < 0.001$ . On the map, dots were separated slightly within a small range to reduce the overlap of nearby sites.

**Figure 2.** Examples of species having traits representing wind (A-D) and animal (E-G) pollination used to classify plant species according to pollination mode in this study. A, *Ambrosia artemisiifolia* L. (Asteraceae) a diclinous wind pollinated plant with minute flowers that produce a large amount of pollen (Photo by Andrew Butko via Wiki images), B, *Mercurialis annua* L. (Euphorbiaceae) a dioecious wind-pollinated plant species; the circle highlights the female flower and, in detail at the right side of the image, we show the inflorescence of a male individual (Photo by Hasan Yldirin), C, *Olyra ciliatifolia* Raddi (Poaceae) female flower with feathery stigmas (Photo by Pedro Viana), D, *Paspalum notatum* Flügge (Poaceae) showing the flexible white filaments; the arrow is pointing to the wide anther aperture (Photo by Pedro Viana), E, *Turnera ulmifolia* L. (Passifloraceae) a 3D heterostyle species and a butterfly, one of its animal pollinators, F, *Couroupita guianensis* Aubl. (Lecythidaceae) with food (colourful) and pollination (white) specialised stamens (Photo by Hipolito Neto), G, *Gongora bufonia* Lindl. (Orchidaceae) being pollinated by a male euglossine bee (*Eufrisea violacea*) when collecting scented oil from its labellum (Photo by Carlos Eduardo Coquinho).

**Figure 3.** The relationship between the proportion of animal pollinated plant species per community and the most important and consistent predictors, a) plant richness; b) temperature; c) precipitation. Open symbols illustrate open vegetation types whereas filled symbols

illustrate forest communities. All relationships significant in simple univariate correlations are shown: all vegetation types (n = 67; dashed line); open vegetation types (n = 51, dotted line); and forest (n = 16, full line). See Table 2 and Tables S1-S2 for statistics.

Appendix Table S1. Site descriptions for each community analysed in the study: “The macroecology of animal *versus* wind pollination”. Each number in the first column corresponds to a reference below.

| N° | Study site                          | Mainland = 0 | Forest: 1 | Total   | N° wind      | % animal    | Latitude    | Longitude   |
|----|-------------------------------------|--------------|-----------|---------|--------------|-------------|-------------|-------------|
|    |                                     | Island = 1   | Other = 0 | plants: | pol. species | pollination | Decimalised | Decimalised |
| 1  | Juan Fernandez Islands, Chile       | 1            | 0         | 149     | 70           | 53.02       | -33.64      | -78.84      |
| 2  | Coastal scrub, UK                   | 1            | 0         | 37      | 9            | 75.68       | 54.40       | -0.51       |
| 3  | Degraded tropical scrub/secondary   | 0            | 1         | 82      | 3            | 96.34       | 22.28       | 114.2       |
| 4  | Serra da Bocaina - Grassland        | 0            | 0         | 179     | 55           | 69.27       | -22.73      | -44.61      |
| 5  | Cerrado, Brazil                     | 0            | 0         | 294     | 52           | 82.31       | -22.87      | -48.49      |
| 6  | semi-arid scrub - Spain             | 0            | 0         | 31      | 6            | 80.65       | 37.01       | -6.55       |
| 7  | Amami Island, Japan                 | 1            | 0         | 103     | 6            | 94.17       | 28.32       | 129.42      |
| 8  | All habitats across New Caledonia   | 1            | 0         | 97      | 3            | 96.91       | -21.17      | 165.25      |
| 9  | La Selva, Costa Rica                | 0            | 1         | 283     | 13           | 95.41       | 10.45       | -84.00      |
| 10 | Caatinga, Brazil                    | 0            | 0         | 147     | 3            | 97.96       | -8.45       | -36.76      |
| 11 | various localities, Faroe Islands   | 1            | 0         | 70      | 19           | 72.86       | 62.00       | 6.783       |
| 12 | Guyana Highlands, Venezuela         | 0            | 0         | 55      | 6            | 89.09       | 5.58        | -61.71      |
| 13 | Rainforest, Malaysia                | 1            | 1         | 262     | 0            | 100.00      | 4.20        | 114.5       |
| 14 | Alpine pavement plain, California   | 0            | 0         | 17      | 4            | 76.47       | 34.30       | -116.86     |
| 15 | Woodland, USA                       | 0            | 1         | 208     | 57           | 72.60       | 42.1        | -111.59     |
| 16 | various localities, St Kilda        | 1            | 0         | 92      | 39           | 57.61       | 57.81       | -8.58       |
| 17 | Palm swamp, Venezuela               | 0            | 1         | 33      | 5            | 84.85       | 8.93        | -67.25      |
| 18 | Prairie, Illinois                   | 0            | 0         | 409     | 14           | 96.58       | 39.46       | -89.90      |
| 19 | Flooded/Upland Rainforest, Colombia | 0            | 1         | 90      | 2            | 98.51       | -0.61       | -72.33      |
| 20 | Lower alpine, Scotland              | 1            | 0         | 153     | 8            | 94.77       | 56.50       | -3.11       |
| 21 | Serra do Cipo                       | 0            | 0         | 64      | 5            | 92.19       | -23.336667  | -45.14      |
| 22 | Botucatu                            | 0            | 0         | 34      | 3            | 91.18       | -22.844167  | -48.68      |
| 23 | Santa Virginia                      | 0            | 1         | 70      | 8            | 88.57       | -19.334722  | -43.58      |

|    |                                     |   |   |     |    |       |            |             |
|----|-------------------------------------|---|---|-----|----|-------|------------|-------------|
| 24 | Picinguaba - restinga               | 0 | 0 | 47  | 5  | 89.36 | -23.358056 | -44.85      |
| 25 | Picinguaba - Atlantic forest        | 0 | 1 | 43  | 2  | 95.35 | -23.338611 | -44.83      |
| 26 | La Floresta - Canelones - Uruguay - | 0 | 0 | 24  | 10 | 58.33 | -34.760706 | -55.69      |
| 27 | Quebrada de los Cuervos - Uruguay - | 0 | 0 | 42  | 14 | 66.67 | -32.935091 | -54.46      |
| 28 | Kumu - Guyana - rainforest          | 0 | 1 | 59  | 3  | 94.92 | 3.266667   | -59.75      |
| 29 | Kumu - Guyana - savannah            | 0 | 0 | 43  | 6  | 86.05 | 3.266667   | -59.77      |
| 30 | Wahroonga - South Africa            | 0 | 0 | 73  | 4  | 94.52 | -29.616667 | 30.13       |
| 31 | Mantanay - Peru                     | 0 | 0 | 148 | 6  | 95.95 | -13.2      | -72.08      |
| 32 | Scrub Field - Northampton - UK      | 1 | 0 | 162 | 35 | 78.40 | 52.26      | -0.88       |
| 33 | Bahia de Patano - Venezuela         | 0 | 0 | 78  | 12 | 84.62 | 10.466667  | -67.75      |
| 34 | Guimar Badlands, Tenerife           | 1 | 0 | 18  | 2  | 88.89 | 28.33      | -16.42      |
| 35 | Mean JasperRidge Plant Covariates - | 0 | 0 | 109 | 24 | 78.98 | 37.417     | -122.19     |
| 36 | Mean JasperRidge Plant Covariates - | 0 | 1 | 78  | 16 | 77.85 | 37.417     | -122.19     |
| 37 | JasperRidge Plant Covariates -      | 1 | 0 | 16  | 7  | 56.25 | 37.701067  | -123.00     |
| 38 | MatherPlantCovariatesPUPPLEEsEt -   | 0 | 0 | 92  | 31 | 66.30 | 37.8774    | -119.25     |
| 39 | MatherPlantCovariatesPUPPLEEsEt -   | 0 | 0 | 58  | 6  | 89.66 | 37.8374    | -120.30     |
| 40 | MatherPlantCovariatesPUPPLEEsEt -   | 0 | 1 | 132 | 19 | 85.61 | 37.8774    | -119.29     |
| 41 | MatherPlantCovariatesPUPPLEEsEt -   | 0 | 0 | 134 | 33 | 75.37 | 38.0095    | -123.00     |
| 42 | TimberlinePlantDataPurple - Dore    | 0 | 0 | 60  | 19 | 68.33 | 37.9683    | -119.30     |
| 43 | TimberlinePlantDataPurple -         | 0 | 0 | 95  | 47 | 50.53 | 37.9382    | -119.25     |
| 44 | TimberlinePlantDataPurple -         | 0 | 1 | 91  | 34 | 62.64 | 37.9612    | -119.29     |
| 45 | TimberlinePlantDataPurple - Talus   | 0 | 0 | 126 | 38 | 69.84 | 37.9462    | -119.25     |
| 46 | Virginia Basin (Colorado)           | 0 | 0 | 64  | 6  | 90.63 | 38.98      | -106.966667 |
| 47 | Mean Hangklip, South Africa         | 0 | 0 | 124 | 41 | 65.55 | -34.25     | 18.75       |
| 48 | Brattnesdalen, Norway               | 0 | 0 | 18  | 8  | 55.56 | 70.25      | 22.07       |
| 49 | Fjorddalen, Norway                  | 0 | 0 | 31  | 10 | 67.74 | 70.19      | 22.10       |
| 50 | various localities, Australia       | 0 | 1 | 148 | 8  | 94.59 | -19.18     | 146.75      |
| 51 | Torrey Pines - California           | 0 | 0 | 85  | 21 | 75.29 | 32.89      | -117.24     |
| 52 | Japatul Valley - California         | 0 | 0 | 91  | 15 | 83.52 | 32.78      | -116.68     |
| 53 | Echo Valley - California            | 0 | 0 | 56  | 4  | 92.86 | 32.89      | -116.65     |



|    |                               |   |   |     |    |       |        |             |
|----|-------------------------------|---|---|-----|----|-------|--------|-------------|
| 54 | Mount Laguna - California     | 0 | 1 | 65  | 15 | 76.92 | 32.86  | -116.41     |
| 55 | Ocotillo - California         | 0 | 0 | 80  | 14 | 82.50 | 32.74  | -115.99     |
| 56 | Papudo - Chile                | 0 | 0 | 95  | 23 | 75.79 | -32.55 | -71.44      |
| 57 | Fundo Santa Laura - Chile     | 0 | 0 | 106 | 18 | 83.02 | -33.26 | -70.85      |
| 58 | Cerro Potrerillo - Chile      | 0 | 0 | 47  | 11 | 76.60 | -30.29 | -70.57      |
| 59 | El Tofo - Chile               | 0 | 0 | 44  | 5  | 88.64 | -29.88 | -71.20      |
| 60 | Alpine - montane Colorado     | 0 | 0 | 68  | 17 | 75.00 | 38.68  | -107.115532 |
| 61 | Aspen - montane Colorado      | 0 | 1 | 56  | 15 | 73.21 | 38.73  | -106.772653 |
| 62 | Sage - montane Colorado       | 0 | 0 | 45  | 9  | 80.00 | 38.73  | -106.823376 |
| 63 | Grassland - montane Colorado  | 0 | 0 | 103 | 24 | 76.70 | 38.95  | -106.988824 |
| 64 | Spruce-fir - montane Colorado | 0 | 1 | 50  | 11 | 78.00 | 38.86  | -107.101081 |
| 65 | Salt marsh, Canada            | 0 | 0 | 18  | 6  | 66.67 | 49.08  | -125.85     |
| 66 | Sphagnum bog, Canada          | 0 | 0 | 33  | 13 | 60.61 | 49.08  | -125.86     |
| 67 | Subalpine meadow, Canada      | 0 | 0 | 45  | 11 | 75.56 | 49.11  | -120.84     |

---

## Appendix Table S1 - References

- 1- Bernadello, G. et al. 2001. A survey of floral traits, breeding systems, floral visitors, and pollination systems of the angiosperms of the Juan Fernandez Islands (Chile). - *Bot. Rev.* 67: 255 - 308.
- 2- Burkill, I. H. 1897. Fertilization of some spring flowers on the Yorkshire coast. - *J. Bot.* 35: 92 - 189.
- 3- Corlett, R. T. 2001. Pollination in a degraded tropical landscape: a Hong Kong case study. - *J. Trop. Ecol.* 17: 155-161.
- 4- Freitas, L. and Sazima, M. 2006. Pollination biology in a tropical high-altitude grassland in Brazil: interactions at the community level. *Ann.Missouri Bot.Gard.* 93: 465-516.
- 4- Martinelli, G. 1989. *Campos de Altitude*. Editora Index, Rio de Janeiro.
- 5- Gottsberger, G. and Silberbauer-Gottsberger, I. 2006. *Life in the Cerrado*. Reta Verlag.
- 6- Herrera, J. 1988. Pollination relationships in southern Spanish mediterranean shrublands. *J. Ecol.* 76: 274 - 287.
- 7- Kato, M. 2000. Anthophilous insect community and plant-pollinator interactions on Amami Islands in the Ryukyu Archipelago, Japan. - *Contrib. Biol. Lab. Kyoto Univ.* 29: 157 – 252.
- 8- Kato, M. and Kawakita, A. 2004. Plant-pollinator interactions in New Caledonia influenced by introduced honey bees. *Am. J. Bot.* 91: 1814-1827.

- 9- Kress, W. J. and Beach, J. H. 1994. Flowering plant reproductive systems. - *In*: McDade, L. A. et al. (eds), *La Selva: ecology and natural history of a Neotropical rainforest*. Univ. Chicago Press, pp. 161 - 182.
- 10- Machado, I. C. and Lopes, A. V. 2004. Floral traits and pollination systems in the caatinga, a Brazilian tropical dry forest. *Ann. Bot.* 94: 365-376.
- 11 - 12- Ollerton, J. Unpublished data, available upon request from the author via [Jeff.Ollerton@northampton.ac.uk](mailto:Jeff.Ollerton@northampton.ac.uk)
- 13- Momose, K. et al. 1998. Pollination biology in a lowland dipterocarp forest in Sarawak, Malaysia. Characteristics of the plant-pollinator community in a lowland dipterocarp forest. *Am. J. Bot.* 85: 1477-1501.
- 14- O'Brien, M. H. 1980. The pollination biology of a pavement plain: pollinator visitation patterns. - *Oecologia* 47: 213-218
- 15- Ostler, W. K. and Harper, K. T. 1978. Floral ecology in relation to plant species diversity in the Wasatch Mountains of Utah and Idaho. *Ecology* 59:848-861.
- 16 and 48-49- Holmes, N. 2014. Unpublished data, available upon request from the author via [Naomi.Holmes@northampton.ac.uk](mailto:Naomi.Holmes@northampton.ac.uk)
- 17- Ramirez, N. and Brito, Y. 1992. Pollination biology in a palm swamp community in the Venezuelan Central Plains. *Bot. J. Linn. Soc.* 110: 277-302.

- 18- Robertson, C. 1928. Flowers and insects: lists of visitors of four hundred and fifty-three flowers. Privately published, Carlinville, Illinois.
- 19- van Dulmen, A. 2001. Pollination and phenology of flowers in the canopy of two contrasting rain forest types in Amazonia, Colombia. - Plant Ecol. 153: 73-85.
- 20- Willis, J. C. and Burkill, I. H. 1895. Flowers and insects in Great Britain. Part I. Ann. Bot. 9: 227-273.
- 20- Willis, J. C. and Burkill, I. H. 1903a. Flowers and insects in Great Britain. Part II. Ann. Bot. 17: 313-349
- 20- Willis, J. C. and Burkill, I. H. 1903b. Flowers and insects in Great Britain. Part III. Ann. Bot. 17: 539-570.
- 20- Willis, J. C. and Burkill, I. H. 1908. Flowers and insects in Great Britain. Part IV. Ann. Bot. 22: 603-649.
- 21-25- Ollerton, J. and Rech, A.R. 2013. unpublished data, available upon request from the authors via [Jeff.Ollerton@northampton.ac.uk](mailto:Jeff.Ollerton@northampton.ac.uk)
- 26-27- Rech, A.R. 2014. unpublished data, available upon request from the author via [andrerodrigorech@gmail.com](mailto:andrerodrigorech@gmail.com)
- 28-34- Ollerton, J. Unpublished data, available upon request from the author via [Jeff.Ollerton@northampton.ac.uk](mailto:Jeff.Ollerton@northampton.ac.uk)
- 35-36- Moldenke, D. Unpublished data, requested from the author.
- 37-46- Moldenke, D. Unpublished data, requested from the author.

- 47 and 50- Ollerton, J. Unpublished data, available upon request from the author via [Jeff.Ollerton@northampton.ac.uk](mailto:Jeff.Ollerton@northampton.ac.uk)
- 51-59- Moldenke, A. R. 1979. Pollination ecology as an assay for ecosystemic organization: convergent evolution in Chile and California. *Phytologia* 42: 415-454.
- 60-64- Moldenke, A. R. and Lincoln, P. G. 1979. Pollination ecology in montane Colorado: a community analysis. *Phytologia* 42: 349 - 379.
- 65-67- Pojar, J. 1974. Reproductive dynamics of four plant communities of southwestern British Columbia. *Can. J. Bot.* 52: 1819 - 1834.

Appendix Table S2. Correlations between the proportion of animal pollinated plant species and all predictor variables for all vegetation types (n = 67, except for velocities which had one data point less as we were unable to extract it for St Kilda Island, Scotland) above the diagonal, and separately for open vegetation types (n = 51, except for velocities which had one data point less as we were unable to extract it for St Kilda Island, Scotland) below the diagonal.

|                                 | % animal<br>pollinated | Plant<br>richness   | MAT                 | MAP                      | MAP<br>seasonality  | Topography               | MAT<br>anomaly      | MAP<br>anomaly      | MAT<br>velocity          | MAP<br>velocity     |
|---------------------------------|------------------------|---------------------|---------------------|--------------------------|---------------------|--------------------------|---------------------|---------------------|--------------------------|---------------------|
| % animal pollinated             |                        | <b>+0.32**</b>      | <b>+0.57*</b>       | <b>+0.29<sup>†</sup></b> | +0.09 <sup>NS</sup> | -0.11 <sup>NS</sup>      | -0.14 <sup>NS</sup> | +0.03 <sup>NS</sup> | +0.07 <sup>NS</sup>      | +0.13 <sup>NS</sup> |
| Plant richness                  | <b>+0.28*</b>          |                     | +0.11 <sup>NS</sup> | +0.11 <sup>NS</sup>      | -0.05 <sup>NS</sup> | -0.11 <sup>NS</sup>      | +0.06 <sup>NS</sup> | +0.23 <sup>NS</sup> | +0.16 <sup>NS</sup>      | +0.05 <sup>NS</sup> |
| MAT (Mean Annual Temperature)   | <b>+0.42*</b>          | +0.09 <sup>NS</sup> |                     | <b>+0.44*</b>            | <b>+0.35*</b>       | <b>-0.45**</b>           | <b>-0.50*</b>       | +0.01 <sup>NS</sup> | +0.05 <sup>NS</sup>      | <b>+0.37**</b>      |
| MAP (Mean Annual Precipitation) | -0.02 <sup>NS</sup>    | -0.04 <sup>NS</sup> | +0.24 <sup>NS</sup> |                          | -0.17 <sup>NS</sup> | <b>-0.25<sup>†</sup></b> | -0.20 <sup>NS</sup> | <b>+0.33**</b>      | +0.14 <sup>NS</sup>      | <b>+0.25*</b>       |
| MAP seasonality                 | +0.14 <sup>NS</sup>    | +0.02 <sup>NS</sup> | <b>+0.39*</b>       | <b>-0.18*</b>            |                     | +0.12 <sup>NS</sup>      | <b>-0.53*</b>       | -0.05 <sup>NS</sup> | <b>-0.36<sup>†</sup></b> | +0.20 <sup>NS</sup> |
| Topography                      | +0.04 <sup>NS</sup>    | -0.08 <sup>NS</sup> | <b>-0.36**</b>      | -0.17 <sup>NS</sup>      | +0.19 <sup>NS</sup> |                          | -0.04 <sup>NS</sup> | -0.16 <sup>NS</sup> | <b>-0.71**</b>           | <b>-0.47**</b>      |
| MAT anomaly                     | -0.08 <sup>NS</sup>    | +0.12 <sup>NS</sup> | <b>-0.51*</b>       | -0.10 <sup>NS</sup>      | <b>-0.60**</b>      | -0.13 <sup>NS</sup>      |                     | -0.17 <sup>NS</sup> | <b>+0.50**</b>           | -0.21 <sup>NS</sup> |
| MAP anomaly                     | -0.22 <sup>NS</sup>    | +0.08 <sup>NS</sup> | -0.17 <sup>NS</sup> | +0.22 <sup>NS</sup>      | +0.06 <sup>NS</sup> | -0.01 <sup>NS</sup>      | -0.13 <sup>NS</sup> |                     | +0.16 <sup>NS</sup>      | +0.10 <sup>NS</sup> |
| MAT velocity                    | +0.01 <sup>NS</sup>    | +0.16 <sup>NS</sup> | -0.04 <sup>NS</sup> | +0.10 <sup>NS</sup>      | <b>-0.41*</b>       | <b>-0.70**</b>           | <b>+0.58**</b>      | +0.05 <sup>NS</sup> |                          | <b>+0.40*</b>       |
| MAP velocity                    | +0.02 <sup>NS</sup>    | -0.01 <sup>NS</sup> | <b>+0.29**</b>      | +0.14 <sup>NS</sup>      | +0.17 <sup>NS</sup> | <b>-0.42**</b>           | -0.15 <sup>NS</sup> | -0.05 <sup>NS</sup> | <b>+0.43*</b>            |                     |

\*\*P < 0.01; \*P < 0.05 when P-values based on degrees of freedom corrected for spatial autocorrelation using Dutilleul's (1993) method;

†significant when using traditional non-spatial statistics, but non-significant when corrected for spatial autocorrelation; <sup>NS</sup>non-significant.

Appendix Table S2, continued. Correlations between predictor variables separately for forest (n = 16).

|                                 | % animal<br>pollinated | Plant<br>richness   | MAT                      | MAP                      | MAP<br>seasonality  | Topography               | MAT<br>anomaly           | MAP<br>anomaly           | MAT<br>velocity          | MAP<br>velocity          |
|---------------------------------|------------------------|---------------------|--------------------------|--------------------------|---------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| % animal pol.                   |                        | +0.30 <sup>NS</sup> | <b>+0.82<sup>†</sup></b> | <b>+0.84<sup>*</sup></b> | -0.01 <sup>NS</sup> | <b>-0.59<sup>†</sup></b> | -0.46 <sup>NS</sup>      | +0.37 <sup>NS</sup>      | +0.32 <sup>NS</sup>      | <b>+0.50<sup>†</sup></b> |
| Plant richness                  |                        |                     | +0.05 <sup>NS</sup>      | +0.38 <sup>NS</sup>      | -0.29 <sup>NS</sup> | -0.22 <sup>NS</sup>      | -0.29 <sup>NS</sup>      | <b>+0.57<sup>*</sup></b> | +0.21 <sup>NS</sup>      | +0.26 <sup>NS</sup>      |
| MAT (Mean Annual Temperature)   |                        |                     |                          | +0.74 <sup>†</sup>       | +0.32 <sup>NS</sup> | <b>-0.73<sup>†</sup></b> | <b>-0.67<sup>†</sup></b> | +0.19 <sup>NS</sup>      | +0.34 <sup>NS</sup>      | <b>+0.58<sup>†</sup></b> |
| MAP (Mean Annual Precipitation) |                        |                     |                          |                          | -0.13 <sup>NS</sup> | <b>-0.52<sup>†</sup></b> | <b>-0.70<sup>*</sup></b> | +0.42 <sup>NS</sup>      | +0.33 <sup>NS</sup>      | <b>+0.58<sup>†</sup></b> |
| MAP seasonality                 |                        |                     |                          |                          |                     | -0.14 <sup>NS</sup>      | -0.26 <sup>NS</sup>      | -0.30 <sup>NS</sup>      | -0.13 <sup>NS</sup>      | +0.26 <sup>NS</sup>      |
| Topography                      |                        |                     |                          |                          |                     |                          | +0.55 <sup>†</sup>       | <b>-0.56<sup>*</sup></b> | <b>-0.78<sup>*</sup></b> | <b>-0.61<sup>†</sup></b> |
| MAT anomaly                     |                        |                     |                          |                          |                     |                          |                          | -0.41 <sup>NS</sup>      | -0.17 <sup>NS</sup>      | <b>-0.64<sup>*</sup></b> |
| MAP anomaly                     |                        |                     |                          |                          |                     |                          |                          |                          | <b>+0.56<sup>*</sup></b> | +0.44 <sup>NS</sup>      |
| MAT velocity                    |                        |                     |                          |                          |                     |                          |                          |                          |                          | +0.36 <sup>NS</sup>      |
| MAP velocity                    |                        |                     |                          |                          |                     |                          |                          |                          |                          |                          |

\*\*P < 0.01; \*P < 0.05 when P-values based on degrees of freedom corrected for spatial autocorrelation using Dutilleul's (1993) method;

†significant when using traditional non-spatial statistics, but non-significant when corrected for spatial autocorrelation; <sup>NS</sup>non-significant.

Appendix Table S3. Contemporary and historical determinants (precipitation and temperature velocities) of the proportion of animal-pollinated plant species in plant communities worldwide (n=66). The standardized regression coefficients are reported both for ordinary least square (OLS)



and logistic regression, and reported for both an averaged model based on weighted  $w_i$  and minimum adequate models (MAMs), as in Diniz-Filho *et al.* (2008). For all MAMs based on OLS, we give AIC<sub>c</sub>, Condition Number, Moran's I, and coefficients of determination ( $R^2$ ). Finally, " $R^2_{\text{species richness}}$ ", " $R^2_{\text{current climate}}$ " and " $R^2_{\text{historical climate}}$ " reflect the unique variation explained by species richness, current climate and historical climate, respectively. Notice that historical climate stability is represented by temperature and precipitation velocity between 21000 years ago and now, and that topography is not included as strongly correlated with velocities. See Table 2 for similar calculations using precipitation and temperature anomalies as historical climate stability measures.

|                                 | OLS          |          |                  | Logistic     |          |       |
|---------------------------------|--------------|----------|------------------|--------------|----------|-------|
|                                 | $\Sigma w_i$ | Averaged | MAM <sup>†</sup> | $\Sigma w_i$ | Averaged | MAM   |
| Insularity                      | 0.25         | +0.05    |                  | 0.75         | -0.24    | -0.25 |
| Plant species richness          | 0.85         | +0.24    | +0.24            | 0.85         | +0.21    | +0.20 |
| Open vegetation vs forest       | 0.44         | -0.24    |                  | 1.00         | +0.31    | +0.29 |
| MAP x Open vegetation vs forest | 0.67         | +0.39    | +0.20            | 1.00         | +1.22    | +1.23 |
| MAP (Mean Annual Precipitation) | 0.27         | -0.07    |                  | 1.00         | +0.03    | +0.02 |
| MAT (Mean Annual Temperature)   | 1.00         | +0.50    | +0.47            | 1.00         | +0.82    | +0.84 |

|  |      |       |        |      |       |       |
|--|------|-------|--------|------|-------|-------|
| MAP seasonality                              | 0.31 | -0.10 |        | 0.87 | -0.26 | -0.27 |
| MAT velocity                                 | 0.23 | -0.00 |        | 1.00 | +0.42 | +0.40 |
| MAP velocity                                 | 0.29 | -0.09 |        | 0.22 | -0.01 |       |
| Akaike Information Criteria <sub>c</sub>     |      |       | -69.94 |      |       | 675.9 |
| Moran's Index                                |      |       | ≤0.14* |      |       |       |
| Condition Number                             |      |       | 1.5    |      |       |       |
| R <sup>2</sup>                               |      |       | 0.43   |      |       |       |
| R <sup>2</sup> <sub>species richness</sub>   |      |       | 0.05   |      |       |       |
| R <sup>2</sup> <sub>current climate</sub>    |      |       | 0.32   |      |       |       |
| R <sup>2</sup> <sub>historical climate</sub> |      |       | 0.00   |      |       |       |

---

\*\*P < 0.01; \*P < 0.05; <sup>NS</sup>non-significant. † five other models were equally fit (i.e.  $\Delta AIC_c \leq 2$ ) containing the following variables, 1) Open vegetation vs forest, plant species richness, MAT, MAP x Open vegetation versus forest; 2) plant species richness, MAT; 3) plant species

richness, MAT, MAP x Open vegetation versus forest, MAP velocity; 4) plant species richness, MAT, MAP x Open vegetation versus forest, MAP seasonality; 5) Open vegetation vs forest, plant species richness, MAT.