

Kent Academic Repository

Full text document (pdf)

Citation for published version

De Coninck, Kyra and Hambly, Karen and Passfield, Louis and Dickinson, John W. and Muthumayandi, Karthikeyan (2015) Inter-observer agreement of thoracolumbar fascia morphology: an exploratory analysis of ultrasound images. *Journal of Bodywork and Movement Therapies*, 19 (4). pp. 668-669. ISSN 1360-8592.

DOI

<http://doi.org/10.1016/j.jbmt.2015.07.005>

Link to record in KAR

<http://kar.kent.ac.uk/58489/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Inter-observer agreement of thoracolumbar fascia morphology: an exploratory analysis of ultrasound images

Kyra De Coninck, Karen Hambly, Louis Passfield, John W Dickinson, Karthik Muthumayandi

School of Sport and Exercise Sciences, University of Kent, Chatham ME4 4AG.

Phone +44 1634 88 88 58 Fax+44 1634 88 88 90 Email: K.De-Coninck@kent.ac.uk

BACKGROUND: Ultrasound imaging (USI) has been shown to be a valid method to investigate the morphology of the thoracolumbar fascia (TLF) [1]. A USI-based study has demonstrated that the TLF of subjects with chronic lower back pain (LBP) is on average 25% thicker and more disorganised compared to a control group [1]. The aim of this study is to explore inter-observer agreement between a range of clinicians on (dis)organisation of TLF in ultrasound images. There are currently no validated methods for the evaluation of USI of TLF.

METHODS: Design: an exploratory analysis using a fully crossed design of inter-observer agreement. This study was approved by the University of Kent's School of Sport and Exercise Sciences Research and Ethics Committee (Prop. 163 – 2013). Participants: Thirty observers consisting of 21 (70%) Medical Doctors, 7 (23%) physiotherapists and 2 (6%) radiologists, with a combined total average of 13 years of clinical experience (\pm SD 9.4). 57% had no experience in USI, 36% had experience ranging from monthly to daily evaluations of USI, no observers had experience in evaluating USI of TLF.

Protocol: A sub-set of thirty ultrasound scans of TLF were randomly selected from a data set of 308 scans of subjects with and without LBP (from a larger study conducted by the first author). All scans were anonymised and displayed on a desktop computer, or projected on a screen. All observers viewed and rated each of the 30 scans independently on a Likert-type scale from 1 (very disorganised) to 10 (very organised). Inter-observer agreement was assessed using a two-way mixed, consistency, average measures intra-class correlation (ICC), the Cronbach's Alpha, to assess consistency among observers. The Krippendorff's Alpha (Kalpha) [2] reliability estimate was used to assess agreement.

RESULTS: The resulting ICC was in the excellent range, ICC = 0.98, indicating that observers had a high degree of consistency, suggesting that (dis)organisation was rated similarly across observers. Observers without USI experience scored an ICC = 0.96, observers with USI experience scored an ICC = 0.95, again both in the excellent range. In this small cohort, experience in USI does not appear to impact on consistency. The Krippendorff's ordinal alpha α was .621, indicating a modest degree of agreement.

CONCLUSIONS: The high ICC and modest Kalpha suggest that a minimal amount of measurement error was introduced by the independent observers, and therefore statistical power for subsequent analyses is not substantially reduced. This will allow for further analysis of USI images of TLF in terms of morphology and classification. This could ultimately, lead to a meaningful evaluation of treatments of TLF.

REFERENCES[1] Langevin, HM, Stevens-Tuttle, D, Fox, JR, Badger, GJ, Bouffard, NA, Krag, MH, Wu, J, Henry, SM. Ultrasound evidence of altered lumbar connective tissue structure in human subjects with chronic low back pain. BMC Musculoskeletal Disorders, 10:151–159, 2009.

[2] Hayes, AF, Krippendorff, K. Answering the call for a standard reliability measure for coding data. Communication Methods and Measures, 1, 77-89, 2007.