

Article (refereed)

Dowding, Claire V.; Shore, Richard F.; Worgan, Andrew; Baker, Philip J.; Harris, Stephen. 2010 Accumulation of Anticoagulant Rodenticides in a Non-target Insectivore, the European hedgehog (*Erinaceus europaeus*). *Environmental Pollution*, 158 (1). 161-166. [10.1016/j.envpol.2009.07.017](https://doi.org/10.1016/j.envpol.2009.07.017)

Crown copyright © 2009 Published by Elsevier Ltd.

This version available <http://nora.nerc.ac.uk/7856/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the authors and/or other rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.

www.elsevier.com

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 Accumulation of Anticoagulant Rodenticides in a Non-target Insectivore,
2 the European hedgehog (*Erinaceus europaeus*)

3
4 Claire V. Dowding^{1*}, Richard F. Shore², Andrew Worgan², Philip J. Baker³ and
5 Stephen Harris

6
7 *School of Biological Sciences, University of Bristol, Woodland Road, Bristol BS8*
8 *1UG, UK*

9
10
11
12
13
14 ¹ Current address: Natural England, 11 Fenlock Court, Blenheim Office Park, Long
15 Hanborough, Oxfordshire OX29 8LN, UK

16
17 ² NERC Centre for Ecology and Hydrology, Lancaster Environment Centre, Library
18 Avenue, Bailrigg, Lancaster LA1 4AP, UK

19
20 ³ Current address: Harborne Building, School of Biological Sciences, University of
21 Reading, Whiteknights, Reading, Berkshire RG6 6AS, UK

22
23 * Corresponding author

24
25
26
27
28
29
30 Address for correspondence: Dr Claire Dowding
31 Natural England
32 11 Fenlock Court
33 Blenheim Office Park
34 Long Hanborough
35 Oxford
36 OX29 8LN, UK

37
38
39 Tel: (+44) 0300 060 1924

40 Fax: (+44) 1993 886541

41 e-mail: Claire.Dowding@naturalengland.org.uk

42 **Abstract**

43 Studies on exposure of non-targets to anticoagulant rodenticides have largely focussed
44 on predatory birds and mammals; insectivores have rarely been studied. We investigated the
45 exposure of 120 European hedgehogs (*Erinaceus europaeus*) from throughout Britain to first-
46 and second-generation anticoagulant rodenticides (FGARs and SGARs) using high
47 performance liquid chromatography coupled with fluorescence detection (HPLC) and liquid-
48 chromatography mass spectrometry (LCMS). The proportion of hedgehogs with liver SGAR
49 concentrations detected by HPLC was 3-13% per compound, 23% overall. LCMS identified
50 much higher prevalence for difenacoum and bromadiolone, mainly because of greater ability
51 to detect low level contamination. The overall proportion of hedgehogs with LCMS-detected
52 residues was 57.5% (SGARs alone) and 66.7% (FGARs and SGARs combined); 27 (22.5%)
53 hedgehogs contained >1 rodenticide. Exposure of insectivores and predators to
54 anticoagulant rodenticides appears to be similar. The greater sensitivity of LCMS suggests
55 that hitherto exposure of non-targets is likely to have been under-estimated using HPLC
56 techniques.

57

58

59 *Keywords:* first- and second-generation anticoagulant rodenticide, insectivore,
60 brodifacoum, bromadiolone, difenacoum, flocoumafen, coumatetralyl, warfarin, non-
61 target

62

63 *Capsule:* Exposure of insectivorous hedgehogs to anticoagulant rodenticides in
64 Britain is similar to predatory birds and mammals that specialise in eating small
65 mammals, and hitherto exposure levels have been underestimated using HPLC
66 techniques.

67

68 **1. Introduction**

69

70 Globally, rodents destroy or spoil substantial amounts of food intended for
71 human or animal consumption (Singleton et al., 1999; Stenseth et al., 2003).
72 Consequently, a range of methods is employed to reduce rodent density and
73 associated damage. This is most commonly done in developed countries using
74 anticoagulant rodenticides, vitamin K antagonists that prevent the synthesis of
75 functional prothombrin and related blood-clotting factors. Extensive use of first-
76 generation anticoagulant rodenticides (FGARs) during the 1950s, however, led to the
77 evolution of genetic resistance in brown rats (*Rattus norvegicus*), with widespread
78 cross-resistance to other compounds (Cowan et al., 1995; Thijssen, 1995). As a
79 result, more potent second-generation anticoagulant rodenticides (SGARs) were
80 developed which have a greater affinity to binding sites, resulting in greater
81 accumulation, persistence and toxicity (Parmar et al., 1987; Huckle and Warburton,
82 1986).

83 Given their mode of action, both FGARs and SGARs are potentially harmful to all
84 vertebrates, and so users are expected to adopt measures that limit direct exposure
85 to non-target species. However, the degree to which these preventive measures are
86 adhered to, particularly by non-professionals, is unknown. For example, in Britain
87 some products are readily available to householders who may be less aware of the
88 risks of non-target poisoning and/or less likely to follow manufacturer's guidelines.
89 Non-target species may also be deliberately poisoned (Barnett et al., 2006).

90 Most studies investigating indirect exposure of non-target species to
91 anticoagulant rodenticides have focussed on the consumption of poisoned rodents by
92 predatory birds and mammals (Newton et al., 1990, 1999a; Berny et al., 1997;

93 McDonald et al., 1998; Shore et al., 1999, 2003a). However, invertebrates can be a
94 route of contamination for insectivorous vertebrates (Spurr and Drew, 1999) and,
95 although exposure of insectivorous birds has been reported (Borst and Couston,
96 2002; Dowding et al., 2006), exposure of insectivorous mammals has not been
97 studied. Potential routes of uptake by invertebrates include: the consumption of
98 rodent faeces (Laas et al., 1985; Craddock, 2002; Eason et al., 2002); the
99 consumption of rodent carcasses; ingestion of soil-bound residues by e.g.
100 earthworms; and direct consumption of poison baits (Spurr and Drew, 1999; Dunlevy
101 et al., 2000; Craddock, 2002). Given that many ecological communities typically
102 contain larger numbers of insectivorous vertebrates relative to predators, the
103 contamination of invertebrates potentially poses the greater risk of non-target
104 poisoning in terms of species and individuals.

105 The European hedgehog (*Erinaceus europaeus*) is a medium-sized (0.8 - 1.2 kg)
106 insectivorous mammal distributed throughout Britain and across Western Europe
107 (Morris and Reeve, 2008). Hedgehogs are of particular interest in terms of exposure
108 to anticoagulant rodenticides, as they are reputed to have declined significantly in the
109 last few decades in Britain, and poisoning by industrial chemicals, including
110 rodenticides, may have been a contributory factor (Battersby and Tracking Mammals
111 Partnership, 2005). Our overall aim in this study was to investigate the scale and
112 severity of exposure of hedgehogs throughout Britain to some of the first-generation
113 (warfarin, coumatetralyl) and all of the second-generation (difenacoum,
114 bromadiolone, brodifacoum, flocoumafen) anticoagulant rodenticides that are
115 licensed for use in Britain; the indandione compounds were not determined using the
116 analytical techniques available to us in this study. The current study is the first to
117 assess anticoagulant rodenticide contamination in Britain of species at this trophic

118 level. Furthermore, we analysed tissue residues using both high performance liquid
119 chromatography coupled with fluorescence detection (hereafter HPLC) and liquid-
120 chromatography mass spectrometry (LCMS). To date, characterisation of exposure
121 of non-target species has mostly used HPLC (for example, McDonald et al., 1998;
122 Shore et al., 2003a, 2006a; Walker et al., 2008) but LCMS is potentially a more
123 sensitive technique and, perhaps more importantly, enables compounds with similar
124 chemical structure to be differentiated with greater confidence since identification is
125 based upon mass rather than elution times. Our specific objectives were: to compare
126 and contrast the (i) frequency of occurrence and (ii) average residue magnitude of
127 FGARs and SGARs in hedgehogs by analysing liver concentrations using both HPLC
128 and LCMS techniques; (iii) to determine whether there were differences in levels of
129 contamination between males and females and between geographical regions; and
130 (iv) on the basis of these results, compare the extent of sub-lethal exposure of
131 hedgehogs in Britain with that of predatory birds and mammals, and assess whether
132 hedgehogs are at risk of acute toxicity from their exposure.

133

134 **2. Materials and Methods**

135

136 During 2004-2006, 20 adult hedgehog carcasses were collected from wildlife
137 rehabilitation hospitals from each of six (Scotland, Wales, Midlands and West, South-
138 Western, South-Eastern, and Eastern) of the seven regions of Britain as defined by
139 the Department for Environment, Food and Rural affairs when assessing rodenticide
140 usage (Dawson et al., 2003); we were unable to obtain samples from the remaining
141 region (Northern England). All 120 hedgehogs used in the study had either died
142 following admission or were euthanased due to their injuries or illness.

143 Each carcass was weighed, sexed and stored at -20°C until dissection, when it
144 was inspected for lesions, injuries or other abnormalities. These observations, along
145 with information collected at admission, were used to determine the cause of death
146 or reason for euthanasia. The whole liver, the primary organ for accumulation of
147 rodenticides (Huckle and Warburton, 1986), was removed, weighed to two decimal
148 places and stored in aluminium foil at -20°C until further analysis.

149

150 *2.1. Residue analyses*

151

152 Anticoagulant rodenticide residues were quantified using both HPLC and LCMS.
153 The four main SGARs licensed for use in the UK (brodifacoum, bromadiolone,
154 difenacoum and flocoumafen) were quantified using both techniques. The two most
155 commonly applied FGARs in the UK, coumatetralyl and warfarin (Dawson and
156 Garthwaite, 2004), were also analysed using LCMS only. All reagents were from
157 Rathburn Chemical Co. Ltd, Walkerburn, Scotland and of a grade suitable for HPLC
158 and LCMS analysis.

159 Extraction procedures for second-generation compounds followed Hunter (1985)
160 and Jones (1996). Samples were analysed in randomised batches of 15. Each liver
161 was defrosted at room temperature and a subsample of approximately 1g (mean wet
162 weight \pm SE=0.98 \pm 0.01g) ground to a homogenous paste using acid-washed furnace-
163 cleaned sand and anhydrous sodium sulphate. A 30ml aliquot of extraction solvent
164 (50:50 acetone/chloroform) was mixed thoroughly with the ground tissue, stood for 1
165 hour, then decanted and collected in a 100ml measuring cylinder through a funnel
166 containing glass wool and anhydrous sodium sulphate. The ground tissue was
167 subsequently washed with 30ml aliquots of extraction solvent and washings were

168 added to the original extraction aliquot until a total volume of 100ml was collected.
169 The mixture was mixed by inversion and left to stand at room temperature for a
170 minimum of 12 hours. Subsequently the extract was divided into 50ml for analysis by
171 HPLC and 30ml was archived at 4°C in the dark for later analysis by LCMS. Both
172 samples were reduced to zero volume by evaporation of solvent in a fume cupboard
173 and the remaining 20ml was poured to waste.

174 The reduced extract was re-dissolved in 1ml of extract solvent and 4ml
175 acetonitrile and cleaned using an SPE Isolute C₁₈ (EC) 1g column (Internation
176 Sorbent Technology, Mid-Glamorgan, UK) connected to an SPE 500-mg NH₂ column
177 solvated with methanol. Columns were conditioned with 5ml methanol followed by
178 5ml acetonitrile. The re-dissolved extract was loaded onto the C₁₈ column and
179 washed with three 5ml aliquots of acetonitrile at <4ml/minute. The C₁₈ column was
180 then removed and 4ml ammoniacal methanol was washed through the NH₂ column
181 (flow <4ml/min). The resulting eluant was combined with 5ml methanol, reduced to
182 near dryness (to remove ammonia) and re-dissolved in 0.5ml methanol. Samples
183 were finally transferred to a chromatography vial via a 4mm syringe filter (Whatman
184 International Ltd, Kent, UK).

185

186 *2.2. High performance liquid chromatography*

187

188 High performance liquid chromatography (HP Series 1100, Agilent Technologies,
189 Bracknell, Berkshire, UK) was performed using a ODS Hypersil 200mm x 4.6mm
190 5µm column (Thermo electron corporation, Runcorn, Cheshire, UK) at 30°C. A 15µl
191 aliquot of cleaned-up extract was injected onto the column using 76:24
192 methanol:water (v/v) supplemented with 0.25% (v/v) acetic acid and 40mM

193 ammonium acetate, as the mobile phase pumped at 1.0ml/min isocratically. SGARs
194 were detected by fluorescence spectrometry (HP 1100 series fluorescence detector)
195 using three excitation wavelengths (313nm, 320nm and 350nm) simultaneously to
196 allow for correction of co-eluting peaks that interfered with the fluorescence of the
197 rodenticides. The emission for each excitation wavelength was measured at 380nm.
198 The excitation wavelength of 313nm gave the greatest emission signal at 380nm and
199 was thus used for quantification. The ratio the emission response elicited by the
200 320nm wavelength to that elicited by 313nm and the ratio elicited by 350nm to that
201 elicited by 313nm were both used to aid identification. A chromatographic peak was
202 identified as a specific SGAR if the ratios of the signals for each excitation
203 wavelength matched the ratios in the standards and if the absolute retention time of
204 the peak fell within the retention time window of the calibration standards.

205

206 *2.3. Liquid chromatography mass spectrometry*

207

208 The archived extraction samples were cleaned using methods previously
209 outlined and analysed by liquid-chromatography tandem mass spectrometry
210 conducted on a Zorbax Eclipse C18 3µm column (150 x 2mm). The analysis was
211 conducted using an isocratic mobile phase consisting of acetonitrile:water containing
212 0.1% formic acid in the ratio 75:25 and at a flow rate of 200µl/min. The column was
213 maintained at 35°C; injection volume was set at 15µl. A Surveyor HPLC system
214 (Thermo Corporation, Hemel Hempstead, Hertfordshire, UK) was used to separate
215 the sample and deliver it to an LCQ Duo, API ion trap mass spectrometer (Thermo
216 Corporation, Hemel Hempstead, Hertfordshire, UK).

217 Analyses were performed using electrospray ionisation in the negative mode.
218 The capillary temperature was set at 270°C with an ionisation voltage of -36.0V. The
219 sheath and auxiliary gasses used were helium and nitrogen maintained at 80psi and
220 20psi respectively. Sensitivity was increased using single ion monitoring, scanning
221 for the molecular ion of each of the rodenticides. Selectivity and conformational
222 analysis was undertaken using tandem mass spectrometry.

223

224 2.4. Quality assurance

225

226 Quantification of residues was carried out by comparison with rodenticide
227 standards (Chemservice, Greyhound Chromatography, Merseyside, UK) for all the
228 FGARS and SGARS that were quantified. For HPLC analysis, the linear calibration
229 range was 50-500ng/ml and the limit of detection (LoD) for peaks identified as
230 SGARs was determined from the linear regression of the multilevel calibration using
231 the equation $Y=Y_0+3S_{y/x}$, where Y is the LoD response, Y_0 is the intercept and $S_{y/x}$ is
232 the standard error of the regression line. The HPLC LoDs for bromadiolone,
233 difenacoum, flocoumafen and brodifacoum based on the standards were 0.03, 0.01,
234 0.01 and 0.02µg respectively, which were analogous to previous analyses of polecat
235 (*Mustela putorius*) livers (Shore et al., 2003a). The LoDs for LCMS were obtained
236 using a similar method and were 0.002µg for all compounds.

237 For LCMS analysis, three concentrations (100, 50 and 10ng/ml) of the standards
238 for all the FGARs and SGARS were run alongside procedural blanks after every eight
239 samples to determine day-to-day quantitation. Calibration curves were obtained
240 using a range of concentrations (500, 400, 200, 100, 50, 20, 10, 5, 1 and 0.1ng/ml) of

241 these standards; the average areas of ten determinations of each standard
242 concentration were used to produce these curves.

243 For both HPLC and LCMS analysis, procedural blanks (reagents only) were
244 analysed alongside samples to detect possible contamination during sample
245 preparation. Chicken liver samples were each spiked with known concentrations of
246 each SGAR and were prepared, stored and analysed in the same way as unknown
247 samples to determine sample matrix recovery and percent recovery data. For HPLC
248 the mean (\pm SE%) recovery, determined from analyses of eight spiked samples, were
249 $108\pm 11.5\%$, $81.6\pm 5.0\%$, $95.2\pm 9.8\%$ and $93.3\pm 9.0\%$ for difenacoum, bromadiolone,
250 flocoumafen and brodifacoum respectively. Corresponding figures for LCMS recovery
251 were $59.2\pm 9.9\%$, $27.3\pm 12.0\%$, $59.2\pm 9.9\%$ and $65.9\pm 7.3\%$, determined from analyses
252 of four samples spiked for each SGAR. The apparently lower recovery associated
253 with LCMS than HPLC may have been an artefact reflecting poor stability of spiked
254 samples when archived. The bromadiolone and difenacoum concentrations in the
255 actual samples of hedgehog livers were not significantly lower when quantified by
256 LCMS than when measured by HPLC (see Results). Concentration data in tissue
257 samples were not recovery-corrected.

258

259 *2.5. Statistical analysis*

260

261 The numbers of samples with detectable and non-detectable rodenticide
262 residues as determined by HPLC and LCMS were compared using Fisher's exact
263 tests. Liver concentrations were not normally distributed and average residue
264 concentrations are given as medians. Median liver concentrations in animals with
265 detectable residues were compared using Mann-Whitney U tests. Wilcoxon matched

266 pairs tests were used to compare residue concentrations detected by the two
267 techniques within the same individual. Binary logistic regression was used to
268 examine the effect of region, batch number and sex on the presence/absence of
269 contamination; batch was included as a factor to confirm that batching samples for
270 analysis did not introduce any analytical biases. All analyses were conducting using
271 SPSS, Release 15.0 (Field, 2005).

272

273 **3. Results**

274

275 Reasons cited by wildlife hospitals for admission of the hedgehogs used in this
276 study were: injury ($n=55$); unknown ($n=46$); natural causes ($n=18$); and suspected
277 poisoning ($n=1$), although this diagnosis was not confirmed clinically or chemically.
278 No obvious signs of haemorrhage other than that associated with trauma were found
279 during *post-mortem* examinations ($n=120$).

280 Using HPLC, detectable liver concentrations of brodifacoum, bromadiolone,
281 difenacoum and flocoumafen were found in four, 13, 16 and zero animals
282 respectively (Table 1); in total, SGARs were detected in 27 individuals (23% of the
283 animals analysed: Table 2). In contrast, SGARs were detected in 69 (57.5%)
284 hedgehogs when the analysis was conducted by LCMS (Table 2). FGARs (only
285 determined by LCMS) were detected in 27 (22.5%) animals (Table 2). Overall,
286 residues of at least one FGAR or SGAR were detected in two thirds of hedgehogs
287 when samples were analysed by LCMS. Fifty-three (44%) individuals had liver
288 residues of one compound; 21 (18%), five (4%) and one (1%) animal contained
289 residues of two, three and four compounds respectively.

290 The greater frequency of detection of SGARs by LCMS than HPLC was largely
291 because more instances of difenacoum and bromadiolone contamination were
292 detected by LCMS (Table 2); the difference in frequency of detection between the
293 analytical methods was significant for difenacoum (Fisher's Exact test, $P < 0.001$) and
294 approached significance for bromadiolone (two-tailed Fisher's Exact test, $P = 0.10$).
295 Much of this higher frequency of detection was due to the greater sensitivity of the
296 LCMS. Liver difenacoum and bromadiolone concentrations below $0.025 \mu\text{g/g}$ wet
297 weight (ww) and $0.05 \mu\text{g/g}$ ww, respectively, were not detected by HPLC, whereas
298 these concentrations comprised 25-50% of the LCMS detections for these
299 compounds (Fig. 1). Overall, detection of these low level difenacoum and
300 bromadiolone residues by LCMS accounted for an extra 30 hedgehogs (25% of the
301 sample) being identified as containing rodenticide.

302 The average magnitude of residues (Table 3), not just the frequency of
303 occurrence, also varied with analytical technique. When only hedgehogs with HPLC
304 and/or LCMS detectable residues were included in the statistical analysis, the
305 median liver bromadiolone concentration was lower when determined by LCMS than
306 by HPLC (Mann Whitney U test: $U = 61.0$, $n_1 = 23$, $n_2 = 13$, $P < 0.01$; Fig. 2). This reflected
307 the presence of low-level bromadiolone concentrations (typically $< 0.1 \mu\text{g/g}$ ww; Fig.
308 1) that were detected by LCMS but not by HPLC (and so were not included in the
309 HPLC dataset of animals with detected residues). When the statistical analysis was
310 further restricted to a matched pair comparison of just animals with bromadiolone
311 residues detected by *both* analytical methods, there was no significant difference
312 between LCMS and HPLC measurements (Wilcoxon matched pairs test: $n = 10$, $Z = -$
313 0.663 , $P > 0.05$). This again suggested that differences between HPLC- and LCMS-
314 determined measurements were solely due to detection of low-level concentrations

315 by LCMS. However, this was not true for difenacoum. Median liver concentrations of
316 difenacoum in animals with detectable residues did not differ with the method of
317 determination ($U=427.5$, $n_1=16$, $n_2=57$, $P>0.05$), despite the presence of a relatively
318 large number of low-level difenacoum residues in the LCMS sample (Fig. 1). This
319 may reflect differential responses (involving enhancement or quenching of response)
320 of the two techniques, as matched-pair analysis indicated that residues were higher
321 in animals when measured by LCMS ($n=9$, $Z=-2.429$, $P<0.05$).

322 Analyses of potential differences in residue magnitude with sex and region were
323 based on LCMS data. Geographical region was not significantly associated with the
324 presence/absence of (i) FGARs (coumatetralyl and warfarin), (ii) bromadiolone and
325 difenacoum combined (the most commonly found SGARs), (iii) all four SGARs, or (iv)
326 all FGARs and SGARs combined (Table 4). Sex did, however, approach significance
327 in two of the four models (bromadiolone and difenacoum combined, $P=0.052$; all
328 SGARs, $P=0.072$; Table 4), with a greater frequency of occurrence of contamination
329 in males than females.

330

331 **4. Discussion**

332

333 The major proportion of hedgehog diet consists of invertebrates, particularly
334 molluscs, beetles and earthworms (Wroot, 1984). Invertebrates have different blood-
335 clotting mechanisms to vertebrates and so are less susceptible to anticoagulant
336 rodenticides than birds and mammals (Shirer, 1992; Pain et al., 2000; Craddock,
337 2002; Johnston et al., 2005). However, ground-dwelling invertebrates can access
338 and feed on rodenticides, including those placed in bait stations (Spurr and Drew,
339 1999; Dunlevy et al., 2000; Craddock, 2002), and retain ingested compound in their

340 bodies for four weeks or longer (Booth et al., 2001; Craddock, 2002). Additional
341 exposure of invertebrates to rodenticides may also arise through ingesting
342 contaminated soil (where baits have not been protected or have been displaced or
343 removed from bait stations), rodent food caches and rodent carcasses. Thus,
344 predation of contaminated invertebrates is likely to be a major pathway by which
345 hedgehogs are exposed to anticoagulant rodenticides. However, hedgehogs will
346 consume small mammal carcasses if they are available (Yalden, 1976) and may also
347 access spilt, cached or unprotected baits directly, and these may be alternative
348 secondary and primary exposure routes.

349 Whatever the route of exposure, it is clear from our results that contamination of
350 hedgehogs with anticoagulant rodenticides is commonplace. These compounds may
351 therefore similarly pose a risk to other species at the same trophic level, such as
352 insectivorous birds (Rammell et al., 1984; Empson and Miskelly, 1999; Robertson
353 and Colbourne, 2001). The frequencies with which we detected SGAR residues by
354 HPLC were towards the mid (brodifacoum, bromadiolone) or low (difenacoum) end of
355 the spectrum documented for predatory birds and mammals in Britain (Table 1), but
356 were comparable in some instances to prevalence rates in species considered to be
357 specialist predators of small mammals, such as the polecat (Shore et al., 2003a),
358 barn owl (*Tyto alba*) (Newton et al., 1999b) and tawny owl (*Strix aluco*) (Walker et al.,
359 2008). Likewise, the magnitudes of residues were also broadly similar to those
360 measured in predatory birds and mammals in Britain (Table 3). Thus, hedgehogs in
361 Britain appear to be at similar risk of exposure and effects from anticoagulant
362 rodenticides as non-target predatory birds and mammals.

363 Our data also suggest that exposure of hedgehogs is geographically widespread.
364 The absence of any significant difference between the proportion of individuals with

365 residues and region indicates that the scale of exposure of hedgehogs does not vary
366 markedly across Britain, consistent with studies of polecats (Shore et al., 2003a),
367 even though the apparent use of rodenticides in arable regions varies geographically
368 (Dawson et al., 2003). In part, however, the likelihood of detecting correlated patterns
369 between prevalence rates in animals and regional patterns of use will be affected by
370 exactly which specific compounds are used. This is because compounds, and
371 particularly FGARs and SGARs, differ in their biological half-life and toxicity (Eason
372 et al., 2002). Furthermore, geographical variation in arable use of rodenticides is
373 unlikely to be of relevance to those animals that were from urban areas. There are no
374 published data for rodenticide use in urban areas in Britain and so it is not possible to
375 assess how urban use may relate to exposure of hedgehogs. Finally, our finding that
376 male hedgehogs tended to be more likely to accumulate rodenticides than females
377 may also have a spatial, albeit small scale, explanation. Males have a greater
378 ranging behaviour than females (Reeve, 1994) and this is likely to increase the
379 likelihood of individuals finding baits and contaminated forage.

380 The overall similarity between hedgehogs and specialist avian and mammalian
381 predators of small mammals was unexpected. This may simply indicate that
382 secondary exposure is more common than previously anticipated for food chains in
383 which small mammals are not a major component. However, this similarity may mask
384 other factors, such as differences in the likely exposure of non-target species in
385 urban and rural areas. We had no information on the exact location in which our
386 hedgehogs were found. Our reliance on analysing the carcasses of animals admitted
387 to wildlife hospitals may have biased the sample towards urban areas because their
388 relatively high human population density may mean that sick/injured hedgehogs are
389 more likely to be found. In contrast, most UK studies on secondary exposure in

390 predatory birds and mammals have analysed animals that are predominantly from
391 rural areas. It is not clear whether an urban-biased sample would tend to increase or
392 decrease the likelihood of detecting exposure. Rodenticides are widely used on
393 farms in rural Britain but are also commonly used throughout urban and suburban
394 landscapes by both professional practitioners and the general public. The density of
395 baits and contaminated prey relative to population numbers of non-target species in
396 rural and urban areas is completely unknown. Furthermore, it is possible that
397 hedgehogs may be particularly susceptible to exposure in urban areas where
398 untrained domestic users may be prone to unintentional misuse. Animals may also
399 be more likely to suffer traumatic injuries in human-dominated habitats through
400 collisions with motor vehicles or injuries arising from misadventure (Reeve and
401 Huijser, 1999). If such injuries occur independently of levels of rodenticide uptake,
402 such a sample would give a reliable indication of levels of sub-lethal contamination in
403 those areas, but if rodenticide uptake increases the likelihood of injury (Fournier-
404 Chambrillon et al., 2004), then urban samples in particular may over-estimate
405 exposure rates. Comparison of exposure rates of hedgehogs or other species from
406 known urban and rural locations is merited.

407 The analysis of our sample of hedgehog tissues using LCMS as well as HPLC
408 has shown that exposure, particularly low-level exposure, is markedly
409 underestimated by HPLC. The proportion of hedgehogs exposed to SGARs
410 increased by two- to three-fold when the analysis was conducted by LCMS. We
411 postulate that current estimates of the exposure of predatory birds and mammals to
412 SGARs have been similarly under-estimated where they have been determined using
413 HPLC measurements.

414 Although exposure of hedgehogs to anticoagulants may be widespread, there is
415 no evidence from our study that this commonly causes lethal poisoning. The *post*
416 *mortem* examination of the animals in our study did not identify any instances of
417 haemorrhage that appeared consistent with rodenticide poisoning. Although there is
418 no precise liver concentration in hedgehogs or other species that is diagnostic of
419 lethal poisoning, SGAR residues in excess of 0.2µg/g ww are considered to be of
420 concern in barn owls (Newton et al., 1999a) and residues of >1µg/g ww are generally
421 considered to be very high. Irrespective of the measurement technique in our study,
422 the percentage of hedgehogs with summed SGAR residues above 0.2µg/g ww and
423 1µg/g ww was <11% and <5% respectively. The detection of liver residues exceeding
424 1µg/g ww suggests that lethal poisoning by rodenticides is likely to occur in some
425 hedgehogs, but the lack of haemorrhaging and relatively low magnitude of most
426 residues suggests that, for animals in our study, contamination with rodenticides was
427 generally not a contributory factor in their admission to wildlife hospitals. Overall,
428 however, poisoning of non-target wild animals by anticoagulant rodenticides is
429 difficult to monitor and studies such as ours may underestimate poisoning events
430 because animals with fatal doses may become lethargic some hours before death
431 and die in cryptic locations (Newton et al., 1999a). Furthermore, there is a general
432 lack of knowledge about whether sub-lethal exposure, as appears to be common in
433 hedgehogs, may be associated with any sub-lethal impacts or an increased
434 susceptibility to toxicity following repeated exposures.

435

436 **5. Conclusion**

437 This study has shown that the European hedgehog, an insectivorous species,
438 has similar rates of exposure (judged from the proportion of animals with HPLC-

439 detected liver concentrations and the size of those residues) to those of specialist
440 predators of small mammals. Given that hedgehogs only rarely eat rodents, these
441 results indicate that anticoagulant rodenticides are finding their way into ecosystems
442 via transfer pathways other than through consumption of contaminated rodents.
443 Furthermore, our data indicate that analysis of samples using LCMS can increase the
444 estimate of exposure by two- to three-fold, largely through the detection of low-level
445 residues, and that the use of HPLC may have markedly under-estimated the true
446 scale of exposure of other non-target species to anticoagulant rodenticides.

447

448 **Acknowledgements**

449

450 We thank the RSPCA Westhatch, RSPCA Stapeley Grange, RSPCA East
451 Winch, RSPCA Mallydams Wood, The Gower Bird Hospital, St Tiggywinkles and
452 Hessilhead Wildlife Rescue Trust for supplying hedgehog carcasses and the
453 Dulverton Trust (C.V. Dowding and S. Harris) for financial support .

454

455 **References**

456

457 Barnett, E.A., Fletcher, M.R., Hunter, K., Sharp, E.A. 2006. Pesticide poisoning of
458 animals in 2005: investigations of suspected incidents in the United Kingdom.
459 Pesticide Services Division, Department for Environment, Food and Rural Affairs,
460 London, UK.
461 Battersby, J. (Ed.), Tracking Mammals Partnership 2005. UK mammals: species
462 status and population trends. First report by the Tracking Mammals Partnership.

463 Joint Nature Conservation Committee/Tracking Mammals Partnership,
464 Peterborough, UK.

465 Berny, P.J., Buronfosse, T., Buronfosse, F., Lamarque, F., Lorgue, G. 1997. Field
466 evidence of secondary poisoning of foxes (*Vulpes vulpes*) and buzzards (*Buteo*
467 *buteo*) by bromadiolone, a 4-year survey. *Chemosphere* 35, 1817-1829.

468 Booth, L.H., Eason, C.T., Spurr, E.B. 2001. Literature review of the acute toxicity and
469 persistence of brodifacoum to invertebrates. *Science for Conservation* 177, 1-9.

470 Borst, G.H.A., Counotte, G.H.M. 2002. Shortfalls using second-generation
471 anticoagulant rodenticides. *Journal of Zoo and Wildlife Medicine* 33, 85.

472 Carter, I., Burn, A. 2000. Problems with rodenticides: the threat to red kites and other
473 wildlife. *British Wildlife* 11, 192-197.

474 Cowan, D., Dunsford, G., Gill, E., Jones, A., Kerins, G., MacNicoll, A., Quay, R. 1995.
475 The impact of resistance on the use of second-generation anticoagulants against
476 rats on farms in southern England. *Pesticide Science* 43, 83-93.

477 Craddock, P. 2002. Aspects of the ecology of forest invertebrates and the use of
478 brodifacoum. PhD thesis, University of Auckland, New Zealand.

479 Dawson, A., Garthwaite, D. 2004. Pesticide usage survey report 185: rodenticide
480 usage by local authorities in Great Britain 2001. Department for Environment,
481 Food and Rural Affairs, London, UK.

482 Dawson, A., Bankes, J., Garthwaite, D. 2003. Pesticide usage survey report 175:
483 rodenticide usage on farms in Great Britain growing arable crops 2000. Department for
484 Environment, Food and Rural Affairs, London, UK.

485 Dowding, J.E., Lovegrove, T.G., Ritchie, J., Kast, S.N., Puckett, M. 2006. Mortality of
486 northern New Zealand dotterels (*Charadrius obscurus aquilonius*) following an aerial
487 poisoning operation. *Notornis* 53, 235-239.

488 Dunlevy, P.A., Campbell, E. W., Lindsey, G.D. 2000. Broadcast application of a
489 placebo rodenticide bait in a native Hawaiian forest. *International Biodeterioration*
490 & *Biodegradation* 45, 199-208.

491 Eason, C.T., Murphy, E.C., Wright, G.R.G., Spurr, E.B. 2002. Assessment of risks of
492 brodifacoum to non-target birds and mammals in New Zealand. *Ecotoxicology* 11,
493 35-48.

494 Empson, R.A., Miskelly, C.M. 1999. The risks, costs and benefits of using
495 brodifacoum to eradicate rats from Kapiti Island, New Zealand. *New Zealand*
496 *Journal of Ecology* 23, 241-254.

497 Field, A. P. 2005. *Discovering statistics using SPSS*. SAGE Publications, London,UK.

498 Fournier-Chambrillon, C., Berny, P.J., Coiffier, O., Barbedienne, P., Dassé, B., Delas,
499 G., Galineau, H., Mazet, A., Pouzenc, P., Rosoux, R., Fournier, P. 2004.
500 Evidence of secondary poisoning of free-ranging riparian mustelids by
501 anticoagulant rodenticides in France: implications for conservation of European
502 mink (*Mustela lutreola*). *Journal of Wildlife Diseases* 40, 688-695.

503 Huckle, K.R., Warburton, P.A. 1986. Elimination, metabolism and disposition of ¹⁴C-
504 WL 108366 in the Fischer 344 rat following repeated oral administration.
505 Sittingbourne Research Centre, Shell Research Ltd, Sittingbourne, Kent, UK.

506 Hunter, K. 1985. High-performance liquid chromatographic strategies for the
507 determination and confirmation of anticoagulant rodenticide residues in animal
508 tissues. *Journal of Chromatography* 321, 255-272.

509 Johnston, J.J., Pitt, W.C., Sugihara, R.T., Eisemann, J.D., Primus, T.M., Holmes,
510 M.J., Crocker, J., Hart, A. 2005. Probabilistic risk assessment for snails, slugs,
511 and endangered honeycreepers in diphacinone rodenticide baited areas on
512 Hawaii, USA. *Environmental Toxicology and Chemistry* 24, 1557-1567.

513 Jones, A. 1996. HPLC determination of anticoagulant rodenticide residues in animal
514 livers. *Bulletin of Environmental Contamination and Toxicology* 56, 8-15.

515 Laas, F.J., Forss, D.A., Godfrey, M.E.R. 1985. Retention of brodifacoum in sheep
516 tissues and excretion in faeces. *New Zealand Journal of Agricultural Research*
517 28, 357-359.

518 McDonald, R.A., Harris, S., Turnbull, G., Brown, P., Fletcher, M. 1998. Anticoagulant
519 rodenticides in stoats (*Mustela erminea*) and weasels (*Mustela nivalis*) in
520 England. *Environmental Pollution* 103, 17-23.

521 Morris, P.A., Reeve, N.J. 2008. Hedgehog *Erinaceus europaeus*. In: Harris, S.,
522 Yalden, D.W. (Eds.), *Mammals of the British Isles: Handbook*. Fourth ed. The
523 Mammal Society, Southampton, UK, pp. 241-249.

524 Newton, I., Wyllie, I., Freestone, P. 1990. Rodenticides in British barn owls.
525 *Environmental Pollution* 68, 101-117.

526 Newton, I., Shore, R.F., Wyllie, I., Birks, J.D.S., Dale, L. 1999a. Empirical evidence of
527 side-effects of rodenticides on some predatory birds and mammals. In: Cowan,
528 D.P., Feare, C.J. (Eds.), *Advances in vertebrate pest management*. Filander,
529 Fürth, Germany, pp. 347-367.

530 Newton, I., Dale, L., Finnie, J.K., Freestone, P., Wright, J., Wyatt, C., Wyllie, I. 1999b.
531 *Wildlife and pollution: 1997/98 annual report*. Joint Nature Conservation
532 Committee Report No. 285, Peterborough, UK.

533 Pain, D.J., Brooke, M. de L., Finnie, J.K., Jackson, A. 2000. Effects of brodifacoum
534 on the land crab of Ascension island. *Journal of Wildlife Management* 64, 380-
535 387.

536 Parmar, G., Bratt, H., Moore, R., Batten, P.L. 1987. Evidence for common binding
537 site in vivo for the retention of anticoagulants in rat liver. *Human Toxicology* 6,
538 431-432.

539 Rammell, C.G., Hoogenboom, J.J.L., Cotter, M., Williams, J.M., Bell, J. 1984.
540 Brodifacoum residues in target and non-target animals following rabbit poisoning
541 trials. *New Zealand Journal of Experimental Agriculture* 12, 107-111.

542 Reeve, N. 1994. *Hedgehogs*. Poyser, London, UK.

543 Reeve, N.J., Huijser, M.P. 1999. Mortality factors affecting wild hedgehogs: a study
544 of records from wildlife rescue centres. *Lutra* 42, 7-24.

545 Robertson, H.A., Colbourne, R.M. 2001. Survival of little spotted kiwi exposed to the
546 rodenticide brodifacoum. *Journal of Wildlife Management* 65, 29-34.

547 Shirer, M. 1992. In poison's defence. *Terra Nova* 17, 3.

548 Shore, R.F., Birks, J.D.S., Freestone, P. 1999. Exposure of non-target vertebrates to
549 second-generation rodenticides in Britain, with particular reference to the polecat
550 *Mustela putorius*. *New Zealand Journal of Ecology* 23, 199-206.

551 Shore, R.F., Afsar, A., Horne, J.A., Wright, J. 2000. Rodenticide and lead
552 concentrations in red kites (*Milvus milvus*). Centre for Ecology and Hydrology
553 Contract Report for English Nature, Peterborough, UK.

554 Shore, R.F., Birks, J.D.S., Afsar, A., Wienburg, C.L., Kitchener, A.C. 2003a. Spatial
555 and temporal analysis of second-generation anticoagulant rodenticide residues in
556 polecats (*Mustela putorius*) from throughout their range in Britain, 1992-1999.
557 *Environmental Pollution* 122, 183-193.

558 Shore, R.F., Fletcher, M.R., Walker, L.A. 2003b. Agricultural pesticides and
559 mammals in Britain. In: Tattersall, F., Manley, W. (Eds.), *Conservation and*

560 conflict: mammals and farming in Britain. Westbury Publishing, Otley, UK, pp. 37-
561 50.

562 Shore, R.F., Malcom, H.M., McLennan, D., Turk, A., Walker, L.A., Wienburg, C.L.,
563 Burn, A.J. 2006a. Did foot-and-mouth disease-control operations affect
564 rodenticide exposure in raptors? *Journal of Wildlife Management* 70, 588-593.

565 Shore, R.F., Walker, L.A., Turk, A., Wienburg, C.L., Wright, J., Murk, A., Wanless, S.
566 2006b. *Wildlife and pollution: 2003/04 annual report*. Joint Nature Conservation
567 Committee Report No. 391, Peterborough, UK.

568 Singleton, G.R., Hinds, L.A., Leirs, H., Zhang, Z. 1999. Ecologically-based
569 management of rodent pests. Australian Centre for Agricultural Research,
570 Canberra, Australia.

571 Spurr, E.B., Drew, K.W. 1999. Invertebrates feeding on baits used for vertebrate pest
572 control in New Zealand. *New Zealand Journal of Ecology* 23, 167-173.

573 Stenseth, N.C., Leirs, H., Skonhott, A., Davis, S.A., Pech, R.P., Andreassen, H.P.,
574 Singleton, G.R., Lima, M., Machang'u, R.S., Makundi, R.H., Zhang, Z., Brown,
575 P.R., Shi, D., Wan, X. 2003. Mice, rats, and people: the bio-economics of
576 agricultural rodent pests. *Frontiers in Ecology and the Environment* 1, 367-375.

577 Thijssen, H.H.W. 1995. Warfarin-based rodenticides: mode of action and mechanism
578 of resistance. *Pesticide Science* 43, 73-78.

579 Walker, L.A., Turk, A., Long, S.M., Wienburg, C.L., Best, J., Shore, R.F. 2008.
580 Second generation anticoagulant rodenticides in tawny owls (*Strix aluco*) from
581 Great Britain. *Science of the Total Environment* 392, 93-98.

582 Wroot, A.J. 1984. Feeding ecology of the European hedgehog *Erinaceus europaeus*.
583 PhD thesis, Royal Holloway College, University of London, UK.

584 Yalden, D.W. 1976. The food of the hedgehog in England. *Acta Theriologica* 30, 401-
585 424.
586

587 **Figure legends**

588

589 Fig. 1. Frequency distribution of bromadiolone and difenacoum liver concentrations in
590 hedgehogs detected by HPLC and LCMS.

591

592 Fig. 2. Median and interquartile ranges of liver concentrations of first- and second-
593 generation anticoagulant rodenticides in hedgehogs with detectable residues as
594 quantified using HPLC and LCMS. Sample sizes are given in Table 2.

595

596

597 Table 1

598 Percentage occurrence of the residues of the first-generation anticoagulant
 599 rodenticide coumatetralyl (coum) and the second-generation anticoagulant
 600 rodenticides brodifacoum (brod), bromadiolone (brom), difenacoum (difen) and
 601 flocoumafen (floc) in the livers of predatory birds and mammals in British wildlife as
 602 identified using high performance liquid chromatography. ND indicates residue not
 603 detected; - indicates chemical was not investigated
 604

Species	n	Coum	Brod	Brom	Difen	Floc	Total^a	Ref^b
Hedgehog (<i>Erinaceus europaeus</i>)	120	-	3.3	10.8	13.3	ND	22.5	1
Polecat (<i>Mustela putorius</i>)	100	-	3.0	12.0	22.0	ND	36.0	2
Stoat (<i>Mustela erminea</i>)	40	15.0	2.5	6.7	-	-	22.5	3
Weasel (<i>Mustela nivalis</i>)	10	30.0	-	10.0	-	-	30.0	3
Red fox (<i>Vulpes vulpes</i>)	92	7.6	5.4	26.1	16.3	-	45.7	4
Barn owl (<i>Tyto alba</i>)	717	-	3.9	11.0	16.7	1.1	26.1	5
Barn owl (<i>Tyto alba</i>)	52	-	5.8	28.8	30.8	ND	42.3	6
Buzzard (<i>Buteo buteo</i>)	40	-	2.5	5.0	32.5	2.5	37.5	6
Tawny owl (<i>Strix aluco</i>)	172	-	4.7	11.6	5.8	ND	19.2	7
Red kite (<i>Milvus milvus</i>)	20	-	-	-	-	-	70.0	8
Kestrel (<i>Falco tinnunculus</i>)	36	-	-	-	-	-	67.0	8
Kestrel (<i>Falco tinnunculus</i>)	40	-	15.0	40.0	72.5	ND	84.6	9

605 ^a Total percentage of individuals positive for one or more chemicals. ^b Reference: 1 - present study; 2 - Shore et
 606 al. (2003a); 3 - McDonald et al. (1998); 4 - Shore et al. (2003b); 5 - Newton et al. (1999b); 6 - Shore et al. (2006a);
 607 7 - Walker et al. (2008), 8 - Shore et al. (2000); 9 - Shore et al. (2006b).

608 Table 2
 609 Number and percentage (out of sample of 120) of hedgehogs with first- (FGAR) and
 610 second-generation anticoagulant rodenticides (SGAR) detected using high
 611 performance liquid chromatography (HPLC) and liquid-chromatography mass
 612 spectrometry (LCMS)
 613

	Hedgehogs with residues detected by			
	HPLC		LCMS	
	%	<i>n</i>	%	<i>n</i>
Coumatetralyl (FGAR)			14.2	17
Warfarin (FGAR)			8.3	10
Brodifacoum (SGAR)	3.3	4	5.0	6
Bromadiolone (SGAR)	10.8	13	19.2	23
Difenacoum (SGAR)	13.3	16	47.5	57
Flocoumafen (SGAR)	0	0	0.8	1
Total SGARs only	22.5	27	57.5	69
Total FGARs and SGARs	-		66.7	80

614 Coumatetralyl and warfarin only determined using LCMS

615

616

617 Table 3
 618 Mean \pm SE (*n*) concentration ($\mu\text{g/g ww}$) of second-generation anticoagulant
 619 rodenticide residues in British wildlife identified using high performance liquid
 620 chromatography. Figures are the concentrations only for those animals where
 621 residue was detected
 622

Species ^a	<i>n</i>	Brodifacoum	Bromadiolone	Difenacoum	Ref ^b
Hedgehog	120	0.05 \pm <0.01 (4)	0.59 \pm 0.24 (13)	0.10 \pm 0.03 (16)	1
Polecat	50	0.06 \pm 0.01 (3)	0.12 \pm 0.03 (12)	0.30 \pm 0.07 (22)	2
Stoat	9	0.12	0.20 \pm 0.10 (3)	-	3
Weasel	3	-	0.25 (1)	-	3
Barn owl	88	0.02 \pm <0.01 (9)	0.09 \pm 0.02 (23)	0.03 \pm 0.01 (35)	4
Kestrel	40	0.08 \pm 0.03 (6)	0.18 \pm 0.04 (16)	0.08 \pm 0.02 (29)	4
Red kite	8	0.35 \pm 0.22 (5)	0.11 \pm 0.01 (3)	0.20 (1)	5
Tawny owl	172	0.25 \pm 0.14 (8)	0.21 \pm 0.05 (20)	0.06 \pm 0.02 (10)	6

623 ^a For Latin names, see Table 1; ^b reference: 1 present study; 2 - Shore et al. (2003a); 3 - McDonald et al. (1998);
 624 4 - Shore et al. (2006b); 5 - Carter and Burn (2000), 6 - RF Shore (unpubl. data).
 625

626

627 Table 4

628 Binary logistic regression models examining the relationship between region, batch
 629 and sex and the presence/absence of (a) first-generation anticoagulant rodenticides
 630 (coumatetralyl and warfarin), (b) the second-generation anticoagulant rodenticides
 631 bromadiolone and difenacoum, (c) all second-generation anticoagulant rodenticides
 632 (brodifacoum, bromadiolone, difenacoum and flocoumafen) and (d) all first- and
 633 second-generation anticoagulant rodenticides in hedgehogs from across Britain
 634 ($n=120$)

635


Model	Variable	B	S.E.	Wald	d.f.	P
a	Batch			3.426	8	0.905
	Region			6.885	5	0.226
	Sex	0.154	0.506	0.092	1	0.762
	Constant	-1.997	1.255	2.531	1	0.112
b	Batch			5.494	8	0.704
	Region			4.353	5	0.500
	Sex	0.831	0.428	3.775	1	0.052
	Constant	-1.130	0.951	1.412	1	0.235
c	Batch			2.821	8	0.945
	Region			3.474	5	0.627
	Sex	0.760	0.423	3.233	1	0.072
	Constant	-0.561	0.931	0.363	1	0.547
d	Batch			1.182	8	0.997
	Region			3.602	5	0.608
	Sex	0.730	0.450	2.635	1	0.105
	Constant	0.388	0.985	0.155	1	0.693

636 Male:female ratio for hedgehogs from different regions were: South-Eastern 12:8; South-Western
 637 15:5; Eastern 11:9; Midlands and West 8:12; Wales 12:8; Scotland 7:13

638

639 Fig. 1


640



641

642 Fig. 2

643



644

645

646