

# Stature Estimation in a Central Patagonian Prehispanic Population: Development of New Models Considering Specific Body Proportions

M. BÉGUELIN\*

*División Antropología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, La Plata (1900), Argentina*

**ABSTRACT** Stature estimation of individuals from extinct human populations is a classic topic in anthropology. The estimations, using regression formulae generated from different reference samples, display different results. This fact is related to inter-populational differences in body proportions, which is a phenotypic trait mainly correlated with climatic parameters. The aim of this paper is to address the problem of stature estimation of an archaeological skeletal sample from Patagonia – a region for which there are no specific models available – using different methods and considering differences in body proportions between reference and target populations. The sample used in this analysis is composed of 35 Late Holocene adults of both sexes recovered in central Patagonia (Argentina). The stature of each individual was first reconstructed using the anatomical method [Fully G. 1956. Une nouvelle méthode de détermination de la taille. *Annales Medicine Legale* 35: 266–273], which has no assumptions on body proportions. The results were compared with estimations based on 32 different regression formulae [Trotter M, Gleser G. 1958. A re-evaluation of estimation of stature based on measurements taken during life and the long bones after death. *American Journal of Physical Anthropology* 16: 79–124. 10.1002/ajpa.1330160106] and three femur/stature ratios [Feldesman MR, Fountain RL. 1996. Race specificity and the femur/stature ratio. *American Journal of Physical Anthropology* 100: 207–224. 10.1002/(SICI)1096-8644(199606)]. The average reconstructed stature was 160.8 cm for females (95% confidence band = 155.6–166.2 cm), and 170.5 cm for males (95% confidence band = 168.8–172.2 cm). Most of the comparisons of the regression formulae and femur/stature ratios showed significant differences, which are explained by differences in body proportions between the Patagonian sample and the ones chosen as reference. Finally, a set of new equations was developed using simple regression techniques. It is suggested that whenever possible, population-specific formulae should be used in archaeological studies. In any other situation, the choice of a reference population should be made by taking into account its geographic (latitudinal) provenance. Copyright © 2009 John Wiley & Sons, Ltd.

*Key words:* stature estimation; body proportions; extinct hunter-gatherers; Patagonia

## Introduction

The estimation of living stature from human skeletal remains has major relevance to biological anthropology. It provides important information regarding issues such as health and nutrition, sexual dimorphism and general trends in body size (Stini, 1969; Tanner, 1988; Bush, 1991; Auerbach & Ruff, 2004). Beginning in the 19th century (Rollet, 1888 in Telkkä, 1950), physical

anthropologists have developed several stature estimation methods based on different segments of the human body, mostly single long bones, using skeletal samples from different parts of the world (Trotter & Gleser, 1952; Genovés, 1967; Olivier, 1976; Feldesman *et al.*, 1990; Sciulli & Giesen, 1993; Feldesman & Fountain, 1996; Formicola & Franceschi, 1996; Hens *et al.*, 2000; Özaslan *et al.*, 2003; Işcan, 2005 and the literature cited therein; El-Meligy *et al.*, 2005; Hauser *et al.*, 2005; Kondo *et al.*, 2005; Celbis & Agritmis, 2006).

It has been stated that formulae (i.e. regression equations intended to predict stature from long bones) are population specific, subsequently raising caution about the applicability of such models to individuals

\* Correspondence to: División Antropología, Museo de La Plata, Universidad Nacional de La Plata, Paseo del Bosque s/n La Plata (1900), Argentina.  
e-mail: mbeguelin@fcnym.unlp.edu.ar

drawn from populations different from the reference one (Pearson, 1899; Telkkä, 1950). The estimation methods based upon one or two long bones, i.e. regression methods (Trotter & Gleser, 1958) and femur/stature ratio (Feldesman & Fountain, 1996) work under the assumption that the individuals of both the targeted and the reference populations have the same long bone-stature ratio. Nevertheless, this assumption is not always true or it cannot be reliably corroborated, such is the case of extinct populations. It is virtually impossible in most cases to have population-specific standards, particularly when ancient populations are the research target. This is an omnipresent problem. In fact, the general lack of comparative skeletal samples of known living statures is a major obstacle found in developing reliable stature estimation methods useful for the study of ancient populations (Işcan, 2005). However, this is a problem that can be, if not totally circumvented, at least controlled if careful attention is paid to the differential body proportions that populations display. The use of universal standards to estimate any biological parameter is progressively discarded when local values are known. Thus, it becomes necessary to approximate the living stature with the least error possible by applying population-specific models instead of universal standards or those from populations completely different from the one being studied.

Although body proportions vary in a spatiotemporal pattern as a function of climate (Ruff, 2002) and that this variation has an effect on the estimation of stature, there is a lack of stature estimation models in many geographic regions of the world. Southern South America is one of these regions. As a consequence, population-specific regression equations are often used to estimate stature of individuals whose body proportions may be significantly different (Barrientos, 1997; Onaha *et al.*, 2002; Méndez *et al.*, 2003). This, in turn, can have ulterior implications if it is transferred to further analysis of other models (i.e. body mass assessment, biomass, population density, etc.).

The aim of this study is to develop, evaluate and discuss stature estimation formulae of adult individuals from archaeological samples in central Patagonia, belonging to Late Holocene populations with no present representatives. The stature is anatomically reconstructed (Fully, 1956) in a central Patagonian sample. Then, the application of different existing techniques of stature estimation is assessed and compared, taking the anatomically reconstructed stature as a reference. Finally, new stature estimation equations are presented from the values drawn upon this sample.

## Materials and methods

### Samples

The skeletal material comes from central Patagonia burials and belongs to hunter-gatherer Late Holocene populations (*ca.* 2500–400 BP). The individuals come from two close geographical regions. The archaeological locality named Sierra Colorada (SAC) lies on the Salitroso Lake Basin (about 30 km west of the eastern slopes of the Andes) ( $47^{\circ}25'S$ ,  $71^{\circ}29'W$ ) in the province of Santa Cruz in southwestern Argentina (Figure 1) (Goñi & Barrientos, 2004). The sample from SAC used in this work is restricted to those adult individuals whose remains were complete enough to perform the anatomical method outlined below. This selection resulted in a sub-sample of 16 individuals (i.e. 10 males and 6 females). The second sub-sample, housed at the División Antropología of the Museo de La Plata in Buenos Aires, Argentina, was collected in



Figure 1. Geographic localisation of the samples analysed.

the last decades of the 19th century by various museum expeditions (Lehmann-Nitsche, 1911) in the lower valley of the Chubut River, in the Chubut Province of southeastern Argentina ( $43^{\circ}15'S$ ,  $65^{\circ}18'W$ ) (Figure 1). The sample consists of 17 male and 2 female complete adult skeletons.

The human osteological collection of the Museo de la Plata is the largest and most important in the country, especially for its post-cranial Patagonian samples. Regretfully, complete and reliable field notes from the early 19th century excavations are virtually non-existent. Therefore, the information contained within them is considered within the appropriate context. Despite these restrictions, it is possible to research this collection using more modern diagnostic tools and methods.

Since the early 20th century a number of studies have examined the limited samples of human remains recovered from sites in central and southern Patagonia (Guichón, 2000). Although in the last few years the quantity of artefactual evidence found in Patagonia has increased, the discovery of human skeletal remains has not increased accordingly. Thus, the difficulty is not the quantity of recovered elements but more related to issues of taphonomy, archaeological visibility and population density (Barrientos, 2002). Most human burials are isolated findings and/or contain highly fragmentary bones with pitiful preservation. Subsequently, such limited remains are often quite inadequate for the kind of studies performed in this work (Prieto, 1991, 1993–1994; Salceda *et al.*, 1999–2001; L'Heureux *et al.*, 2003; Castro *et al.*, 2004; Hauser *et al.*, 2005; Gómez Otero, 2006; Raxter *et al.*, 2006).

Both male sub-samples were compared with regard to body proportions in order to evaluate the possibility of analysing them altogether. The Mann–Whitney *U*-test was applied with a 0.05 probability in the following ratios: crural index; talus-calcaneus articulated height/tibial length; sum of vertebrae (trunk)/lower limb length (tibial length + femur length); basion-bregma cranial height/sum of vertebrae and physiological femur length/sum of vertebrae. As there were no significant differences found in any of the ratios, samples from Chubut and Santa Cruz were pooled for the analysis. In addition, several craniometric studies (e.g. Perez *et al.*, 2004) support the biological homogeneity of these populations. This resulted in a larger single sample of 27 individuals. However, this sample size must be considered to evaluate the following analyses.

#### Age and sex determination

Age was estimated according to changes in the auricular surface of the ilium (Lovejoy *et al.*, 1985)

Table 1. Descriptive data on the sample and stature reconstructions (in centimetres) of Patagonian individuals by means of the anatomical method (Fully, 1956 with corrections suggested by Raxter *et al.*, 2006)

Sample	Male		Female	
	Age	Stature	Age	Stature
Chubut	20	167.69	21	158.37
	33	175.06	26	161.43
	20	175.23		
	37	168.08		
	28	167.48		
	20	168.97		
	39	167.38		
	28	168.25		
	23	176.34		
	34	162.51		
	30	172.57		
	20	171.02		
	39	170.42		
	41	171.60		
	29	167.37		
	30	173.06		
	45	171.20		
Santa Cruz	30	176.42	22	150.57
	47	177.17	22	154.43
	22	162.77	42	165.86
	50	172.76	42	170.23
	25	169.37	37	164.30
	30	177.27	30	161.53
	50	172.30		
	47	161.11		
	55	170.56		
	50	169.42		
<i>n</i>		27	8	
Mean		170.50	160.84	

and in the pubic symphysis (Brooks & Suchey, 1990; Ghidini, 2008 personal communication for Chubut individuals; García Guraieb, 2006 for Santa Cruz individuals). Mean range stature was used in the following analyses. More details about the samples are shown in Table 1. Sex was determined based on morphological observations and metric assessment of the pelvis, cranium and long bones. The sexing criteria considered in the innominates were ventral arc, sub-pubic angle, ischiopubic ramus (Phenice, 1969), greater sciatic notch width and preauricular sulcus (Buikstra & Ubelaker, 1994). The morphological traits considered in the cranium were those suggested by Buikstra & Ubelaker (1994). Head diameter of femur and humerus were analysed as sex indicators in long bones (Béguelin & Gonzalez, 2009).

#### Osteometrics

The skeletal measurements used in this research are cranial height (basion-bregma), maximum height of the vertebral body from second cervical (C2) through

fifth lumbar (L5); maximum height of the first sacral vertebra; physiological and maximum femoral length (FLB and FLM correspondingly); tibial medial malleolus-lateral condyle length (TL); talus-calcaneus articulated height (T-CH); humerus and radius maximum length (HLM, RLM) (Fully, 1956; Martin & Saller, 1957; Raxter *et al.*, 2006). These variables were measured with a sliding caliper and an osteometric board.

#### *Intra-observer error*

The same observer (M.B.) who has previous experience in osteometric analyses performed all the measurements (recorded in millimetres) in order to avoid inter-observer error. Assessment of intra-observer error was performed using a sample composed by individuals from Patagonia. Operator inconsistency was evaluated by measuring the same set of variables in two events 2 weeks apart from each other. All the measurements show an acceptable level of consistency between observational series (Béguelin, 2009).

#### *Stature estimation methods*

The individual statures were first reconstructed through the anatomical method (Fully, 1956, following Raxter *et al.*, 2006, p. 5, Eq. 1). This technique consists of adding the length or height of every element that makes an individual's stature. A soft-tissue correction factor is added to this sum, and if the individual has a known age, it can be added as another correction factor. It is considered a reconstruction rather than an estimation given that all the bones that account for living stature are considered. Consequently, there are no assumptions on differences in body proportions when using this technique. Thus, an important advantage of the anatomical method is the relatively small difference between the predicted stature and the living stature. However, this is only the case when almost complete skeletons are available. The equation used includes a correction coefficient for age. Major changes in stature related to ageing, such as compression of vertebral bodies, are incorporated in the method by measuring the vertebrae themselves (see Raxter *et al.*, 2006 for further discussion on age changes).

Following this, two stature estimation methods – mathematical equations derived from the length of one or more long bones – were used to estimate stature in the Patagonian sample. The classical long bone regression equations developed by Trotter & Gleser (1952) were applied to the sample. Long bones produce the most precise estimations, given that they are highly correlated with stature (Damuth & MacFadden, 1990). This

is the reason why most equations used to estimate stature are based on such bones. There is a wide variety of regression models built on different anatomical elements and on diverse human populations (Sciulli & Giesen, 1993; Özaslan *et al.*, 2003; Hauser *et al.*, 2005; Işcan, 2005). Stature estimation from regression techniques represents a single mathematical operation: the replacement of the measure in the corresponding equation. Regression equations used in such studies were generated with the least squares method. This method assumes random values only in the dependent variable, stature in this case. The female stature was assessed employing eight long bone regression formulae based on African-American and U.S. Whites' populations. The male stature was calculated using 19 long bone regression equations based on Mongoloids, Mexicans, African-American, and U.S. Whites' populations. An age-correction factor suggested by Trotter & Gleser (1951) was applied to the values obtained with both long bone equations and femur/stature ratio methods.

The statures were also estimated using 3 of the 55 published femur/stature ratios. These include Mesoamerican (sex-discriminated), Asiatic and generic (Feldesman & Fountain, 1996). The Mesoamerican and Asiatic ratios were chosen on the basis of the genetic closeness of the reference and the studied populations. The stature estimation method derived from the femur/stature ratio introduced by Feldesman *et al.* (1990), is a special case of a line equation, which intercept is equal to zero. This method, which is grounded on the fact that there is a higher correlation between femur length and body size than for any other skeletal bone (Damuth & MacFadden, 1990), consists of the calculation of stature from the femur length and a chosen ratio:

$$\begin{aligned} F/S R &= \frac{\text{Femur length}}{\text{stature}} \times 100 \\ \Rightarrow \text{Stature} &= \frac{\text{Femur length}}{F/S R} \end{aligned} \quad (1)$$

Feldesman & Fountain (1996) calculated the ratios for 55 populations around the world, many of them sex specific, and found an almost constant femur-stature ratio, regardless of sex or race. These authors also calculated a generic ratio that they found very useful due to its conservative nature. The generic ratio consists of the average of all the population's ratios' values.

#### *Comparison of estimates and development of new models*

Non-parametric Wilcoxon paired sample tests were used to compare each of the stature estimation methods

with the anatomical method. To further estimate stature from different independent variables, simple regression equations derived from the male group were performed from the values reconstructed with the anatomical method. The assumptions of the regression method were formerly evaluated (i.e. homoscedasticity, with the Bartlett test, and normality, linearity and independence were evaluated by means of graphic methods).

The female sample was used to compare different methods but not to develop new models, given its small size.

## Results

Statures obtained by means of the anatomical method are shown in Table 1; the average female stature is 160.8 cm (95% confidence band: 155.6–166.2 cm;  $n = 8$ ) and 170.5 cm for the male group (95% confidence band: 168.8–172.2 cm;  $n = 27$ ). Table 2 shows mean statures

Table 2. Mean stature estimation in centimetres calculated through different femur/stature ratios (Feldesman & Fountain, 1996) and regression equations on limbs long bones (Trotter & Gleser, 1952, 1958)

Method	Male	Female
	Mean stature	Mean stature
Anatomical	170.50	160.84
C F/S Ma	175.12***	166.34*
C F/S As	174.32***	163.50*
C F/S Ge	172.49*	161.79
Regr W H	169.85	159.90
Regr W R	176.36***	168.04*
Regr W F	171.80	160.92
Regr W T	176.72***	167.22*
Regr W F + T	173.78***	164.00*
Regr AA H	167.32***	158.09
Regr AA R	169.61	160.04
Regr AA F	167.54**	158.35*
Regr AA T	171.23	161.90
Regr AA F + T	168.75**	160.17
Regr MH	169.64	
Regr M R	173.15**	
Regr M F	171.61	
Regr M T	174.48***	
Regr M F + T	174.04***	
Regr Mx H	168.16**	
Regr Mx R	172.12	
Regr Mx F	171.12	
Regr Mx T	172.47**	

F/S R = femur/stature ratio; Ma = Mesoamerican; As = Asiatic; Ge = Generic; Regr = regression; H = humerus; R = radius; F = femur; T = tibia; W = U.S. whites; AA = African-American; M = Mongoloid; Mx = Mexican.

\*  $p < 0.05$ .

\*\*  $p < 0.01$ .

\*\*\*  $p < 0.001$  for non-parametric Wilcoxon matched pairs test between each method and the anatomical one.

derived from F/S ratios and regression methods published for other populations as well as the probabilities of Wilcoxon matched pairs tests for male and female. The results of each stature estimation method were compared with the results of the anatomical method. Several significant differences were found within both groups. The outstanding result obtained from this analysis is that in different models derived from the same populations, none of the cases functions as adequate estimators of stature for the Patagonian samples used in this study.

The male statures obtained with the F/S R method significantly exceeds the values reconstructed for Patagonian samples. The equations for U.S. Whites overestimate significantly the Patagonian values for distal limb segments, i.e. radius and tibia as well as for the tibia + femur models. Instead, estimations for proximal elements do not differ from the anatomical method. Mongoloids' models show the same pattern described for U.S. Whites. Stature estimation from the Mexican femur equation does not differ significantly from that obtained with Fully's method. Tibia and humerus equations significantly overestimate and underestimate, respectively. The radius shows marginally significant differences ( $p = 0.051$ ). The regression equations for African-American populations significantly underestimate Patagonian statures in proximal segments, i.e. humerus and femur, as well as tibia + femur models.

The female group presents many significant differences. The estimations done on the F/S R for Mesoamericans and Asiatic appear to be significantly larger, whereas the generic F/S R is not significant ( $p = 0.123$ ). The regression equation for U.S. Whites based on the distal limb bones (tibia and radio) overestimates significantly the values obtained for Patagonia with the anatomical method. Significant differences were found between African-American equations and the anatomical method only for the femur equation.

Simple regression equations were developed from the values reconstructed with the anatomical method (Table 3). The regression equations are presented in a descending order of precision according to the standard estimation error (SEE). The stature estimation formula using TL and TCH + T + FLB showed the largest accuracy, while the formulae of FLB and HL provided the least accuracy.

## Discussion

The results of this work show that all the F/S R overestimate the values obtained from the anatomical

Table 3. Stature regression models from different skeletal remains derived from Patagonian samples

	Formula (variables in cm.)	$R^2$	$p$	SEE
1	$71.60 + 2.54 \times TL$	0.687	0.0000	2.11
2	$27.89 + 1.55 \times (TCH + T + FLB)$	0.741	0.0000	2.25
3	$52.15 + .79 \times (TCH + TL + FLB + HL + RL)$	0.605	0.0000	2.34
4	$64.95 + 1.25 \times (TL + FLB)$	0.612	0.0000	2.35
5	$59.72 + 2.90 \times PL$	0.656	0.0000	2.50
6	$75.48 + 2.06 \times FLM$	0.500	0.0001	2.72
7	$140.68 + 0.55 \times (SUM VERT)$	0.464	0.0002	2.77
8	$90.18 + 1.38 \times (HL + RL)$	0.360	0.0025	2.98
9	$103.11 + 2.61 \times RL$	0.336	0.0024	3.20
10	$105.83 + 2.31 \times UL$	0.355	0.0027	3.28
11	$69.38 + 2.21 \times FLB$	0.437	0.0002	3.32
12	$99.74 + 2.19 \times HL$	0.288	0.0069	3.39

method. This would imply that the individuals under study have longer femurs in relation with Asiatic and Mesoamerican populations, as well as the generic mean. Notwithstanding this, stature estimates of regression methods using femur as a predictive variable are consistent with the results of the anatomical method in three out of four cases. Estimating stature on the basis of femur length is generally more accurate than using measurements of other bones. The F/S ratios were calculated for the Patagonian sample accounting for the anatomical method, being 26.95 the female ratio and 27.13 the male one. Significant differences between males and females were not detected ( $t$ -test  $p=0.436$ ). Among the 55 published F/S ratios (Feldesman & Fountain, 1996), those of African-American populations are the most similar to Patagonian ratios for females (26.80) as well as for males (27.46).

The male stature overestimation resulting from the regression equations of U.S. Whites shows that Patagonian long bones (except for the humerus) are relatively longer than in U.S. Whites' populations. Given that models including both tibia and femur overestimate the stature of the sample under study, it is reasonable to assume relatively shorter trunks in Patagonian individuals relative to individuals of U.S. origin. Similarly, the distal upper and lower limbs of Mongoloids are relatively shorter than the homologous limb of Patagonian individuals. Even though the results derived from Mexicans' regressions do not show a clear pattern, the equation involving femora seems to be consistent with Patagonian proportions. Besides this fact, there is an apparent slight discordance in the limb proportions of Patagonian and Mexican samples. The equation for the humerus tends to underestimate the stature, and those involving radius and tibia tend to overestimate it (estimation with radio is not significant, but there is a tendency  $p=0.051$ ). Regarding the African-American sample, Patagonian individuals present

shorter proximal upper and lower limbs relative to stature. It is worth noting that even when a model derived from a certain population limb bone may display accurate results (e.g. male Mongoloid humerus), the rest of the bones from that population may not necessarily match in relation to the stature of the population under study (e.g. male Mongoloid tibia; see Table 3). In fact, this is reflecting the difference in body proportions among the populations.

Differences in results between regression and F/S R estimates (e.g. African-American ratio is similar to Patagonians' one, but stature is underestimated by the regression model based on femur length of the same population) can be explained by a detailed analysis of the statistical models. F/S R can be viewed as a regression model that includes the slope alone, neglecting the elevation of the line (intercept). Linear regression in turn, considers both, slope and intercept and encloses more information. Therefore, linear regression models could be preferable for estimating stature.

With regard to the female group, the estimations derived from the U.S. Whites' regression equations (Trotter & Gleser, 1958) overestimate the stature when using the distal segment of both upper and lower limbs. This means that the Patagonian female sample has a relatively longer upper and lower distal limb segments than U.S. Whites' populations. Instead, the African-American proportions seem similar to that of Patagonia except for the femur that underestimates statures. Still, these statements need to be proven using a larger sample.

Different human populations vary in their body proportions (Roberts, 1953; Stinson, 1990). At least some of these differences could be accounted for as climatic adaptations. For example, populations adapted to cold climate have short limbs relative to stature, while populations adapted to warm climate have longer limbs (Ruff, 1993, 1994, 2002; Holliday, 1997a).

These adaptations also affect intra-limb proportions (e.g. crural and brachial indices), a phenomenon that seems to be highly correlated with mean annual temperature (Trinkaus, 1981). The limb/stature and intra-limb proportions show a geographic arrangement following a latitudinal gradient (linked to climatic features, mainly temperature). In particular, Béguelin & Barrientos (2006) studied samples of Late Holocene hunter-gatherer populations from Patagonia and neighbouring regions, finding a clear geographic variation pattern in a latitudinal orientation. The samples, composed of adult male and female individuals, were arranged into four geographic groups corresponding to latitudinal bands of 5° widths each, beginning at 40° south. On the basis of femur maximum length and the brachial and crural indices, several intra- and inter-group statistical comparisons were performed. The results suggest that there is an intelligible pattern of geographic variation in the body proportions, which can be explained by the convergence of different factors and processes such as climatic adaptation and migratory movements. Thus the samples used for the present work differ in body proportions from the northernmost ones studied by Béguelin & Barrientos (2006), probably due to the influence of climatic factors. Therefore, the results of this work show that the differences in body proportions between Patagonian and other American and Asian populations greatly affect the stature estimations.

Method choice depends largely on the particular problem. The anatomical method requires a generally complete skeleton and is a highly time-consuming task. The regression and F/S methods can be easily applied having the length of a single bone, which represents a substantial saving of time, *albeit* in detriment of estimation accuracy. Thus, the anatomical method offers advantages when estimating stature because it tends to more closely approximate living stature, based on the fact that the technique is developed from actual body proportions. This method is highly recommended when all of the bones needed to perform it are available (Raxter *et al.*, 2006). Unfortunately most archaeological skeletons are incomplete, so other methods are required such as regression equations linking a single long bone to the stature built on the same population under study. In many cases this is not obtainable, so regression formulae from other populations can be used, considering the geographical region as a criterion to choose the reference population.

In this work, a number of simple regression equations have been developed from a central Patagonian archaeological sample. These models are recommended to estimate the stature of a population geographically

close or with similar body proportions instead of other commonly used methods. The regression equations developed from Patagonian samples (see Table 3) are, for the moment, the only ones made from autochthonous populations in southern South America. Considering the issues mentioned above, like the misleading results of methods derived from one population and applied to another, the models presented here are more reliable for local samples than other methods commonly used (e.g. Trotter & Gleser, 1958; Genovés, 1967).

## Conclusions

In this work it has been established from a Late Holocene Patagonian sample that the body stature estimation, using formulae calculated on a different population, may produce misleading results. Depending on the bone used to predict the stature, there is quite a significant level of variation. Thus, the results obtained reinforce the statement that there is a remarkable influence of body proportions on stature estimation. This is not surprising since stature is a general parameter of body dimensions, recognisably linked to biogeographical aspects (Baker, 1988; Holliday, 1997b; Gustafsson & Lindenfors, 2004; Weinstein, 2005; Pfeiffer & Sealy, 2006). In Patagonia, the body proportions of ancient populations vary at least in relation to latitude, a proxy measure of climatic variation (Béguelin & Barrientos, 2006). All this evidence suggests that a biogeographically (environmental) informed selection of the method is critical in order to improve the predictions of ancient population statures.

## Acknowledgements

I want to acknowledge the careful reading, assistance and encouragement from Ivan Perez and Fernando Archuby. I am grateful to Hector M. Pucciarelli (División Antropología, Facultad de Ciencias Naturales y Museo of La Plata, Argentina) and Rafael Goñi (Instituto Nacional de Antropología y Pensamiento Latinoamericano, Argentina) for granting access to the human skeletal collections under their care. I also thank Tyler O'Brien and the staff of Grammar 101 for their kind help with the English version of the manuscript. I appreciate the Editor George Maat and two anonymous reviewers for their comments and suggestions on an earlier version of this manuscript.

## References

- Auerbach BM, Ruff CB. 2004. Human body mass estimation: a comparison of 'morphometric' and 'mechanical' methods. *American Journal of Physical Anthropology* **125**: 331–342. 10.1002/ajpa.20032.
- Baker PT. 1988. Human adaptability. In *Human Biology. An Introduction to Human Evolution, Variation, Growth and Adaptability*, Harrison GA, Tanner JM, Pilbeam DR, Baker PT (eds). Science Press: Oxford; 437–547.
- Barrientos G. 1997. Nutrición y Dieta de las Poblaciones Aborígenes Prehispánicas del Sudeste de la Región Pampeana. *PhD dissertation*. Facultad de Ciencias Naturales y Museo, Universidad Nacional La Plata, Argentina.
- Barrientos G. 2002. The archaeological analysis of death-related behaviors from an evolutionary perspective: exploring the bioarchaeological record of early American huntergatherers. In *Perspectivas Integradoras entre Arqueología y Evolución. Teoría, Método y Casos de Aplicación*, Martínez G, Lanata JL (eds). INCUAPA, Facultad de Ciencias Sociales (UNCPBA): Olavarría; 221–253.
- Béguelin M. 2009. Variación Geográfica en la Morfología del Esqueleto Postcranial de las Poblaciones Humanas de Pampa y Patagonia durante el Holoceno Tardío: Una Aproximación Morfométrica. *PhD dissertation*. Universidad Nacional de La Plata, Argentina.
- Béguelin M, Barrientos G. 2006. Variación morfométrica postcranial en muestras tardías de restos humanos de Patagonia: una aproximación biogeográfica. *Intersecciones en Antropología* **7**: 49–62.
- Béguelin M, Gonzalez PN, 2008. Estimación del sexo a partir del fémur en muestras arqueológicas de Pampa y Patagonia. *Revista Argentina de Antropología Biológica* **10**: 55–70.
- Brooks S, Suchey J. 1990. Skeletal age determination based on the os pubis: a comparison of the Acsádi-Nemeskéri and Suchey-Brooks Methods. *Human Evolution* **5**: 227–238.
- Buikstra J, Ubelaker D. 1994. *Standards for Data Collection from Human Skeletal Remains*. Arkansas Archeological Survey: Fayetteville, AR.
- Bush E. 1991. Concepts of health and stress. In *Health in Past Societies: Biocultural Interpretations of Human Skeletal Remains in Archaeological Context*, Bush E, Zvebil M (eds). BAR International Series 567: Oxford; 3–9.
- Castro A, Gómez Otero J, Arrigoni G, Moreno JE. 2004. Prospección macrorregional comparativa a las lomerías de la Costa Atlántica continental de Patagonia: algunas claves sobre el uso del espacio y de otros recursos. In *Contra Viento y Marea. Arqueología de Patagonia*, Civalero M, Fernández P, Guraieb A (eds). Instituto Nacional de Antropología y Pensamiento Latinoamericano. Sociedad Argentina de Antropología: Buenos Aires; 197–215.
- Celbis O, Agritmis H. 2006. Estimation of stature and determination of sex from radial and ulnar bone lengths in a Turkish corpse sample. *Forensic Science International* **158**: 135–139.
- Damuth J, MacFadden BJ. 1990. Introduction: body size and its estimation. In *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*, Damuth J, MacFadden BJ (eds.). Cambridge University Press: Cambridge; 1–10.
- El-Meligy M, Abdel-Hady R, Abdel-Maaboud RM, Mohamed ZT. 2005. Estimation of human body built in Egyptians. *Forensic Science International* **159**: 27–31.
- Feldesman MR, Fountain RL. 1996. Race specificity and the femur/stature ratio. *American Journal of Physical Anthropology* **100**: 207–224. 10.1002/(SICI)1096-8644(199606).
- Feldesman MR, Kleckner JG, Lundy JK. 1990. Femur/stature ratio and estimates of stature in mid- and late-Pleistocene fossil hominids. *American Journal of Physical Anthropology* **83**: 359–372. 10.1002/ajpa.1330830309.
- Formicola V, Franceschi M. 1996. Regression equations for estimating stature from long bones of early Holocene European samples. *American Journal of Physical Anthropology* **100**: 83–88. 10.1002/(SICI)1096-8644(199605)100:1<83::AID.-AJPA8>3.0.CO;2-E.
- Fully G. 1956. Une nouvelle méthode de détermination de la taille. *Annales Medicine Legale* **35**: 266–273.
- García Guraieb S. 2006. Salud y enfermedad en cazadores-recolectores del Holoceno tardío en la cuenca del Lago Salitroso (Santa Cruz). *Intersecciones en Antropología* **7**: 37–48.
- Genovés S. 1967. Proportionality of the long bones and their relation to stature among Mesoamericans. *American Journal of Physical Anthropology* **26**: 67–78. 10.1002/ajpa.1330260109.
- Gómez Otero J. 2006. Dieta, uso del espacio y evolución en poblaciones cazadoras-recolectoras de la costa centroseptentrional de Patagonia durante el Holoceno medio y Tardío. *PhD dissertation*. Facultad de Filosofía y Letras, Universidad de Buenos Aires, Argentina.
- Goñi R, Barrientos G. 2004. Poblamiento tardío y movilidad en la cuenca del lago Salitroso. In *Contra Viento y Marea. Arqueología de Patagonia*, Civalero T, Fernández P, Guraieb G (eds). Instituto Nacional de Antropología y Pensamiento Latinoamericano. Sociedad Argentina de Antropología: Buenos Aires; 313–324.
- Guichón RA. 2000. Agenda para una evaluación en la Antropología Biológica de Patagonia Austral. In *Desde el País de los Gigantes. Perspectivas Arqueológicas en Patagonia*, Belardi J, Carvallo M, Espinosa S (eds). Universidad Nacional de la Patagonia Austral: Río Gallegos; 99–108.
- Gustafsson A, Lindenfors P. 2004. Human size evolution: no evolutionary allometric relationship between male and female stature. *Journal of Human Evolution* **47**: 253–266.
- Hauser R, Smolinski J, Gos T. 2005. The estimation of stature on the basis of measurements of the femur. *Forensic Science International* **147**: 185–190.
- Hens SM, Konigsberg LW, Jungers WL. 2000. Estimating stature in fossil hominids: Which regression model and reference sample to use? *Journal of Human Evolution* **38**: 767–784.
- Holliday TW. 1997a. Postcranial evidence of cold adaptation in European Neandertal. *American Journal of Physical Anthropology* **104**: 245–258.



- Holliday TW. 1997b. Body proportions in Late Pleistocene Europe and modern human origins. *Journal of Human Evolution* **32**: 423–447.
- Işcan MY. 2005. Forensic anthropology of sex and body size. *Forensic Science International* **147**: 107–112.
- Kondo O, Dodo Y, Akazawa T, Muhesen S. 2005. Estimation of stature from the skeletal reconstruction of an immature Neandertal from Dederiyeh Cave, Syria. *Journal of Human Evolution* **38**: 457–473.
- LeHeureux C, Guichón R, Barberena R, Borrero L. 2003. Durmiendo bajo el faro. Estudio de un entierro humano en Cabo Vírgenes (C.V.17), provincia de Santa Cruz, República Argentina. *Intersecciones en Antropología* **4**: 87–97.
- Lehmann-Nitsche R. 1911. Catálogo de la Sección Antropología del Museo de La Plata. 1910. Buenos Aires, Coni Hnos.
- Lovejoy O, Meindl R, Pryzbeck T, Mensforth R. 1985. Chronological metamorphosis of the auricular surface of the ilium: a new method for the determination of adult skeletal age at death. *American Journal of Physical Anthropology* **68**: 15–28. 10.1002/ajpa.1330680103.
- Martin R, Saller K. 1957. *Lehrbuch der Anthropologie. Band 1*. Gustav Fischer Verlag: Stuttgart.
- Méndez M, Calandra H, Ferrarini S, Salceda SA, Tobisch A. 2003. Arqueología Chaqueña 2. Nota preliminar sobre un hallazgo de restos óseos en urna. In *Actas del XIII Congreso Nacional de Arqueología Argentina*, 91–103.
- Olivier G. 1976. The stature of Australopithecus. *Journal of Human Evolution* **5**: 529–534.
- Onaha ME, Tobisch A, Pan MF, Padula G, Drube H, Vassallo M. 2002. Enterratorio humano en la mesada del Carrizal, Valle de Hualfín (Dto. De Belén, pcia. de Catamarca). In *XIII Congreso Nacional de Arqueología Argentina*, Córdoba, 497–509.
- Özaslan A, Işcan M, Özaslan I, Tuğcu H, Koç S. 2003. Estimation of stature from body parts. *Forensic Science International* **132**: 40–45.
- Pearson K. 1899. Mathematical contributions to the theory of evolution. V. On the reconstruction of the stature of prehistoric races. *Philosophical Transactions Royal Society of London* **192**: 169–244.
- Perez SI, Béguelin M, Del Papa M. 2004. Evaluación de relaciones biológicas regionales y extra regionales de muestras del N.O. de Santa Cruz. In *Contra viento y marea*, Civalero MT, Fernández PM, Guráieb AGA (eds). Arqueología de Patagonia, 359–437. 832. Instituto Nacional de Antropología y Pensamiento Latinoamericano, Sociedad Argentina de Antropología.
- Pfeiffer S, Sealy J. 2006. Body size among Holocene Foragers of the Cape Ecozone, Southern Africa. *American Journal of Physical Anthropology* **129**: 1–111. 10.1002/ajpa.20231.
- Phenice T. 1969. A newly developed visual methods of sexing os pubis. *American Journal of Physical Anthropology* **30**: 297–301. 10.1002/ajpa.1330300214.
- Prieto A. 1991. Cazadores tempranos y tardíos en la Cueva 1 del Lago Sofía. *Anales del Instituto de la Patagonia. Serie Ciencias Humanas* **20**: 75–99.
- Prieto A. 1993–1994. Algunos datos en torno a los enterratorios humanos de la región continental de Magallanes. *Anales del Instituto de la Patagonia. Serie Ciencias Humanas* **22**: 91–100.
- Raxter M, Auerbach B, Ruff C. 2006. Revision of the fully technique for estimating statures. *American Journal of Physical Anthropology* **130**: 374–384. 10.1002/ajpa.20361.
- Roberts DF. 1953. Body weight, race and climate. *American Journal of Physical Anthropology* **11**: 533–558. 10.1002/ajpa.1330110404.
- Ruff CB. 1993. Climatic adaptation and hominid evolution: the thermoregulatory imperative. *Evolutionary Anthropology* **2**: 53–60. 10.1002/evan.1360020207.
- Ruff CB. 1994. Morphological adaptation to climate in modern and fossil hominids. *Yearbook of Physical Anthropology* **37**: 65–107. 10.1002/ajpa.1330370605.
- Ruff CB. 2002. Variation in human body size and shape. *Annual Review of Anthropology* **31**: 211–232.
- Salceda SA, Méndez MG, Castro AS, Moreno JE. 1999–2001. Enterratorios indígenas de Patagonia: el caso del sitio Heupel-Caletta Olivia Santa Cruz (Argentina). *Xama* **12–14**: 161–171.
- Sciulli PW, Giesen MJ. 1993. Brief communication: an update on stature estimation in prehistoric native Americans of Ohio. *American Journal of Physical Anthropology* **92**: 395–399. 10.1002/ajpa.1330920309.
- Stini WA. 1969. Nutritional stress and growth: sex difference in adaptive response. *American Journal of Physical Anthropology* **31**: 417–426. 10.1002/ajpa.1330310316.
- Stinson S. 1990. Variation in body size and shape among South American Indians. *American Journal of Human Biology* **2**: 37–51.
- Tanner J. 1988. Human growth and constitution. In *Human Biology. An Introduction to Human Evolution, Variation, Growth and Adaptability*, Harrison G, Tanner J, Pilbeam D, Baker P (eds). Oxford Science Press: Oxford, 339–432.
- Telkkä A. 1950. On the prediction of human stature from long bones. *Acta Anatomica* **9**: 103–117.
- Trinkaus E. 1981. Neanderthal limb proportions and cold adaptation. In *Aspects of Human Evolution*, Stringer CB (ed.). Taylor & Francis: London, 187–224.
- Trotter M, Gleser G. 1951. The effect of ageing on stature. *American Journal of Physical Anthropology* **9**: 311–324. 10.1002/ajpa.1330090307.
- Trotter M, Gleser G. 1952. Estimation of stature from long bones of American Whites and Negroes. *American Journal of Physical Anthropology* **10**: 469–514. 10.1002/ajpa.1330100407.
- Trotter M, Gleser G. 1958. A re-evaluation of estimation of stature based on measurements taken during life and the long bones after death. *American Journal of Physical Anthropology* **16**: 79–124. 10.1002/ajpa.1330160106.
- Weinstein K. 2005. Body proportions in ancient Andeans from high and low altitudes. *American Journal of Physical Anthropology* **128**: 559–585. 10.1002/ajpa.20137.