

Earth, Moon, and Planets

LAVRAS DO SUL: A NEW EQUILIBRATED ORDINARY CHONDRITE FROM RIO GRANDE DO SUL, BRAZIL

--Manuscript Draft--

Manuscript Number:	MOON607R1
Full Title:	LAVRAS DO SUL: A NEW EQUILIBRATED ORDINARY CHONDRITE FROM RIO GRANDE DO SUL, BRAZIL
Article Type:	Manuscript
Keywords:	Brazilian Meteorite; Chondrite; Electron Microprobe; Mössbauer Spectroscopy
Corresponding Author:	Rosa B Scorzelli, Ph.D Rio de Janeiro, Rio de Janeiro BRAZIL
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	M E Zucolotto, Dr.
First Author Secondary Information:	
All Authors:	M E Zucolotto, Dr. L L Antonello, Dr. M E M.E.Varela, Dr. Rosa B Scorzelli, Ph.D Isabel P Ludka, Dr. P Munayco, Ph.D E dos Santos
All Authors Secondary Information:	
Manuscript Region of Origin:	
Abstract:	<p>The new Brazilian chondrite, Lavras do Sul was found in 1985 at Lavras do Sul, Rio Grande do Sul State, Brazil. (33°30'48"S; 53°54'65"W). It consists of one mass weighting about 1kg and being covered by a black fusion crust with grayish interior. Four polished thin sections were prepared from a slice weighing 67g on deposit at Museu Nacional/UFRJ. It consists mostly of chondrules and chondrule fragments dispersed in a recrystallized matrix. Most chondrules are poorly defined and range in size from 300µm to 2000 µm, although some of them show distinct outlines, particularly when viewed under cross-polarized transmitted and reflected light. The texture of chondrules varies from: non-porphyritic (e.g., barred-olivine, radial-pyroxene); to porphyritic ones (e.g., granular olivine as well as olivine-pyroxene). The meteorite contains mainly olivine (Fa24.9), low-Ca pyroxene (Fs22.6) and metal phases, with minor amount of plagioclase, chromite and magnetite. Mössbauer Spectroscopy studies indicate that the metal phase is kamacite, tetrataenite and antitaenite. Veins of secondary iddingsite crosscut the thin section and some ferromagnesian silicates. The chemical composition indicates that Lavras do Sul is a member of the low iron L chondrite group. The poorly delineated chondritic texture with little quantity of well-defined chondrules, the occurrence of few clinopyroxene and plagioclase (and maskelinite) with apparent diameters ranging from 5µm to 123µm in length, led us to classify Lavras do Sul in an equilibrated petrologic type 5. The shock features of some minerals, suggests a shock stage S3, and the presence of a small amount of secondary minerals such as iddingsite and goethite, a degree of weathering W1. The meteorite name was approved by the Nomenclature Committee (Nom Com) of the</p>

M. E. Zucolotto², L. L. Antonello¹, M.E.Varela³, R. B. Scorzelli¹, Isabel P. Ludka⁴
P. Munayco¹, E. dos Santos¹.

**LAVRAS DO SUL: A NEW EQUILIBRATED ORDINARY L5 CHONDRITE FROM RIO
GRANDE DO SUL, BRAZIL.**

¹Centro Brasileiro de Pesquisas Físicas (CBPF/MCT), Brazil.

²Museu Nacional /UFRJ, RJ, Brazil.

³ICATE- CONICET, San Juan, Argentina.

⁴Instituto de Geociências/UFRJ, RJ, Brazil

Corresponding Author:

Rosa B scorzelli

scorza@cbpf.br

Telephone: +55-21-21417188

Fax: +55-21-21417540

LAVRAS DO SUL: A NEW EQUILIBRATED ORDINARY L5 CHONDRITE FROM RIO GRANDE DO SUL, BRAZIL.

M. E. Zucolotto², L. L. Antonello¹, M.E.Varela³, R. B. Scorzelli¹, Isabel P. Ludka⁴
P. Munayco¹, E. dos Santos¹.

ABSTRACT

The new Brazilian chondrite, Lavras do Sul, was found in 1985 at Lavras do Sul, Rio Grande do Sul State, Brazil (33°30'48"S; 53°54'65"W). It consists of a single mass weighing about 1kg, covered by a black fusion crust with grayish interior. Four polished thin sections were prepared from a slice weighing 67g on deposit at the Museu Nacional/UFRJ. It consists mostly of chondrules and chondrule fragments dispersed in a recrystallized matrix. Most chondrules are poorly defined and range in size from 300µm to 2000 µm, although some of them show distinct outlines, particularly when viewed under cross-polarized transmitted and reflected light. The texture of chondrules varies from non-porphyritic (e.g., barred-olivine, radial-pyroxene) to porphyritic ones (e.g., granular olivine as well as olivine-pyroxene). The meteorite contains mainly olivine (Fa_{24.9}), low-Ca pyroxene (Fs_{22.6}) and metal phases, with minor amounts of plagioclase, chromite and magnetite. Mössbauer Spectroscopy studies indicate that the metal phase is kamacite, tetrataenite and antitaenite. Veins of secondary iddingsite crosscut the thin section and some ferromagnesian silicates. The chemical composition indicates that Lavras do Sul is a member of the low iron L chondrite group. The poorly delineated chondritic texture with few well-defined chondrules, the occurrence of rare clinopyroxene and plagioclase (and maskelinite) with apparent diameters ranging from 5µm to 123µm led us to classify Lavras do Sul as an equilibrated petrologic type 5. The shock features of some minerals suggest a shock stage S3, and the presence of a small amount of secondary minerals such as iddingsite and goethite, a degree of weathering W₁. The meteorite name was approved by the Nomenclature Committee (Nom Com) of the Meteoritical Society (Meteoritic Bulletin N°99).

Keywords: Brazilian Meteorite, Chondrite, Electron Microprobe, Mössbauer Spectroscopy.

INTRODUCTION

The Lavras do Sul meteorite was found in 1985 by Professor Picada from the Geosciences Institute/UFRGS, in a stream pebble field bed in the vicinity of Lavras do Sul, Rio Grande do Sul State, Brazil. It is a single stone of about 1 kg, with mean dimensions of 10 x 5 x 6 cm, and covered by a black fusion crust. A mass of 67g was donated to Museu Nacional/UFRJ and is housed at the Department of Geology and Paleontology (Inventory number 142 MT). In hand specimen, it shows a grayish interior with some slightly reddish-brown spots due to weathering, also visible in thin-section. Metal shows very few weathering suggesting that the recovery was done shortly after the fall (Fig. 1). In this paper a detailed petrographic, chemical and Mössbauer spectroscopy study of this new meteorite is reported.

MATERIALS AND METHODS

Four polished thin-sections were etched with 2% nital to reveal the metallographic microstructures and studied using optical, scanning electron microscopy (SEM) electron microprobe and Mössbauer spectroscopy. Optical microscopy was performed by transmitted and reflected light, using a Petrographic Microscope Carl Zeiss, with Axionvision Release 4.7 software. The electron microscopy was performed by back-scattering electron imaging (BSEi) on a Digital Scanning Microscope, (DSM 940A Zeiss), equipped with a system for energy-dispersive X-ray micro-analysis (LINK/Oxford-exLII).

Major element chemical compositions were obtained by an ARL-SEMQ (WDS) electron microprobe with analyses acquired at conditions of 15 kV acceleration potential and 15 nA sample currents. Estimated precision for major and minor elements is better than 3% and for Na about 10%. Natural and synthetic standards were used for calibration and an online ZAF correction was applied to the data.

The ^{57}Fe Mössbauer spectroscopy (^{57}Fe -MS) experiments were performed at room temperature in standard transmission geometry, using a 25mCi $^{57}\text{Co}/\text{Rh}$ radioactive source in sinusoidal mode. The NORMOS code was used for the spectrum analysis.

PETROGRAPHY

TEXTURE

Under optical inspection, Lavras do Sul shows a texture that is similar to other equilibrated ordinary chondrites. Both matrix and chondrules have undergone considerable recrystallization. The chondrules are embedded in a recrystallized and clastic matrix with most grains ranging in size between 50 μm to 150 μm and down to the limits of microscopic resolution. There are patchy areas with different colors and veins of secondary minerals (mainly iddingsite) that crosscut the matrix. Some chondrules show a poorly defined interface between the chondrules and matrix, in particular some radial and barred chondrules with irregular boundaries. Metal grains and troilite are scattered through the matrix as well as in the chondrules.

THE CHONDRULES

The chondrules in Lavras do Sul are mainly poorly defined, although some show distinct outlines, particularly when viewed under crossed polarized transmitted and reflected light (Fig. 2). The chondrules are more or less rounded in outline, irregular or fragmented. Chondrules range in apparent diameter between 200 μm to 2000 μm , with three average population sizes: small ($\sim 300\mu\text{m}$), medium ($\sim 700\mu\text{m}$) and large ($\sim 1.700\mu\text{m}$). Here, the size frequency and apparent dimensions of chondrules measured in thin-sections were determined by a combination of measurement techniques (Eisenhour 1996; King and King 1978). Chondrules with non-porphyritic and porphyritic textures are observed, including: Barred Olivine (BO), Radial Pyroxene (RP), Granular olivine-pyroxene (GOP), Porphyritic olivine-pyroxene (POP), Microcrystalline chondrule and a Chromite-rich chondrule.

Barred olivine (BO) chondrules range in size between 500 μm to 1200 μm in apparent diameter. In most of them the olivine bars as well as the rims occur in parallel orientation (e.g., showing simultaneous extinction under cross-polarized transmitted light) suggestive of a single crystal. Some of these objects are bounded by a rim of olivine. The bars of the defined chondrules vary widely in size from 18 μm to 20 μm up to 80 μm to 120 μm (the latter measured in a fragment of a BO). The skeletal crystals of olivine commonly are associated with interstitial aggregates of orthopyroxene. The mesostasis is most glassy, with incipient recrystallization with microcrystalline thin pyroxene bars (Figs. 2b)

Radial Pyroxene (RP) chondrules are less common than those of the Barred Olivine type, having sizes varying from 500 μm to 900 μm with a sub-rounded shape. They have thin lamellae of pyroxene with a radiating fan distribution as a result of an eccentric nucleation point near the surface. The main mineral is orthopyroxene, with fine lamellae of clinopyroxene. In some of these objects euhedral skeletal olivine grains were observed.

Granular olivine-pyroxene chondrules (GOP) have sizes varying from 500 μm to 600 μm . The anhedral crystals of olivine and pyroxene have sizes around 25 μm in a recrystallized mesostasis.

Porphyritic olivine-pyroxene chondrules (POP) are the largest objects, with a size range from 700 to 1900 μm . They are composed of large euhedral to subhedral olivine or pyroxene crystals with interstitial aggregates of pyroxene and sometime plagioclase. They have sizes up to 1800 μm and are immersed in a slightly devitrified glassy mesostasis sometimes are surrounded by large opaque minerals. Crystals show undulatory extinction, planar structure and numerous cracks sometimes filled with iddingsite.

Microcrystalline chondrules have sizes around 700 μm with anhedral small crystals of olivine and pyroxene in a recrystallized mesostasis.

Among the different types of chondrules described above, one particular object has called attention. It is a chromite-rich chondrule (600 μm in apparent diameter) composed of large olivine bars and glass (Fig. 2d). Immersed in the Na-rich glass there are small euhedral olivine crystals and euhedral chromite crystals that contribute to the dark aspect of the object in transmitted light.

MINERAL CHEMISTRY

MAJOR ELEMENT PHASE COMPOSITIONS

Representative and averaged electron microprobe analyses (EMPA) of the phases are given in Tables 1 – 7. Olivine ($\text{Fa}_{24,9}$) typically has FeO contents between 22.1 and 23.4 wt% (Table 1). The MnO and CaO contents vary between (0.65 – 0.77 wt %) and (0.02–to 0.07 wt %), respectively. Pyroxenes are low-Ca pyroxene ($\text{Fs}_{22,6}$) with FeO contents varying from 13.8 to 14.5 wt% (Table 2), TiO_2 and CaO in the range of 0.11 – 0.27 wt% and 0.7 – 0.97 wt%, respectively, with Cr_2O_3 from 0.07 to 0.13wt%. Na and K are mostly below the detection limit of the electron microprobe. In the chromite-rich chondrule, the composition of the large and small euhedral olivine crystals show no significant variation in their major elements. Small euhedral chromites in the mesostasis have TiO_2 , Al_2O_3 and MgO contents around 4.3; 6 and 3, 7 wt%, respectively. The glassy mesostasis is rich in Na_2O and K_2O (~2 and 1 wt%, respectively). Representative analyses of the different phases are shown in Table 3.

ACCESSORY MINERALS

In order of decreasing abundance the accessory minerals are: kamacite, troilite, taenite, plagioclase, chromite and magnetite.

Plagioclase occurs as tiny anhedral crystals, occasionally showing twinning. However, most of them are transformed into maskelinite, ranging in size from 5 μm to 123 μm (the majority of the grains are < 50 μm) (Table 4).

Troilite is very abundant, comprising about 60 % of the opaque minerals. It is commonly observed surrounding large chondrules, but can also occur isolated or associated with metal. Several large troilite crystals exhibit well developed polycrystalline grains with 120° intersections (triple junctions). Representative analyses are given in Table 5.

Metal grains are typically irregular in shape; some are elongated, others subhedral (Fig. 3). They are less abundant than troilite and some grains are composed of kamacite and taenite. In etched samples the taenite shows a cloudy core and a tetrataenite rim that can reach 1-3 μm thick (Fig. 3b). Under optical microscopy the cloudy core is seen to be composed of tetrataenite and sometimes martensite. A compositional profile (P1, Fig 2a) performed near the edge of a large metal grain included in troilite shows the high variation in Ni contents in kamacite, tetrataenite and cloudy taenite (Table 6, Fig 4). Kamacite (α nickel-iron, b.c.c) grains are polycrystalline and exhibits Neumann lines, shock-produced twin lamellae, and sometimes displays Widmanstätten-like patterns. Individual kamacite appears commonly with irregular shapes either isolated or associated with troilite. Kamacite can also form large veins (~ 7 mm x 1 mm), that are observed crosscutting a thin-section (Fig. 3c). The kamacite crystals show irregular borders towards silicates. A step profile, performed across the vein, indicates that the chemical composition (e.g. Ni content) is quite constant (P2, Fig. 3c, Table 7). Metal can also develop zoneless plessite texture as pearlitic plessite (Sears and Axon 1975), composed of

roughly equal parts of kamacite and taenite (Table 8 Fig. 3d,e,f).

Small FeNi metal spherical within silicate melt pockets and veins (Sears and Dodd 1998) are also observed.. Magnetite appears associated with troilite and is poikilitic in the silicate phases. Occasionally silicate minerals (e.g., olivine, pyroxenes, plagioclase) occur as inclusions in metal. Chromite occurs as subhedral to euhedral unzoned crystals, associated with silicate phases.

The Mössbauer analysis of the Lavras do Sul bulk sample showed a spectrum composed of overlapping paramagnetic and magnetic phases. The spectra at room temperature exhibit two Fe^{2+} doublets attributed to olivine and pyroxene, and additionally two magnetic phases associated to Fe-Ni phases (kamacite and taenite) and troilite. The Mössbauer spectrum of the metal particles extracted from the bulk sample (Fig. 5), free of silicates and troilite, exhibits only kamacite and small amounts of the intergrowth tetrataenite/antitaenite (Rancourt and Scorzelli.1995).

SECONDARY MINERALS

The products of terrestrial weathering are iron oxide hematite (Fe_2O_3) and hydrated iron oxide minerals as iddingsite, goethite, lepidocrocite and others. These products are generally fine-grained and intimately mixed. In Lavras do Sul, iddingsite fills fractures and veins and tarnishes the silicate minerals with reddish and yellowish colors. Metal, however, is relatively fresh with only few weathering features, suggesting a recovery shortly after the fall.

DISCUSSIONS AND CONCLUSIONS

Lavras do Sul consists of chemically homogeneous major (and minor) phases (e.g., Table 1 and 2), with plagioclase showing a uniform composition (Table 3). The progressive stages of shock metamorphism in ordinary chondrites (Stöffler et al, 1991) are mainly based on shock effects in olivine and plagioclase. The presence of deformation features such as planar fractures and undulatory extinction in plagioclase and olivine, with fractures in most silicate grains, determine the shock stage as S3.

Homogenous plagioclase – such as those observed in Lavras do Sul - should, however, be expected in unshocked ordinary chondrites. But, because of the slow diffusion rates of NaSi-CaAl substitutions (Hart 1981), uniform compositions can also be expected in highly shocked chondrites.

The presence of ambiguous features indicating different shock stages in the same chondrite is not uncommon, and could point towards shock and subsequent post-shock annealing, that could preserve features normally associated with high degrees of shock metamorphism. Shock metamorphic effects can be partially removed by annealing resulting in a shock classification that could be lower than the maximum stage experienced by the chondrite.

Troilite appears to be a sensitive indicator of shock (Bennett and McEwen 1996); it becomes

polycrystalline with 120° triple junctions with shock loading between 35GPa-60GPa. This corresponds to a peak pressure much higher (e.g., S4: 20-35 GPA, S5: 35-55 GPA) than those estimated based in shock effects in olivine and plagioclase (S3).

Zoneless plessite is common in type 5 and 6 low shock chondrites (S1-S2) (Bennett and McEwen 1996). Also perlitic plessite is only observed in L chondrites of low shock stage below S4 (Sears and Axon 1975). However, Stöffler et al (1991) suggest that the presence of coarse-grained perlitic plessite could be interpreted as having been formed by a post-shock annealing after the chondrite was affected by a high shock stage above S4.

Metal is a phase that should be easier to equilibrate than all other phases, but is commonly not equilibrated. The presence of melt pockets indicates temperatures > 800 °C (e.g., partial or *in situ* melting above 950°C (Sears and Axon 1975) and this should have homogenized all metal grains. In such a scenario kamacite and taenite might produce composite grains with equal amount of both phases. This is rarely observed in chondrites (Urey and Mayeda 1959) as well as in Lavras. The zoned taenite grains with tetrataenite and cloudy taenite core with the classical “M” shaped Ni profile (Figs. 4) are interpreted to be the result of diffusion-controlled transformation taenite → taenite + kamacite during very slow cooling (Heymann 1967). This indicates that Lavras do Sul had a variable metallographic cooling rate. This result is in agreement with previous work (Willis and Goldstein 1983) that shows that metallographic cooling rates are rarely unique in a given chondrite.

The new Brazilian meteorite, Lavras do Sul, is thus classified as an ordinary chondrite (L5) (Van Schmus and Wood 1967; Rubin 1990) (Fig. 6) with shock stage S3 (Stöffler et al 1991) and a degree of weathering W1 (Wlotzka 1993).

ACKNOWLEDGMENTS

E. Zucolotto acknowledges FAPERJ for financial support. M.E. Varela is grateful to Agencia PICT 212 and CNPq for financial support during scientific visits to CBPF. R.B. Scorzelli would like to thank FAPERJ and CNPq for financial support; P. Munayco and E. dos Santos are grateful to FAPERJ and CAPES and L. L. Antonello to CBPF/MCT for their fellowships.

REFERENCES:

1. M. E. Bennett and H. Y. Jr. McSween. *Meteoritics & Planetary Science* **31**:783-792. (1996).
2. A. J. Brearley and R. H. Jones. Chondritic meteorites. In *Planetary materials*, edited by Papike J. J. Washington D.C.: Mineralogical Society of America. pp. 3-1–3-398. (1998).
3. D. D. Eisenhour. *Meteoritics & Planetary Science* **31**:243-248. (1996).
4. S. R. Hart. *Geochimica Cosmochimica Acta*, **45**: 279-291. (1981).
5. D. Heymann. *Icarus* **6**: 189 - 221. (1967).
6. T.V. V. King and E. A. King. *Meteoritics* **31**:47-72. (1978).
7. D. G. Rancourt and R. B. Scorzelli. *Journal of Magnetism and Magnetic Materials* **150**: 30-36. (1995).

8. A. E. Rubin, *Geochimica et Cosmochimica Acta* **54**:1217–1232. (1990). H. Y. Jr.
9. A. E. Rubin. *Geochimica et Cosmochimica Acta*, **56**: 1705-1714. (1992).
10. D. W. G. Sears and H. J. Axon. *Meteoritics* **10**: 486-487. (1975)
11. D. W.G. Sears and R. T. Dodd. In J. F. Kerridge and M. S. Matthews, eds., *Meteorites and the Early Solar System*. pp. 3-31. Tucson, AZ: University of Arizona Press. (1998).
12. D. Stöffler, K. Keil, and E.R.D Scott. *Geochimica et Cosmochimica Acta* **55**: 3845-3867. (1991).
13. H. C. Urey and Mayeda. *Geochimica et Cosmochimica Acta*, **17**: Issue 1-2, pp.113-116. (1959).
14. J. Willis and J.I. Goldstein In *14th Lunar & Planetary Science Conference*. (*J. Geophys. Res. Suppl.* **88**) B 287- B 292. (1983).
15. F. Wlotzka. *Meteoritics* **28**: 460. (1993).
16. W. R. Van Schmus and J. A. Wood. *Geochimica et Cosmochimica Acta* **31**: 747-65. (1967).
17. M. E Zucolotto et al. *Meteoritics & Planetary Science*. **45**:226. the 73 rd Annual Meeting of the Meteoritical Society (2010).

Figure 1



Figure 2




Figure 3




Figure 4




Figure 5




figure 6
[Click here to download high resolution image](#)




FIGURE CAPTIONS

Figure 1

Panoramic photomicrograph in plane (a) and crossed transmitted light (b) of the whole thin-section, showing the distribution of objects.

Figure 2

(a) Photomicrograph of Lavras do Sul meteorite in plane polarized transmitted light showing the chondritic texture with chondrules poorly defined. Length of the field: 5.2mm. (b) Plane transmitted light photomicrographs showing a barred olivine chondrule with their unmistakable texture composed of one set of elongated parallel bars of olivine with a glassy mesostasis between bars. Length of the field: 1,3mm. (c) Plane polarized transmitted light photomicrographs of microcrystalline chondrule showing anhedral small crystals of olivine and pyroxene in a glassy mesostasis. (d) Photomicrographs in reflected light showing a chromite-rich chondrule (600 μm in apparent diameter), composed of large olivine bars and glass. Immersed in the Na-rich glassy mesostasis there are small euhedral olivine crystals and euhedral chromite crystals,

Figure 3

(a) BSE image of kamacite with cloudy taenite inside bordered by tetrataenite. Dark line indicates location of the profile P1 (Table 6). The white rectangle in dotted line shows the location of figure 3b. (b) Reflected light photomicrograph showing the aspect of phases after being etched by nital2%. The cloudy taenite (black) is bordered by tetrataenite (bright silver) and surrounding by polycrystalline kamacite. (c) Reflected light photomicrograph showing a metal vein. Black line indicates profile P2 (Table 7). The white rectangle in dotted line shows the location of figure 3d. (d) BSE image of metal with plessitic texture composed of roughly equal parts of kamacite and taenite. (e) Reflected light photomicrograph showing metal with plessitic texture, Length of the field 2,6mm. (f) BSE image of metal. Numbers indicate location of the EMP analysis (Table 8).

Figure 4

Point step profile showing the classical M shape in the Ni contents of P1 (see Fig. 3a).

Figure 5

The ^{57}Fe Mössbauer spectra performed at room temperature, the bulk sample shows kamacite, troilite, olivine and pyroxene; after magnetic separation it shows kamacite, tetrataenite and antitaenite.

Figure 6

Classification of Lavras do Sul considering the average of Low-Ca pyroxene and olivine. Chondrites fields from Brearley and Jones, (1998)

Table 1: Representative EMP analysis of olivine (wt %)

SiO ₂	38,4	38,2	38,4	37,7	38,5	38,0	38,6	39,2
TiO ₂	0,05		0,04	0,05	0,05	0,05		0,02
FeO	22,7	22,7	22,9	23,4	22,4	22,1	22,7	22,5
MnO	0,68	0,65	0,74	0,74	0,77	0,68	0,71	0,65
MgO	38,4	37,7	38,6	37,7	38,2	39,0	37,9	38,0
CaO		0,02	0,06	0,06	0,05	0,04	0,07	
Total	100,2	99,2	100,8	99,7	100,0	99,9	100,0	100,3

Table 2: Representative EMP analysis of pyroxene (wt %)

SiO ₂	55,4	54,8	54,8	55,2	56,0	55,6	55,3	55,6
TiO ₂	0,19	0,11	0,23	0,22	0,22	0,27	0,23	0,27
Al ₂ O ₃	0,19	0,15						
Cr ₂ O ₃	0,11	0,09	0,04	0,11	0,13	0,07	0,13	0,09
FeO	14,1	13,9	14,5	14,3	14,2	14,5	14,0	13,8
MnO	0,67	0,79	0,55	0,74	0,76	0,74	0,67	0,74
MgO	28,5	28,3	28,6	29,0	28,7	28,7	28,6	28,6
CaO	0,82	0,97	0,79	0,99	0,80	0,74	0,93	0,70
Total	100,0	99,1	99,5	100,6	100,8	100,6	99,9	99,8

Table 3: EMP analysis of the olivine-chromite-rich chondrule (wt %)

	OI (B)	OI (B)	OI (euh)	OI (euh)	Chromite	Chromite	Chromite	Glass	Glass
SiO ₂	36,4	39,4	39,7	39,8	0,50	0,53	0,64	68,5	68,1
TiO ₂	0,07	0,07	0,00	0,00	4,47	4,29	4,24	0,00	0,07
Al ₂ O ₃	0,42	0,06	0,25	0,14	5,99	5,97	6,13	25,3	25,2
Cr ₂ O ₃	0,50	0,46	0,31	0,46	53,7	53,6	53,3	0,33	0,00
FeO	29,8	22,6	21,8	22,0	30,7	30,9	31,8	0,37	0,52
MnO	0,34	0,38	0,51	0,47	0,48	0,60	0,64	0,04	0,04
MgO	32,2	36,9	35,3	35,8	3,61	3,81	3,56	0,02	0,02
CaO	0,11	0,02	0,05	0,02	0,07		0,05	2,40	2,13
Na ₂ O								2,21	2,70
K ₂ O								1,03	1,07

Table 4: Representative EMP analysis of plagioclase (wt %)

SiO ₂	67,0	67,9	65,9	65,8	67,5
TiO ₂	0,07	0,04	0,05	0,04	0,00
Al ₂ O ₃	22,1	21,7	21,3	21,7	22,1
Cr ₂ O ₃			0,03		0,07
FeO	0,45	0,58	0,36	0,36	0,75
CaO	2,09	2,12	2,15	2,14	2,12
Na ₂ O	7,84	6,82	8,02	7,52	6,90
K ₂ O	0,90	1,35	1,32	1,20	1,11
Total	100,5	100,6	99,2	98,8	100,5
<i>Ab</i>	81,8	76,8	79,6	79,2	78,4
<i>An</i>	12	13,2	11,8	12,5	13,3
<i>Or</i>	6,2	10	8,6	8,3	8,3

Table 5: Representative analysis of Troilite (wt %)

Fe	61,6	61,2	62,0	62,8	62,8
Ni	2,86	1,10	0,05	0,09	0,44
S	36,5	37,6	38,4	37,8	37,7
Total	100,9	100,1	100,5	100,7	100,9

Table 6: Representative analysis in metal (P1) (wt %)

	T					T				T			
Fe	93,6	93,6	93,5	92,9	51,9	80,4	80,2	81,3	80,5	50,6	49,4	92,0	92,0
Ni	6,4	6,4	6,0	6,5	44,7	19,5	19,5	17,8	19,0	50,1	50,2	6,5	6,8
Co	0,62	0,60	0,58	0,63	0,08	0,55	0,51	0,62	0,53	0,04	0,03	0,55	0,57
Total	100,7	100,7	100,1	100,1	96,8	100,5	100,3	99,8	100,1	100,8	99,6	99,2	99,4

Table 7: Representative analysis in metal vein (P2) (wt %)

Fe	93,1	93,1	93,3	93,3	93,2
Ni	6,8	6,6	6,5	6,6	6,8
Co	0,93	0,94	0,92	0,90	0,90
Total	100,9	100,6	100,7	100,9	100,9

Table 8: EMP analysis of a metal grain with plessitic texture

	1	2	3	4	5	6	7	8	9	10	11	12
Fe	46.9	47.3	50.1	48.2	48.0	67.1	92.1	68.8	69.0	69.6	92.5	69.7
Ni	53.0	52.7	50.5	51.2	52.2	31.2	5.4	30.5	29.3	29.7	5.6	29.7
Co	0.05	0.16	0.12	0.08	0.04	0.37	0.61	0.27	0.25	0.26	0.49	0.34
Total	100.1	100.3	100.7	99.6	100.3	98.8	98.3	99.7	98.6	99.5	98.6	99.8

Tables captions

Table 1: Representative EMP analysis of olivine (wt %)

Table 2: Representative EMP analysis of pyroxene (wt %)

Table 3: EMP analysis of the olivine-chromite-rich chondrule (wt %)

Table 4: Representative EMP analysis of plagioclase (wt %)

Table 5: Representative analysis of Troilite (wt %)

Table 6: Representative analysis in metal (P1) (wt %)

Table 7: Representative analysis in metal vein (P2) (wt %)

Table 8: EMP analysis of a metal grain with plessitic texture