

Direct Interaction of Serotonin Type 3 Receptor Ligands with Recombinant and Native $\alpha 9\alpha 10$ -Containing Nicotinic Cholinergic Receptors

CARLA V. ROTHLIN, MARIA I. LIUDYNO, ANA F. SILBERING, PAOLA V. PLAZAS, MARÍA E. GOMEZ CASATI, ELEONORA KATZ, PAUL S. GUTH, and A. BELÉN ELGOYHEN

Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad de Buenos Aires, (C.V.R., A.F.S., P.V.P., M.E.G.C., E.K., A.B.E.), Departamento de Biología, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina (E.K.); and Department of Pharmacology, Tulane University, New Orleans, Louisiana (M.I.L., P.S.G.).

Received August 5, 2002; accepted January 24, 2002

This article is available online at <http://molpharm.aspetjournals.org>

ABSTRACT

In the present work, we characterized the effects of serotonin type 3 receptor ligands on recombinant and native $\alpha 9\alpha 10$ -containing nicotinic acetylcholine receptors (nAChRs). Our results indicate that the recombinant $\alpha 9\alpha 10$ nAChR shares striking pharmacological properties with 5-HT₃ ligand-gated ion channels. Thus, 5-HT₃ receptor antagonists block ACh-evoked currents in $\alpha 9\alpha 10$ -injected *Xenopus laevis* oocytes with a rank order of potency of tropisetron (IC₅₀, 70.1 ± 0.9 nM) > ondansetron (IC₅₀, 0.6 ± 0.1 μM) = MDL 72222 (IC₅₀, 0.7 ± 0.1 μM).

Although serotonin does not elicit responses in $\alpha 9\alpha 10$ -injected oocytes, it blocks recombinant $\alpha 9\alpha 10$ receptors in a noncompetitive and voltage-dependent manner (IC₅₀, 5.4 ± 0.6 μM). On the other hand, we demonstrate an in vivo correlate of these properties of the recombinant receptor, with those of the $\alpha 9\alpha 10$ -containing nAChR of frog saccular hair cells. The possibility that the biogenic amine serotonin might act as a neuromodulator of the cholinergic efferent transmission in the vestibular apparatus and in the organ of Corti is discussed.

Nicotinic acetylcholine receptors (nAChRs) are complexes of protein subunits that coassemble to form an ion channel gated through the binding of the neurotransmitter ACh to its ligand-binding site (Karlin, 2002). A diversity of subunits have been cloned in recent years. The nAChR at the neuromuscular junction mediates fast synaptic transmission and is thought to have a ($\alpha 1$)₂ $\beta 1\gamma\delta$ stoichiometry (Galzi et al., 1991). Ten genes that encode neuronal nAChR subunits have been identified in the vertebrate central or peripheral nervous system: $\alpha 2$ – $\alpha 8$ and $\beta 2$ – $\beta 4$ (Le Novère and Changeux, 1999). In heterologous expression systems, the neuronal nAChRs may assemble from single α -subunits, from multiple α - and β -subunits, and heteromeric nAChRs formed via pair-wise combinations of $\alpha 2$, $\alpha 3$, $\alpha 4$, or $\alpha 6$ with either the $\beta 2$ or $\beta 4$ subunits (for references, see Elgoyhen et al., 2001). Neuronal nAChRs preserve the structural motif of muscle nAChR, with

a pentameric structure that includes two α and three β subunits (Anand et al., 1991; Cooper et al., 1991).

The cloning of the $\alpha 9$ and $\alpha 10$ subunits added two peculiar members to the family of nAChRs (Elgoyhen et al., 1994, 2001). They are distant members of the family; whereas neuronal nAChR α subunits and the muscle $\alpha 1$ subunit share sequence homologies ranging from 48 to 70%, the sequence identity between $\alpha 9$ and all known nAChR subunits is less than 40%. When expressed in *Xenopus laevis* oocytes, $\alpha 9$ and $\alpha 10$ subunits form a heteromeric receptor-channel complex that is activated by ACh but not by nicotine and displays a very distinct pharmacological profile that falls into neither the nicotinic nor the muscarinic subdivision of the pharmacological classification scheme of cholinergic receptors. However, the properties of the recombinant $\alpha 9\alpha 10$ receptor are strikingly similar to those described for the cholinergic receptor that mediates synaptic transmission between efferent cholinergic fibers and cochlear outer hair cells (Fuchs, 1996; Elgoyhen et al., 2001). Moreover, the $\alpha 9$ and $\alpha 10$ gene subunits exhibit a unique and restricted expression pattern. Whereas $\alpha 9$ and $\alpha 10$ message has not been found in the central nervous system, it is present in the cochlear and

This work was supported by an International Research Scholar grant from the Howard Hughes Medical Institute, the Beca Ramón Carrillo-Arturo Oñativia, the Agencia Nacional de Promoción Científica y Tecnológica (to A.B.E.), and National Institutes of Health grant DC00303 (to P.S.G.). C.V.R. and P.V.P. are supported by a Consejo Nacional de Investigaciones Científicas y Técnicas predoctoral fellowship, M.E.G.C. by a fellowship from ANPCyT, and A.F.S. by a Fundación Antorchas fellowship for undergraduates.

ABBREVIATIONS: nAChR, nicotinic acetylcholine receptor; ACh, acetylcholine; 5-HT, 5-hydroxytryptamine; BAPTA-AM, 1,2-bis(O-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid; MDL 72222, 3-tropanyl-3,5-dichlorobenzoate; I-V, current-voltage; 5HTQ, trimethyl serotonin; ICS 205,930, tropisetron.

vestibular hair cells (Elgoyhen et al., 1994, 2001; Hiel et al., 1996; Morley et al., 1998). This has led to the proposal that efferent modulation of cochlear and vestibular hair cell function occurs, at least in part, via heteromeric nAChRs assembled from both $\alpha 9$ and $\alpha 10$ subunits (Elgoyhen et al., 2001).

Nicotinic AChRs are members of a family of neurotransmitter-gated ion channels that also includes, GABA_A, GABA_C, glycine, 5-HT₃ and some invertebrate anionic glutamate receptors (Le Novère and Changeux, 1999). The subunits of these receptors have similar sequences and distributions of hydrophobic, membrane-spanning segments. Each subunit contains in its ligand-binding N-terminal half two cysteine residues separated by 13 other residues that are presumably disulfide-linked, thus giving this family the name of the Cys-loop receptors. We have demonstrated (Rothlin et al., 1999) that homomeric $\alpha 9$ nAChRs share several pharmacological properties with GABA_A, glycine, and 5-HT₃ receptors. The aim of the present work was to perform an extensive characterization of the effects of 5-HT₃ receptor ligands on recombinant and native $\alpha 9\alpha 10$ -containing nicotinic cholinergic receptors. Our results indicate that the recombinant $\alpha 9\alpha 10$ nAChR shares striking pharmacological properties with type 3 serotonin ligand-gated ion channels. Moreover, it demonstrates an in vivo correlate of these properties of the recombinant receptor, with those of the $\alpha 9\alpha 10$ -containing nAChR of frog saccular hair cells.

Materials and Methods

Expression of Recombinant Receptors in *X. laevis* Oocytes.

For expression studies, $\alpha 4$, $\alpha 7$, $\alpha 9$, $\alpha 10$, and $\beta 2$ rat nAChR subunits were subcloned into a modified pGEMHE vector (Liman et al., 1992). Capped cRNAs were in vitro transcribed from linearized plasmid DNA templates using the mMessage mMachine T7 transcription kit (Ambion Corporation, Austin, TX). The maintenance of *X. laevis* and the preparation and cRNA injection of stage V and VI oocytes have been described in detail elsewhere (Katz et al., 2000). Typically, oocytes were injected with 50 nl of RNase-free water containing 0.01 to 1.0 ng of cRNA (at a 1:1 molar ratio when pair-wise combined) and maintained in Barth's solution at 17°C.

Electrophysiological recordings were performed 2 to 6 days after cRNA injection under two-electrode voltage clamp with a Geneclamp 500 amplifier (Axon Instruments Corp., Union City, CA). Both voltage and current electrodes were filled with 3 M KCl and had resistances of ~1 to 2 M Ω . Data acquisition was performed using a Digidata 1200 and the pClamp 7.0 software (Axon Instruments). Data were analyzed using Clampfit from the pClamp 6.1 software. During electrophysiological recordings, oocytes were continuously superfused (~10 ml/min) with normal frog saline composed of 115 mM NaCl, 2.5 mM KCl, 1.8 mM CaCl₂, and 10 mM HEPES buffer, pH 7.2. Unless otherwise indicated, the membrane potential was clamped to -70 mV. Drugs were applied in the perfusion solution of the oocyte chamber. To minimize activation of the endogenous Ca²⁺-sensitive chloride current (Elgoyhen et al., 2001), all experiments were performed in oocytes incubated with the Ca²⁺ chelator BAPTA-AM (100 μ M) for 3 to 4 h before electrophysiological recordings.

Concentration-response curves were normalized to the maximal agonist response in each oocyte. For the inhibition curves, antagonists were added to the perfusion solution for 2 min before the addition of 10 μ M ACh and then were coapplied with this agonist. Responses were referred to as a percentage of the response to ACh. The mean and S.E.M. of peak current responses are represented. Agonist concentration-response curves were iteratively fitted with the equation: $I/I_{\max} = A^{n_H}/(A^{n_H} + EC_{50}^{n_H})$, where I is the peak inward

current evoked by agonist at concentration A ; I_{\max} is current evoked by the concentration of agonist eliciting a maximal response; EC_{50} is the concentration of agonist inducing half-maximal current response, and n_H is the Hill coefficient. An equation of the same form was used to analyze the concentration dependence of antagonist-induced blockage. The parameters derived were the concentration of antagonist producing a 50% block of the control response to ACh (IC_{50}) and the associated interaction coefficient (n_H). Current-voltage relationships were obtained by applying 2-s voltage ramps from -120 to +50 mV, 10 s after the peak response to 10 μ M ACh from a holding potential (V_{hold}) of -70 mV. Leakage correction was performed by digital subtraction of the current-voltage (I-V) curve obtained by the same voltage ramp protocol before the application of ACh. Generation of voltage protocols and data acquisition were performed using a Digidata 1200 and the pClamp 6.1 or 7.0 software (Axon Instruments). Data were analyzed using Clampfit from the pClamp 6.1 software.

Statistical significance was evaluated by the Student's t test (two-tailed, unpaired samples). Multiple comparisons of IC_{50} values were performed with a one-way analysis of variance followed by Tukey's test. $p < 0.05$ was considered significant.

Isolation and Patch Clamp Recording from Saccular Hair Cells.

Frog saccular hair cells were isolated enzymatically. The dissociation protocol, optimized for the purpose of observing reliable ACh responses, was described previously (Holt et al., 2001). Briefly, leopard frogs (*Rana pipiens*) were chilled, pithed,¹ and decapitated. Each side of the head was placed into a perilymph-like standard external solution composed of 105 mM NaCl, 2.5 mM KCl, 0.81 mM MgCl₂, 1.8 mM CaCl₂, 3.4 mM NaHCO₃, 0.5 mM NaH₂PO₄, 2.5 mM Na₂HPO₄, 1 mM ascorbate, 4 mM glucose, and 5 mM pyruvate. The otic capsule from each side was opened to gain access into the inner ear. The sacculus was dissected, and the macula was excised from the sacculus. The macula was then incubated in 0.05% trypsin in Hanks' balanced salt solution (Invitrogen, Carlsbad, CA) for 8 to 10 min, after 30 s of rinsing in a low-calcium dissociation solution (105 mM NaCl, 2.5 mM KCl, 0.81 mM MgCl₂, 0.1 mM CaCl₂, 3.4 mM NaHCO₃, 0.5 mM NaH₂PO₄, and 2.5 mM Na₂HPO₄, 1 mM ascorbate, 4 mM glucose, and 5 mM pyruvate), which contained 10% fetal calf serum (Invitrogen). It was then washed in dissociation solution containing 500 μ g/ml bovine serum albumin for 5 to 10 min. The macula was then transferred to the recording chamber, previously filled with external solution, and hair cells were gently separated from the epithelium with the help of a thin, hook-shaped glass wisp. The hair cells settled to the base of the recording chamber and adhered firmly to the bottom, which was precoated with 2 mg/ml concanavalin A. The bath was perfused with external solution at a rate of 1 ml/min. All solutions had an osmolality of 220 mOsm and a pH of 7.2.

The recording chamber was placed on the stage of an inverted microscope (IMT-2; Olympus, Tokyo, Japan) equipped with a 40 \times objective with modulation-contrast optics. Continuous gravity perfusion (1 ml/min) was used for exchange of recording solutions. The drugs were applied at a rate of ~3.6 μ l/s using a gravity-driven microperfusion pipette placed ~300 to 600 μ m from the cell of interest. Experiments were performed at room temperature (20–22°C). Currents and voltages were recorded with the perforated patch-clamp technique. The antibiotic amphotericin B (0.6 mg/ml) was added to the internal solution, containing 75 mM KCl, 2 mM MgCl₂, 30 mM K₂SO₄, 6 mM glucose, and 10 mM HEPES. Borosilicate glass pipettes were pulled from 1.5 mm o.d. \times 0.75 mm i.d. capillary tubing (Longreach Scientific Resources, Orr's Island, ME) with a Flaming/Brown micropipette puller (model P-97; Sutter Instruments, Novato, CA) and heat-polished (Narashige Scientific Instrument La, Tokyo, Japan). The tip potential between the internal solution of the pipette and the bath was nulled before seal formation. The junction potential

¹ A pithed frog has had its central nervous system destroyed (its spinal cord has been severed). It is dead, but some of its organs continue to function for a brief period.

was calculated to be approximately -7 mV. The zero current potential was -48 ± 7 mV and ranged from -37 to -68 mV; the series resistance was 16 ± 4 M Ω and was partially compensated (80%) during voltage-clamp recordings using the compensation circuitry of the amplifier. Average cell capacitance was 15 ± 3 pF. Series resistance and cell capacitance compensations were updated continuously throughout all recordings.

The Axopatch 1D patch-clamp amplifier (Axon Instruments) was used for all voltage-clamp experiments in recording from saccular hair cells. Saccular hair cells were stepped from -60 mV (holding potential) to -10 mV during the voltage-clamp protocol. This protocol was used because -10 mV coincides with the peak of ACh-induced calcium activated outward K⁺ currents, as determined from previously reported I-V relationships (Erstegui et al., 1994). Voltage-clamp data were sampled at 50- to 300- μ s intervals. Records were low-pass filtered at 5 to 10 kHz with a four-pole Bessel filter. Stimuli were generated and data were sampled with a 12-bit digital-to-analog and analog-to-digital converter (DigiData 1200 series interface; Axon Instruments) and controlled by the data acquisition software package pClamp7 (Axon Instruments). Voltage-clamp data were stored digitally and later analyzed offline using the pClamp6 Clampfit (Axon Instruments), Excel 97 (Microsoft Corp., Redmond, WA), and Origin (ver. 5; OriginLab Corp., Northampton, MA) programs. Dose-response curves were normalized to the maximal agonist response in each hair cell. For the inhibition dose-response curves, cells were superfused with compounds for 1.5 to 2.5 min before coinjection with 20 μ M ACh (\sim EC₅₀), and responses were calculated as a percentage of the response elicited by the agonist.

Materials. ACh chloride or iodide was bought from Sigma Chemical Co. (St. Louis, MO). Serotonin hydrochloride, 1-(*m*-chlorophenyl)-biguanide hydrochloride, trimethyl serotonin iodide (5HTQ), 3-tropanyl-indole-3-carboxylate hydrochloride (tropisetron; ICS 205,930), and 3-tropanyl-3,5-dichlorobenzoate (MDL 72222) were obtained from RBI/Sigma (Natick, MA). Ondansetron hydrochloride was kindly donated by Raffo Laboratories (Buenos Aires, Argentina). Drugs were dissolved in distilled water as 10 mM stocks and stored aliquoted at -20°C . BAPTA-AM (Molecular Probes, Eugene, OR) was stored at -20°C as aliquots of a 100 mM solution in dimethyl sulfoxide, thawed, and diluted 1000-fold into saline solution shortly before incubation of the oocytes.

All experimental protocols were carried out in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals (<http://oacu.od.nih.gov/regs/guide/guidex.htm>).

Results

Effect of Serotonin on ACh-Evoked Currents through Recombinant $\alpha 9\alpha 10$ nAChRs. Figure 1A shows representative responses to 10 μ M ACh of *X. laevis* oocytes injected with $\alpha 9$ and $\alpha 10$ cRNAs. As expected for a nAChR, serotonin did not evoke currents through the $\alpha 9\alpha 10$ receptor. However, currents elicited by ACh were blocked by serotonin in a concentration-dependent manner (Fig. 1, A and B) with an IC₅₀ of 5.4 ± 0.6 μ M (mean \pm S.E.M.; $n = 6$). Block by serotonin was reversible, because initial control responses to ACh were recovered after washes of the oocytes with frog saline (data not shown).

To further characterize the mechanism underlying the blocking action of serotonin on the $\alpha 9\alpha 10$ receptor, the effect of this drug was analyzed at increasing concentrations of ACh (V_{hold} , -70 mV). As shown in Fig. 2A, 10 μ M serotonin produced a significant reduction of the agonist maximal response (percentage of maximal response, 58.1 ± 5.4 , $n = 4$) and no change in the EC₅₀ value (7.9 ± 1.0 and 11.6 ± 5.3 μ M in the absence and presence of serotonin, respectively). This result is compatible with a noncompetitive mechanism of

block, because the effect of serotonin is reversible. Moreover, as shown in the representative I-V curves of Fig. 2B, block by serotonin of ACh-evoked responses was voltage-dependent. Thus, whereas responses were not modified at positive depolarized potentials, they were diminished at negative hyperpolarized membrane holding potentials (I/I_{max} , 39 ± 4.7 and $20.1 \pm 0.4\%$ of control values at -90 mV for 10 and 30 μ M serotonin, respectively, $p < 0.01$, $n = 3$).

Effect of Serotonin Type 3 Receptor Agonists on the Recombinant $\alpha 9\alpha 10$ nAChR. To analyze whether the effect observed with serotonin could be extended to other serotonergic agonists, the effect of 5HTQ and 1-(*m*-chlorophenyl)-biguanide, selective for 5-HT₃ receptors, was studied on $\alpha 9\alpha 10$ -injected *X. laevis* oocytes. Contrary to that observed with serotonin, both 5-HT₃ agonists elicited inward currents through the $\alpha 9\alpha 10$ receptor (Fig. 3A). However, both compounds behaved as weak partial agonists, because maximal responses were only $10.8 \pm 0.8\%$ ($n = 3$) for 5HTQ and $3.1 \pm 0.7\%$ ($n = 3$) for 1-(*m*-chlorophenyl)-biguanide of the maximum obtained with ACh (Fig. 3B).

The effect of increasing concentrations of 1-(*m*-chlorophenyl)-biguanide was further analyzed at a fixed 10 μ M ACh concentration (V_{hold} , -70 mV). As shown in Fig. 4, A and B, responses to ACh were potentiated by low concentrations of 1-(*m*-chlorophenyl)-biguanide (3 – 10 μ M) that did not elicit currents per se, whereas inhibition of ACh-evoked currents predominated at higher concentrations of the compound. Potentiation reached $43.8 \pm 5.7\%$ ($n = 4$) at 3 μ M 1-(*m*-chlorophenyl)-biguanide and $21 \pm 5.5\%$ ($n = 6$) at 10 μ M. When analyzed at different holding potentials, both potentiation by 10 μ M 1-(*m*-chlorophenyl)-biguanide as well as block by 100 μ M 1-(*m*-chlorophenyl)-biguanide resulted in voltage independence (Fig. 4C). Thus, potentiation of responses at -90 mV ($32 \pm 13\%$), did not significantly differ from that at $+40$ mV ($51 \pm 15\%$; $n = 3$), and inhibition of responses at -90 mV ($65 \pm 6\%$) did not differ from that at $+40$ mV ($61 \pm 4\%$; $n = 4$). However, if the blocking effect of 1-(*m*-chlorophenyl)-biguanide starts at concentrations lower than 3 to 10 μ M, both potentiation and blockage will overlap at 10 μ M 1-(*m*-chloro-




Fig. 1. Effect of serotonin on ACh-evoked currents through recombinant $\alpha 9\alpha 10$ nAChRs. A, representative traces to ACh either alone or in the presence of serotonin. B, inhibition curve performed by the coapplication of 10 μ M ACh and increasing concentrations of serotonin. Oocytes were incubated with each concentration of serotonin for 2 min before the addition of ACh. Peak current values are plotted, expressed as the percentage of the peak control current evoked by ACh. The mean and S.E.M. of six experiments are shown.

phenyl)-biguanide. Therefore, the shape of the I-V curves at low concentrations of 1-(*m*-chlorophenyl)-biguanide might not solely reflect the potentiating action of the drug. The mechanism underlying potentiation of ACh responses in the presence of 1-(*m*-chlorophenyl)-biguanide was not further analyzed. As shown in Fig. 4D, block of ACh-evoked responses

in the presence of 100 μ M 1-(*m*-chlorophenyl)-biguanide was noncompetitive, because although the effect was reversible, block was not surmounted at high concentrations of ACh (V_{hold} , -70 mV). Thus, 100 μ M 1-(*m*-chlorophenyl)-biguanide produced a significant reduction of the agonist maximal response (percentage of maximal response, 32.6 ± 2.9 , $n = 10$)




Fig. 2. Mechanism of blockage by serotonin of ACh-evoked responses. A, concentration-response curves to ACh performed either alone or in the presence of 10 μ M serotonin. Peak current values were normalized and referred to the maximal peak response to ACh. The mean and S.E.M. of three to four experiments per group are shown. B, representative I-V curves obtained upon application of 2-s voltage ramps (-120 to +50 mV) 10 s after the peak response to 10 μ M ACh from a holding potential (V_{hold}) of -70 mV, either alone or in the presence of serotonin ($n = 3$ for each curve).




Fig. 3. Effect of serotonin type 3 receptor agonists on the recombinant $\alpha 9\alpha 10$ nAChR. A, representative current responses to 5-HT₃ receptor agonists. B, concentration-response curves to ACh, 5HTQ, and 1-(*m*-chlorophenyl)-biguanide. Peak current values were normalized and referred to the maximal peak response to ACh. The mean and S.E.M. of three experiments per group are shown.




Fig. 4. Effect of 1-(*m*-chlorophenyl)-biguanide on ACh-evoked currents. A, representative traces to ACh either alone or in the presence of 10 μ M or 100 μ M 1-(*m*-chlorophenyl)-biguanide. Oocytes were preincubated for 2 min with 1-(*m*-chlorophenyl)-biguanide before the addition of 10 μ M ACh. B, responses to 10 μ M ACh at different concentrations of 1-(*m*-chlorophenyl)-biguanide ($n = 4-6$); *, $p < 0.05$; **, $p < 0.01$, Tukey's test. C, representative I-V curves obtained upon application of 2-s voltage ramps (-120 to +50 mV) 10 s after the peak response to 10 μ M ACh from a holding potential (V_{hold}) of -70 mV, either alone or in the presence of 1-(*m*-chlorophenyl)-biguanide ($n = 3$ for each curve). D, concentration-response curves to ACh performed either alone or in the presence of 100 μ M 1-(*m*-chlorophenyl)-biguanide. Peak current values were normalized and referred to the maximal peak response to ACh. The mean and S.E.M. of 10 experiments per group are shown.

and no change in the EC₅₀ value [17.1 ± 1.0 and 19.8 ± 8.9 μM in the absence and presence of 1-(*m*-chlorophenyl)-biguanide, respectively].

Effect of Serotonin on ACh-Evoked Currents in Isolated Saccular Hair Cells. The effect of serotonin on a native hair cell cholinergic receptor preparation was analyzed on isolated frog vestibular hair cells. It has been described previously that hyperpolarization of the frog saccular hair cells in response to the exogenous application of ACh is caused by the activation of $\alpha 9$ -containing nicotinic cholinergic receptors and the subsequent activation of the small-conductance, Ca²⁺-activated (SK) potassium current (Athas et al., 1997; Lioudyno et al., 2000; Holt et al., 2001). The $\alpha 9$ -containing nAChR in frog saccular hair cells is pharmacologically indistinguishable from the $\alpha 9\alpha 10$ nAChR of the cochlear hair cells and the recombinant $\alpha 9\alpha 10$ nAChR, suggesting that both subunits are involved in the ACh-evoked response (Fuchs, 1996; Elgoyhen et al., 2001; Holt et al., 2001). As shown in the representative traces of Fig. 5A, the application of ACh to isolated saccular hair cells typically produces an outward current. The concentration of ACh that produces a half-maximal current response (EC₅₀) has been reported previously to be 18.4 μM (Holt et al., 2001).

The addition of the neurotransmitter serotonin to voltage-clamped saccular hair cells did not elicit a detectable response. However, the coapplication of serotonin and 20 μM ACh blocked the ACh-evoked currents in a concentration-dependent manner (Fig. 5, A and B). The concentration of serotonin required to block 50% of the response to ACh (7.5 ± 6.9 μM , $n = 4$) was in the same order of magnitude as that derived from Fig. 1B for recombinant $\alpha 9\alpha 10$ receptors.

Effect of Serotonin Type 3 Receptor Antagonists on ACh-Evoked Currents through Recombinant $\alpha 9\alpha 10$




Fig. 5. Effect of serotonin on ACh-evoked currents in isolated saccular hair cells. A, representative traces of ACh-evoked currents recorded from isolated frog saccular hair cells either alone or in the presence of serotonin (black traces). The membrane holding potential was stepped from -60 to -10 mV during the voltage-clamp protocol (gray, control traces). B, inhibition curve performed by the coapplication of 20 μM ACh and increasing concentrations of serotonin. Cells were incubated with each concentration of serotonin for 2 min before the addition of ACh. Current values are plotted, expressed as the percentage of the control current evoked by ACh. The data shown are the mean \pm S.E.M. of four experiments.

nAChRs. Tropisetron, an established blocker of 5-HT₃ receptors (Vanner and Sruprenant, 1990; Maricq et al., 1991), is one of the most potent antagonists described so far for the $\alpha 9\alpha 10$ nAChR (Fig. 6) (Elgoyhen et al., 2001). High sensitivity of $\alpha 9\alpha 10$ to 5-HT₃ antagonists seems to be a pharmacological property of this receptor and is not a peculiarity restricted to tropisetron. Thus, ACh-evoked currents in $\alpha 9\alpha 10$ -injected oocytes were also blocked by the 5-HT₃ antagonists ondansetron and MDL 72222 (Fig. 6, A and B). In all cases, the effect was concentration-dependent, with a rank order of potency of tropisetron (IC₅₀, 70.1 ± 0.9 nM, $n = 8$) > ondansetron (IC₅₀, 0.6 ± 0.1 μM , $n = 6$) = MDL 72222 (IC₅₀, 0.7 ± 0.1 μM , $n = 4$). Block by antagonists was reversible, because initial control responses to ACh were recovered after washes of the oocytes with frog saline (not shown). Moreover, antagonists did not elicit per se responses in oocytes expressing $\alpha 9\alpha 10$ receptors. On the other hand, ketanserin (3 μM), an antagonist of 5-HT_{2A} and 5-HT_{2C} metabotropic serotonin receptors, did not significantly block ACh-evoked currents ($n = 3$ oocytes).

Tropisetron is known to interact with 5-HT₃ receptor-binding sites and to block agonist-evoked responses in a competitive manner (Maricq et al., 1991). To further characterize the mechanism underlying the blocking effect of this compound on the $\alpha 9\alpha 10$ nAChR, block by tropisetron was analyzed at increasing concentrations of ACh. As shown in Fig. 6C, tropisetron produced an almost parallel rightward shift of ACh-evoked currents. A significant increase in the ACh EC₅₀ from 13.6 ± 2.9 ($n = 10$) to 31.1 ± 3.5 μM ($n = 10$; $p < 0.001$) was observed, with no changes in agonist maximal responses. This result is compatible with a competitive type of block. However, an increase in the Hill coefficient from 1.1 ± 0.1 to 1.8 ± 0.1 in the presence of tropisetron was observed ($p < 0.001$), suggesting that the underlying inhibitory mechanism could be more complex than just a competitive block. Schild plots were not constructed because high concentrations of ACh produced an increased desensitization of the receptor, resulting in nonreproducible responses.

Effect of Serotonin Type 3 Receptor Antagonists on ACh-Evoked Currents in Isolated Saccular Hair Cells. The native vestibular and cochlear hair cell cholinergic receptor and the recombinant $\alpha 9\alpha 10$ nAChR share similar pharmacological properties (for references, see Elgoyhen et al., 2001). To analyze whether this also holds true in the case of 5-HT₃ receptor antagonists, the effect of these compounds was studied on a native hair cell cholinergic receptor preparation, such as the isolated frog saccular hair cells.

Figure 7A shows representative traces to 20 μM ACh and block of these responses in the presence of either tropisetron, ondansetron, or MDL 72222. As shown previously for the $\alpha 9\alpha 10$ nAChR, block by antagonists was concentration-dependent (Fig. 7B), with a rank order of potency of tropisetron (IC₅₀, 0.33 ± 0.01 μM , $n = 4$) > ondansetron (IC₅₀, 0.64 ± 0.14 μM , $n = 4$) = MDL 72222 (IC₅₀, 0.8 ± 0.2 μM , $n = 4$).

Effect of Tropisetron on $\alpha 4\beta 2$ and $\alpha 7$ nAChRs. To analyze whether the block by tropisetron is selective for the $\alpha 9\alpha 10$ nAChR or is a general feature common to other nAChRs, the effect of this compound was studied on recombinant $\alpha 4\beta 2$ and $\alpha 7$ nAChRs. Although $\alpha 4\beta 2$ nAChRs were blocked by tropisetron (Fig. 8A), the IC₅₀ value (5.9 ± 0.6 μM , $n = 5$) derived from the concentration-response curves, was 2 orders of magnitude higher than that required to block $\alpha 9\alpha 10$

nAChRs. On the other hand, tropisetron elicited inward currents in $\alpha 7$ -injected oocytes, albeit with a maximal response that reached only $15.4 \pm 1.1\%$ of the maximum obtained with nicotine (Fig. 8B).

Discussion

The present study contributes to the pharmacological characterization of the $\alpha 9\alpha 10$ nAChR and indicates that this

receptor shares striking properties with type 3 serotonin ligand-gated ion channels. Moreover, it demonstrates an *in vivo* correlate of these properties of the recombinant receptor, with those of the $\alpha 9\alpha 10$ -containing nAChR of frog saccular hair cells. Finally, it poses the possibility that the biogenic amine serotonin might act as a neuromodulator of the cholinergic efferent transmission in the vestibular apparatus and in the organ of Corti.

The IC_{50} values found for tropisetron, ondansetron, and




Fig. 6. Effect of serotonin type 3 receptor antagonists on ACh-evoked currents through recombinant $\alpha 9\alpha 10$ nAChRs. A, representative traces to $10 \mu\text{M}$ ACh, either alone or in the presence of tropisetron, ondansetron, or MDL 72222. B, inhibition curves performed by the coapplication of $10 \mu\text{M}$ ACh and increasing concentrations of the antagonists. Oocytes were incubated with each concentration of antagonists for 2 min before the addition of ACh. Peak current values are plotted, expressed as the percentage of the peak control current evoked by ACh. C, concentration-response curves to ACh performed either alone or in the presence of $0.3 \mu\text{M}$ tropisetron. Peak current values were normalized and referred to the maximal peak response to ACh. The mean and S.E.M. of four to ten experiments per group are shown in B and C.




Fig. 7. Effect of serotonin type 3 receptor antagonists on ACh-evoked currents in isolated saccular hair cells. A, representative traces to $20 \mu\text{M}$ ACh, either alone or in the presence of tropisetron, ondansetron, or MDL 72222 (black traces). The cells were held at -60 mV and the command potential was stepped from -60 to -10 mV during the voltage-clamp protocol (gray, control traces). B, inhibition curves performed by the coapplication of $20 \mu\text{M}$ ACh and increasing concentrations of the antagonists. Cells were incubated with each concentration of antagonist for 2 min before the addition of ACh. Peak current values are plotted, expressed as the percentage of the peak control current evoked by ACh. The mean \pm S.E.M. of four to five experiments per antagonist concentration are shown.

MDL 72222 to block ACh-evoked currents on the $\alpha 9\alpha 10$ and the native hair cell receptor are in the same order of magnitude as those required for both recombinant (Maricq et al., 1991; Dubin et al., 1999) and native 5-HT₃ receptors present in the guinea pig submucosal plexus and the rabbit heart (Vanner and Sruprenant, 1990; Turconi et al., 1991). Moreover, among all the compounds tested on the $\alpha 9\alpha 10$ nAChR (Elgoyhen et al., 2001; Sgard et al., 2002), the 5-HT₃ antagonist tropisetron/as well as the glycinergic antagonist strychnine and the nicotinic antagonist α -bungarotoxin, have the highest blocking potencies. Thus, the heteromeric $\alpha 9\alpha 10$ receptor has peculiar pharmacological properties within the nAChR family, and resembles what has been described previously for the homomeric $\alpha 9$ recombinant receptor, which preserves pharmacological properties that are characteristic of other members of the Cys-loop family of ligand-gated ion channels (Rothlin et al., 1999). Moreover, this finding is in accordance with the fact that $\alpha 9$ shares with $\alpha 10$ the highest amino acid sequence identity, being more distantly related to other members of the nAChR family (Elgoyhen et al., 2001).

The binding of 5-HT₃ antagonists such as tropisetron to 5-HT₃ receptors is known to require several tryptophan residues in the extracellular amino terminus, probably involved in cation- π interactions with the positive amine in the tropane of these antagonists (Venkataraman et al., 1999; Yan et al., 1999; Spier and Lummis, 2000). Interestingly, in the $\alpha 9$ and $\alpha 10$ nAChR subunits, most residues aligned with the critical tryptophans are conserved. Thus the high-affinity site of interaction of tropisetron with the $\alpha 9\alpha 10$ nAChR could be extracellular and resemble its site on the 5-HT₃ receptor. This is supported by the finding that blockage of the $\alpha 9\alpha 10$ receptor by tropisetron is surmounted at high concentrations of ACh, compatible with a competitive type of block. Competitive antagonists of members of the Cys-loop family of ligand-gated ion channels are most probably binding with an extracellular site of the receptor (Karlin, 2002). Moreover, tropisetron turns to a high-affinity agonist of an $\alpha 9\alpha 10$ receptor bearing a leucine-to-threonine mutation at position L9' within transmembrane region II (P. V. Plazas, E. Katz,

and A. B. Elgoyhen, unpublished observations), further reinforcing the hypothesis of an extracellular amino-terminal binding site for this compound.

Within the nAChR family, high-affinity (nanomolar) block by the 5HT₃ antagonist tropisetron seems to be a distinctive feature of $\alpha 9$ (Rothlin et al., 1999) and $\alpha 9\alpha 10$ receptors. As derived from Fig. 8, micromolar concentrations of this antagonist are required to block $\alpha 4\beta 2$ receptors expressed in *X. laevis* oocytes. Moreover, affinity constants for tropisetron on $\alpha 4\beta 2$ and $\alpha 1\beta 1\gamma\delta$ derived from binding assays are all in the micromolar range (Macor et al., 2001). On the other hand, although tropisetron binds with high affinity to $\alpha 7$ nAChRs (Macor et al., 2001), it behaves as a weak partial agonist of this receptor subtype (Fig. 8) (Macor et al., 2001).

From a pharmacological standpoint, ondansetron and tropisetron are widely prescribed drugs for chemotherapy-induced emesis (Hesketh, 2000). A myriad of signaling pathways lead from the periphery to the emetic center, including inputs from the vestibular apparatus, of special importance during motion sickness (Pasricha, 2001). Because of its high affinity for the $\alpha 9\alpha 10$ receptor, tropisetron will block the $\alpha 9\alpha 10$ -containing vestibular cholinergic receptor at the concentrations used in therapeutics. This effect might (or might not) contribute to its antiemetic action, a question that will only be elucidated when a better understanding of the function of the efferent system to the vestibular apparatus is gained.

The observation that $\alpha 9\alpha 10$ and the native saccular hair cell receptors were blocked by serotonin at low micromolar concentrations and by an apparent noncompetitive mechanism was unexpected. Block of the closely related homomeric $\alpha 9$ receptor by serotonin seems to be competitive and requires high micromolar to millimolar concentrations (IC₅₀, 251 μ M; Rothlin et al., 1999), thus suggesting that serotonin probably binds to different sites within the $\alpha 9$ and the $\alpha 9\alpha 10$ receptor proteins. Moreover, the fact that block of ACh-evoked currents through $\alpha 9\alpha 10$ was dependent upon variations of the membrane holding potential, being evident only at hyperpolarized potentials, might suggest that serotonin is acting at a site within the channel lumen. An interaction of serotonin with the ion channel lumen has been reported for other nAChRs (Arias, 1998). However, although serotonin also blocks neuronal $\alpha 7$, $\alpha 3\beta 4$, $\alpha 4\beta 4$, $\alpha 2\beta 4$, and $\alpha 3\beta 2$ (Nakazawa et al., 1995; Palma et al., 1996), the apparent affinity for these receptor subtypes is 1 or 2 orders of magnitude lower than that reported here for the $\alpha 9\alpha 10$ nAChR. Therefore, within the nAChR family of receptors, the antagonist specificity of serotonin is highest for the $\alpha 9\alpha 10$ receptor.

Serotonin differs from other 5-HT₃ receptor agonists such as 5HTQ and 1-(*m*-chlorophenyl)-biguanide in that the latter behave as weak partial agonists of the recombinant $\alpha 9\alpha 10$ receptor. However, as observed in Fig. 4, the effect of 1-(*m*-chlorophenyl)-biguanide is more complex than just a partial agonism, because both potentiation and inhibition of ACh-evoked responses were observed, probably indicating more than one site of interaction of the compound with the receptor. As reported for the effect of atropine on $\alpha 4\beta 4$ nicotinic receptors and curare-like drugs on neuronal nicotinic receptors containing the $\beta 4$ subunit, where both potentiation and blockage of ACh-evoked responses are observed (Cachelin and Rust, 1994; Zwart and Vivjerberg, 1997), potentiation might result from the interaction with the agonist binding




Fig. 8. Effect of tropisetron on $\alpha 4\beta 2$ and $\alpha 7$ nAChRs. A, inhibition curve performed by the coapplication of 10 μ M ACh and increasing concentrations of tropisetron to $\alpha 4\beta 2$ -injected oocytes. Peak current values are plotted, expressed as the percentage of the peak control current evoked by ACh. B, concentration-response curves to tropisetron and nicotine in $\alpha 7$ -injected oocytes. Peak current values were normalized and referred to the maximal peak response to nicotine. The mean and S.E.M. of three to five experiments per group are shown.

site of the receptor, where simultaneous occupation of the two binding sites, one agonist recognition site with ACh and the other agonist recognition site with 1-(*m*-chlorophenyl)-biguanide, leads to subsequent ion channel activation. On the other hand, blockage was noncompetitive and independent of the membrane holding potential, thus suggesting an interaction with a regulatory site independent of the ion channel pore. As for all experiments in which more than one receptor subunit is expressed, one cannot preclude the possibility of assembly of different receptors subtypes with different stoichiometries to account for the different effects of 1-(*m*-chlorophenyl)-biguanide. Further studies that go beyond the scope of the present work will be required to fully explain the complex effects of 1-(*m*-chlorophenyl)-biguanide on the $\alpha 9\alpha 10$ receptor.

Does the interaction of serotonin with the $\alpha 9\alpha 10$ recombinant and the saccular hair cell cholinergic receptor underlie a physiological implication? The function of a serotonergic innervation to the inner ear is unknown. Intracochlear injection of serotonin reduces the compound action potential of the auditory nerve (Bobbin and Thompson, 1978). Recent evidence suggests the presence of serotonin and its metabolite 5-hydroxyindole-3-acetic acid in the mammalian cochlea and vestibule (Gil-Loyzaga et al., 1997a,b, 2000) and subunit A of 5-HT₃ receptors has been identified by in situ hybridization in the cochlear and vestibular sensory epithelium of the developing rat (Johnson and Heinemann, 1995). In the case of the organ of Corti, serotonergic fibers reach the cochlea accompanying the olivocochlear efferent system. At the periphery, these fibers form glomerulus-like structures within the inner spiral bundle and could directly contact inner hair cells. They reach the Corti's tunnel and then follow a spiral distribution, branching at least on the first row of outer hair cells (Gil-Loyzaga et al., 2000). The fact that serotonergic fibers are present at the base of the outer hair cells, together with the finding that low micromolar concentrations of serotonin block ACh-evoked currents through the $\alpha 9\alpha 10$ receptor, opens the possibility that serotonin might function as a neuromodulator of the efferent cholinergic innervation to the inner ear.

Acknowledgments

We thank Dr Jim Boulter (UCLA) for sharing the $\alpha 4$, $\alpha 7$, and $\beta 2$ rat nAChR subunits cDNAs and Raffo Laboratories (Buenos Aires, Argentina) for the kind donation of ondansetron hydrochloride.

References

- Anand R, Conroy W, Schoepfer R, Whiting P, and Lindstrom J (1991) Neuronal nicotinic acetylcholine receptors expressed in *Xenopus* oocytes have a pentameric quaternary structure. *J Biol Chem* **266**:11192–11198.
- Arias H (1998) Noncompetitive inhibition of nicotinic acetylcholine receptors by endogenous molecules. *J Neurosci Res* **52**:369–379.
- Athas GB, Norris CH, Guth PS, and Garcia MM (1997) Expression profiling of vestibular hair cells from the frog. *Assoc Res Otolaryng Abs* **20**:37.
- Bobbin RP and Thompson MH (1978) Effects of putative transmitters on afferent cochlear transmission. *Ann Otol Rhinol Laryngol* **87**:185–190.
- Cachelin AB and Rust G (1994) Unusual pharmacology of (+)-tubocurarine with rat neuronal nicotinic acetylcholine receptors containing $\beta 4$ subunits. *Mol Pharmacol* **46**:1168–1174.
- Cooper E, Couturier S, and Ballivet M (1991) Pentameric structure and subunit stoichiometry of a neuronal nicotinic acetylcholine receptor. *Nature (Lond)* **350**:235–238.
- Dubin AE, Huvar R, D'Andrea MR, Pyati J, Zhu JY, Joy KC, Wilson SJ, Galindo JE, Glass CA, Luo L, et al. (1999) The pharmacological and functional characteristics of the serotonin 5-HT_{3B} receptor are specifically modified by a 5-HT_{3B} receptor subunit. *J Biol Chem* **274**:30799–30810.
- Elgoyhen AB, Johnson DS, Boulter J, Vetter DE, and Heinemann S (1994) Alpha9: an acetylcholine receptor with novel pharmacological properties expressed in rat cochlear hair cells. *Cell* **79**:705–715.
- Elgoyhen AB, Vetter D, Katz E, Rothlin C, Heinemann S, and Boulter J (2001) Alpha 10: A determinant of nicotinic cholinergic receptor function in mammalian vestibular and cochlear mechanosensory hair cells. *Proc Natl Acad Sci USA* **98**:3501–3506.
- Erostegei C, Nenov AP, Norris CH, and Bobbin RP (1994) Acetylcholine activates a K⁺ conductance permeable to Cs⁺ in guinea pig outer hair cells. *Hearing Res* **81**:119–129.
- Fuchs P (1996) Synaptic transmission at vertebrate hair cells. *Curr Opin Neurobiol* **6**:514–519.
- Galzi J-L, Revah F, Bessis A, and Changuex JP (1991) Functional architecture of the nicotinic acetylcholine receptor: from electric organ to brain. *Annu Rev Pharmacol* **31**:37–72.
- Gil-Loyzaga P, Bartolomé MV, and Vicente-Torres MA (1997a) Serotonergic innervation of the organ of Corti of the cat cochlea. *Neuroreport* **8**:3519–3522.
- Gil-Loyzaga P, Bartolomé V, Vicente-Torres A, and Carricondo F (2000) Serotonergic innervation of the organ of Corti. *Acta Otolaryngol* **120**:128–132.
- Gil-Loyzaga P, Vicente-Torres MA, Garcia-Bonacho M, and Esquifino A (1997b) Presence of catecholamines and serotonin in the rat vestibule. *Brain Res* **746**:265–268.
- Hesketh PJ (2000) Comparative review of 5-HT₃ receptor antagonists in the treatment of acute chemotherapy-induced nausea and vomiting. *Cancer Investig* **18**:163–173.
- Hiel H, Elgoyhen AB, Drescher D, and Morley B (1996) Expression of nicotinic acetylcholine receptor mRNA in the adult rat peripheral vestibular system. *Brain Res* **738**:347–352.
- Holt JC, Lioudyno M, Athas G, Garcia MM, Perin P, and Guth PS (2001) The effect of proteolytic enzymes on the $\alpha 9$ -nicotinic receptor-mediated response in isolated frog vestibular hair cells. *Hearing Res* **152**:25–42.
- Johnson DS and Heinemann SF (1995) Embryonic expression of the 5-HT₃ receptor subunit, 5-HT_{3A}, in the rat: an in situ hybridization study. *Mol Cell Neurosci* **6**:122–138.
- Karlin A (2002) Ion channel structure: emerging structure of the nicotinic acetylcholine receptors. *Nat Rev Neurosci* **3**:102–114.
- Katz E, Verbitsky M, Rothlin C, Vetter D, Heinemann S, and Elgoyhen AB (2000) High calcium permeability and calcium block of the $\alpha 9$ nicotinic acetylcholine receptor. *Hearing Res* **141**:117–128.
- Le Novère N and Changeux JP (1999) The ligand gated channel database. *Nucleic Acids Res* **27**:340–342.
- Liman ER, Tytgat J, and Hess P (1992) Subunit stoichiometry of a mammalian K⁺ channel determined by construction of multimeric cDNAs. *Neuron* **9**:861–871.
- Lioudyno M, Verbitsky M, Holt J, Elgoyhen AB, and Guth P (2000) Morphine inhibits an $\alpha 9$ -acetylcholine nicotinic receptor-mediated response by a mechanism which does not involve opioid receptors. *Hearing Res* **149**:167–177.
- Macor JE, Gurley D, Lanthorn T, Loch J, Mack RA, Mullen G, Tran O, Wright N, and Gordon JC (2001) The 5-HT₃ antagonist tropisetron (ICS 205–930) is a potent and selective nicotinic receptor partial agonist. *Bioorg Med Chem Lett* **11**:319–321.
- Marić A, Peterson A, Brake A, Myers R, and Julius D (1991) Primary structure and functional expression of the 5HT₃ receptor, a serotonin-gated ion channel. *Science (Wash DC)* **254**:432–436.
- Morley B, Li H, Hiel H, Drescher D, and Elgoyhen AB (1998) Identification of the subunits of the nicotinic cholinergic receptors in the rat cochlea using RT-PCR and in situ hybridization. *Mol Brain Res* **53**:78–87.
- Nakazawa K, Akiyama T, and Inoue K (1995) Block by 5-hydroxytryptamine of neuronal acetylcholine receptor channels expressed in *Xenopus* oocytes. *Cell Mol Neurobiol* **15**:495–500.
- Palma E, Miledi AM, Eusebi F, and Miledi R (1996) Threonine-for-leucine mutation within domain M2 of the neuronal $\alpha 7$ nicotinic receptor converts 5-hydroxytryptamine from antagonist to agonist. *Proc Natl Acad Sci USA* **93**:11231–11235.
- Pasricha PJ (2001) Prokinetic agents, antiemetics and agents used in irritable bowel syndrome, in *Goodman and Gilman's The Pharmacological Basis of Therapeutics* (Hardman JG and Limbird LE eds) pp 1021–1036, McGraw Hill, New York.
- Rothlin C, Verbitsky M, Katz E, and Elgoyhen AB (1999) The $\alpha 9$ nicotinic acetylcholine receptor shares pharmacological properties with type A γ -aminobutyric acid, glycine and type 3 serotonin receptors. *Mol Pharmacol* **55**:248–254.
- Spier AD and Lummis SCR (2000) The role of tryptophan residues in the 5-Hydroxytryptamine₃ receptor ligand binding domain. *J Biol Chem* **275**:5620–5625.
- Turconi M, Donetti A, Schiavone A, Sagrafa A, Montagna E, Nicola M, Cesana R, Rizzi C, and Micheletti R (1991) Pharmacological properties of a novel class of 5-HT₃ receptor antagonists. *Eur J Pharmacol* **203**:203–211.
- Vanner S and Sruprenant A (1990) Effects of 5-HT₃ receptor antagonists on 5-HT and nicotinic depolarizations in guinea-pig submucosal neurones. *Br J Pharmacol* **99**:840–844.
- Venkataraman P, Ma C, Kirschbaum KS, and Schulte MK (1999) Tryptophan 89 of the 5HT₃ receptor forms a cation- π interaction with the amino group of 5HT_{3R}. *Soc Neurosci Abstr* **25**:202.
- Yan D, Schulte MK, Bloom KE, and White MM (1999) Structural features of the ligand-binding domain of the serotonin 5HT₃ receptor. *J Biol Chem* **274**:5537–5541.
- Zwart R and Vjiverberg HPM (1997) Potentiation and inhibition of neuronal nicotinic receptors by atropine: competitive and non-competitive effects. *Mol Pharmacol* **52**:886–895.

Address correspondence to: Ana Belén Elgoyhen, Instituto de Investigaciones en Ingeniería Genética y Biología Molecular (CONICET-UBA), Vuelta de Obligado 2490, 1428 Buenos Aires, Argentina. E-mail: elgoyhen@dna.uba.ar