



Mood Symptom Dimensions and Developmental Differences in Neurocognition in Adolescence

Journal:	<i>Clinical Psychological Science</i>
Manuscript ID	CPX-21-0181.R2
Manuscript Type:	Empirical Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Kaiser, Roselinde; University of Colorado Boulder, Psychology and Neuroscience; University of Colorado Boulder, Institute of Cognitive Science; University of Colorado Boulder, Renee Crown Wellness Institute</p> <p>Moser, Amelia; University of Colorado Boulder, Psychology and Neuroscience; University of Colorado Boulder, Institute of Cognitive Science</p> <p>Neilson, Chiara; University of Colorado Boulder, Psychology and Neuroscience; University of Colorado Boulder, Institute of Cognitive Science</p> <p>Peterson, Elena; University of Colorado Boulder, Psychology and Neuroscience; University of Colorado Boulder, Renee Crown Wellness Institute</p> <p>Jones, Jenna; University of Colorado Boulder, Institute of Cognitive Science</p> <p>Hough, Christina; University of California Los Angeles, Psychology</p> <p>Rosenberg, Benjamin; University of California Los Angeles, Psychology</p> <p>Sandman, Christina; University of California Los Angeles, Psychology</p> <p>Schneck, Christopher; University of Colorado Denver School of Medicine, Psychiatry</p> <p>Miklowitz, David; University of California Los Angeles, Psychiatry</p> <p>Friedman, Naomi; University of Colorado Boulder, Institute for Behavioral Genetics</p>
Keywords:	reward sensitivity, executive functioning, mania, anhedonia, adolescence
Abstract:	<p>Adolescence is critical period of neurocognitive development as well as increased prevalence of mood pathology. This cross-sectional study replicated developmental patterns of neurocognition and tested whether mood symptoms moderated developmental effects. Participants were 419 adolescents (n=246 with current mood disorders) who completed reward learning and executive functioning tasks, and reported on age, puberty, and mood symptoms. Structural equation modeling revealed a quadratic relationship between puberty and reward learning performance that was moderated by symptom severity: in early puberty, adolescents reporting higher manic symptoms exhibited heightened reward learning performance (better maximizing of rewards on learning tasks), whereas adolescents reporting elevated anhedonia showed blunted reward learning performance. Models also showed a linear relationship between age and executive functioning that was moderated by manic symptoms:</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	adolescents reporting higher mania showed poorer executive functioning at older ages. Findings suggest neurocognitive development is altered in adolescents with mood pathology and suggest directions for longitudinal studies.



1
2
3
4
5
6
7 **Mood Symptom Dimensions and Developmental Differences in Neurocognition in**
8
9 **Adolescence**

10
11 Roselinde H. Kaiser^{1,2,3}, Amelia D. Moser^{1,2}, Chiara Neilson^{1,2}, Elena C. Peterson¹, Jenna
12 Jones^{1,2}, Christina Hough⁴, Benjamin Rosenberg⁴, Christina F. Sandman⁴, Christopher D.
13 Schneck⁵, David J. Miklowitz⁶, & Naomi P. Friedman^{1,2,7}
14

15
16 ¹ Department of Psychology and Neuroscience, University of Colorado Boulder

17 ² Institute of Cognitive Science, University of Colorado Boulder

18 ³ Renée Crown Wellness Institute, University of Colorado Boulder

19 ⁴ Department of Psychology, University of California Los Angeles

20 ⁵ Department of Psychiatry, University of Colorado Denver

21 ⁶ Department of Psychiatry, University of California Los Angeles

22 ⁷ Institute of Behavioral Genetics, University of Colorado Boulder
23
24

25 *Corresponding author:

26 Roselinde Kaiser, Ph.D.

27 Research on Affective Disorders and Development (RADD) Lab

28 Department of Psychology and Neuroscience

29 University of Colorado Boulder

30 Muenzinger 345 UCB

31 Boulder, CO 80309.

32 Email: Roselinde.Kaiser@colorado.edu
33
34

35 Keywords: reward sensitivity, executive functioning, age, puberty, mania, anhedonia
36
37

38 Abstract: 150 words

39 Article body (including references): ~9,780 words

40 Figures: 2

41 Tables: 2

42 Supplements: 1
43
44

45 Short title: Developmental differences, mood, and neurocognition
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Adolescence is critical period of neurocognitive development as well as increased prevalence of mood pathology. This cross-sectional study replicated developmental patterns of neurocognition and tested whether mood symptoms moderated developmental effects. Participants were 419 adolescents ($n=246$ with current mood disorders) who completed reward learning and executive functioning tasks, and reported on age, puberty, and mood symptoms. Structural equation modeling revealed a quadratic relationship between puberty and reward learning performance that was moderated by symptom severity: in early puberty, adolescents reporting higher manic symptoms exhibited heightened reward learning performance (better maximizing of rewards on learning tasks), whereas adolescents reporting elevated anhedonia showed blunted reward learning performance. Models also showed a linear relationship between age and executive functioning that was moderated by manic symptoms: adolescents reporting higher mania showed poorer executive functioning at older ages. Findings suggest neurocognitive development is altered in adolescents with mood pathology and suggest directions for longitudinal studies.

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Adolescence is a critical period for the onset of mood disorders (Paus et al., 2008). The majority of first-episodes of mania and depression occur in adolescence (Kessler et al., 2005) and mood symptoms in adolescence precede and are associated with adult mood disorders (Bertha & Balazs, 2013). Adolescence is also characterized by remarkable changes in neurocognitive functioning, and specifically in neurocognitive domains in which abnormalities have been observed in mood disorders (Nielson et al., 2020; Nusslock & Alloy, 2017; Zald & Treadway, 2017). These converging patterns suggest that abnormal development of neurocognition may be a marker for bipolar and unipolar mood disorders (Alloy & Nusslock, 2019; Kujawa et al., 2020). In light of the translational implications of such work (Gaffrey et al., 2013; Van Rhee et al., 2021), exploration of abnormal neurocognitive development in mood pathology is needed.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Two areas of neurocognitive functioning are especially relevant to adolescent development and mood pathology: *reward sensitivity*, defined as an individual's behavioral, physiological, or emotional responsiveness to rewards (Berridge, 2018), and *executive functioning*, which includes higher-order cognitive abilities that support goal-directed behavior and adaptation (Friedman & Miyake, 2017; Miyake et al., 2000). Both reward sensitivity and executive functioning undergo changes during adolescence, but prior research indicates that developmental timing differs between these neurocognitive domains. Reward sensitivity follows a curvilinear pattern, increasing steeply in early adolescence, and peaking in mid-adolescence before declining or reaching asymptote in late adolescence into young adulthood (Casey et al., 2008; Luna & Wright, 2016; Shulman et al., 2016). In contrast, executive functioning shows linear improvement across adolescence into young adulthood (Luna et al., 2015; Luna & Wright, 2016), with peak levels of executive functioning not achieved until mid-twenties or later (Best & Miller, 2010). These differences in trajectory are paralleled by differences in the developmental

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

4

factors believed to contribute to changes in each domain: Puberty and pubertal hormones are related to reward-approach behavior (although the nature of such associations may depend on sex (Braams et al., 2015; Harden et al., 2018)), whereas age is associated with executive functioning (Best & Miller, 2010) (also see (Blakemore et al., 2010; Forbes & Dahl, 2010)). Together, this research has motivated a set of dual-systems models (Geier & Luna, 2009; Shulman et al., 2016; Steinberg et al., 2008) proposing that adolescent behavior is shaped by the distinct developmental trajectories of reward sensitivity and executive functioning, with the former more strongly related to pubertal changes and the latter driven by factors related to aging (e.g., maturation of prefrontal cortical regions) (Casey et al., 2008).

Reward sensitivity and executive functioning are also key domains of neurocognitive dysfunction in mood disorders. Reward sensitivity is conceptually related to symptom dimensions of mood pathology, including depressive anhedonia (loss of interest and pleasure) and mania/hypomania. Indeed, blunted reward sensitivity has been associated with anhedonia across mood diagnoses (Forbes & Dahl, 2012; Huys et al., 2013; Pizzagalli, 2014; Pizzagalli et al., 2008; Zald & Treadway, 2017). In contrast, abnormalities in reward sensitivity have been observed in bipolar disorders and at higher levels of manic symptom severity (Alloy & Nusslock, 2019), including increased corticostriatal reward response (Nusslock et al., 2012; Phillips & Swartz, 2014). However, evidence is mixed regarding the direction of reward processing abnormalities in bipolar disorders. For example, some research has indicated reward hyposensitivity in the form of blunted reward response bias (Pizzagalli et al., 2008) or lower self-reported reward response (Biuckians et al., 2007). Other studies provide evidence that reward anomalies in bipolar or highly manic youth may depend on sex, developmental processes, or co-recruitment of other cognitive abilities (Maresh et al., 2019; Urošević, Luciana, et al., 2016;

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

5

1
2
3 Urošević, Youngstrom, et al., 2016). Therefore, the nature of reward abnormalities associated
4
5 with mood disorders and symptoms is complex and may depend on the form of reward
6
7 sensitivity that is targeted.
8
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Meanwhile, global and overlapping deficits in executive functioning have been observed across unipolar and bipolar mood disorders (Lee et al., 2014; Rock et al., 2014; Snyder, 2013) including adolescent samples (Horn et al., 2011; Joseph et al., 2008; Wagner et al., 2015), and the severity of such deficits is shown to scale with severity of illness (Dixon et al., 2004; Scult et al., 2017). Executive deficits may contribute to cognitive symptoms of mood pathology (e.g., problems concentrating, racing thoughts), but also to difficulties regulating affective processes that characterize mood disorders (Lima et al., 2018). Together, this research suggests that overlapping executive functioning deficits but (possibly) distinctive reward processing anomalies characterize mood symptom dimensions of anhedonia and mania/hypomania (Nusslock & Alloy, 2017).

The convergence of dual systems changes and heightened mood symptoms in adolescence has led to integrative models that aim to describe how mood pathology shapes — or is shaped by — developmental factors (Alloy et al., 2016; Alloy & Nusslock, 2019; Forbes et al., 2021; Forbes & Dahl, 2012). However, there are several key gaps in empirical work into such models. First, prior work on mood pathology in adolescence has typically explored executive functioning and reward processing separately, limiting our ability to compare across neurocognitive domains or link results to dual systems models informed by developmental theory. Second and relatedly, although these separate lines of research into executive functioning and reward processing have revealed abnormalities in adolescents with mood disorders (Horn et al., 2011; Wagner et al., 2015), mixed evidence regarding the presence or magnitude of such

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

6

1
2
3 effects (Joseph et al., 2008; Reyes et al., 2017) makes their reliability unclear. One source of
4
5 unreliability in these studies may be heterogeneity in the tasks used to evaluate reward sensitivity
6
7 or executive functioning. Given their unique characteristics, any individual task paradigm is
8
9 contaminated by perceptual or other task-specific factors (Friedman et al., 2008) and
10
11 neuropsychological tasks often draw upon multiple cognitive functions (Snyder et al., 2015).
12
13 Research that adopts a multi-measure, latent variable approach to estimate reward response and
14
15 executive functioning in clinical samples may provide more reliable evaluation of these
16
17 constructs.
18
19

20
21
22 Third, the majority of research examining adolescent mood pathology and neurocognitive
23
24 functioning has taken a categorical, diagnostic approach. This approach has limitations, given
25
26 that within both unipolar and bipolar diagnostic categories there is significant heterogeneity in
27
28 phenotype (Cuthbert & Insel, 2010; McIntyre et al., 2015) and neurocognitive functioning (Van
29
30 Rheenen et al., 2020). In turn, symptom dimensions of anhedonia and mania may be as or more
31
32 strongly related to neurocognitive indices than diagnoses, and subclinical symptoms have been
33
34 related to functionally significant impairments (Ayuso-Mateos et al., 2010; Copeland et al.,
35
36 2015; Vrieze et al., 2013). Investigation of symptom dimensions in transdiagnostic samples can
37
38 provide important insight that complements information from diagnoses.
39
40

41
42 Fourth, few studies have incorporated developmental differences into the investigation of
43
44 mood pathology and neurocognitive functioning (Gaffrey et al., 2013). For example, although
45
46 pubertal status has been linked to reward processing (Harden et al., 2018) and puberty onset is
47
48 associated with both depression and mania (Mendle, 2014; Yazici et al., 2013), these factors
49
50 have not been well integrated. Prior clinical research has viewed puberty and adolescence as a
51
52
53
54
55
56
57
58
59
60

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

7

1
2
3 window of risk, but there is a lack of conclusive evidence for whether or how developmental
4
5 differences in neurocognitive functioning are altered in individuals with mood pathology.
6

7
8 The present study aimed to address the above gaps by investigating reward sensitivity
9
10 and executive functioning in relation to developmental variables and mood symptom dimensions
11
12 in a large sample of adolescents. This study was a cross-sectional investigation, an approach that
13
14 can test developmental differences in neurocognitive functioning but cannot answer questions
15
16 about within-person development over time. Neurocognitive functioning was measured with a
17
18 methodologically robust multi-task design. For the present study, we chose to operationalize
19
20 reward sensitivity as performance across a set of widely used reward learning tasks based on
21
22 prior research indicating that this dimension of reward sensitivity is altered in mood disorders
23
24 (Zald & Treadway, 2017) and undergoes adolescent development (Nussenbaum & Hartley,
25
26 2019). In turn, we chose to operationalize executive functioning in the form of performance
27
28 across gold-standard executive functioning tasks based on evidence that common executive
29
30 functions are broadly disrupted in psychopathology (Friedman et al., 2018) and improve over
31
32 development (Friedman et al., 2016).
33
34
35
36

37
38 Structural equation modeling (SEM) tested the following hypotheses in a single model.
39
40 First, we predicted that there would be a significant curvilinear association between pubertal
41
42 stage and a latent reward learning performance factor. Second, we predicted that the association
43
44 between pubertal stage and reward learning performance would be moderated by manic
45
46 symptom severity and by anhedonia severity, and that these patterns of moderation would differ
47
48 from one another. We did not have specific predictions for the directions of moderation, but we
49
50 predicted that adolescents reporting high severity of manic symptoms would show elevations in
51
52 reward learning performance that varied by pubertal stage, whereas adolescents reporting high
53
54
55
56
57
58
59
60

1
2
3 severity of anhedonic symptoms would show reductions in reward learning performance that
4 varied by pubertal stage. Third, we predicted a significant linear association between age and a
5 latent executive functioning factor. Fourth, we predicted that the association between age and
6 executive functioning would be moderated by manic and anhedonic symptom severity, but we
7 predicted an overlapping pattern, in which age-related increases in executive functioning would
8 be weaker at (any) higher symptom severity. Together, the SEM was designed to replicate
9 developmental dual-systems research in a large sample with a robust methodological approach,
10 and extend this research to address the question of how dual-systems developmental differences
11 are altered in adolescents reporting disturbances in specific mood dimensions. Below and in
12 Supplement, we report how we determined our sample size, all data exclusions, all
13 manipulations, and all measures in the study.

31 Method

33 Participants

34 [Table 1 about here]

35 We recruited 419 participants to research testing at two sites (University of Colorado
36 Boulder, University of California Los Angeles) (Tables 1-2; also see Supplement and Table S1).
37 To be eligible for these studies, participants were fluent in English or native English speakers,
38 had normal or corrected-to-normal vision, and reported no history of head injury, neurological
39 conditions, or other conditions that would interfere with cognitive testing. Recruitment was
40 aimed at individuals in the developmental period of adolescence, which can be defined as
41 beginning when an individual enters puberty, and ending when the individual transitions to
42 independence in social-emotional functioning (Forbes & Dahl, 2010). In practice, recruitment
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

9

1
2
3 across studies was anchored to age (ages 13 to 24) and social role (e.g., middle, high-school, or
4 college students who had not yet entered the workforce). Recruitment was also aimed at high
5 variance in clinical mood symptoms, including those with bipolar or unipolar mood disorders (in
6 final sample, $n=34$ with full or subthreshold bipolar disorders, $n=184$ with full or subthreshold
7 unipolar disorders, $n=28$ with schizoaffective or other mood disorders, $n=173$ with no mood
8 disorders; Table 2). To achieve this sample, recruitment included distribution of research
9 advertisements and outreach in community, school, healthcare, and outpatient psychiatric
10 settings in the Boulder and Los Angeles metropolitan areas. Research protocols for each study
11 were approved by the Institutional Review Boards at each site. Consent was obtained from legal
12 adults (ages 18 and older) and parental consent and child assent was obtained from legal minors
13 (ages 17 and younger) participating in the study.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 [Table 2 about here]

30 Procedures

31
32
33 Participants were recruited for an in-person research session that included a series of
34 behavioral tests and surveys. Participants were informed that they would receive a monetary
35 bonus based on their task performance and winnings to enhance motivation. Participants did not
36 know the specific bonus amount and did not receive the bonus until experimental procedures
37 were completed. After behavioral testing, clinical evaluation was performed (along with
38 procedures that will be reported elsewhere, see Supplement).

46 Measures

48 *Neurocognitive Functioning*

49
50
51 To evaluate neurocognitive functioning, six behavioral tasks were electronically
52 administered with a research staff member present in the room to assist in instruction and data
53
54
55
56
57
58
59
60

1
2
3 collection (Figure S1). Tasks varied between 7-12 minutes per task, and administration of all six
4
5 tasks (including instructions and breaks between tasks) took approximately one hour. Task
6
7 administration was conducted using a Mac Mini, a 27" high-resolution monitor, and a button box
8
9 with millisecond accuracy to record responses and reaction time (RT). Verbal responses (to the
10
11 antisaccade task) were recorded by the research assistant. Tasks were programmed in PsychoPy
12
13 version 1.85.1. Stimuli within tasks were counterbalanced and randomized, and the order of
14
15 stimuli within each task was standardized across participants. Each task included practice trials
16
17 or description of stimuli at the beginning, to confirm that participants understood instructions and
18
19 could perform the task. The order of task administration was standardized to be one of two
20
21 sequences, and participants were randomly assigned to a sequence (there were no significant
22
23 differences in performance between task sequences).

24
25
26
27
28 **Reward Learning Tasks.** The learning tasks used to index reward sensitivity were
29
30 selected because they are widely-used and well-validated in developmental and clinical samples
31
32 (Andre Der-Avakian et al., 2016; Lawlor et al., 2020; Master et al., 2020; Nussenbaum &
33
34 Hartley, 2019; Pizzagalli et al., 2005). Reward tasks featured overlapping instrumental learning
35
36 demands and monetary rewards, but differed in perceptual characteristics and task structure,
37
38 making this set of tasks eligible for latent variable analysis in which the shared process across
39
40 tasks was behavioral ability to maximize reward by learning stimulus-response associations.
41
42
43

44
45 **Two-armed Bandit Task.** The two-armed bandit is a probability learning task. In each
46
47 trial the participant was presented with a pair of stimuli (symbols) and instructed to make a
48
49 choice that elicited a desired outcome, using a button to select either the top or the bottom
50
51 stimulus. For one stimulus pair the desired outcome was to gain monetary reward, for a second
52
53 stimulus pair the desired outcome was to avoid monetary loss, and for the third stimulus pair the
54
55
56
57
58
59
60

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

11

1
2
3 desired outcome was to view a special graphic. Unbeknownst to the participant, within each
4
5 stimulus pair, one stimulus had an 80% probability of a desired outcome, and the other stimulus
6
7 had a 20% probability of a desired outcome. Participants had up to 4 seconds to respond for each
8
9 stimulus pair; after making a choice, the outcome (reward, loss, special neutral outcome, or null
10
11 outcome) was displayed for 2 seconds. The task consisted of 72 trials, with 24 trials for each
12
13 condition.
14
15

16
17 ***Probabilistic Reward Task.*** The probabilistic reward task (Pizzagalli et al., 2005) is a
18
19 probability learning task that takes a signal detection approach in which participants are
20
21 instructed to discriminate between two closely-matched stimuli (McCarthy & Davison, 1979).
22
23 For each trial, the participant was presented (for 500ms) with a cartoon face, and either a short or
24
25 long mouth stimulus (for 100ms) appeared on the face. The participant was asked to indicate
26
27 which stimulus was displayed as quickly as possible by pressing a button. Correct responses
28
29 either resulted in reward feedback (“Correct! You win \$0.05!”) or null feedback (blank screen).
30
31 The reward reinforcement schedule was asymmetrical: One “rich” stimulus was rewarded for
32
33 correct responses at a rate that was three times higher than the reward rate for the “lean” stimulus
34
35 (the mouth stimulus that was “rich”, long vs. short, was counterbalanced across participants).
36
37 The task included two blocks of 100 trials each, and in each block 30 of the 50 rich trials (60%)
38
39 and 10 of the 50 lean trials (20%) was followed by reward feedback.
40
41
42
43

44
45 ***Instrumental Learning Task.*** This instrumental learning task (modeled on (Collins &
46
47 Frank, 2012)) is a stimulus-response learning task in which all correct responses earn rewards.
48
49 Participants were instructed to learn which of three response buttons corresponded with each
50
51 stimulus in a set of sequentially-presented neutral images. Before each block, participants were
52
53 shown the set of images that would be used in that block. During the task, each stimulus image
54
55
56
57
58
59
60

1
2
3 was presented for up to 2 seconds, and after responding, feedback on accuracy was displayed
4
5 (“Correct” or “Try Again!”). Participants were instructed that a monetary bonus would be
6
7 calculated based on accuracy. Stimuli were blocked, and in each block the participant had to
8
9 learn stimulus-response mappings for two, three, four, or five images. Blocks consist of 11 trials
10
11 per image and there were two blocks per type (two blocks each of 22, 33, 44, and 55 trials).
12
13

14
15 **Executive Functioning Tasks.** Executive functioning was measured using a set of well-
16
17 validated tasks that have been administered in developmental and clinical samples, and used in
18
19 prior latent variable analysis (Friedman et al., 2008, 2016). The tasks featured overlapping
20
21 demands for general executive control but differed in perceptual characteristics and specific task
22
23 demands.
24
25

26
27 **Antisaccade task.** The antisaccade task is a measure of response inhibition. For each trial,
28
29 the participant viewed a fixation point presented at the center of the monitor. After a variable
30
31 amount of time (delay ranging from 1500 to 3500ms, at increments of 250ms) a visual distractor
32
33 was shown on one side of the screen for 150ms, after which a target stimulus appeared on the
34
35 opposite side of the screen (target displayed for 250, 233, or 200 ms, then masked). The visual
36
37 distractor was a small black square (1/8 inch) and the target stimulus was a number symbol
38
39 (between 1 and 9). The participant was instructed to keep their eyes on the fixation point and not
40
41 saccade to the visual distractor, instead saccading to the target stimulus in time to see it before it
42
43 was masked, and then report the numeric value out loud to the research assistant. The participant
44
45 completed 36 trials in which the distractor and stimulus were on the same side (target displayed
46
47 for 200 ms), and 12 practice trials in which the distractor and stimulus were on opposite sides
48
49 (target displayed for 250 ms). Then, the task consisted of three blocks (24 blocked trials of each
50
51
52
53
54
55
56
57
58
59
60

1
2
3 target duration in ascending difficulty) of antisaccade trials in which the distractor and stimulus
4
5 were on opposite sides.
6

7
8 ***Spatial N-Back Task.*** The spatial n-back is a test of working memory updating ability. In
9
10 each block of this task, the participant was presented with a display of twelve open squares
11
12 distributed across the screen. For each trial, one square in the display turned black (“flashed”) for
13
14 500ms, followed by a 1500ms inter-trial interval. The participant was instructed to respond by
15
16 pressing a button to indicate whether the square that flashed for that trial was the same spatial
17
18 location as the square that flashed two trials earlier. Each block consisted of 24 trials, and within
19
20 each trial there were six match trials (on which the correct answer was “yes”) and 18 reject trials
21
22 (on which the correct answer was “no”; among these reject trials, there were five lures on which
23
24 the same spatial location was flashed three trials earlier). Participants completed a practice block
25
26 of 20 trials. The task consisted of four blocks (24 trials each).
27
28
29

30
31 ***Color-Shape Task.*** The color-shape task is a test of mental set shifting. In this task, for
32
33 each trial, a shape stimulus (red or blue, circle or triangle) appeared in the center of the screen
34
35 accompanied by a cue letter (C or S) to indicate whether the participant should categorize the
36
37 stimulus based on color or shape. The cue appeared 350 ms before the target and remained on the
38
39 screen until the participant responded using one of two buttons (left for red or circle, right for
40
41 blue or triangle). The next cue appeared 350 ms after the end of the trial. The participant started
42
43 with 12 trials practicing identification of color, 12 trials practicing identification of shape, and
44
45 then 24 trials practicing a mixed sequence in which a trial of one cue type could be preceded by
46
47 the same (“repeat” trials) or the other (“switch” trials) cue type. The task consisted of two blocks
48
49 (52 trials each) of mixed trials. The first four trials of each block were discarded, yielding 48
50
51 trials (half repeat, half switch trials) for each of the two task blocks for analysis.
52
53
54
55
56
57
58
59
60

Developmental Differences

Self-report measures were administered using either Qualtrics or REDCap software. A research staff member was available for questions outside the experiment room.

Age. Participants reported age in years.

Pubertal Development Scale (PDS, (Petersen et al., 1988)). The PDS is a self-report measure of physical pubertal changes which shows good reliability and correlates well with hormone measures (Shirtcliff et al., 2009). Individual items were summed and averaged to yield a PDS score that ranged from 1 to 4.

Gender. Participants reported on sex and gender according to a two-step procedure that asks about current gender identity and sex assigned at birth. Gender was contrast-coded as cisgender male (-1), cisgender female (+1), or other, including transgender, non-binary, or gender-fluid (0).

Mood Pathology

See Supplement for exploratory analyses that considered diagnostic variables.

Mood and Anxiety Symptom Questionnaire- Anhedonia (MASQ-AD, (Watson, Clark, Weber, & Assenheimer, 1995; Watson, Clark, Weber, Assenheimer, et al., 1995)). The 8-item MASQ-AD loss of interest scale was administered to evaluate anhedonic depression in the past week. Each item was rated on a scale of 1 (not at all) to 5 (extremely). The summed score provided a measure of anhedonic symptom severity.

General Behavior Inventory- Mania/Hypomania (GBI-MH, (Depue et al., 1989)). The 10-item GBI-MH scale was administered to measure symptoms of mania or hypomania in the

past week¹. This measure has been used to assess current mania severity (Moriarity et al., 2020) in clinical and community samples (e.g., in the Adolescent Brain and Cognitive Development Study, (Barch et al., 2018)). Each item was rated on a scale of 0 (never or hardly ever) to 3 (very often or constantly). The summed score indexed severity of manic symptoms.

Analyses

Neurocognitive Functioning

The reward learning and executive functioning performance parameters selected for this study are widely used and well-validated (Friedman et al., 2016; Pizzagalli et al., 2005). These parameters were chosen based on evidence that these measures are sensitive to individual differences or development (Friedman et al., 2016; Master et al., 2020; Nussenbaum & Hartley, 2019) and to symptoms of mood pathology (Lawlor et al., 2020; Robinson & Chase, 2017; Zald & Treadway, 2017). The executive function performance parameters are the same as those used in prior latent variable research (Friedman & Miyake, 2017).

Performance Parameters. Performance parameters from each task are described below with additional details in Supplement (Tables S2-3).

Two-armed Bandit Task. Performance on the reward condition was modeled with a prediction error computational model (Daw, 2011; Frank et al., 2007; Rescorla & Wagner, 1972), programmed in R. See (Frank et al., 2007) for formulas and modeling approach. This model fits the participant's trial-by-trial sequence of responses to the pair of stimuli that can elicit reward. The expected value of selecting one of the stimuli i (here, either the stimulus with

¹ We note that due to researcher error, the instructional text preceding GBI items was inconsistent across sites in reminding the participant to consider the past-week timeframe when reporting on symptoms of mania. Sensitivity analyses were performed to ensure that this error did not engender site differences that influenced results, including testing site as a moderator of effects. No site-related differences were observed.

an 80% probability of reward, or the stimulus with a 20% probability of reward) was estimated using the following equation for each participant:

$$Q_i(t + 1) = Q_i(t) + \alpha * [r(t) - (Q_i(t))]$$

Where t is trial number, and $r(t)$ is the reward value for that trial (either 1=received reward, or 0=did not receive reward). (Note that $r(t) - Q_i(t)$ reflects the prediction error for that trial). Q is initialized at 0 at the beginning of the model. The learning rate parameter α that best fits each participant's sequence of responses over time is selected to reflect the degree to which prior reinforcement outcomes affect subsequent Q values. In addition, the probability of choosing one stimulus (choosing A instead of B) was calculated using this equation for each participant:

$$P_A(t) = \left(\frac{e^{(Q_A(t) * \beta)}}{e^{(Q_A(t) * \beta)} + e^{(Q_B(t) * \beta)}} \right)$$

Where β is the inverse temperature parameter reflecting the participant's tendency to randomly choose a stimulus (explore) or to choose the stimulus with the currently-highest Q value (exploit). The model was fit to each participant's sequence of responses, searching through the parameter space (1000 values and 200 random starts for each parameter). Model fit was estimated for each participant according to the log likelihood estimate fit of the model to each participant's choices over time:

$$\text{fit} = -1 * (\sum \log(P))$$

Guided by prior clinical and developmental research (Nussenbaum & Hartley, 2019; Zald & Treadway, 2017), we selected learning rate (α) as the primary performance parameter, and the quality of the model was assessed with the fit parameter (see below). Learning rate can be interpreted as the speed at which the highest reward association is learned for Q .

Probabilistic Reward Task. Performance measures for this task (Pizzagalli et al., 2005) were computed using Python. The primary performance parameter was discriminability, a measure of the participant's ability to accurately discriminate between the two stimuli (long and short mouths). The formula for computing discriminability is described in (Pizzagalli et al., 2005). Discriminability was calculated as:

$$DIS = \frac{1}{2} \log \left(\frac{(Rich_{correct} + 0.5) * (Lean_{correct} + 0.5)}{(Rich_{incorrect} + 0.5) * (Lean_{incorrect} + 0.5)} \right)$$

Discriminability reflects the extent to which the participant is able to accurately discriminate between stimuli overall, reaping the maximum possible rewards for both trial types. We also computed a second parameter (response bias) that reflects the extent to which the participant develops an accuracy bias to respond more accurately to "rich" compared with "lean" stimuli over time (Supplement). Of note, both of these performance parameters have been associated with anhedonic depression (Lawlor et al., 2020; Pizzagalli et al., 2005). However, discriminability was selected as the primary performance parameter for this study because of superior covariance with reward learning parameters from other tasks (Supplement).

Instrumental Learning Task. Performance on this task was operationalized as accuracy across trial blocks, calculated by estimating proportion accuracy for each block and averaging across blocks for a global measure.

Antisaccade Task. The performance measure for the antisaccade task was proportion of correct responses across the antisaccade blocks (Friedman et al., 2016).

Spatial N-Back Task. The performance measure for the spatial n-back task was proportion of correct responses across the task (Friedman et al., 2016).

Color-Shape Task. The performance measure for the color-shape task was switch cost, or the difference between average reaction time on switch trials and average reaction time on repeat

1
2
3 trials for each participant (Friedman et al., 2016). Reaction times for incorrect responses, for
4
5 trials immediately following incorrect responses, below 200ms, and that were >3.32 times the
6
7 median absolute deviation in each condition were removed before calculating condition averages
8
9
10 (Wilcox & Keselman, 2003).

11
12 **Data Quality Assurance and Checks.** Data quality was checked for each task and
13
14 measure. Participants who failed quality assurance steps were removed from analyses involving
15
16 that parameter (Table S2) (Friedman et al., 2016; Pizzagalli et al., 2005). For the probabilistic
17
18 reward task, participants were removed who failed to discriminate above-chance level between
19
20 stimulus types ($<55\%$ accuracy), who failed to achieve an overall reinforcement schedule of
21
22 approximately 3:1 (reward ratio of rich:lean), or who showed high frequency of reaction times
23
24 indicating invalid responses or outlier responses ($>20\%$ trials with $RT < 150\text{ms}$ or $RT > 2500\text{ms}$).
25
26 These performance checks are standard for this task (Pizzagalli et al. 2005). For the two-armed
27
28 bandit task, participants were removed on the basis of poor fit for the prediction error model (fit
29
30 < 2). For other tasks, participants were removed who showed chance-level accuracy: instrumental
31
32 learning task ($<32\%$ accuracy), antisaccade task ($<20\%$ accuracy), spatial n-back ($<60\%$
33
34 accuracy), and color-shape task ($<60\%$ accuracy). The distributions of all measures were
35
36 evaluated (Table S2); all performance, developmental, and mood measures showed acceptable
37
38 normality (skew < 2 , kurtosis < 6) (Hair, 2010; Kline, 2011). Outlier checks failed to reveal
39
40 measurements more than ± 3 standard deviations from mean scores.
41
42
43
44
45

46 ***Site Checks***

47
48
49 There were no significant differences between sites in experimental variables or symptom
50
51 severity, $ps > 0.13$. There were significant differences between sites in average age and pubertal
52
53 stage, $ps < 0.05$, which was expected given the younger ages eligible for recruitment at the CU
54
55
56
57
58
59
60

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

19

1
2
3 Boulder site (Table 1, Supplement). Supplementary analyses were performed to further evaluate
4
5 potential site differences, including testing invariance in the factor structures across sites and
6
7 repeating models with site as a covariate or moderator. There were no significant differences in
8
9 factor structure or experimental effects between sites. (Supplement).
10
11

Modeling

12
13
14
15 All continuous variables were standardized for analyses. Task performance parameters
16
17 that loaded negatively on a factor were reversed for SEM and Figures to enhance interpretability.
18

19 We conducted modeling in Mplus (Muthen & Muthen, 2017) using full information
20
21 maximum likelihood estimation (i.e., including individuals missing subsets of data) with robust
22
23 standard error estimates (robust maximum likelihood estimator; MLR). Neurocognitive factors
24
25 were estimated with confirmatory factor analyses across performance parameters from reward
26
27 processing tasks (reward learning performance factor), and across performance parameters from
28
29 the executive function tasks (executive functioning factor). The structural equation model (SEM)
30
31 tested *a priori* hypotheses regarding developmental patterns in neurocognitive factors and
32
33 moderation of these developmental patterns by mood symptoms. The following predicted the
34
35 reward learning performance factor: linear and quadratic effects of puberty, linear effects of
36
37 anhedonic and manic symptom severity, and interactions between symptom measures and
38
39 puberty variables (covariates were the linear and quadratic effects of age). In the same model, the
40
41 following predicted the executive functioning factor: linear and quadratic effects of age, linear
42
43 effects of anhedonic and manic symptom severity, and interactions between symptom measures
44
45 and age variables (covariates were the linear and quadratic effects of puberty). Model fit was
46
47 evaluated on the basis of confirmatory fit index (CFI) > 0.9 and root-mean-square error of
48
49 approximation (RMSEA) < 0.06, (Hair, 2010). Model comparison was performed using the chi-
50
51
52
53
54
55
56
57
58
59
60

squared difference test using the Satorra-Bentler scaled chi-square (Satorra, 2000). Associations between developmental or clinical variables and neurocognitive factors are reported as standardized estimates, with confidence intervals and (two-tailed) significance reported. To ensure the interaction effects were appropriately standardized, we standardized the components of the interactions prior to multiplying them, then used the STDY estimates from Mplus, which standardize the estimates with respect to the dependent variables. Thus, the products were not standardized and the interaction effects (e.g., of $X1*X2$) are interpretable as the change in $X1$'s standardized effect on Y with one standard deviation change in $X2$.

Results

Confirmatory Factor Analysis

Confirmatory factor analysis modeling the reward learning performance and executive functioning factors is reported in the Supplement (Figure S2). Significant indicator loadings supported the latent variable approach for evaluating individual differences in reward learning performance and executive functioning. Of note, reward learning performance and executive functioning were correlated.

Structural Equation Model

The SEM evaluated direct and moderated effects of puberty and mood symptom severity on the reward learning performance factor, and direct and moderated effects of age and mood symptom severity on the executive functioning factor (Figure 1). The model showed adequate fit, $\chi^2(72) = 89.87, p > 0.05$, RMSEA = 0.03 [90% CI 0.00 to 0.04], CFI = 0.92. The chi-square difference test showed this model was superior to the constrained model in which paths were fixed at zero, $\chi^2_{diff}(20) = 81.40, p < 0.05$.

[Figure 1 about here]

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

21

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

There was a significant quadratic effect of pubertal status on reward learning performance, (standardized) estimate = -0.11 [90% CI -0.20 to -0.02], $p=0.037$, in which participants at early or late pubertal stages exhibited relatively lower reward learning performance than participants mid-puberty. The quadratic effect of pubertal status on reward learning performance was moderated by both anhedonic symptoms, (standardized) estimate = -0.20 [90% CI -0.35 to -0.05], $p=0.027$, and manic symptoms (standardized) estimate = 0.27 [90% CI 0.13 to 0.42], $p=0.002$. Adolescents who were more severely anhedonic exhibited lower reward learning performance especially at early pubertal stages, whereas adolescents who were more severely manic exhibited higher reward learning performance at early pubertal stages that declined at mid-to-late pubertal stages.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

The linear and quadratic effects of age on executive functioning were significant, (standardized) linear estimate = 0.42 [90% CI 0.30 to 0.55], $p<0.001$, and (standardized) quadratic estimate = -0.16 [90% CI -0.25 to -0.06], $p=0.008$. Adolescents at older ages showed better executive functioning, and age-related improvements leveled off in the oldest adolescents. Linear age differences in executive functioning were moderated by manic symptom severity, (standardized) estimate = -0.16 [90% CI -0.28 to -0.05], $p=0.022$. Adolescents reporting higher mania showed worse executive functioning, especially at older ages. Anhedonia did not significantly moderate age differences in executive functioning.

45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Although we had no *a priori* hypotheses regarding gender, we repeated the above model including self-reported gender as a predictor (Supplement). Developmental and psychopathology effects were not significantly altered by the addition of gender (changes in standardized estimates <0.06 , changes in $ps<0.02$).

1
2
3 Together, results showed pubertal differences in reward learning performance that were
4 moderated by anhedonic or manic symptom severity, and age differences in executive
5 functioning that were moderated by manic symptom severity (Figure 2).
6
7
8
9

10 Discussion

11
12 The goals of this study were to replicate developmental differences in reward sensitivity
13 and executive functioning, and test whether such developmental differences followed distinctive
14 patterns in adolescents with high levels of anhedonic or manic symptoms. Results support a
15 model in which dual-systems developmental differences emerge differently for adolescents with
16 mood problems, but the nature of the developmental abnormalities depended on the nature of
17 mood symptoms. Higher levels of manic symptoms were associated with higher reward learning
18 performance at early pubertal stages but declining reward learning performance later in puberty,
19 and poorer executive functioning especially at older ages. In contrast, higher levels of anhedonic
20 symptoms were associated with lower reward learning performance at early pubertal stages.
21 These findings inform models in which abnormal developmental trajectories correspond with
22 specific dimensions of mood pathology, and motivate future longitudinal research to evaluate
23 developmental and clinical changes over time.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 [Figure 2 about here]

41
42 In this cross-sectional adolescent sample, we replicated previous research demonstrating
43 puberty-related curvilinear differences in reward sensitivity, and age-related linear (and
44 curvilinear) differences in executive functioning (Geier & Luna, 2009; Shulman et al., 2016).
45 Consistent with prior work, in this study mid-pubertal adolescents showed the highest reward
46 learning performance (ability to maximize rewards across learning tasks), but reward learning
47 performance was lower at early or late pubertal stages (Braams et al., 2015; Davidow et al.,
48
49
50
51
52
53
54
55
56
57
58
59
60

2016). In contrast, adolescents at older ages showed better executive functioning (Steinberg et al., 2008). In the present study, reward learning performance and executive functioning were evaluated using a robust latent variable approach. Performance indexed by a single task parameter derives from a number of sources (Collins & Frank, 2012) and may reflect a mixture of neurocognitive abilities, contributing to mixed findings in developmental or clinical samples (Snyder et al., 2015). Latent variable approaches take advantage of shared processes across a set of conceptually-related tasks to derive a more reliable estimate of a common dimension (Friedman et al., 2016; Harden et al., 2018), and the present study leveraged this approach for replication.

The main contribution of this study is to show that developmental differences in reward sensitivity and executive functioning are altered in adolescents with currently elevated mood symptoms. These findings have implications for conceptual models of mood pathology and clinical application. First, results highlight the complexity of neurocognitive abnormalities as they manifest over development. Here, reward learning performance was elevated in more highly manic, early-puberty adolescents, but decreased over pubertal development as a function of symptom severity. This may reflect a developmental shift from reward hypersensitivity to hyposensitivity in this particular domain of reward processing, or changes in learning and performance ability that cause mania-related reward abnormalities to manifest differently over development. To explore these interpretations, it will be valuable to test how reward response changes over puberty or the course of illness. Second, these results point to developmental periods within adolescence when neurocognitive functioning may be altered in mood pathology, and therefore periods in which it is especially valuable to identify neurocognitive abnormalities that may inform risk prediction and intervention (Kujawa et al., 2020). Third, these results point

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

24

1
2
3 to distinctive patterns of neurocognitive abnormality that may inform differential diagnosis in
4
5 adolescence. Because early-pubertal reward hypersensitivity was related to manic symptom
6
7 severity over and above elevated anhedonia (and even in the absence of bipolar diagnoses),
8
9 reward processing abnormalities may be risk markers for bipolar illness (Alloy & Nusslock,
10
11 2019; Nusslock & Alloy, 2017). In light of the serious challenges distinguishing between
12
13 unipolar and bipolar disorders in adolescence (Birmaher & Axelson, 2006), differential
14
15 neurocognitive markers of illness are a needed advancement for clinical treatment.
16
17
18

19
20 These findings point to several important next steps in this research. One direction for
21
22 this work will be to evaluate neurobiological and physiological mechanisms of mood-related
23
24 neurocognitive anomalies (Galvan, 2010). It may be that among adolescents with mood
25
26 pathology, abnormal reward learning performance in early puberty is driven by emerging
27
28 abnormalities in pubertal hormone functioning: for example, elevated testosterone and lower
29
30 estradiol in adolescents are associated with higher levels of reward sensitivity (Harden et al.,
31
32 2018) and maturation of dopaminergic projections to medial prefrontal systems (Delevich et al.,
33
34 2021). Individual differences in sex hormones have been linked with mania and depression,
35
36 although effects are mixed across studies and commonly examined in adults (Johnson et al.,
37
38 2013; Lombardo et al., 2021). Research that evaluates how changes in pubertal hormones track
39
40 with striatal development and emerging mood symptoms may address these ideas. In turn, given
41
42 evidence that adolescent improvements in executive functioning are related to maturation of
43
44 prefrontal systems and synchrony among frontoparietal brain networks (Luna et al., 2015),
45
46 research that explores differences in structural and functional connectivity of prefrontal regions
47
48 may clarify and extend these results. For example, in the present study (and contrary to our
49
50 hypotheses) anhedonia did not significantly moderate developmental patterns of executive
51
52
53
54
55
56
57
58
59
60

1
2
3 functioning; neuroimaging may identify subtle differences in prefrontal development that do not
4
5 emerge behaviorally.
6

7
8 A second important direction for this work is to extend to longitudinal research designs.
9
10 The key limitation of the present study is that, in this cross-sectional design, it was only possible
11
12 to test for *differences* related to developmental variables, but not to test developmental *processes*
13
14 that unfold over time. Longitudinal evaluation of neurocognitive functioning is needed to
15
16 characterize the temporal associations between, e.g., pubertal development and reward
17
18 processing (Van Leijenhorst et al., 2010). Relatedly, longitudinal assessment would clarify the
19
20 extent to which neurocognitive developmental trajectories correspond with onset of or
21
22 fluctuations in mood pathology within a person. Cross-sectional mood symptoms reflect both
23
24 state-dependent information about a given episode and trait-like information about overall illness
25
26 severity (Sarapas et al., 2012). Only repeated measurement of mood over time can disentangle
27
28 state- versus trait-like features, and address key questions about the timing of neurocognitive
29
30 anomalies as related to the course of mood pathology. Finally, longitudinal assessment is a
31
32 superior approach for disentangling neurocognitive changes associated with age versus puberty,
33
34 as these variables tend to be correlated in cross-sectional research (Braams et al., 2015). In the
35
36 present study, age and pubertal stage were correlated ($r=0.52$), but neurocognitive differences
37
38 related to age versus puberty were significant when controlling for the other developmental
39
40 variable. However, longitudinal studies can provide nuanced investigation e.g., of puberty-
41
42 related changes in neurocognition that occur within age groups.
43
44
45
46
47
48

49 Several other limitations and future directions are noted. First, we chose to focus on
50
51 common dimensions of reward learning performance and executive functioning, but these are
52
53 non-unitary constructs that can also be decomposed into subdimensions (e.g., within reward
54
55
56
57
58
59
60

1
2
3 learning performance, exploration). Second, there are other domains of reward processing and
4 executive function that are not captured in the present study (e.g., reward liking, decision
5 making, selective attention). Future research that expands to capture additional domains of
6 reward and higher-order cognition, or that targets subdimensions of these neurocognitive
7 domains, may provide complementary insight. Third, we note that although reward learning
8 performance and executive functioning had different associations with developmental and
9 symptom variables, these dimensions were highly correlated at the factor and indicator levels
10 (Table S3). These patterns indicate that latent variables may have captured both independent
11 processes unique to reward learning performance versus executive functioning, and shared
12 processes that influence behavior across neurocognitive domains (e.g., motivation).

13
14 Confirmatory factor analyses should be replicated. Fourth, this study relied on self-report
15 measurement of puberty, and did not include pre-pubertal participants or the transition to
16 puberty. Future research in younger samples and that captures other (e.g., hormone) measures of
17 puberty is needed. Together, next steps for this work include replication and extension of the
18 present findings in large-scale and well-powered longitudinal developmental samples.

19
20 In conclusion, this study used a latent-variable approach to evaluate age- or puberty-
21 related differences in neurocognitive functioning in adolescence, and showed that youths with
22 elevated symptoms of mania or anhedonia exhibited altered developmental patterns. Our findings
23 replicate and extend research on reward and executive functioning in adolescent mood disorders,
24 and suggest that mood-related abnormalities in neurocognition may manifest differently over
25 development. More broadly, this work lies at the intersection of clinical, developmental and
26 neurocognitive science, and highlights the value of interdisciplinary models to inform efforts to
27 understand — and intervene in — adolescent mood disorders. Future longitudinal investigation

1
2
3 will take the important next step of evaluating trajectories of neurocognitive development in
4 adolescents with mood pathology.
5
6

7 **Author Contributions**

8
9
10 R.H.K. developed the study concept and design. A.D.M., C.N., E.C.P., J.J., C.H., B.R., and
11 C.F.S performed data collection and clinical evaluation, under R.H.K.'s supervision. C.D.S. and
12 D.M. provided consultation on recruitment and clinical evaluation. R.H.K. performed analyses
13 with consultation from N.P.F. The manuscript was drafted by R.H.K., C.N., and J.J., and all
14 other co-authors provided critical revisions. All authors approved the final version of the
15 manuscript before submission.
16
17
18
19
20
21
22
23
24
25

26 **Declaration of Conflicting Interests**

27
28 The authors declared no conflicts of interest with respect to the authorship or the publication of
29 this article.
30
31
32
33
34

35 **Acknowledgements**

36
37 With sincere thanks to Daniel Dillon for sharing a Python-based analytic pipeline for scoring the
38 probabilistic reward task. Research supported by NARSAD Young Investigator Award (24879)
39 and NIMH grant MH117131 (R.H.K.).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Alloy, L. B., & Nusslock, R. (2019). Future Directions for Understanding Adolescent Bipolar Spectrum Disorders: A Reward Hypersensitivity Perspective. *Journal of Clinical Child & Adolescent Psychology*, 48(4), 669–683. <https://doi.org/10.1080/15374416.2019.1567347>
- Alloy, L. B., Olino, T., Freed, R. D., & Nusslock, R. (2016). Role of Reward Sensitivity and Processing in Major Depressive and Bipolar Spectrum Disorders. *Behavior Therapy*, 47(5), 600–621. <https://doi.org/10.1016/j.beth.2016.02.014>
- Andre Der-Avakian, Samuel A. Barnes, Athina Markou, & Diego A. Pizzagalli. (2016). Translational Assessment of Reward and Motivational Deficits in Psychiatric Disorders. In *Translational Neuropsychopharmacology* (Vol. 28). Springer International Publishing.
- Ayuso-Mateos, J. L., Nuevo, R., Verdes, E., Naidoo, N., & Chatterji, S. (2010). From depressive symptoms to depressive disorders: The relevance of thresholds. *British Journal of Psychiatry*, 196(5), 365–371. <https://doi.org/10.1192/bjp.bp.109.071191>
- Barch, D. M., Albaugh, M. D., Avenevoli, S., Chang, L., Clark, D. B., Glantz, M. D., Hudziak, J. J., Jernigan, T. L., Tapert, S. F., Yurgelun-Todd, D., Alia-Klein, N., Potter, A. S., Paulus, M. P., Prouty, D., Zucker, R. A., & Sher, K. J. (2018). Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Developmental Cognitive Neuroscience*, 32, 55–66. <https://doi.org/10.1016/j.dcn.2017.10.010>
- Berridge, K. C. (2018). Evolving Concepts of Emotion and Motivation. *Frontiers in Psychology*, 9, 1647. <https://doi.org/10.3389/fpsyg.2018.01647>
- Bertha, E. A., & Balazs, J. (2013). Subthreshold depression in adolescence: A systematic review. *European Child & Adolescent Psychiatry*, 22(10), 589–603. <https://doi.org/10.1007/s00787-013-0411-0>
- Best, J. R., & Miller, P. H. (2010). A Developmental Perspective on Executive Function. *Child Development*, 81(6), 1641–1660. <https://doi.org/10.1111/j.1467-8624.2010.01499.x>

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

29

- 1
2
3 Birmaher, B., & Axelson, D. (2006). Course and outcome of bipolar spectrum disorder in children and
4 adolescents: A review of the existing literature. *Development and Psychopathology*, *18*(04).
5
6 <https://doi.org/10.1017/S0954579406060500>
7
8
9 Biuckians, A., Miklowitz, D. J., & Kim, E. Y. (2007). Behavioral activation, inhibition and mood
10 symptoms in early-onset bipolar disorder. *Journal of Affective Disorders*, *97*(1–3), 71–76.
11
12 <https://doi.org/10.1016/j.jad.2006.07.005>
13
14
15 Blakemore, S.-J., Burnett, S., & Dahl, R. E. (2010). The role of puberty in the developing adolescent
16 brain. *Human Brain Mapping*, *31*(6), 926–933. <https://doi.org/10.1002/hbm.21052>
17
18
19 Braams, B. R., van Duijvenvoorde, A. C. K., Peper, J. S., & Crone, E. A. (2015). Longitudinal Changes in
20 Adolescent Risk-Taking: A Comprehensive Study of Neural Responses to Rewards, Pubertal
21 Development, and Risk-Taking Behavior. *Journal of Neuroscience*, *35*(18), 7226–7238.
22
23 <https://doi.org/10.1523/JNEUROSCI.4764-14.2015>
24
25
26 Casey, B. J., Jones, R. M., & Hare, T. A. (2008). The Adolescent Brain. *Annals of the New York Academy*
27 *of Sciences*, *1124*, 111–126. <https://doi.org/10.1196/annals.1440.010>
28
29
30 Collins, A. G. E., & Frank, M. J. (2012). How much of reinforcement learning is working memory, not
31 reinforcement learning? A behavioral, computational, and neurogenetic analysis: Working
32 memory in reinforcement learning. *European Journal of Neuroscience*, *35*(7), 1024–1035.
33
34 <https://doi.org/10.1111/j.1460-9568.2011.07980.x>
35
36
37 Copeland, W. E., Wolke, D., Shanahan, L., & Costello, E. J. (2015). Adult Functional Outcomes of
38 Common Childhood Psychiatric Problems: A Prospective, Longitudinal Study. *JAMA Psychiatry*,
39 *72*(9), 892. <https://doi.org/10.1001/jamapsychiatry.2015.0730>
40
41
42 Cuthbert, B., & Insel, T. (2010). The Data of Diagnosis: New Approaches to Psychiatric Classification.
43 *Psychiatry*, *5*.
44
45
46 Davidow, J. Y., Foerde, K., Galván, A., & Shohamy, D. (2016). An Upside to Reward Sensitivity: The
47 Hippocampus Supports Enhanced Reinforcement Learning in Adolescence. *Neuron*, *92*(1), 93–
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Daw, N. (2011). Trial-by-trial data analysis using computational modelx. In *Decision Making, Affect, and*
4
5 *Learning: Attention and Performance XXIII* (pp. 3–37). Oxford University Press.
6
7 Delevich, K., Klinger, M., Okada, N. J., & Wilbrecht, L. (2021). Coming of age in the frontal cortex: The
8
9 role of puberty in cortical maturation. *Seminars in Cell & Developmental Biology, 118*, 64–72.
10
11 <https://doi.org/10.1016/j.semcdb.2021.04.021>
12
13 Depue, R. A., Krauss, S., Spont, M. R., & Arbisi, P. (1989). General Behavior Inventory Identification
14
15 of Unipolar and Bipolar Affective Conditions in a Nonclinical University Population. *Journal of*
16
17 *Abnormal Psychology, 98*(2), 117–126.
18
19 Dixon, T., Kravariti, E., Frith, C., Murray, R. M., & McGUIRE, P. K. (2004). Effect of symptoms on
20
21 executive function in bipolar illness. *Psychological Medicine, 34*(5), 811–821.
22
23 <https://doi.org/10.1017/S0033291703001570>
24
25 Forbes, E. E., & Dahl, R. E. (2010). Pubertal development and behavior: Hormonal activation of social
26
27 and motivational tendencies. *Brain and Cognition, 72*(1), 66–72.
28
29 <https://doi.org/10.1016/j.bandc.2009.10.007>
30
31 Forbes, E. E., & Dahl, R. E. (2012). Research Review: Altered reward function in adolescent depression:
32
33 what, when and how?: Reward function and adolescent depression. *Journal of Child Psychology*
34
35 *and Psychiatry, 53*(1), 3–15. <https://doi.org/10.1111/j.1469-7610.2011.02477.x>
36
37
38 Forbes, E. E., Eckstrand, K. L., Rofey, D. L., & Silk, J. S. (2021). A Social Affective Neuroscience
39
40 Model of Risk and Resilience in Adolescent Depression: Preliminary Evidence and Application
41
42 to Sexual and Gender Minority Adolescents. *Biological Psychiatry: Cognitive Neuroscience and*
43
44 *Neuroimaging, 6*(2), 188–199. <https://doi.org/10.1016/j.bpsc.2020.07.020>
45
46
47 Frank, M. J., Moustafa, A. A., Haughey, H. M., Curran, T., & Hutchison, K. E. (2007). Genetic triple
48
49 dissociation reveals multiple roles for dopamine in reinforcement learning. *Proceedings of the*
50
51 *National Academy of Sciences, 104*(41), 16311–16316. <https://doi.org/10.1073/pnas.0706111104>
52
53
54
55
56
57
58
59
60

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

31

1
2
3 Friedman, N., & Miyake, A. (2017). Unity and diversity of executive functions: Individual differences as
4 a window on cognitive structure. *Cortex*, *86*, 186–204.

5
6
7 <https://doi.org/10.1016/j.cortex.2016.04.023>

8
9 Friedman, N. P., du Pont, A., Corley, R. P., & Hewitt, J. K. (2018). Longitudinal Relations Between
10 Depressive Symptoms and Executive Functions From Adolescence to Early Adulthood: A Twin
11 Study. *Clinical Psychological Science*, *6*(4), 543–560.

12
13
14
15
16 <https://doi.org/10.1177/2167702618766360>

17
18 Friedman, N. P., Miyake, A., Altamirano, L. J., Corley, R. P., Young, S. E., Rhea, S. A., & Hewitt, J. K.
19 (2016). Stability and change in executive function abilities from late adolescence to early
20 adulthood: A longitudinal twin study. *Developmental Psychology*, *52*(2), 326–340.

21
22
23
24 <https://doi.org/10.1037/dev0000075>

25
26 Friedman, N. P., Miyake, A., Young, S. E., DeFries, J. C., Corley, R. P., & Hewitt, J. K. (2008).

27 Individual differences in executive functions are almost entirely genetic in origin. *Journal of*
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Experimental Psychology-General, *137*(2), 201–225. [https://doi.org/10.1037/0096-](https://doi.org/10.1037/0096-3445.137.2.201)

3445.137.2.201

35 Gaffrey, M. S., Luby, J. L., & Barch, D. M. (2013). Towards the study of functional brain development in
36 depression: An Interactive Specialization approach. *Neurobiology of Disease*, *52*, 38–48.

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
<https://doi.org/10.1016/j.nbd.2012.06.012>

Galvan. (2010). Adolescent development of the reward system. *Frontiers in Human Neuroscience*.

<https://doi.org/10.3389/neuro.09.006.2010>

Geier, C., & Luna, B. (2009). The maturation of incentive processing and cognitive control.

Pharmacology, Biochemistry, and Behavior, *93*(3), 212–221.

<https://doi.org/10.1016/j.pbb.2009.01.021>

Hair, J. F. (Ed.). (2010). *Multivariate data analysis* (7th ed). Prentice Hall.

Harden, K. P., Mann, F. D., Grotzinger, A. D., Patterson, M. W., Steinberg, L., Tackett, J. L., & Tucker-

Drob, E. M. (2018). Developmental differences in reward sensitivity and sensation seeking in

- adolescence: Testing sex-specific associations with gonadal hormones and pubertal development. *Journal of Personality and Social Psychology*, *115*(1), 161–178.
<https://doi.org/10.1037/pspp0000172>
- Horn, K., Roessner, V., & Holtmann, M. (2011). Neurocognitive performance in children and adolescents with bipolar disorder: A review. *European Child & Adolescent Psychiatry*, *20*(9), 433–450.
<https://doi.org/10.1007/s00787-011-0209-x>
- Huys, Q. J., Pizzagalli, D. A., Bogdan, R., & Dayan, P. (2013). Mapping anhedonia onto reinforcement learning: A behavioural meta-analysis. *Biology of Mood & Anxiety Disorders*, *3*(1), 12.
<https://doi.org/10.1186/2045-5380-3-12>
- Johnson, J. M., Nachtigall, L. B., & Stern, T. A. (2013). The Effect of Testosterone Levels on Mood in Men: A Review. *Psychosomatics*, *54*(6), 509–514. <https://doi.org/10.1016/j.psych.2013.06.018>
- Joseph, M. F., Frazier, T. W., Youngstrom, E. A., & Soares, J. C. (2008). A Quantitative and Qualitative Review of Neurocognitive Performance in Pediatric Bipolar Disorder. *Journal of Child and Adolescent Psychopharmacology*, *18*(6), 595–605. <https://doi.org/10.1089/cap.2008.064>
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*(6), 593.
<https://doi.org/10.1001/archpsyc.62.6.593>
- Kline, R. B. (2011). *Principles and practice of structural equation modeling* (3rd ed). Guilford Press.
- Kujawa, A., Klein, D. N., Pegg, S., & Weinberg, A. (2020). Developmental trajectories to reduced activation of positive valence systems: A review of biological and environmental contributions. *Developmental Cognitive Neuroscience*, *43*, 100791. <https://doi.org/10.1016/j.dcn.2020.100791>
- Lawlor, V. M., Webb, C. A., Wiecki, T. V., Frank, M. J., Trivedi, M., Pizzagalli, D. A., & Dillon, D. G. (2020). Dissecting the impact of depression on decision-making. *Psychological Medicine*, *50*(10), 1613–1622. <https://doi.org/10.1017/S0033291719001570>

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

33

- 1
2
3 Lee, R. S. C., Hermens, D. F., Scott, J., Redoblado-Hodge, M. A., Naismith, S. L., Lagopoulos, J.,
4
5 Griffiths, K. R., Porter, M. A., & Hickie, I. B. (2014). A meta-analysis of neuropsychological
6
7 functioning in first-episode bipolar disorders. *Journal of Psychiatric Research*, *57*, 1–11.
8
9 <https://doi.org/10.1016/j.jpsychires.2014.06.019>
10
11 Lima, I. M. M., Peckham, A. D., & Johnson, S. L. (2018). Cognitive deficits in bipolar disorders:
12
13 Implications for emotion. *Clinical Psychology Review*, *59*, 126–136.
14
15 <https://doi.org/10.1016/j.cpr.2017.11.006>
16
17 Lombardo, G., Mondelli, V., Dazzan, P., & Pariante, C. M. (2021). Sex hormones and immune system: A
18
19 possible interplay in affective disorders? A systematic review. *Journal of Affective Disorders*,
20
21 *290*, 1–14. <https://doi.org/10.1016/j.jad.2021.04.035>
22
23 Luna, B., Marek, S., Larsen, B., Tervo-Clemmens, B., & Chahal, R. (2015). An Integrative Model of the
24
25 Maturation of Cognitive Control. *Annual Review of Neuroscience*, *38*(1), 151–170.
26
27 <https://doi.org/10.1146/annurev-neuro-071714-034054>
28
29 Luna, B., & Wright, C. (2016). Adolescent brain development: Implications for the juvenile criminal
30
31 justice system. In *APA handbook of psychology and juvenile justice* (pp. 91–116). American
32
33 Psychological Association. <https://doi.org/10.1037/14643-005>
34
35 Maresh, E. L., Stim, J. J., Van Voorhis, A. C., Kang, S. S., Luciana, M., Sponheim, S. R., & Urošević, S.
36
37 (2019). Neurophysiological correlates of cognitive control and approach motivation abnormalities
38
39 in adolescent bipolar disorders. *Cognitive, Affective, & Behavioral Neuroscience*, *19*(3), 677–691.
40
41 <https://doi.org/10.3758/s13415-019-00719-x>
42
43
44 Master, S. L., Eckstein, M. K., Gotlieb, N., Dahl, R., Wilbrecht, L., & Collins, A. G. E. (2020).
45
46 Distangling the systems contributing to changes in learning during adolescence.
47
48 *Developmental Cognitive Neuroscience*, *41*, 100732. <https://doi.org/10.1016/j.dcn.2019.100732>
49
50 McCarthy, D., & Davison, M. (1979). Signal probability, reinforcement, and signal detection. *Journal of*
51
52 *the Experimental Analysis of Behavior*, *32*, 373–386.
53
54
55
56
57
58
59
60

1
2
3 McIntyre, R. S., Soczynska, J. K., Cha, D. S., Woldeyohannes, H. O., Dale, R. S., Alsuwaidan, M. T.,

4
5 Gallaughier, L. A., Mansur, R. B., Muzina, D. J., Carvalho, A., & Kennedy, S. H. (2015). The
6 prevalence and illness characteristics of DSM-5-defined “mixed feature specifier” in adults with
7 major depressive disorder and bipolar disorder: Results from the International Mood Disorders
8 Collaborative Project. *Journal of Affective Disorders*, *172*, 259–264.

9
10
11
12 <https://doi.org/10.1016/j.jad.2014.09.026>

13
14
15
16 Mendle, J. (2014). Why Puberty Matters for Psychopathology. *Child Development Perspectives*, *8*(4),
17 218–222. <https://doi.org/10.1111/cdep.12092>

18
19
20 Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The
21 unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks:
22 A latent variable analysis. *Cognitive Psychology*, *41*(1), 49–100.

23
24
25 <https://doi.org/10.1006/cogp.1999.0734>

26
27
28 Moriarity, D. P., Ng, T., Titone, M. K., Chat, I. K.-Y., Nusslock, R., Miller, G. E., & Alloy, L. B. (2020).
29 Reward Responsiveness and Ruminative Styles Interact to Predict Inflammation and Mood
30 Symptomatology. *Behavior Therapy*, *51*(5), 829–842. <https://doi.org/10.1016/j.beth.2019.11.007>

31
32
33 Muthen, L. K., & Muthen, B. O. (2017). *Mplus user's guide* (8th ed.). Muthen & Muthen.

34
35
36
37 Nielson, D. M., Keren, H., O'Callaghan, G., Jackson, S. M., Douka, I., Zheng, C. Y., Vidal-Ribas, P.,

38
39 Pornpattananangkul, N., Camp, C. C., Gorham, L. S., Wei, C., Kirwan, S., & Stringaris, A.

40
41 (2020). Great Expectations: A Critical Review of and Recommendations for the study of Reward
42 Processing as a Cause and Predictor of Depression. *Biological Psychiatry*, S0006322320317005.

43
44
45 <https://doi.org/10.1016/j.biopsych.2020.06.012>

46
47
48 Nussenbaum, K., & Hartley, C. A. (2019). Reinforcement learning across development: What insights can
49 we draw from a decade of research? *Developmental Cognitive Neuroscience*, *40*, 100733.

50
51
52 <https://doi.org/10.1016/j.dcn.2019.100733>

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

35

- 1
2
3 Nusslock, R., & Alloy, L. B. (2017). Reward processing and mood-related symptoms: An RDoC and
4 translational neuroscience perspective. *Journal of Affective Disorders, 216*, 3–16.
5
6 <https://doi.org/10.1016/j.jad.2017.02.001>
7
8
9 Nusslock, R., Almeida, J. R., Forbes, E. E., Versace, A., Frank, E., LaBarbara, E. J., Klein, C. R., &
10
11 Phillips, M. L. (2012). Waiting to win: Elevated striatal and orbitofrontal cortical activity during
12 reward anticipation in euthymic bipolar disorder adults. *Bipolar Disorders, 14*(3).
13
14 <https://doi.org/10.1111/j.1399-5618.2012.01012.x>
15
16
17 Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during
18 adolescence? *Nature Reviews Neuroscience, 9*(12), 947–957. <https://doi.org/10.1038/nrn2513>
19
20
21 Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of pubertal status:
22 Reliability, validity, and initial norms. *Journal of Youth and Adolescence, 17*(2), 117–133.
23
24 <https://doi.org/10.1007/BF01537962>
25
26
27 Phillips, M. L., & Swartz, H. A. (2014). A Critical Appraisal of Neuroimaging Studies of Bipolar
28 Disorder: Toward a New Conceptualization of Underlying Neural Circuitry and a Road Map for
29 Future Research. *American Journal of Psychiatry, 171*(8), 829–843.
30
31 <https://doi.org/10.1176/appi.ajp.2014.13081008>
32
33
34 Pizzagalli, D. A. (2014). Depression, Stress, and Anhedonia: Toward a Synthesis and Integrated Model.
35 *Annual Review of Clinical Psychology, Vol 10, 10*, 393–423. <https://doi.org/10.1146/annurev-clinpsy-050212-185606>
36
37
38
39 Pizzagalli, D. A., Goetz, E., Ostacher, M., Iosifescu, D. V., & Perlis, R. H. (2008). Euthymic Patients
40 with Bipolar Disorder Show Decreased Reward Learning in a Probabilistic Reward Task.
41 *Biological Psychiatry, 64*(2), 162–168. <https://doi.org/10.1016/j.biopsych.2007.12.001>
42
43
44 Pizzagalli, D. A., Jahn, A. L., & O’Shea, J. P. (2005). Toward an objective characterization of an
45 anhedonic phenotype: A signal detection approach. *Biological Psychiatry, 57*(4), 319–327.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Reyes, A. N., Cardoso, T. A., Jansen, K., Mondin, T. C., Souza, L. D. M., Magalhães, P. V. S.,
4
5 Kapczynski, F., & Silva, R. A. (2017). Functional impairment and cognitive performance in mood
6
7 disorders: A community sample of young adults. *Psychiatry Research*, *251*, 85–89.
8
9 <https://doi.org/10.1016/j.psychres.2017.01.069>
- 11 Robert Rescorla & Allan Wagner. (1972). A theory of Pavlovian conditioning: Variations in the
12
13 effectiveness of reinforcement and nonreinforcement. In *Classical conditioning: Current*
14
15 *research and theory* (pp. 64–99). Appleton-Century-Crofts.
- 17 Robinson, O., & Chase, H. (2017). Learning and Choice in Mood Disorders: Searching for the
18
19 Computational Parameters of Anhedonia. *Computational Psychiatry*, *1*, 208–233.
20
21 https://doi.org/10.1162/CPSY_a_00009
- 23 Rock, P. L., Roiser, J. P., Riedel, W. J., & Blackwell, A. D. (2014). Cognitive impairment in depression:
24
25 A systematic review and meta-analysis. *Psychological Medicine*, *44*(10), 2029–2040.
26
27 <https://doi.org/10.1017/S0033291713002535>
- 29 Sarapas, C., Shankman, S. A., Harrow, M., & Goldberg, J. F. (2012). Parsing trait and state effects of
30
31 depression severity on neurocognition: Evidence from a 26-year longitudinal study. *Journal of*
32
33 *Abnormal Psychology*, *121*(4), 830–837. <https://doi.org/10.1037/a0028141>
- 35 Satorra, A. (2000). Scaled and adjusted restricted tests in multi-sample analysis of moment structures. In
36
37 *Innovations in multivariate statistical analysis. A Festschrift for Heinz Neudecker* (pp. 223–247).
38
39 Kluwer Academic Publishers.
- 41 Scult, M. A., Paulli, A. R., Mazure, E. S., Moffitt, T. E., Hariri, A. R., & Strauman, T. J. (2017). The
42
43 association between cognitive function and subsequent depression: A systematic review and
44
45 meta-analysis. *Psychological Medicine*, *47*(1), 1–17.
46
47 <https://doi.org/10.1017/S0033291716002075>
- 49 Shirliff, E. A., Dahl, R. E., & Pollak, S. D. (2009). Pubertal Development: Correspondence Between
50
51 Hormonal and Physical Development: Hormonal Correlates of Pubertal Stage. *Child*
52
53 *Development*, *80*(2), 327–337. <https://doi.org/10.1111/j.1467-8624.2009.01263.x>

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

37

- 1
2
3 Shulman, E. P., Smith, A. R., Silva, K., Icenogle, G., Duell, N., Chein, J., & Steinberg, L. (2016). The
4
5 dual systems model: Review, reappraisal, and reaffirmation. *Developmental Cognitive*
6
7 *Neuroscience*, *17*, 103–117. <https://doi.org/10.1016/j.dcn.2015.12.010>
8
9
10 Snyder, H. R. (2013). Major Depressive Disorder is associated with broad impairments on
11
12 neuropsychological measures of executive function: A meta-analysis and review. *Psychological*
13
14 *Bulletin*, *139*(1), 81–132. <https://doi.org/10.1037/a0028727>
15
16 Snyder, H. R., Miyake, A., & Hankin, B. L. (2015). Advancing understanding of executive function
17
18 impairments and psychopathology: Bridging the gap between clinical and cognitive approaches.
19
20 *Frontiers in Psychology*, *6*, 24. <https://doi.org/10.3389/fpsyg.2015.00728>
21
22 Steinberg, L., Albert, D., Cauffman, E., Banich, M., Graham, S., & Woolard, J. (2008). Age differences in
23
24 sensation seeking and impulsivity as indexed by behavior and self-report: Evidence for a dual
25
26 systems model. *Developmental Psychology*, *44*(6), 1764–1778. <https://doi.org/10.1037/a0012955>
27
28 Urošević, S., Luciana, M., Jensen, J. B., Youngstrom, E. A., & Thomas, K. M. (2016). Age associations
29
30 with neural processing of reward anticipation in adolescents with bipolar disorders. *NeuroImage:*
31
32 *Clinical*, *11*, 476–485. <https://doi.org/10.1016/j.nicl.2016.03.013>
33
34 Urošević, S., Youngstrom, E. A., Collins, P., Jensen, J. B., & Luciana, M. (2016). Associations of age
35
36 with reward delay discounting and response inhibition in adolescents with bipolar disorders.
37
38 *Journal of Affective Disorders*, *190*, 649–656. <https://doi.org/10.1016/j.jad.2015.11.005>
39
40
41 Van Leijenhorst, L., Zanolie, K., Van Meel, C. S., Westenberg, P. M., Rombouts, S. A. R. B., & Crone,
42
43 E. A. (2010). What Motivates the Adolescent? Brain Regions Mediating Reward Sensitivity
44
45 across Adolescence. *Cerebral Cortex*, *20*(1), 61–69. <https://doi.org/10.1093/cercor/bhp078>
46
47
48 Van Rheeën, T. E., Lewandowski, K. E., Bauer, I. E., Kapczynski, F., Miskowiak, K., Burdick, K. E., &
49
50 Balanzá-Martínez, V. (2020). Current understandings of the trajectory and emerging correlates of
51
52 cognitive impairment in bipolar disorder: An overview of evidence. *Bipolar Disorders*, *22*(1),
53
54 13–27. <https://doi.org/10.1111/bdi.12821>
55
56
57
58
59

- 1
2
3 Van Rheenen, T. E., Miskowiak, K., & Burdick, K. E. (2021). Recognising the relevance of cognitive
4
5 dysfunction in the clinical management of bipolar disorder. *Bipolar Disorders*, bdi.13063.
6
7 <https://doi.org/10.1111/bdi.13063>
8
9
10 Vrieze, E., Pizzagalli, D. A., Demyttenaere, K., Hompes, T., Sienaert, P., de Boer, P., Schmidt, M., &
11
12 Claes, S. (2013). Reduced Reward Learning Predicts Outcome in Major Depressive Disorder.
13
14 *Biological Psychiatry*, 73(7), 639–645. <https://doi.org/10.1016/j.biopsych.2012.10.014>
15
16 Wagner, S., Müller, C., Helmreich, I., Huss, M., & Tadić, A. (2015). A meta-analysis of cognitive
17
18 functions in children and adolescents with major depressive disorder. *European Child &*
19
20 *Adolescent Psychiatry*, 24(1), 5–19. <https://doi.org/10.1007/s00787-014-0559-2>
21
22 Watson, D., Clark, L. A., Weber, K., & Assenheimer, J. S. (1995). Testing a tripartite model II: Exploring
23
24 the symptom structure of anxiety and depression in student, adult, and patient samples. *Journal of*
25
26 *Abnormal Psychology*, 104(1), 15–25. <https://doi.org/10.1037/0021-843x.104.1.15>
27
28
29 Watson, D., Clark, L. A., Weber, K., Assenheimer, J. S., Strauss, M. E., & McCormick, R. A. (1995).
30
31 Testing a tripartite model I: Evaluating the convergent and discriminant validity of anxiety and
32
33 depression symptom scales. *Journal of Abnormal Psychology*, 104(1), 3–14.
34
35 <https://doi.org/10.1037/0021-843x.104.1.3>
36
37 Yazici, E., Bursalioglu, F. S., Aydin, N., & Yazici, A. B. (2013). Menarche, puberty and psychiatric
38
39 disorders. *Gynecological Endocrinology*, 29(12), 1055–1058.
40
41 <https://doi.org/10.3109/09513590.2013.829447>
42
43 Zald, D. H., & Treadway, M. T. (2017). Reward Processing, Neuroeconomics, and Psychopathology.
44
45 *Annual Review of Clinical Psychology*, 13(1), 471–495. [https://doi.org/10.1146/annurev-clinpsy-](https://doi.org/10.1146/annurev-clinpsy-032816-044957)
46
47 [032816-044957](https://doi.org/10.1146/annurev-clinpsy-032816-044957)
48
49
50
51
52
53
54
55
56
57
58
59
60

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

39

Table 1. Demographics

	CU Boulder	UCLA	All
	<i>n</i> = 264	<i>n</i> = 155	<i>n</i> = 419
	M (SD)	M (SD)	M (SD)
Age	17.37 (2.23)	20.34 (2.21)	18.46 (2.65)
PDS*	3.56 (0.49)	3.82 (0.24)	3.66 (0.44)
GBI-MH	4.00 (4.22)	4.48 (5.90)	4.16 (4.85)
MASQ-AD	16.60 (6.65)	18.99 (8.63)	17.49 (7.53)
	%	%	%
Gender			
Cisgender Female	54.75%	69.03%	60.03%
Cisgender Male	38.78%	29.03%	35.17%
Other	6.46%	1.94%	4.79%
Ethnicity			
Hispanic or Latinx	11.85%	19.75%	14.77%
Non-Hispanic and Non-Latinx	84.44%	75.93%	81.29%
Other	3.70%	4.32%	3.93%
Race			
Asian	4.43%	19.14%	9.87%
Biracial or More than One Race	9.96%	14.81%	11.75%
Black or African American	1.84%	3.10%	2.31%
Native Hawaiian	0.00%	0.00%	0.00%
Native American	0.00%	0.00%	0.00%
White	80.07%	49.38%	68.72%
Other	3.69%	13.58%	7.35%
Parent Education (Highest Completed)			
8 th Grade or Less	0.00%	6.17%	2.28%
Partial High School	2.95%	3.09%	3.00%
High School/GED	8.49%	10.49%	9.23%
Vocational/Trade	1.85%	0.62%	1.39%
Partial College or 2-year Degree	12.55%	17.90%	14.53%
College or 4-year Degree	30.63%	35.80%	32.54%
Graduate Degree	39.48%	21.60%	32.87%
Not reported	4.06%	3.10%	3.70%
Family Income (yr)			
<10,000	5.54%	10.49%	7.37%
~10,000-25,000	8.12%	9.26%	8.54%
~25,000-50,000	12.18%	12.96%	12.47%
~50,000-75,000	17.34%	15.43%	16.63%
~75,000-100,000	18.82%	14.20%	17.11%
>100,000	34.32%	33.33%	33.95%
Not reported	3.69%	4.32%	3.92%

Note: Age = age in years, CU = University of Colorado Boulder, GBI-MH = General Behavior Inventory, Mania/Hypomania subscale, MASQ-AD = Mood and Anxiety Symptom Questionnaire, Anhedonic Loss of Interest subscale, PDS = Pubertal Development Scale *on *n*=402, UCLA = University of California Los Angeles

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

40

Table 2. Clinical Characteristics

	CU Boulder	UCLA	All
	<i>n</i> = 264	<i>n</i> = 155	<i>n</i> = 419
	%	%	%
Mood Disorders			
Unipolar Disorders			
Major Depressive Disorder	21.40%	23.46%	22.16%
Persistent Depressive Disorder	5.90%	22.84%	12.17%
Subclinical Depressive Disorder	12.55%	4.32%	9.51%
Bipolar Disorders			
Bipolar I Disorder	1.85%	4.32%	2.76%
Bipolar II Disorder	2.58%	3.09%	2.77%
Bipolar Disorder Not Otherwise Specified	1.11%	1.85%	1.38%
Subthreshold Bipolar Disorder	1.48%	0.62%	1.16%
Other Mood Disorder Not Otherwise Specified	5.90%	6.79%	6.23%
Schizoaffective Disorder	0.00%	1.23%	0.46%
No Mood Disorders	47.23%	31.48%	41.40%
Secondary Disorders			
Anxiety Disorders	14.77%	27.10%	19.33%
Substance Use Disorders			
Mild	5.30%	12.26%	7.87%
Moderate or Severe	1.14%	8.39%	3.82%
Eating Disorders	1.51%	5.16%	2.86%
Trauma and Stress-Related Disorders (Lifetime)	6.43%	14.84%	9.54%
Attention Deficit Hyperactivity Disorder	8.33%	14.19%	10.50%
Unknown	3.03%	7.10%	4.54%
Medications			
Anticonvulsant	1.14%	8.39%	3.82%
Anxiolytic (not within 48 hours of session)	0.76%	1.93%	1.19%
Atypical antipsychotic	0.38%	5.16%	2.15%
Benzodiazepine (not within 48 hours of session)	0.76%	3.22%	1.67%
Lithium	0.00%	1.93%	0.71%
Beta-blockers (not within 48 hours of session)	1.51%	1.93%	1.67%
Norepinephrine-dopamine reuptake inhibitor	0.38%	7.74%	3.10%
Selective serotonin-norepinephrine reuptake inhibitor	0.76%	3.87%	1.91%
Serotonin antagonist and reuptake inhibitor	1.14%	2.58%	1.67%
Selective serotonin reuptake inhibitor	8.71%	17.42%	11.93%
Stimulant (not within 48 hours of session)	4.54%	5.16%	4.77%
Tetracyclics	0.00%	0.38%	0.14%
Any psychoactive medication	20.07%	32.26%	24.58%


Note. Reported are lifetime mood diagnoses. Current mood symptom severity is reflected in primary dimensional measures reported in main text. Comorbid psychiatric disorders secondary to current mood disorders were permitted in this study and are reported above, based on standard DSM5 criteria and with extended reporting on Substance Use Disorders and Eating Disorders (i.e., remitted but recent diagnoses – past one year for SUDs, past two years for EDs – are reported for those categories) and Trauma and Stress Related Disorders (lifetime history). Current use of psychoactive medications is reported, and eligibility criteria required that participants were on a stable medication and dose (no changes in past six weeks) and abstained from use of stimulant, beta-blocker, anxiolytic, or benzodiazepine medication for 48+ hours prior to the research session. Analyses were repeated excluding individuals with secondary disorders or for whom secondary disorder data was incomplete/unknown, or covarying use of psychoactive medications, but these exclusions did not influence the nature or significance of effects therefore the full sample was retained for analyses.

DEVELOPMENTAL DIFFERENCES, MOOD, AND NEUROCOGNITION

41

Figure 1. Structural Equation Model. A structural equation model evaluated developmental differences in reward learning performance and executive functioning, and moderation of developmental differences by mood symptom severity. The reward learning performance factor was estimated across probabilistic reward task discriminability (PRT-DIS), instrumental learning task accuracy (ILT-ACC), and bandit task learning rate (BAN-LRN). The executive functioning factor was estimated across antisaccade accuracy (ANT-ACC), color-shape task reaction time switch cost (reversed for display, CSS-SWC), and spatial 2-back accuracy (S2B-ACC). The model tested the predictive linear and quadratic effects of pubertal stage (Pubertal Development Scale scores, PDS) and age on reward learning performance, and moderation of pubertal effects on reward learning performance by severity of manic symptoms (General Behavior Inventory-Mania/Hypomania scale, GBI-MH) and severity of anhedonic symptoms (Mood and Anxiety Symptom Questionnaire- Anhedonic Loss of Interest subscale, MASQ-AD). The model also tested predictive linear and quadratic effects of pubertal stage and age on executive functioning, and moderation of age effects on executive functioning by severity of manic or anhedonic symptoms. *Note:* Heavy lines with standardized estimates indicate significant path effects at $*p < 0.05$.

1
2
3
4
5 **Figure 2. Mood symptom severity moderates developmental differences in reward learning**
6 **performance and executive functioning.** (A) Quadratic association between pubertal stage and
7 reward learning performance factor scores. (B) Higher severity of anhedonic depression was
8 associated with stronger quadratic pubertal differences in reward learning performance,
9 including lower reward learning performance at earlier pubertal stages. (C) Higher severity of
10 mania was associated with weaker or reversed quadratic pubertal differences in reward learning
11 performance, including higher reward learning performance at earlier pubertal stages. (D)
12 Curvilinear (linear and quadratic effects) association between age and executive functioning
13 factor scores. (E) Severity of anhedonic depression did not significantly moderate age
14 differences in executive functioning. (F) Higher severity of mania moderated linear age
15 differences in executive functioning, in the form of poorer executive functioning especially at
16 older ages. *Note:* Plots show extracted factor scores on the y axes for ease of presentation, but
17 statistical associations among variables were estimated using structural equation models in which
18 reward learning performance and executive functioning were modeled as latent variables.
19 Displayed are lines fitted to models at low (-1.5 SD, blue) or high (+1.5 SD, red) levels of
20 symptom severity, and point color is scaled to low (blue) to high (red) symptom severity.
21 * $p < 0.05$ developmental (A, D) or moderated (B-C, E-F) effects.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Moderated Developmental Patterns

