

1
2
3 **Uptake and safety of pneumococcal vaccination in adults with immune mediated**
4 **inflammatory diseases: a UK wide observational study**
5
6
7
8
9

10 Georgina Nakafero¹, Matthew J. Grainge², Tim Card^{2,3}, Christian D. Mallen⁴, Jonathan
11 S. Nguyen Van-Tam², Abhishek Abhishek¹.
12
13
14

15
16
17 ¹Academic Rheumatology, School of Medicine, University of Nottingham, Nottingham,
18 UK. ²Lifespan and Population Health, School of Medicine, University of Nottingham,
19 Nottingham, UK. ³Nottingham Digestive Diseases Centre, Translational Medical
20 Sciences, School of Medicine, University of Nottingham, Nottingham, UK. ⁴Primary
21 Care Centre Versus Arthritis, School of Medicine, Keele University, Keele, UK.
22
23
24
25
26
27
28
29

30
31 **Corresponding author:** Dr Georgina Nakafero
32

33 **Address for correspondence:**
34

35 Dr Georgina Nakafero, Senior Research Fellow
36

37 A23, Academic Rheumatology
38

39 Clinical Sciences Building
40

41 School of Medicine
42

43 The University of Nottingham,
44

45 Nottingham NG5 1PB
46

47 UK
48

49 Email: georgina.nakafero@nottingham.ac.uk
50

51 ORCiD iD: <https://orcid.org/0000-0002-3859-7354>
52
53
54
55
56

57 © The Author(s) 2024. Published by Oxford University Press on behalf of the British Society for Rheumatology.
58 This is an Open Access article distributed under the terms of the Creative Commons Attribution License
59 (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium,
60 provided the original work is properly cited.

Abstract

Objective The uptake and safety of pneumococcal vaccination in people with immune mediated inflammatory diseases (IMIDs) is poorly understood. We investigated the UK wide pneumococcal vaccine uptake in adults with IMIDs and explored the association between vaccination and IMID flare.

Methods Adults with IMIDs diagnosed on or before 01/09/2018, prescribed steroid-sparing drugs within the last 12 months and contributing data to the Clinical Practice Research Datalink Gold were included. Vaccine uptake was assessed using a cross-sectional study design. Self-controlled case series (SCCS) analysis investigated the association between pneumococcal vaccination and IMID flare. The SCCS observation period was up-to six-month before and after pneumococcal vaccination. This was partitioned into a 14-day pre-vaccination induction, 90-days post-vaccination exposed, and the remaining unexposed periods.

Results We included 32,277 patients, 14,151 with RA, 13,631 with IBD, 3,804 with axial spondyloarthritis and 691 with SLE. Overall, 57% were vaccinated against pneumococcus. Vaccine uptake was lower in those younger than 45 years (32%), with IBD (42%), and without additional indication(s) for vaccination (46%). In the vaccine-safety study, data for 1,067, 935, and 451 vaccinated patients with primary-care consultations for joint pain, AIRD flare and IBD flare respectively were included. Vaccination against pneumococcal pneumonia was not associated with primary-care consultations for joint pain, AIRD flare and IBD flare in the exposed period with incidence rate ratios (95% Confidence Interval) 0.95 (0.83-1.09), 1.05 (0.92-1.19), and 0.83 (0.65-1.06) respectively.

1
2
3 **Conclusion** Uptake of pneumococcal vaccination in UK patients with IMIDs was
4 suboptimal. Vaccination against pneumococcal disease was not associated with IMID
5
6
7 flare.
8
9

10
11
12 **Keywords:** Pneumococcal vaccination, rheumatoid arthritis, psoriatic arthritis,
13 vaccine safety, vaccine uptake.
14
15
16

17 18 19 **Key messages**

- 20
21
22 • The uptake of pneumococcal vaccination in people with immune mediated
23 inflammatory diseases is suboptimal.
- 24
25
26 • Vaccination against pneumococcal disease is safe.
- 27
28
29 • Pneumococcal vaccination should be actively promoted in people with
30 inflammatory conditions.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Immunosuppressed adults with immune-mediated inflammatory diseases (IMIDs) such as rheumatoid arthritis (RA), inflammatory bowel disease (IBD), and systemic lupus erythematosus (SLE) are at an increased risk of pneumonia and its complications including death (1, 2, 3, 4). Consequently, pneumococcal vaccination is recommended for this at-risk population (5, 6). Despite long-standing recommendations for vaccinating the high risk groups since the year 1992 and the over 65s since the year 2003 (5), the uptake of pneumococcal vaccination in the at-risk populations was suboptimal (7). In a previous study from the UK, the uptake of pneumococcal vaccination among patients with RA was reported to be 50% overall, and 43% in those younger than 65 years in age (8). The uptake of pneumococcal vaccination across a broad range of IMIDs in a UK wide cohort has not been evaluated to the best of our knowledge. Understanding vaccine uptake across a range of conditions is important as the uptake of pneumococcal vaccination in people with IBD was noted to be lower in North America and Europe at 10.3%-38% (9, 10).

Belief that the vaccination could trigger an IMID flare, and cause other IMIDs e.g., vasculitis (11, 12) are key barriers to vaccination (9, 13, 14). The association between pneumococcal vaccination and IMID flare has not been evaluated in an adequately controlled study. There is some evidence from small studies restricted to a few conditions that vaccination against pneumococcal disease does not cause a flare of IMIDs (15, 16). In a systematic review and meta-analysis of pneumococcal vaccine immunogenicity studies in patients with SLE, disease activity did not worsen up to eight weeks after pneumococcal vaccination (17).

In this study we evaluated the uptake and safety of pneumococcal vaccination in UK adults with IMIDs.

Methods

Data source: Data from the Clinical Practice Research Datalink (CPRD) Gold were used in this study. Incepted in the year 1987, CPRD Gold is an anonymised longitudinal database of electronic health records of >14 million people in the UK. CPRD participants are representative of the UK population in terms of age, sex, and ethnicity (18). CPRD includes information on demographics, lifestyle factors, diagnoses stored as Read codes – a coded thesaurus of clinical terms, primary-care prescriptions, and immunisations. Vaccination and date of vaccination are also recorded.

Approval/patient consent This study was approved by Clinical Practice Research Datalink's Research Data Governance (Reference 21_000614), which has overarching research ethics committee approval for research studies using anonymous data. Practices that contributed data to the Clinical Practice Research Datalink consented to using anonymized patient data for approved research projects and additional consent was not required prior to individual studies (cprd.com).

Study design Cross-sectional and self-controlled case series (SCCS) study designs were used to examine the uptake and safety of pneumococcal vaccination.

Population Adults aged ≥ 18 years on the 1st September 2018, with at-least one primary-care record of an IMID (i.e., rheumatoid arthritis (RA), inflammatory bowel disease (IBD), axial spondyloarthritis (Ax-SpA), systemic lupus erythematosus (SLE)) and with at least one prescription of a steroid sparing drug (i.e., either methotrexate, azathioprine, 6-mercaptopurine, sulfasalazine, 5-aminosalicylates, mycophenolate, leflunomide, ciclosporin, tacrolimus or sirolimus) within the previous 12 months were eligible to be included in this study.

Pneumococcal vaccination Pneumococcal vaccination was defined using both product codes and Read codes (Supplementary Table S1, available at *Rheumatology* online). Dates of vaccination were extracted from CPRD. Data on vaccination from inception of CPRD in 1987 to 1st September 2018 were considered in the vaccine uptake study. Vaccinations recorded in the CPRD as not administered in primary care e.g., vaccination from hospitals, pharmacy were included in the vaccine uptake study but were excluded from the safety study because the date of administration is not reliably recorded in the CPRD. Similarly in the vaccine safety study we only considered the first vaccination in those with two or more records of vaccination as people that experience an adverse-event with a vaccination are less likely to agree to have a second dose of the vaccine if offered for any clinical indication.

Outcomes

Vaccine uptake Vaccination against pneumococcal pneumonia, defined as any pneumococcal vaccination up to 1st September 2018.

Vaccine safety:

A. Auto-immune rheumatic disease (AIRD)

[1] Primary care consultation for joint pain. This was defined using Read codes (Supplementary Table S2, available at *Rheumatology* online). Consultations for joint pain within 14 days of each other were considered as part of the same episode.

[2] AIRD flare. This was defined as present when there was a primary-care prescription of oral corticosteroid without another corticosteroid prescription in the preceding sixty days. The patient was also required to not have consulted for an alternate condition that could justify corticosteroid prescription on the same date. For this, all relevant primary care consultations were retrieved and reviewed by AA (General Medicine and Rheumatology expertise) for conditions that might explain the corticosteroid

1
2
3 prescribed and such participants were excluded from the analysis as there was
4 considerable uncertainty whether they experienced IMID flare or another illness
5 (Supplementary Table S3, available at *Rheumatology* online). The corticosteroid
6 prescription free period used to define consultation for AIRD flare was increased to at-
7 least 120 days in a sensitivity analysis as this time-period has been validated for the
8 IBD flare(19) (See below).
9

10
11
12 [3] RA flare. This was defined as present when there was either a Read code for RA
13 flare or a primary-care prescription of oral corticosteroid without another corticosteroid
14 prescription in the preceding sixty days in patients diagnosed with RA. The patient was
15 also required to not have consulted for an alternate condition that could justify
16 corticosteroid prescription on the same date. This condition was applied following the
17 same procedure as for AIRD flare described above.
18
19

20 B: IBD

21 [1] IBD flare. This was defined as present when there was primary-care prescription
22 of corticosteroid without another corticosteroid prescription in the preceding 120
23 days(19). The patient was also required to not have consulted for an alternate
24 condition that could justify corticosteroid prescription on the same date defined as a
25 new primary care prescription of corticosteroid(19).
26
27

28 **Covariates**

29 *Vaccine uptake* Age, sex, type of IMID and presence of additional indication(s) for
30 vaccination as per the UK Health Security Agency (5). Briefly, the additional indications
31 for vaccination included chronic heart diseases, chronic respiratory diseases, chronic
32 kidney diseases, chronic liver diseases, immunosuppression (defined as either solid
33 organ transplant, bone marrow transplant, HIV/AIDS, lymphoma, leukaemia,
34 myeloma, or chemotherapy), diabetes and asplenia. The vaccine uptake study was a
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 cross-sectional study that included patients alive on the 1st September 2018. Thus,
4
5 covariates were ascertained at 01/09/2018.
6

7
8 *Vaccine safety* Season defined in line with the Meteorological Office. According to the
9
10 Meteorological Office, winter spans from the 1st December to the 28th February of the
11
12 next year, spring spans from 1st March to 31st May, summer spans from 1st June to
13
14 31st August, and autumn spans from 1st September to 31st November.
15

16 17 **Statistical analyses**

18
19 *Vaccine uptake* The percentage and 95% CI of the study population alive on 1st
20
21 September 2018 that were vaccinated was calculated. The proportion vaccinated was
22
23 stratified according to their age (<45, 45–64, ≥65 years) on the 1st of September 2018,
24
25 sex, presence of other indications for vaccination, and type of IMID (RA, IBD, SLE,
26
27 Ax-SpA). Poisson regression with robust standard error was used to examine mutually
28
29 adjusted associations between pneumococcal vaccination and age group, sex, IMID
30
31 type and presence of additional at-risk condition for vaccination.
32
33

34
35 *Vaccine safety* Patients vaccinated against pneumococcal pneumonia and who also
36
37 consulted their GP for at-least one IMID flare in the six-month period before and the
38
39 six-month period after vaccination were included in a self-controlled case-series
40
41 (SCCS) analysis. SCCS is an established study design for assessing vaccine safety.
42
43 By including patients with both an exposure and an outcome, and undertaking within
44
45 person comparisons, SCCS analysis removes between person time-fixed
46
47 confounding, a key confounder in vaccine safety studies. The baseline period
48
49 extended from the latest of current registration date, first IMID diagnosis date recorded
50
51 in the CPRD, and 165 days preceding vaccination to 15-days pre-vaccination, and
52
53 from 90 days post-vaccination to the earliest date of six-months post vaccination,
54
55 leaving GP surgery, death, or last data collection from the GP surgery. The exposed
56
57
58
59
60

1
2
3 period extended from vaccination date to 90 days later and, was further categorised
4 as 0-14 days, 15-30 days, 31-60 days, and 61-90 days (Supplementary Figure S1,
5 available at *Rheumatology* online). The first cut-off was selected at 14-days post-
6 vaccination as it takes two weeks for the serological response and, we hypothesised
7 this period of immune reconstitution would be most likely to associate with disease
8 activity. The 15-day period immediately preceding vaccination was excluded from the
9 baseline period to minimise confounding due to healthy vaccinee effect or due to active
10 promotion of vaccination in those consulting for a disease flare(20).
11

12 A Poisson model conditioned on the number of events adjusted for the four UK
13 seasons as categories was fitted to calculate incidence rate ratios (aIRR) and 95%
14 confidence interval (CI) for each exposure period compared to the baseline period.
15 This approach was also followed to assess the association between pneumococcal
16 vaccination and AIRD flare, defined as a ≥ 4 -month gap between corticosteroid
17 prescriptions in people with autoimmune rheumatic diseases during the observation
18 period as a sensitivity analysis.
19

20
21
22 Data management and analysis were performed in Stata v17, Stata Corp LLC,
23 Texas, USA.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Uptake Data from 32,277 people with IMIDs were included in this study (Figure 1). Their mean age (SD) on 1st of September 2018 was 58 (16) years and 59% were female. 14,151 (43.8%) had RA, 13,631 (42.2%) IBD, 3,804 (11.8%) Ax-SpA, and 691 (2.1%) SLE.

Overall uptake of pneumococcal vaccination was 56.5% (95%CI 55.9-57.0%). Pneumococcal vaccinations occurred between 01/01/1992 and 01/09/2018. Vaccination uptake was significantly lower in people with IBD (42.4%, 95% CI (41.6-43.2%)) than in those with AIRDs (66.8%, 95% CI (66.1-67.4%)) with adjusted incidence rate ratio (aIRR) 0.75 (95%CI 0.73- 0.76) . Increasing age, female sex and presence of additional at-risk conditions were independently associated with the uptake of pneumococcal vaccination (Table 1). Vaccination uptake was higher in people aged at-least 65 years or with an additional at-risk condition than in those less than 65-years in age and without an additional at-risk condition for vaccination (proportion vaccinated (95% CI); 0.72 (0.72-0.73) and 0.37(0.36-0.38) respectively). On adjusting for gender and type of inflammatory condition, people not considered at additional risk of pneumococcal pneumonia (i.e., age < 65 years and without another at-risk condition) were 46% less likely to get vaccinated than those considered at additional risk of pneumococcal pneumonia (i.e., aged ≥65-year and/or with additional at-risk condition) with aIRR 0.54 (95% CI 0.53-0.56)).

Vaccine Safety Data for 1,067, 935, 778 and 451 participants with primary care consultations for joint pain, AIRD flare, RA flare and IBD flare respectively were included. 1,838 participants had either AIRD flare or a primary care consultation for joint pain and of these, 1,412 (76.8%) had RA, 281 (15.3%) had SpA and 145 (7.9%) had SLE. The majority were female (71.6%) and their mean (SD) age was 55 (12)

1
2
3 years. Of the participants with IBD flare, 240 (53.2%) had ulcerative colitis, 160
4 (35.5%) had Crohn's disease, 51 (11.3%) had IBD without any specific coding for
5
6 subtype.
7
8

9
10 Vaccination against pneumococcal pneumonia was not associated with joint pain
11 consultation, AIRD flare or IBD flares respectively in the 90-days post vaccination
12 (Tables 2, 3). The 15-day pre-vaccination period associated with significantly more
13 primary care consultations for joint pain, AIRD flare and IBD flare (Tables 2, 3).
14
15
16
17
18
19
20

21 **Discussion**

22
23 This UK wide study has shown that approximately one in two immunosuppressed
24 adults with IMIDs in the UK is vaccinated against pneumococcal pneumonia. This is
25 similar to the vaccine uptake of 54% to 56% reported in people with chronic respiratory
26 disease, chronic kidney disease, and diabetes requiring insulin or oral hypoglycaemic
27 medication (21). The vaccine uptake was even lower at 31.8% in those less than 45-
28 years in age and at 46.4% in IMID patients without an additional indication for
29 vaccination. The vaccine uptake ranged from 42.4% to 69.7% in IBD and RA
30 respectively.
31
32
33
34
35
36
37
38
39
40

41 This study did not find an association between pneumococcal vaccination and IMID
42 flare requiring primary-care consultation and/or treatment. An increased risk in flare of
43 underlying disease was observed 15-days pre-vaccination, which could be attributed
44 to opportunistic vaccine promotion to people consulting for an IMID flare resulting in
45 vaccination.
46
47
48
49
50
51
52

53 It is difficult to compare our findings on vaccine uptake in IMIDs with those of previous
54 studies, since, to our knowledge, this is the first study to assess pneumococcal
55 vaccination uptake across many inflammatory conditions and to compare uptake
56 between different inflammatory conditions. These low vaccination rates are
57
58
59
60

1
2
3 concerning given the increased risk of pneumococcal disease in this at-risk population
4 for whom vaccination is recommended and indicates that they would benefit from
5 targeted measures to increase pneumococcal vaccine uptake. We observed
6 substantial variation in vaccination rates across different inflammatory diseases. This
7 may reflect differential advice from different specialties. A 2013 UK primary care study
8 reported 50% pneumococcal vaccine uptake in RA patients which was lower than the
9 70% pneumococcal vaccine uptake reported in the current study (8). This
10 improvement in pneumococcal vaccination uptake over a 5-year period is remarkable
11 and may be related to clear guidance from the British Society of Rheumatology to offer
12 vaccination against pneumococcal pneumonia in patients with RA (22, 23). Similarly,
13 more and more patients with RA are being treated with potent combination DMARDs
14 and this has resulted in vaccination being promoted more actively in people with this
15 condition(24). Improvement in pneumococcal vaccination uptake in RA patients has
16 also been reported in a multi-centre prospective study of 1,679 patients in Greece
17 (25). In our study patients with IBD had a low vaccine uptake. A similar low
18 pneumococcal vaccine uptake of 38% has been reported from a French online survey
19 of IBD patients (10) while a lower rate of 10.3% was reported from a gastro-enterology
20 clinic in Canada (9). Pneumococcal vaccine uptake was higher in people with Ax-SpA
21 in the current study than has been reported in other countries in Europe. This may be
22 because Ax-SpA patients treated with NSAIDs alone were not included in this study.
23 For comparison, in Switzerland, the pneumococcal vaccine uptake was reported to be
24 32.5% in an online survey of Ax-SpA patients that was not restricted to patients on
25 immunosuppressive treatment (26). The multinational COMORA study reported higher
26 uptake of pneumococcal vaccine in some countries e.g., France, but lower uptake in
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 most other countries (27). These questionnaire studies are prone to bias from self-
4 reported vaccination uptake and should be interpreted with caution.
5
6

7
8 Consistent with previous research on factors associated with increased vaccine
9 uptake, female sex, increasing age and other indications for vaccination significantly
10 associated with pneumococcal vaccination (13, 28, 29).
11
12

13
14 Barriers to vaccination have included the fear that vaccines may trigger an IMID flare
15 (14, 30). This study did not find a significant association between pneumococcal
16 vaccination and flare of the underlying inflammatory disease. Similarly, a systematic
17 review and meta-analysis of pneumococcal vaccine immunogenicity studies in
18 patients with SLE did not find an association between the vaccination and increased
19 disease activity (17). Safety studies in the general population have shown that
20 pneumococcal vaccine is well tolerated(31). Similarly, there was no association
21 between vaccination with the seasonal influenza vaccine and AIRD flare, and no
22 association between vaccination against COVID-19 and AIRD, IBD, and psoriasis flare
23 in previous studies (32, 33, 34, 35). A meta-analysis of prior uncontrolled studies
24 reported a 2% pooled prevalence of IBD flare after vaccination, however, it is not
25 known if the flares were temporally related or coincidental (36).
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Strengths of this study included a large nationally representative sample of people with
43 IMIDs in the UK given the almost universal registration with a GP for all UK residents
44 (18). We studied a wide range of IMIDs improving the generalizability of the findings.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
The use of primary-care prescription and consultation data minimised recall bias on
the association between vaccination and disease flares. To improve the validity of our
case definition, we used a combination of diagnostic and prescription codes to
ascertain people with IMIDs. Additionally, we defined IBD flares according to a
validated definition (19) and we undertook a sensitivity analysis for the association

1
2
3 between pneumococcal vaccination and AIRD flare using a validated IBD flare
4 definition. Furthermore, to improve the outcome fidelity, we excluded participants with
5
6 diagnoses that could potentially explain corticosteroid prescribed on the same date as
7
8 the AIRD or IBD flare. Finally, our use of SCCS methodology controlled for between-
9
10 person confounding which is a serious problem in observational studies of vaccine
11
12 safety.

13
14
15
16
17 There are some limitations to our study. Firstly, some vaccinations that were
18
19 administered outside of primary-care for example in hospital or at the workplace as for
20
21 health care professionals may not have been recorded in the CPRD, reducing
22
23 vaccination uptake estimates. This is unlikely to have a significant impact on our
24
25 results as vaccination is almost exclusively a general practice activity in the UK. Where
26
27 non-primary care administration of the vaccine was recorded, it was excluded from the
28
29 vaccine safety study as it is difficult to be sure of the date of vaccination in such
30
31 instances. Second, the type of vaccine was not assessed as the vast majority of
32
33 vaccinations were with the PPV23 vaccine, which has been universally used in the UK
34
35 for risk groups since the year 1992 (5). Third, we were unable to assess the impact of
36
37 biologics on vaccine safety because their prescription is not recorded in the CPRD.
38
39 We see no reason though to expect more extreme immunologically driven side effects
40
41 in these groups given the possibility of less immunogenic response with biologic use
42
43 (37). Fourth, data on disease activity and flares managed in hospital or specialist
44
45 clinics are not recorded in CPRD. . Fifth, because our definition of AIRD and IBD flare
46
47 required consultation and/or prescription, minor flares not needing drug treatment
48
49 were not considered as an outcome in the vaccine safety study. It is possible that there
50
51 may be an association of vaccination with minor flares that were not ascertained in our
52
53 study. However, such effects would be unlikely to greatly discourage vaccination
54
55
56
57
58
59
60

1
2
3 uptake and it is the more significant flares that we have studied which are of primary
4 concern. Flares managed in hospital or specialist clinics were excluded. Additionally,
5
6 joint pain was considered as an outcome of interest because it is a common symptom
7
8 of inflammatory arthritis. However, joint pain might also be caused by another illness
9
10 such as osteoarthritis reducing the specificity for this outcome.
11
12

13
14 In conclusion, this study provides recent UK-wide population evidence that the uptake
15
16 of pneumococcal vaccination in people with IMIDs is suboptimal particularly in patients
17
18 with IBD, those younger than 65 years in age, and in those without another indication
19
20 for vaccination. It also demonstrated that pneumococcal vaccination does not
21
22 associate with flare of the underlying IMIDs. These data should be used to promote
23
24 pneumococcal vaccination in this at-risk population.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Funding** This work was funded by a grant from the NIHR (Reference number:
4
5 NIHR201973)
6

7
8 **Disclosure statement** A.A. has received personal fees from UpToDate (royalty),
9
10 Springer (royalty), Cadilla Pharmaceuticals (lecture fees), NGM Bio (consulting), and
11
12 Limbic (consulting) unrelated to this work. CDM is Director of the NIHR School for
13
14 Primary Care Research. Keele University has received research funding for CDM from
15
16 NIHR, MRC, Versus Arthritis and BMS. JSN-V-T was seconded to the Department of
17
18 Health and Social Care (DHSC) from October 2017 to March 2022. Since March 2022
19
20 he has received personal fees from CSL Seqirus (lectures, writing and consulting),
21
22 AstraZeneca (lecture) and Sanofi Pasteur (lectures and speaking) all of whom
23
24 manufacture influenza vaccines. He has consulted for Moderna Therapeutics who are
25
26 developing influenza vaccines. He has performed consultancy for MSD, which
27
28 manufactures PPV23, in spring 2023 but unrelated to PPV23. The views expressed in
29
30 this manuscript are those of its authors and not necessarily those of DHSC or any
31
32 other entity mentioned above. The other authors have no conflict of interest to declare.
33
34
35

36
37 **Author contribution** The study was conceived by Prof Abhishek. All authors were
38
39 involved in the design of the study. The analysis was carried out by Dr Nakafero. All
40
41 authors edited the first and all subsequent drafts and approved the final draft for
42
43 submission.
44
45

46
47 **Data sharing statement** Data used in the study are from the Clinical Practice
48
49 Research Datalink (CPRD). Due to CPRD licencing rules, we are unable to share data
50
51 used in this study with third parties. The data used in this study may be obtained
52
53 directly from the CPRD. Study protocol is available from www.cprd.com.
54
55
56
57
58
59
60

References

1. van Aalst M, Lötsch F, Spijker R, van der Meer JTM, Langendam MW, Goorhuis A, et al. Incidence of invasive pneumococcal disease in immunocompromised patients: A systematic review and meta-analysis. *Travel Med Infect Dis*. 2018;24:89-100.
2. Wotton CJ, Goldacre MJ. Risk of invasive pneumococcal disease in people admitted to hospital with selected immune-mediated diseases: record linkage cohort analyses. *J Epidemiol Community Health*. 2012;66(12):1177-81.
3. Shea KM, Edelsberg J, Weycker D, Farkouh RA, Strutton DR, Pelton SI. Rates of pneumococcal disease in adults with chronic medical conditions. *Open Forum Infect Dis*. 2014;1(1):ofu024.
4. Shigayeva A, Rudnick W, Green K, Chen DK, Demczuk W, Gold WL, et al. Invasive Pneumococcal Disease Among Immunocompromised Persons: Implications for Vaccination Programs. *Clin Infect Dis*. 2016;62(2):139-47.
5. UK Health Security Agency. *Pneumococcal: the green book*, chapter 25. 2023.
6. Lopez A, Mariette X, Bachelez H, Belot A, Bonnotte B, Hachulla E, et al. Vaccination recommendations for the adult immunosuppressed patient: A systematic review and comprehensive field synopsis. *J Autoimmun*. 2017;80:10-27.
7. Campling J, Vyse A, Liu H-H, Wright H, Slack M, Reinert R-R, et al. A review of evidence for pneumococcal vaccination in adults at increased risk of pneumococcal disease: risk group definitions and optimization of vaccination coverage in the United Kingdom. *Expert Review of Vaccines*. 2023;22(1):785-800.
8. Costello R, Winthrop KL, Pye SR, Brown B, Dixon WG. Influenza and Pneumococcal Vaccination Uptake in Patients with Rheumatoid Arthritis Treated with Immunosuppressive Therapy in the UK: A Retrospective Cohort Study Using Data from the Clinical Practice Research Datalink. *PLoS One*. 2016;11(4):e0153848.
9. Malhi G, Rumman A, Thanabalan R, Croitoru K, Silverberg MS, Hillary Steinhart A, et al. Vaccination in inflammatory bowel disease patients: attitudes, knowledge, and uptake. *J Crohns Colitis*. 2015;9(6):439-44.
10. Loubet P, Verger P, Abitbol V, Peyrin-Biroulet L, Launay O. Pneumococcal and influenza vaccine uptake in adults with inflammatory bowel disease in France: Results from a web-based study. *Dig Liver Dis*. 2018;50(6):563-7.
11. Birck R, Kaelsch I, Schnuelle P, Flores-Suárez LF, Nowack R. ANCA-associated vasculitis following influenza vaccination: causal association or mere coincidence? *J Clin Rheumatol*. 2009;15(6):289-91.
12. Abdalla A, Sebaoui S, Alraqi S. Diffuse cutaneous reaction following PPV-23 pneumococcal vaccine: an immunisation-associated hypersensitivity vasculitis. *BMJ Case Rep*. 2020;13(3).
13. Loubet P, Kernéis S, Groh M, Loulergue P, Blanche P, Verger P, et al. Attitude, knowledge and factors associated with influenza and pneumococcal vaccine uptake in a large cohort of patients with secondary immune deficiency. *Vaccine*. 2015;33(31):3703-8.
14. Fuller A, Hancox J, Vedhara K, Card T, Mallen C, Van-Tam JSN, et al. Barriers and facilitators to vaccination uptake against COVID-19, influenza, and pneumococcal pneumonia in immunosuppressed adults with immune-mediated inflammatory diseases: A qualitative interview study during the COVID-19 pandemic. *PLoS One*. 2022;17(9):e0267769.
15. Rezende R, Ribeiro F, Albuquerque E, Gayer C, Andrade L, Klumb E. Immunogenicity of pneumococcal polysaccharide vaccine in adult systemic lupus erythematosus patients undergoing immunosuppressive treatment. *Lupus*. 2016;25(11):1254-9.
16. Migita K, Akeda Y, Akazawa M, Tohma S, Hirano F, Ideguchi H, et al. Opsonic and Antibody Responses to Pneumococcal Polysaccharide in Rheumatoid Arthritis Patients Receiving Golimumab Plus Methotrexate. *Medicine (Baltimore)*. 2015;94(52):e2184.
17. Pugès M, Biscay P, Barnetche T, Truchetet M, Richez C, Seneschal J, et al. Immunogenicity and impact on disease activity of influenza and pneumococcal vaccines

- 1
2
3 in systemic lupus erythematosus: a systematic literature review and meta-analysis.
4 *Rheumatology (Oxford)*. 2016;55(9):1664-72.
- 5 18. Herrett E, Gallagher AM, Bhaskaran K, Forbes H, Mathur R, van Staa T, et al.
6 Data Resource Profile: Clinical Practice Research Datalink (CPRD). *International journal*
7 *of epidemiology*. 2015;44(3):827-36.
- 8 19. Lewis JD, Aberra FN, Lichtenstein GR, Bilker WB, Brensinger C, Strom BL.
9 Seasonal variation in flares of inflammatory bowel disease. *Gastroenterology*.
10 2004;126(3):665-73.
- 11 20. Nakafero G, Grainge MJ, Myles PR, Mallen CD, Zhang W, Doherty M, et al.
12 Association between inactivated influenza vaccine and primary care consultations for
13 autoimmune rheumatic disease flares: a self-controlled case series study using data
14 from the Clinical Practice Research Datalink. *Annals of the Rheumatic Diseases*.
15 2019;78(8):1122-6.
- 16 21. Public Health England. Pneumococcal Polysaccharide Vaccine (PPV) coverage
17 report, England, April 2020 to March 2021. In: Report HP, editor. 2022.
- 18 22. van Assen S, Agmon-Levin N, Elkayam O, Cervera R, Doran MF, Dougados M, et
19 al. EULAR recommendations for vaccination in adult patients with autoimmune
20 inflammatory rheumatic diseases. *Ann Rheum Dis*. 2011;70(3):414-22.
- 21 23. Singh JA, Saag KG, Bridges SL, Jr., Akl EA, Bannuru RR, Sullivan MC, et al. 2015
22 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis.
23 *Arthritis Care Res (Hoboken)*. 2016;68(1):1-25.
- 24 24. National Institute for Health and Care Excellence (NICE). Scenario: General
25 principles of managing adults on DMARDs. 2023.
- 26 25. Thomas K, Lazarini A, Kaltsonoudis E, Voulgari PV, Drosos AA, Repa A, et al.
27 Patterns and factors associated with pneumococcal vaccination in a prospective cohort of
28 1,697 patients with rheumatoid arthritis. *Front Med (Lausanne)*. 2022;9:1039464.
- 29 26. Stoffel ST, Colaninno A, Bräm R, Schwenkglens M. Pneumococcal vaccination
30 among adult risk patient with axial spondyloarthritis in Switzerland: Data from the
31 survey of the ankylosing spondylitis association of Switzerland (SVMB). *Vaccine*.
32 2022;40(43):6206-10.
- 33 27. Hmamouchi I, Winthrop K, Launay O, Dougados M. Low rate of influenza and
34 pneumococcal vaccine coverage in rheumatoid arthritis: data from the international
35 COMORA cohort. *Vaccine*. 2015;33(12):1446-52.
- 36 28. Nakafero G, Grainge MJ, Myles PR, Mallen CD, Zhang W, Doherty M, et al.
37 Predictors and temporal trend of flu vaccination in auto-immune rheumatic diseases in
38 the UK: a nationwide prospective cohort study. *Rheumatology (Oxford)*.
39 2018;57(10):1726-34.
- 40 29. Garg M, Mufti N, Palmore TN, Hasni SA. Recommendations and barriers to
41 vaccination in systemic lupus erythematosus. *Autoimmunity reviews*. 2018;17(10):990-
42 1001.
- 43 30. Lawson EF, Trupin L, Yelin EH, Yazdany J. Reasons for failure to receive
44 pneumococcal and influenza vaccinations among immunosuppressed patients with
45 systemic lupus erythematosus. *Semin Arthritis Rheum*. 2015;44(6):666-71.
- 46 31. Musher DM, Manof SB, Liss C, McFetridge RD, Marchese RD, Bushnell B, et al.
47 Safety and antibody response, including antibody persistence for 5 years, after primary
48 vaccination or revaccination with pneumococcal polysaccharide vaccine in middle-aged
49 and older adults. *J Infect Dis*. 2010;201(4):516-24.
- 50 32. Adams L, Nakafero G, Grainge MJ, Card T, Mallen CD, Van-Tam JSN, et al. Is
51 vaccination against COVID-19 associated with psoriasis or eczema flare? Self-controlled
52 case series analysis using data from the Clinical Practice Research Datalink (Aurum). *Br J*
53 *Dermatol*. 2023;188(2):297-9.
- 54 33. Card TR, Nakafero G, Grainge MJ, Mallen CD, Van-Tam JSN, Williams HC, et al. Is
55 Vaccination Against COVID-19 Associated With Inflammatory Bowel Disease Flare? Self-
56 Controlled Case Series Analysis Using the UK CPRD. *Am J Gastroenterol*.
57 2023;118(8):1388-94.
- 58 34. Nakafero G, Grainge MJ, Card T, Mallen CD, Nguyen Van-Tam JS, Williams HC, et
59 al. Is vaccination against COVID-19 associated with autoimmune rheumatic disease
60

1
2
3 flare? A self-controlled case series analysis. *Rheumatology (Oxford)*. 2023;62(4):1445-
4 50.

5 35. Nakafero G, Grainge MJ, Myles PR, Mallen CD, Zhang W, Doherty M, et al.
6 Association between inactivated influenza vaccine and primary care consultations for
7 autoimmune rheumatic disease flares: a self-controlled case series study using data
8 from the Clinical Practice Research Datalink. *Ann Rheum Dis*. 2019;78(8):1122-6.

9 36. Desalermos A, Pimienta M, Kalligeros M, Shehadeh F, Diamantopoulos L,
10 Karamanolis G, et al. Safety of Immunizations for the Adult Patient With Inflammatory
11 Bowel Disease-A Systematic Review and Meta-analysis. *Inflamm Bowel Dis*.
12 2022;28(9):1430-42.

13 37. Hua C, Barnetche T, Combe B, Morel J. Effect of Methotrexate, Anti-Tumor
14 Necrosis Factor α , and Rituximab on the Immune Response to Influenza and
15 Pneumococcal Vaccines in Patients With Rheumatoid Arthritis: A Systematic Review and
16 Meta-Analysis. *Arthritis Care Res (Hoboken)*. 2014;66(7):1016-26.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Uptake and risk factors of pneumococcal vaccination in immune mediated inflammatory diseases (n=32,277)

		Vaccination uptake			Incidence rate ratios (IRR) and 95% confidence interval (CI)	
		Total number	Number vaccinated	Percent (95% CI)	Crude	Adjusted
Overall		32,277	18,227	56.47 (55.93,57.01)	-	-
Age, years						
	< 45	7,190	2,285	31.78 (30.71,32.87)	1	1
	45-64	13,045	6,297	48.27 (47.41,49.13)	1.52 (1.46,1.58)	1.36 (1.31,1.42)
	≥65	12,042	9,645	80.09 (79.37,80.80)	2.52 (2.43,2.61)	2.07 (1.99,2.14)
Sex						
	Male	13,231	7,055	53.32 (52.47,54.17)	1	1
	Female	19,046	11,172	58.66 (58.00,59.36)	1.10 (1.08,1.12)	1.04 (1.02,1.06)
Additional clinical risk group(s)						
	Absent	19,643	9,117	46.41 (45.72,47.11)	1	1
	Present	12,634	9,110	72.11 (71.32,72.88)	1.55 (1.53,1.58)	1.30 (1.28,1.32)
Inflammatory condition						
	Rheumatoid Arthritis	14,151	9,868	69.73 (68.97,70.49)	1	1
	Inflammatory Bowel Disease	13,631	5,779	42.40 (41.57,43.23)	0.61 (0.59,0.62)	0.75 (0.73,0.76)
	Systemic Lupus Erythematosus	691	387	56.01 (52.28,59.67)	0.80 (0.75,0.86)	0.98 (0.92,1.04)
	Axial spondyloarthritis*	3,804	2,193	57.65 (56.07,59.21)	0.83 (0.80,0.85)	0.99 (0.96,1.02)

*Psoriatic arthritis, reactive arthritis, and ankylosing spondylitis

Table 2: The association between pneumococcal vaccination and consultation for autoimmune rheumatic disease (AIRD) flare, rheumatoid arthritis (RA) flare, joint pain, and inflammatory bowel disease (IBD) flare

Outcome	Risk period (days)	Events (n)	Person-time (days)	IRR (95%CI)	Adjusted IRR (95%CI) *	p-value
AIRD flare	Baseline	1,048	373,886	1	1	-/-
	15 days pre-vaccination	96	22,656	1.50	1.52	<0.001
	Post vaccination intervals					
	0 - 90 days	391	138,045	0.99	1.05	0.514
	0 - 14 days	63	22,914	0.97	0.99	0.953
	15 - 30 days	51	22,985	0.78	0.80	0.133
RA flare	Baseline	856	30,5713	1	1	-/-
	15 days pre-vaccination	77	18,521	1.48	1.50	0.001
	Post vaccination intervals					
	0 - 90 days	321	11,3087	0.99	1.04	0.430
	0 - 14 days	54	18,769	1.01	1.04	0.782
	15 - 30 days	42	18,835	0.78	0.81	0.184
Joint pain	Baseline	956	33,1904	1	1	-/-
	15 days pre-vaccination	80	20,153	1.38	1.38	0.006
	Post vaccination intervals					
	0 - 90 days	341	122,959	0.94	0.95	0.434
	0 - 14 days	60	20,371	1.01	1.02	0.859
	15 - 30 days	46	20,469	0.77	0.78	0.107
IBD flare	Baseline	338	125,190	1	1	-/-
	15 days pre-vaccination	35	7,485	1.72	1.79	0.001
	Post vaccination intervals					
	0 - 90 days	126	44,906	1.03	0.83	0.143
	0 - 14 days	18	7,485	0.89	0.89	0.629
	15 - 30 days	16	7,485	0.79	0.75	0.271
	31 - 60 days	53	14,970	1.30	1.07	0.666

	61 - 90 days	39	14,966	0.96 (0.69,1.34)	0.79 (0.55,1.13)	0.193
--	-----------------	----	--------	---------------------	---------------------	-------


*adjusted for season. The baseline period extended from the latest of current registration date, first disease diagnosis date recorded in the CPRD, and six-months preceding vaccination to 15-days pre-vaccination, and from 90 days post-vaccination to the earliest date of six-months post vaccination, leaving GP surgery, death, or last data collection from the GP surgery.

Table 3. The association between pneumococcal vaccination and autoimmune rheumatic disease (AIRD) flare[†]: Sensitivity analysis

Risk period (days)	Events (n)	Person-time (days)	IRR (95%CI)	Adjusted IRR (95%CI) *	p-value
Baseline	225	86,797	1	1	-/-
15 days pre-vaccination	24	5,204	1.78 (1.17,2.71)	1.94 (1.26,2.97)	0.002
Post vaccination intervals					
0 - 90 days	105	31,817	1.25 (0.99,1.58)	1.21 (0.92,1.58)	0.166
0 - 14 days	18	5,204	1.30 (0.80,2.10)	1.41 (0.87,2.29)	0.167
15 - 30 days	13	5,299	0.93 (0.53,1.63)	0.99 (0.56,1.74)	0.967
31 - 60 days	36	10,620	1.28 (0.90,1.82)	1.29 (0.89,1.87)	0.179
61 - 90 days	38	10,620	1.35 (0.96,1.91)	1.34 (0.93,1.94)	0.114

*adjusted for season; [†]AIRD flare defined as a ≥4-month gap between steroid prescriptions in people with autoimmune rheumatic diseases (AIRDs) during the observation period. The baseline period extended from the latest of current registration date, first disease diagnosis date recorded in the CPRD, and six-months preceding vaccination to 15-days pre-vaccination, and from 90 days post-vaccination to the earliest date of six-months post vaccination, leaving GP surgery, death, or last data collection from the GP surgery.

Figure 1: Participant selection criteria



CPRD: Clinical Practice Research Datalink.

Consistent safety profile with over 8 years of real-world evidence, across licensed indications¹⁻³



1,000,000 patients treated globally, and counting*⁴



100+
clinical trials*⁵



8+ years of
real-world evidence¹⁻³



8
indications¹⁻³



Click here to visit our HCP portal and learn more

Real-world evidence shows a consistent safety profile over 6 years^{6,7}

No trend toward increased AE rates over time (pooled PsA, AS, PsO):^{†6}

AEs of select interest (EAIR per 100 PY)	1 year	2 years	3 years	4 years	5 years	6 years	Cumulative rate
Serious infections Cases	2.0 n=149	1.7 n=475	0.7 n=649	1.3 n=1,841	1.3 n=2,285	1.1 n=2,226	1.3 n=8,719
Malignant or unspecified tumours Cases	0.2 n=15	0.2 n=50	0.2 n=225	0.3 n=422	0.3 n=520	0.3 n=573	0.3 n=1,896
MACE Cases	0.2 n=15	0.1 n=39	0.2 n=151	0.2 n=238	0.2 n=264	0.1 n=287	0.2 n=1,031
Total IBD Cases	0.2 n=12	0.2 n=46	0.2 n=185	0.3 n=340	0.2 n=312	0.1 n=261	0.2 n=1,291
Exposure (PY)	7450	28,549	93,744	137,325	182,024	212,636	680,470

No trend towards increased rates of malignancy, MACE or IBD over time⁶

The most frequently reported adverse reactions are upper respiratory tract infections (17.1%) (most frequently nasopharyngitis, rhinitis).^{1,2} Refer to the prescribing information for a summary of adverse events.

Adapted from Novartis Data on File. 2021.⁶

Refer to the Cosentyx Summary of Product Characteristics for full details, dosing and administration, including special populations.

Cosentyx® (secukinumab) licensed indications in rheumatology: Cosentyx, alone or in combination with methotrexate, is indicated for the treatment of active **psoriatic arthritis** in adult patients when the response to previous disease-modifying anti-rheumatic drug therapy has been inadequate; active **ankylosing spondylitis** in adults who have responded inadequately to conventional therapy; active **non-radiographic axial spondyloarthritis** with objective signs of inflammation as indicated by elevated C-reactive protein and/or magnetic resonance imaging evidence in adults who have responded inadequately to non-steroidal anti-inflammatory drugs; active **enthesitis-related arthritis** in patients 6 years and older (alone or in combination with methotrexate) whose disease has responded inadequately to, or who cannot tolerate conventional therapy; active **juvenile psoriatic arthritis** in patients 6 years or older (alone or in combination with methotrexate) whose disease has responded inadequately to, or who cannot tolerate, conventional therapy.^{1,2}

Prescribing information, adverse event reporting and full indication can be found on the next page.

*Patients prescribed Cosentyx for any indication since launch.

[†]Successive time periods of PSUR shown with cumulative rate: 26 Dec 2014 to 25 Dec 2015; 26 Dec 2015 to 25 Dec 2016; 26 Dec 2016 to 25 Dec 2017; 26 Dec 2017 to 25 Dec 2018; 26 Dec 2018 to 25 Dec 2019; 26 Dec 2019 to 25 Dec 2020.⁶

Abbreviations: AE, adverse event; AS, ankylosing spondylitis; EAIR, exposure-adjusted incidence rate; HCP, healthcare professional; IBD, inflammatory bowel disease; MACE, major adverse cardiac event; PsA, psoriatic arthritis; PsO, plaque psoriasis; PY, patient year.

References: **1.** Cosentyx® (secukinumab) GB Summary of Product Characteristics; **2.** Cosentyx® (secukinumab) NI Summary of Product Characteristics; **3.** European Medicines Agency, European public assessment report. Available at: https://www.ema.europa.eu/en/documents/overview/cosentyx-epar-medicine-overview_en.pdf [Accessed February 2024]; **4.** Novartis Data on File. Secukinumab – Sec008. 2023; **5.** Novartis. Novartis Cosentyx® positive 16-week PREVENT results advance potential new indication for patients with axial spondyloarthritis. Available at: <https://www.novartis.com/news/media-releases/novartis-cosentyx-positive-16-week-prevent-results-advance-potential-new-indication-patients-axial-spondyloarthritis> [Accessed February 2024]; **6.** Novartis data on file. Cosentyx Periodic Safety Update Report (PSUR); 26 December 2019 – 25 December 2020. 22 February 2021; **7.** Deodhar A, et al. Arthritis Res Ther 2019;21(1):111.



Cosentyx® (secukinumab) Northern Ireland Prescribing Information.

Please refer to the Summary of Product Characteristics (SmPC) before prescribing.

Indications: Treatment of: moderate to severe plaque psoriasis in adults, children and adolescents from the age of 6 years who are candidates for systemic therapy; active psoriatic arthritis in adults (alone or in combination with methotrexate) who have responded inadequately to disease-modifying anti-rheumatic drug therapy; active ankylosing spondylitis in adults who have responded inadequately to conventional therapy; active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation as indicated by elevated C-reactive protein (CRP) and/or magnetic resonance imaging (MRI) evidence in adults who have responded inadequately to non-steroidal anti-inflammatory drugs; active enthesitis-related arthritis and juvenile psoriatic arthritis in patients 6 years and older (alone or in combination with methotrexate) whose disease has responded inadequately to, or who cannot tolerate, conventional therapy; active moderate to severe hidradenitis suppurativa (acne inversa) in adults with an inadequate response to conventional systemic HS therapy. **Presentations:** Cosentyx 150 mg solution for injection in pre-filled pen; Cosentyx 300 mg solution for injection in pre-filled pen. **Dosage & Administration:** Administered by subcutaneous injection at weeks 0, 1, 2, 3 and 4, followed by monthly maintenance dosing. Consider discontinuation if no response after 16 weeks of treatment. Each 150 mg dose is given as one injection of 150 mg. Each 300 mg dose is given as two injections of 150 mg or one injection of 300 mg. If possible avoid areas of the skin showing psoriasis. **Plaque Psoriasis:** Adult recommended dose is 300 mg monthly. Based on clinical response, a maintenance dose of 300 mg every 2 weeks may provide additional benefit for patients with a body weight of 90 kg or higher. Adolescents and children from the age of 6 years: if weight ≥ 50 kg, recommended dose is 150 mg (may be increased to 300 mg as some patients may derive additional benefit from the higher dose). If weight < 50 kg, recommended dose is 75 mg. However, 150mg solution for injection in pre-filled pen is not indicated for administration of this dose and no suitable alternative formulation is available. **Psoriatic Arthritis:** For patients with concomitant moderate to severe plaque psoriasis see adult plaque psoriasis recommendation. For patients who are anti-TNF α inadequate responders, the recommended dose is 300 mg, 150 mg in other patients. Can be increased to 300 mg based on clinical response. **Ankylosing Spondylitis:** Recommended dose 150 mg. Can be increased to 300 mg based on clinical response. **nr-axSpA:** Recommended dose 150 mg. **Enthesitis-related arthritis and juvenile psoriatic arthritis:** From the age of 6 years, if weight ≥ 50 kg, recommended dose is 150 mg. If weight < 50 kg, recommended dose

Cosentyx® (secukinumab) Great Britain Prescribing Information.

Please refer to the Summary of Product Characteristics (SmPC) before prescribing.

Indications: Treatment of: moderate to severe plaque psoriasis in adults, children and adolescents from the age of 6 years who are candidates for systemic therapy; active psoriatic arthritis in adults (alone or in combination with methotrexate) who have responded inadequately to disease-modifying anti-rheumatic drug therapy; active ankylosing spondylitis in adults who have responded inadequately to conventional therapy; active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation as indicated by elevated C-reactive protein (CRP) and/or magnetic resonance imaging (MRI) evidence in adults who have responded inadequately to non-steroidal anti-inflammatory drugs; active enthesitis-related arthritis and juvenile psoriatic arthritis in patients 6 years and older (alone or in combination with methotrexate) whose disease has responded inadequately to, or who cannot tolerate, conventional therapy; active moderate to severe hidradenitis suppurativa (acne inversa) in adults with an inadequate response to conventional systemic HS therapy. **Presentations:** Cosentyx 75 mg solution for injection in pre-filled syringe; Cosentyx 150 mg solution for injection in pre-filled syringe; Cosentyx 150 mg solution for injection in pre-filled pen; Cosentyx 300 mg solution for injection in pre-filled pen. **Dosage & Administration:** Administered by subcutaneous injection at weeks 0, 1, 2, 3 and 4, followed by monthly maintenance dosing. Consider discontinuation if no response after 16 weeks of treatment. Each 75 mg dose is given as one injection of 75 mg. Each 150 mg dose is given as one injection of 150 mg. Each 300 mg dose is given as two injections of 150 mg or one injection of 300 mg. If possible avoid areas of the skin showing psoriasis. **Plaque Psoriasis:** Adult recommended dose is 300 mg. Based on clinical response, a maintenance dose of 300 mg every 2 weeks may provide additional benefit for patients with a body weight of 90 kg or higher. Adolescents and children from the age of 6 years: if weight ≥ 50 kg, recommended dose is 150 mg (may be increased to 300 mg as some patients may derive additional benefit from the higher dose). If weight < 50 kg, recommended dose is 75 mg. **Psoriatic Arthritis:** For patients with concomitant moderate to severe plaque psoriasis see adult plaque psoriasis recommendation. For patients who are anti-TNF α inadequate responders, the recommended dose is 300 mg, 150 mg in other patients. Can be increased to 300 mg based on clinical response. **Ankylosing Spondylitis:** Recommended dose 150 mg. Can be increased to 300 mg based on clinical response. **nr-axSpA:** Recommended dose 150 mg. **Enthesitis-related arthritis and juvenile psoriatic arthritis:** From the age of 6 years, if weight ≥ 50 kg, recommended dose is 150 mg. If

weight < 50 kg, recommended dose is 75 mg. **Hidradenitis suppurativa:** Recommended dose is 300 mg monthly. Based on clinical response, the maintenance dose can be increased to 300 mg every 2 weeks. **Contraindications:** Hypersensitivity to the active substance or excipients. Clinically important, active infection. **Warnings & Precautions:** **Infections:** Potential to increase risk of infections; serious infections have been observed. Caution in patients with chronic infection or history of recurrent infection. Advise patients to seek medical advice if signs/symptoms of infection occur. Monitor patients with serious infection closely and do not administer Cosentyx until the infection resolves. Non-serious mucocutaneous candida infections were more frequently reported for secukinumab than placebo in the psoriasis clinical studies. Should not be given to patients with active tuberculosis (TB). Consider anti-tuberculosis therapy before starting Cosentyx in patients with latent TB. **Inflammatory bowel disease (including Crohn's disease and ulcerative colitis):** New cases or exacerbations of inflammatory bowel disease have been reported with secukinumab. Secukinumab, is not recommended in patients with inflammatory bowel disease. If a patient develops signs and symptoms of inflammatory bowel disease or experiences an exacerbation of pre-existing inflammatory bowel disease, secukinumab should be discontinued and appropriate medical management should be initiated. **Hypersensitivity reactions:** Rare cases of anaphylactic reactions have been observed. If an anaphylactic or serious allergic reactions occur, discontinue immediately and initiate appropriate therapy. **Vaccinations:** Do not give live vaccines concurrently with Cosentyx; inactivated or non-live vaccinations may be given. Paediatric patients should receive all age appropriate immunisations before treatment with Cosentyx. **Latex-Sensitive Individuals:** The removable needle cap of the 150mg pre-filled pen contains a derivative of natural rubber latex. **Concomitant immunosuppressive therapy:** Combination with immunosuppressants, including biologics, or phototherapy has not been evaluated in psoriasis studies. Cosentyx was given concomitantly with methotrexate, sulfasalazine and/or corticosteroids in arthritis studies. Caution when considering concomitant use of other immunosuppressants. **Interactions:** Live vaccines should not be given concurrently with secukinumab. No interaction between Cosentyx and midazolam (CYP3A4 substrate) seen in adult psoriasis study. No interaction between Cosentyx and methotrexate and/or corticosteroids seen in arthritis studies. **Fertility, pregnancy and lactation: Women of childbearing potential:** Use an effective method of contraception during and for at least 20 weeks after treatment. **Pregnancy:** Preferably avoid use of Cosentyx in pregnancy. **Breast feeding:** It is not known if secukinumab is excreted in human breast milk. A clinical decision should be made on continuation of breast feeding during Cosentyx treatment (and up to 20 weeks after

weight < 50 kg, recommended dose is 75 mg. **Hidradenitis suppurativa:** Recommended dose is 300 mg monthly. Based on clinical response, the maintenance dose can be increased to 300 mg every 2 weeks. **Contraindications:** Hypersensitivity to the active substance or excipients. Clinically important, active infection. **Warnings & Precautions:** **Infections:** Potential to increase risk of infections; serious infections have been observed. Caution in patients with chronic infection or history of recurrent infection. Advise patients to seek medical advice if signs/symptoms of infection occur. Monitor patients with serious infection closely and do not administer Cosentyx until the infection resolves. Non-serious mucocutaneous candida infections were more frequently reported for secukinumab in the psoriasis clinical studies. Should not be given to patients with active tuberculosis (TB). Consider anti-tuberculosis therapy before starting Cosentyx in patients with latent TB. **Inflammatory bowel disease (including Crohn's disease and ulcerative colitis):** New cases or exacerbations of inflammatory bowel disease have been reported with secukinumab. Secukinumab, is not recommended in patients with inflammatory bowel disease. If a patient develops signs and symptoms of inflammatory bowel disease or experiences an exacerbation of pre-existing inflammatory bowel disease, secukinumab should be discontinued and appropriate medical management should be initiated. **Hypersensitivity reactions:** Rare cases of anaphylactic reactions have been observed. If an anaphylactic or serious allergic reactions occur, discontinue immediately and initiate appropriate therapy. **Vaccinations:** Do not give live vaccines concurrently with Cosentyx; inactivated or non-live vaccinations may be given. Paediatric patients should receive all age appropriate immunisations before treatment with Cosentyx. **Latex-Sensitive Individuals:** The removable needle cap of the 75mg and 150 mg pre-filled syringe and 150mg pre-filled pen contains a derivative of natural rubber latex. **Concomitant immunosuppressive therapy:** Combination with immunosuppressants, including biologics, or phototherapy has not been evaluated in psoriasis studies. Cosentyx was given concomitantly with methotrexate, sulfasalazine and/or corticosteroids in arthritis studies. Caution when considering concomitant use of other immunosuppressants. **Interactions:** Live vaccines should not be given concurrently with secukinumab. No interaction between Cosentyx and midazolam (CYP3A4 substrate) seen in adult psoriasis study. No interaction between Cosentyx and methotrexate and/or corticosteroids seen in arthritis studies. **Fertility, pregnancy and lactation: Women of childbearing potential:** Use an effective method of contraception during and for at least 20 weeks after treatment. **Pregnancy:** Preferably avoid use of Cosentyx in pregnancy. **Breast feeding:** It is not known if secukinumab is excreted in human breast milk. A clinical decision should be made on continuation of breast feeding during Cosentyx treatment (and up to 20 weeks after discontinuation) based on benefit of breast feeding to the

child and benefit of breast feeding to the woman. **Fertility:** Effect on human fertility not evaluated. **Adverse Reactions:** *Very Common* ($\geq 1/10$): Upper respiratory tract infection. *Common* ($\geq 1/100$ to $< 1/10$): Oral herpes, headache, rhinorrhoea, diarrhoea, nausea, fatigue. *Uncommon* ($\geq 1/1,000$ to $< 1/100$): Oral candidiasis, lower respiratory tract infections, neutropenia, inflammatory bowel disease. *Rare* ($\geq 1/10,000$ to $< 1/1,000$): anaphylactic reactions, exfoliative dermatitis (psoriasis patients), hypersensitivity vasculitis. *Not known:* Mucosal and cutaneous candidiasis (including oesophageal candidiasis). **Infections:** Most infections were non-serious and mild to moderate upper respiratory tract infections, e.g. nasopharyngitis, and did not necessitate treatment discontinuation. There was an increase in mucosal and cutaneous (including oesophageal) candidiasis, but cases were mild or moderate in severity, non-serious, responsive to standard treatment and did not necessitate treatment discontinuation. Serious infections occurred in a small proportion of patients (0.015 serious infections reported per patient year of follow up). **Neutropenia:** Neutropenia was more frequent with secukinumab than placebo, but most cases were mild, transient and reversible. Rare cases of neutropenia CTCAE Grade 4 were reported. **Hypersensitivity reactions:** Urticaria and rare cases of anaphylactic reactions were seen. **Immunogenicity:** Less than 1% of patients treated with Cosentyx developed antibodies to secukinumab up to 52 weeks of treatment. **Other Adverse Effects:** The list of adverse events is not exhaustive, please consult the SmPC for a detailed listing of all adverse events before prescribing. **Legal Category:** POM. **MA Number & List Price:** EU/1/14/980/005 - 150 mg pre-filled pen x2 £1,218.78; EU/1/14/980/010 - 300 mg pre-filled pen x1 £1,218.78. **PI Last Revised:** May 2023. Full prescribing information, (SmPC) is available from: Novartis Pharmaceuticals UK Limited, 2nd Floor, The WestWorks Building, White City Place, 195 Wood Lane, London, W12 7FQ. Telephone: (01276) 692255.

UK | 284832 | May 2023

Adverse Event Reporting:

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard. Adverse events should also be reported to Novartis via uk.patientsafety@novartis.com or online through the pharmacovigilance intake (PVI) tool at www.novartis.com/report

If you have a question about the product, please contact Medical Information on 01276 698370 or by email at medinfo.uk@novartis.com

child and benefit of Cosentyx therapy to the woman. **Fertility:** Effect on human fertility not evaluated. **Adverse Reactions:** *Very Common* ($\geq 1/10$): Upper respiratory tract infection. *Common* ($\geq 1/100$ to $< 1/10$): Oral herpes, headache, rhinorrhoea, diarrhoea, nausea, fatigue. *Uncommon* ($\geq 1/1,000$ to $< 1/100$): Oral candidiasis, lower respiratory tract infections, neutropenia, inflammatory bowel disease. *Rare* ($\geq 1/10,000$ to $< 1/1,000$): anaphylactic reactions, exfoliative dermatitis (psoriasis patients), hypersensitivity vasculitis. *Not known:* Mucosal and cutaneous candidiasis (including oesophageal candidiasis). **Infections:** Most infections were non-serious and mild to moderate upper respiratory tract infections, e.g. nasopharyngitis, and did not necessitate treatment discontinuation. There was an increase in mucosal and cutaneous (including oesophageal) candidiasis, but cases were mild or moderate in severity, non-serious, responsive to standard treatment and did not necessitate treatment discontinuation. Serious infections occurred in a small proportion of patients (0.015 serious infections reported per patient year of follow up). **Neutropenia:** Neutropenia was more frequent with secukinumab than placebo, but most cases were mild, transient and reversible. Rare cases of neutropenia CTCAE Grade 4 were reported. **Hypersensitivity reactions:** Urticaria and rare cases of anaphylactic reactions were seen. **Immunogenicity:** Less than 1% of patients treated with Cosentyx developed antibodies to secukinumab up to 52 weeks of treatment. **Other Adverse Effects:** The list of adverse events is not exhaustive, please consult the SmPC for a detailed listing of all adverse events before prescribing. **Legal Category:** POM. **MA Number & List Price:** PLGB 00101/1205 - 75 mg pre-filled syringe x 1 - £304.70; PLGB 00101/1029 - 150 mg pre-filled pen x2 £1,218.78; PLGB 00101/1030 - 150 mg pre-filled syringe x2 £1,218.78; PLGB 00101/1198 - 300 mg pre-filled pen x 1 £1,218.78. **PI Last Revised:** June 2023. Full prescribing information, (SmPC) is available from: Novartis Pharmaceuticals UK Limited, 2nd Floor, The WestWorks Building, White City Place, 195 Wood Lane, London, W12 7FQ. Telephone: (01276) 692255.

UK | 290802 | June 2023

Adverse Event Reporting:

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard. Adverse events should also be reported to Novartis via uk.patientsafety@novartis.com or online through the pharmacovigilance intake (PVI) tool at www.novartis.com/report.

If you have a question about the product, please contact Medical Information on 01276 698370 or by email at medinfo.uk@novartis.com