

ACCEPTED MANUSCRIPT • OPEN ACCESS

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X stellarator

To cite this article before publication: Omstavan Samant *et al* 2024 *Nucl. Fusion* in press <https://doi.org/10.1088/1741-4326/ad36d4>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2024 The Author(s). Published by IOP Publishing Ltd on behalf of the IAEA. All rights reserved.



As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 4.0 licence, this Accepted Manuscript is available for reuse under a CC BY 4.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/4.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X stellarator

O Samant¹, R O Dendy¹, S C Chapman^{1,2,3}, D Moseev⁴ and R Ouchoukov⁵

¹Centre for Fusion, Space and Astrophysics, University of Warwick, Coventry, CV47AL, United Kingdom

²Department of Mathematics and Statistics, University of Tromsø, Norway

³ International Space Science Institute, Hallerstrasse 6, 3012 Bern, Switzerland

⁴Max-Planck-Institut für Plasmaphysik, Wendelsteinstr. 1, 17491 Greifswald, Germany

⁵Max-Planck-Institut für Plasmaphysik, Boltzmannstr. 2, 85748 Garching, Germany

E-mail: Omstavan.Samant@warwick.ac.uk / omstavansamant@gmail.com

Abstract.

Measurements of ion cyclotron emission (ICE) are planned for magnetically confined fusion (MCF) plasmas heated by neutral beam injection (NBI) in the Wendelstein 7-X stellarator (W7-X). Freshly injected NBI ions in the edge region, whose velocity-space distribution function approximates a delta-function, are potentially unstable against the magnetoacoustic cyclotron instability (MCI), which could drive a detectable ICE signal. Prediction of ICE from NBI protons in W7-X hydrogen plasmas is challenging, owing to the low ratio of the ions' perpendicular velocity to the local Alfvén speed, $v_{\perp(NBI)}/V_A \simeq 0.14$. We address this from first principles, using the particle-in-cell (PIC) kinetic code EPOCH. This self-consistently solves the Lorentz force equation and Maxwell's equations for tens of millions of computational ions (both thermal majority and energetic NBI minority) and electrons, fully resolving gyromotion and hence capturing the cyclotron resonant phenomenology which gives rise to ICE. Our simulations predict an ICE signal which is predominantly electrostatic while incorporating a significant electromagnetic component. Its frequency power spectrum reflects novel MCI physics, reported here for the first time. The NBI ions relaxing under the MCI first drive broadband field energy at frequencies a little below the lower hybrid frequency ω_{LH} , across the wavenumber range $k\omega_c/V_A = 40$ to 60, where ω_c and V_A denote ion cyclotron frequency and Alfvén velocity. Nonlinear coupling between these waves then excites spectrally structured ICE with narrow peaks, at much lower frequencies, typically the proton cyclotron frequency and its lower harmonics. The relative strength of these peaks depends on the specifics of the NBI ion velocity-space distribution and of the local plasma conditions, implying diagnostic potential for the predicted ICE signal from W7-X.

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X2

1. Introduction

Ion cyclotron emission (ICE) is spontaneously generated strongly suprathermal radiation peaking spectrally at multiple harmonics of the cyclotron frequency of an energetic ion population within a plasma. The narrowband character of the ICE spectral peaks implies that the source is strongly spatially localised. ICE spectra are widely observed in magnetic confinement fusion (MCF) plasmas, and the peak intensities are usually a few orders of magnitude stronger than the thermal radiation.

ICE was early observed from fusion-born ion populations in pure deuterium and DT plasmas in JET[1–4] and TFTR [5, 6], and from neutral beam injected (NBI) ions in TFTR[7]. Since 2017, ICE has been detected and analysed from NBI and fusion-born ions in the KSTAR[8–10], DIII-D[11–16], ASDEX-Upgrade[17–21], NSTX-U[22, 23], JT-60U[24–26], TUMAN-3M[27–29], EAST[30–33] and HL-2A[34, 35] tokamaks, and the LHD heliotron-stellarator[36–41]. The magnetoacoustic cyclotron instability (MCI) has been identified as the excitation mechanism for ICE through analytical studies[6, 7, 42] and simulations[8–10, 12, 38, 39, 43–46]. The MCI can arise at spatial locations where the velocity space distribution of an energetic ion population has a positive slope, $\frac{\partial f}{\partial v_{\perp}} > 0$ for values of $v_{\perp} \gtrsim V_A$, in the velocity space. Under these conditions, waves propagating nearly perpendicular to the magnetic field are excited on the fast Alfvén-cyclotron harmonic branch through wave-particle cyclotron resonance, with the fast ions transferring some of their energy to the excited waves. The spatial location of the population inversion can be inferred by matching the frequency separation between successive ICE spectral peaks to the local cyclotron frequency of the energetic ions, and hence to the local magnetic field strength and the corresponding radial location. The energetic ion species whose free energy has been found to drive ICE through the MCI include fusion born ions[1–3, 5, 8, 10, 14, 24, 25, 47–51], neutral beam injected(NBI) ions[5, 7, 9, 11–14, 19, 24, 38, 48, 49] and ions heated via ion cyclotron resonant heating[52, 53].

In this paper, we focus on the likely prospects, physics basis and spectral character for ICE associated with NBI of energetic ions in future Wendelstein 7-X stellarator plasmas. As noted above, ICE due to NBI ion populations has been previously detected from several tokamaks and the LHD heliotron-stellarator. Comparison between ICE spectra from tokamaks and stellarators sheds light on the relative importance of overall magnetic field structure compared to spatially localised physics. Edge NBI ICE has been simulated in Refs. [9, 12, 38, 39] from first principles, using particle-in-cell (PIC) kinetic codes which solve the Maxwell-Lorentz system of equations self-consistently for tens of millions of gyro-orbit resolved particles. We note that simulations Refs [38, 39] for the LHD stellarator, unlike those presented here, were hybrid simulations with fluid electrons and did not fully resolve electron gyro-dynamics. Here we report the results of PIC simulations using the EPOCH code[54] with fully gyro-resolved thermal ions, energetic ions and electrons. We use a locally uniform (slab geometry) treatment of the plasma physics of ICE emission in these simulations, as in the numerous previous

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X3

PIC simulations for a wide range of MCF plasmas. This local treatment does not incorporate toroidal magnetic field geometry, whether tokamak or stellarator. Its past success in matching observed ICE signals suggests that these spectra are generated by the spatially localised physics that is captured in our code. In particular, the recent success in matching ICE simulations to ICE observations from the LHD heliotron-stellarator [38–40] provides support for adopting here our predictive approach for future ICE physics in NBI-heated plasmas in the W7-X stellarator. These simulations are computationally resource-intensive, partly due to the low ratio of the perpendicular velocity of the NBI ions to the local Alfvén velocity, $V_{NBI}/V_A = 0.14$. The sub-Alfvénic regime of ICE, though uncommon, has been previously observed in TFTR[5], ASDEX-U[19] and LHD[39]. Our simulations capture the full frequency range from ion cyclotron through lower hybrid and beyond, with high resolution. It appears that both the MCI and the lower hybrid drift instability, found in related simulations[55, 56], may operate simultaneously under the Wendelstein 7-X plasma conditions. It appears relevant that turbulence in the lower hybrid frequency range has been observed from NBI-heated plasmas in W7-X [57] and LHD[41]. Also important, and explored here, is the extent to which NBI-driven ICE phenomenology is predominantly electromagnetic or electrostatic. In the stellarator context, we refer again to Refs.[38, 39] on this topic, together with the PIC studies in Refs.[58, 59] which also bear on lower hybrid aspects. The latter papers are in the same spirit as the present work, and differ insofar as in Ref.[58] the higher-frequency excitation is electromagnetic as distinct from electrostatic, and in Ref.[59] the ion cyclotron and lower hybrid frequencies are substantially closer to each other. These two instabilities, which are taken to be well separated in frequency and wavenumber in a linear analytical description, and hence decoupled, appear here as manifestations of a single coupled plasma phenomenon at the level of first principles self-consistent nonlinear kinetics. The development of a predictive capability, in addition to interpretive, for linking the spectral structure of ICE to the velocity-space structure and spatial location of the emitting ion population is important for the diagnostic exploitation of ICE in present and future[60, 61] fusion experiments.

2. Physical and computational approach to W7-X ICE simulations

It is widely accepted that ICE from MCF plasmas is driven by the MCI, whereby a minority non-Maxwellian population of energetic ions excites modes on the fast Alfvén-cyclotron harmonic wave branches, transferring some of its energy also to the kinetic energy of the thermal ions and electrons[42, 62–66]. Here we study the MCI under conditions relevant to NBI populations (freshly ionised, or undergoing prompt loss) in the W7-X edge plasma. This location is the most likely one for excitation of NBI-driven ICE in W7-X, as previously observed in the LHD heliotron-stellarator[38, 49] and the TFTR[5, 7], KSTAR[9], DIII-D[11–14] and ASDEX-Upgrade[21, 48] tokamaks, for example. The reason is that, near the NBI injection point in the edge plasma, the velocity-space distribution of freshly ionised NBI ions, which have not yet slowed

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X4

down through collisions, is closest to a delta-function; and hence most strongly unstable against the MCI which gives rise to the observed ICE signal. We adopt the first principles approach previously used to interpret observations of NBI-driven ICE from other MCF plasmas [9, 12, 38, 39]. This approach involves the use of the particle-in-cell (PIC) code [67, 68] EPOCH [54], which captures the fully gyro-resolved Lorentz dynamics of all particle species, and evolves these dynamics self-consistently with the electric and magnetic fields by solving the full Maxwell's equations. We use EPOCH in 1D3V mode, that is, the simulation domain incorporates one spatial axis, which can be oriented at any angle to the direction of the ambient magnetic field, and all three vector components in velocity-space. It is known that the 1D3V approach, which demands substantial but not unrealistic computational resources, provides a good description of MCI physics for ICE simulation and interpretation [9, 12, 38, 39, 43]. Hence one can be moderately confident in now deploying this capability in predictive mode for W7-X.

EPOCH self consistently evolves the Maxwell equations using a finite-difference time-domain method (FDTD) for all three components of electric and magnetic fields defined on a Yee staggered grid and the gyro orbit resolved charged particle dynamics using a modified version of the Boris algorithm. The charged particles are represented by macroparticles. To reduce noise in the system, we use a fifth order weighing scheme. The electrons and majority thermal ion (protons) populations form the maxwell distributed pseudorandom thermal background. The minority fast ions representing the NBI protons are initialised as a drifting ring-beam distribution, originally modelled with a delta function for the perpendicular component in [42, 65] and later investigated with a spread in the perpendicular component in [69], given by

$$f(v_{\parallel}, v_{\perp}) \propto \exp\left(-\frac{(v_{\parallel} - v_d)^2}{v_r^2}\right) \exp\left(-\frac{(v_{\perp} - u_0)^2}{u_r^2}\right) \quad (1)$$

where v_{\perp} and v_{\parallel} are the velocity components perpendicular and parallel to the background magnetic field, u_0 is initial perpendicular velocity and v_d is initial drift along the background magnetic field, u_r and v_r are the perpendicular and parallel velocity spreads respectively. The system is then allowed to relax.

We ran several simulations using the physical and computational approach described above with one higher (by a factor of 8) resolution simulation used to benchmark the outputs of simulations at lower resolution. For all simulations except the higher resolution simulation, we used 150,000 cells with the time step 0.95 times that required by the CFL condition. The higher resolution simulation used 1,200,000 cells. PIC simulations typically require that the cell size is less than the electron Debye length λ_{D_e} thereby resolving the associated plasma physics. The Nyquist condition in wavenumber space requires that dk , the resolution in wavenumber (k) space, is smaller than ω_c/v_A where ω_c is the ion cyclotron frequency and v_A is the Alfvén velocity. To pre-empt numerical heating of electrons, the cell size must resolve the smallest gyroradius in the system which is the electron gyroradius. The spatial domain should also be long enough to have at least several gyroradii of all species. The data from the simulation

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X5

cannot be saved at every timestep due to memory limits, but must nevertheless be stored at sufficiently high cadence to resolve the physics being explored here. This work requires the resolution in angular frequency, $d\omega$, to be smaller than the ion cyclotron frequencies as the MCI is a cyclotron resonant wave-particle phenomenon, hence the time duration of our simulations must exceed several ion gyroperiods. The simulation Nyquist frequency must exceed the largest frequency present in the system, in our case the electron cyclotron frequency, to avoid aliasing or adding noise in the lower frequencies. The noise is also inversely dependent on the number of macroparticles used to represent the charged particles. Here we use 1,995,000 macro particles for each species. In these simulations, the spatial domain, and hence the allowable wavenumber vectors, was oriented quasiperpendicular (85-89.5 degrees) to the direction of the magnetic field. Other initial parameters are listed in Table 1. In the simulations presented here, we

Parameter	Value
Electron density(n_e)	$6 \times 10^{18} m^{-3}$
Thermal electron and Proton temperature	30eV
Magnetic field	2.25 T
NBI Proton beam Energy	55keV
NBI Proton density	$0.03 \times n_e$
Beam injection Angle w.r.t. B	70°
Initial drift along B(v_d)	$1.11 \times 10^6 m/s$
Initial Perpendicular velocity(u_0)	$3.05 \times 10^6 m/s$
Perpendicular velocity spread(u_r)	$0.001 \times u_0$
Parallel velocity spread(v_r)	$0.01 \times v_d$

Table 1. Physical Parameters used in the simulations

find(see e.g. Fig. 6) that nonlinear three-wave coupling plays a significant role in the flow of energy in the frequency-wavenumber space, and hence in determining the structure of the simulated ICE spectra. This nonlinear coupling is quantified in Sec. 3 using bispectral analysis techniques which are outlined briefly in Appendix A and has previously been used to assist interpretation of ICE simulations in, for example, Fig. 3 of [8].

3. ICE simulation results

Our simulation results are encapsulated in the following six Figures, which we discuss in detail below. To summarise these briefly, Fig. 1 displays the time evolution of the change in energy densities of particles and fields. Figure 2 shows the spatiotemporal Fourier transform of the energy associated with the z-component of the excited magnetic, that is, the distribution of power across different frequencies and wavenumbers. Figure 3 plots simulated ICE power spectra, derived by summing spatiotemporal Fourier transforms over wavenumber, for a range of relevant parameters. We compare the power excited

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X6

in electric versus magnetic field components in Fig.4. The time evolution of energy with respect to wavenumber is shown in Fig.5. Nonlinear coupling of energy across wavenumber space is quantified in the bispectral plots of Fig. 6, for both electric and magnetic fields, at different propagation angles. Figure 1 shows the time evolution of particle and field energy densities from a 1D3V EPOCH PIC simulation of NBI ions relaxing under the MCI for W7-X edge plasma conditions. We emphasise that this reflects spontaneous collective relaxation of millions of simulation particles - NBI ions, thermal ions and electrons - together with the self-consistent electric and magnetic field under the Lorentz-Maxwell system of equations; the MCI is not present in the simulation initial conditions and emerges as the simulation evolves.

MCI involves a collective relaxation of non-Maxwellian ion population, i.e. the beam ions in our simulations. This is evidenced in Fig. 1 which shows the change in energy densities in the field components E_x and B_z and in the different particle populations. In this sub-Alfvénic regime for the NBI ions, it is expected from previous analyses of observations of NBI-driven ICE [7, 9, 38, 39] that the MCI may include a significant electrostatic component. This is seen here in Fig. 1, where the initial rise is at times corresponding to the growth of field energy at $\omega \approx \omega_{LH}$ (cf. Fig.2 and 5). This transitions at $t = 1.25\tau_{gp}$ in Fig.1 (cf. Fig.5, below) to the growth of spectral peaks at lower ion cyclotron harmonics. The system settles into its saturated regime, which presumably corresponds to the state determining the observable ICE spectrum, well before the simulation ends at three ion gyroperiods. In Fig.1, the time traces are flat after two gyroperiods. As we shall see later, when resolved with respect to wavenumber, there is correspondingly little or no time evolution in the upper third of Fig.5 below, covering the time interval between two and three gyroperiods.

Magnetic field data from the simulation is used to calculate the spatiotemporal Fourier transform(2D discrete Fast Fourier Transform), as shown in Fig.2(left) for the case with wave propagation at an angle of 89.5° with respect to the magnetic field B . The B_z component of the magnetic field is chosen for the analysis, because it is perpendicular to the simulation direction and hence the k -vector, thus capturing the electromagnetic physics of the system. Fig. 2(left) shows that the spontaneously excited fields are concentrated in distinct regions of (ω, k) space: spectrally peaked at the second ion cyclotron harmonic; and an extended lower hybrid feature between dimensionless wavenumbers $k = 10$ to 30 , and $k = 50$ to 60 (hereafter the “ELHF”) close to the lower hybrid resonance frequency given by equation (5) in reference [70]

$$\omega^2 = \omega_{LH}^2 \left(1 + \frac{m_i}{m_e} \cos^2 \theta \right). \quad (2)$$

Here m_i , m_e are the ion and electron masses, θ is the angle of propagation and ω_{LH} is the cold plasma lower hybrid frequency given by equation (1) in [70]. Using equation (2) gives 14.96, 25.21, 34.41 and 54.19, normalised to ω_c , for propagation angles $89.5^\circ, 88^\circ, 87^\circ$ and 85° respectively. The ELHFs can be observed around these frequencies also in the spatiotemporal Fourier transforms displayed in Fig. 6 (below),

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X7




Figure 1. Time evolution of change in energy densities of particles, electric field E_x and magnetic field B_z in the 89.5° case. The MCI starts around $t = 1.1\tau_{gp}$, which matches Fig.5 and is predominantly electrostatic, with the energy excited in E_x significantly greater than in B_z .

where they are paired with their corresponding bispectra (discussed below). ELHF might be related to the lower hybrid wave physics studied in [55, 71].

Integrating the squared spatiotemporal FFT of B_z over k -space and multiplying by the constant $\frac{1}{2\mu_0}$ yields the frequency power spectrum, Fig.2(right). Although the exact values of the spectral peak in an experiment can vary depending on various factors such as measuring device, technique, and position of antennae, the relative heights of peaks should follow the same trend in these logarithmic plots. We note that a difference of α in the log scale (sometimes referred to as “prominence”) indicates that the peak has 10^α times the power compared to the base level. Multiple simulations (Fig.3) for quasi-perpendicular propagation at angles between 85° and 89.5° to the magnetic field show that the excitation of spectral peaks at the 2nd and 3rd harmonics is a robust feature. Their amplitude is at least three orders of magnitude above the noise floor, which is plotted in the upper two panels of Fig.3, and these peaks are consistently present (while others are consistently absent) in multiple simulation cases. Comparison with the noise floor (which possesses spectral structure in consequence of the fluctuation dissipation theorem, whereby noise energy is concentrated at normal modes) also shows that the simulated ICE spectrum is strongly suprathermal. This aligns with ICE observations in all experiments, and has been a prime driver of ICE interpretation in terms of collective instability since the earliest measurements, see e.g. Ref.[1]. This is a key prediction of the present work, reinforced by the overall invariance of the phenomenology with respect to the modest variations in electron temperature shown in Fig.3(b) and in velocity-space spread shown in Fig.3(c). The 1D3V EPOCH

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X8




Figure 2. Left: Spatiotemporal Fourier transform of the z-component of the magnetic field (B_z). The logarithmic (base ten) colour scale quantifies the relative distribution of energy across frequency-wavenumber space. Frequency is normalised to the proton cyclotron frequency and wavenumber to the ratio of proton cyclotron frequency to Alfvén velocity. The blue line shows the cold plasma dispersion relation (Eq. (5.49) in [72]). The black line is the lower hybrid resonance frequency from Eq. (2). Right: Corresponding simulated ICE power spectrum obtained by summing, over all k , the power at each frequency in the left panel. The extended lower hybrid feature appears in both panels in the frequency range $11 \lesssim \omega \lesssim 16$.

PIC simulations reported here follow all three vector components of both the electric and the magnetic fields, which are evaluated at each timestep at each grid cell. This enables us to quantify the balance of the simulated ICE spectrum with respect to electrostatic and electromagnetic components, information which is important also from a diagnostic perspective. For example, the field component E_x lies along the spatial domain of the simulation, and is purely electrostatic; whereas the oscillating component of B_z is purely electromagnetic. The frequency power spectrum for the electrostatic component E_x is obtained in the same way as for the electromagnetic component B_z described above, but the multiplicative constant is replaced by $\frac{c_0}{2}$. The two panels of Fig.4 shows examples of these, together with a computation of the ratio of the magnitude of the electrostatic to electromagnetic components of this simulated ICE spectrum. It can be seen that the electrostatic component is dominant across most of the frequency domain, whereas the electromagnetic component is dominant around the 2nd cyclotron harmonic peak. The relative differences in the electric and magnetic field spectra also indicate which probe might be more suitable to detect the signal at different frequency ranges on W7-X. Since the current B-dot probe ICE diagnostic [73] operates primarily inductively, it is unlikely that the predominantly electrostatic high frequency waves which comprise the ELHF will be detected by it. They should however be detectable in the ICRF antenna which will be introduced in the next experimental campaign, OP2.2. The variation in nonlinear energy coupling between the electromagnetic and electrostatic channels is explored further in Fig. 6 using bispectral analysis.

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X9




Figure 3. Variation of simulated ICE power spectra with respect to k vector orientation, electron temperature and NBI ion velocity space distribution. Spectra plotted versus frequency following integration, over wavenumber, of spatiotemporal Fourier transforms of the excited fields. a)(top) Each coloured trace corresponds to a simulation with different quasi-perpendicular orientation of the spatial domain with respect to the magnetic field direction, in the range of 85 to 89.5 degrees. The line labelled thermal background shows the baseline spectrum from a simulation containing no suprathermal ions. b)(middle) each trace corresponds to simulations with slightly different electron temperatures. c)(bottom) shows power spectra from simulations with slightly different spreads in v_d from the ring-beam Eq. (1). The 2nd and 3rd harmonic peaks arise in all cases with a ring-beam, together with harmonics 4, 5 and 6 in some of them. The ELHF spans harmonics 8 to 18 in the 89.5° cases; 20 to 23 for 88°; and is off scale to the right in the 87° and 85° cases. Note that the y-axis range is different in each panel.

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X10




Figure 4. (top) Simulated ICE power spectra of electric field E_x and magnetic field B_z for the 89.5° case, plotted versus frequency following integration, over wavenumber, of spatiotemporal Fourier transforms of the excited fields. (bottom) Frequency dependence of the ratio of the power in B_z to E_x .

The 1D FFTs of magnetic field component B_z in the spatial domain at a given time show the power in different wavenumbers (k) in the system at that instant. Horizontally stacking these k -space spectra taken at successively later times in the simulation yields us Figure 5. The ELHF structure can be observed growing first, before the ICE at lower harmonics is initiated, and then dying out around the time when the lower harmonics gain energy. This suggests the ELHF may drive the ICE in the lower harmonics, with the energy transferred by the nonlinear coupling, which has been tested using bispectral methods as described below.

Fig. 6 shows the dispersion relations and the bicoherence (See Appendix A) of the magnetic field B_z and electric field E_x signals. All the dispersion plots are qualitatively similar with the ELHF located around the lower hybrid frequency and significant power at the lower ICE harmonics. The bicoherence plots have colourmaps ranging from 0 to 1 since bicoherence is a normalised bispectrum, see again in the Appendix. This Figure only informs about the strength of nonlinear coupling between each set of three phase-matched waves, but not the direction of energy transfer among them. The wavenumbers from the bicoherence can be matched to the dispersion plot to find the corresponding frequencies. The change in energy in different wavenumbers from Fig. 5 indicates which modes gain or lose energy. It can be seen that the ELHF is nonlinearly coupled to the lower and middle harmonics for all cases, sometimes indirectly through a third frequency. The bicoherence of the B_z signal is similar, but not identical, to that of E_x .

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X11




Figure 5. Time evolution of 1D spatial FFT of magnetic field component B_z . This shows the variation of the energy distribution across different wavenumbers, k (normalised to the ratio of ion cyclotron frequency to the Alfvén velocity), at different points in time (normalised to the ion gyroperiod). The ELHF, located approximately between wavenumbers 45 and 65, starts around $0.5\tau_{gp}$. This is earlier than the spectral peaks at the 2nd and 3rd harmonics, which rise around $1.25\tau_{gp}$, at which time the ELHF declines.

For example, there is nonlinear wave coupling in a small region of dimensionless k -space, which is visible in the bicoherence of B_z but not in that of E_x , around $(\sim 58, \sim 2)$ for the 89.5° cases, around $(\sim 85, \sim 5)$ for the 88° case, and around $(\sim 115, \sim 5)$ for the 87° case. These differences indicate the importance of self consistently evolving and exploring both electrostatic and electromagnetic components, as here, when simulating ICE phenomenology. Faint diagonal lines which intersect the ELHF are visible in some dispersion plots. These features, seen also in Ref.[58], are a factor of order one thousand weaker than the dominant features of the spatiotemporal Fourier transform, hence are close to the thermal noise floor in these simulations. While it cannot yet be established whether they are numerical or physical in origin, it seems likely that they are an effect, rather than a cause, of the ELHF; they may repay investigation in future studies.

4. Conclusions

We predict the excitation of ICE spectral peaks at the 2nd and 3rd proton cyclotron harmonics in the edge region of hydrogen plasmas heated by proton NBI in the W7-X stellarator. The predominantly electrostatic character that we predict for these ICE peaks suggests that they could be detectable with probes, as previously in TFTR [5, 7], as well as antennas. The lower cyclotron harmonics in our simulation may be driven by nonlinear interactions within the ELHF structure, which is excited at frequencies

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X12




Figure 6. Each row of plots corresponds to the case for the propagation angle stated on the left-hand margin. The first column shows the spatiotemporal FFTs of B_z with the logarithmic colour scale indicating the relative distribution of energies in the (ω, k) combination calculated in similar fashion to that in Fig. 2. The ELHF appears in each case around the LH resonance frequency indicated by the black line. The second and third columns show the bicoherence of the magnetic field B_z and electric field E_x respectively. The colour scale ranges from 0 to 1, with values closer to 1 indicating stronger nonlinear coupling between the two wavenumbers. The first two rows correspond to the same case, but top row data was obtained from the higher resolution simulation used in Fig. 2.

Predicting ion cyclotron emission from neutral beam heated plasmas in Wendelstein7-X13

closer to the lower hybrid frequency. The excitation of the ELHF, involving radiation at frequencies comparable to the lower hybrid frequency, is our second main prediction. Based on the change in power in the wavenumbers with time and the nonlinear coupling revealed by the bispectral analysis of the fields excited in our simulations, cross-scale energy transfer is expected between the ELHF and the lower frequency ion cyclotron harmonic peaks characteristic of ICE. Whilst these predictions are specialised to the W7-X NBI edge plasma scenario specified, we note that the nonlinear coupling between lower hybrid waves, identified here, could also arise in other MCF plasmas irrespective of their overall confining magnetic field geometry, and indeed in solar-terrestrial and astrophysical plasmas where ICE-type phenomena have been observed[74–77] or are anticipated[78, 79]. In summary, the phenomenology predicted by these simulations is not narrowly concentrated in frequency and wavenumber. Instead, energy flows initially into wavenumbers corresponding to frequencies near the lower hybrid frequency, the broadband ELHF. It appears from Figs.5 and 6 that, probably, energy then flows by nonlinear self-interactions into: wavenumbers corresponding to the intermediate frequency range, which would also be expected to be excited by standard MCI and ICE physics; and to the lowest ion cyclotron harmonic peaks.

Acknowledgements

Calculations were performed using the Sulis Tier 2 HPC platform hosted by the Scientific Computing Research Technology Platform at the University of Warwick. Sulis is funded by EPSRC Grant EP/T022108/1 and the HPC Midlands+ consortium.

This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 — EUROfusion). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

SCC acknowledges support via the Johannes Geiss Fellowship of the ISSI, and AFOSR grant FA8655-22-1-7056

Appendix A

We use bispectral analysis to identify nonlinear phase coupling, which is indicative of energy being transferred between waves. In particular, we identify where three waves interact nonlinearly while satisfying the wavenumber and (angular) frequency matching conditions $k_3 = k_1 + k_2$ and $\omega_3 = \omega_1 + \omega_2$; i.e. where waves “1” and “2” interact to produce “3”. We use a Fourier transform based bispectrum b_s , defined as[80]

$$b_s^2(k_1, k_2) = |B(k_1, k_2)|^2 = |\langle F(k_1)F(k_2)F^*(k_1 + k_2) \rangle|^2 \quad (3)$$

where $F(k_1)$ is the complex Fourier transform of the quantity being analysed, in our case the electric field E_x or magnetic field B_z . The angled brackets $\langle \cdot \rangle$ can represent

REFERENCES

14

averaging over time or ensemble. In our case, we implement ensemble averaging by creating multiple records of the signal by using a sliding window of the signal with respect to time. Computing this for the entire space is expensive, in terms of computation time and memory. Therefore, we calculate the bicoherence around the regions with maximum power in the spatiotemporal Fourier transform. The bispectrum can be normalised in a number of ways[81, 82] to obtain the bicoherence, b_c , and we choose to do so using Schwartz's inequality:

$$b_c^2(k_1, k_2) = \frac{|\langle F(k_1)F(k_2)F^*(k_1 + k_2) \rangle|^2}{\langle |F(k_1)F(k_2)|^2 \rangle \langle |F^*(k_1 + k_2)|^2 \rangle} \quad (4)$$

such that the bicoherence is bounded by $0 \leq b_c \leq 1$. A value closer to 1 indicates there is coupling between the two frequencies and their sum while a value close to 0 indicates its absence. For example, if there is an intense region around (a, b) then waves with wavenumbers a, b and $a + b$ are coupled. The value of b_c^2 measures the fraction of power at $a + b$ that is coupled in the three wave coupling. The preceding definitions are in terms of wavenumber(k) which is relevant to our case, but the same definitions and relations hold in (angular) frequency space and k can be substituted by ω . We note that Eq. (4) posses the following symmetries: $B(k_1, k_2) = B(k_2, k_1) = B^*(-k_1, -k_2)$ and $B(k_1, k_2) = B(-k_1 - k_2, k_2) = B(k_1, -k_1 - k_2)$ [80]. These symmetries apply equally to the bicoherence in three wave interactions[83].

References

- [1] G A Cottrell and R O Dendy. "Superthermal radiation from fusion products in JET". *Physical Review Letters* 60.1 (1988), p. 33.
- [2] G A Cottrell et al. "Ion cyclotron emission measurements during JET deuterium-tritium experiments". *Nuclear Fusion* 33.9 (1993), p. 1365.
- [3] P Schild, G A Cottrell, and R O Dendy. "Sawtooth oscillations in ion cyclotron emission from JET". *Nuclear Fusion* 29.5 (1989), p. 834.
- [4] K G McClements et al. "Ion cyclotron emission from JET DT plasmas". *Physical Review Letters* 82.10 (1999), p. 2099.
- [5] S Cauffman et al. "Alfvénic behaviour of alpha particle driven ion cyclotron emission in TFTR". *Nuclear fusion* 35.12 (1995), p. 1597.
- [6] R O Dendy et al. "Ion cyclotron emission due to collective instability of fusion products and beam ions in TFTR and JET". *Nuclear Fusion* 35.12 (1995), p. 1733.
- [7] R O Dendy et al. "A mechanism for beam-driven excitation of ion cyclotron harmonic waves in the Tokamak Fusion Test Reactor". *Physics of Plasmas* 1.10 (1994), pp. 3407–3413.
- [8] Benjamin Chapman et al. "Nonlinear wave interactions generate high-harmonic cyclotron emission from fusion-born protons during a KSTAR ELM crash". *Nuclear Fusion* 58.9 (2018), p. 096027.

REFERENCES

15

- [9] B Chapman et al. "Interpretation of suprathreshold emission at deuteron cyclotron harmonics from deuterium plasmas heated by neutral beam injection in the KSTAR tokamak". *Nuclear Fusion* 59.10 (2019), p. 106021.
- [10] B Chapman et al. "Sub-microsecond temporal evolution of edge density during edge localized modes in KSTAR tokamak plasmas inferred from ion cyclotron emission". *Nuclear Fusion* 57.12 (2017), p. 124004.
- [11] Kathreen E Thome et al. "Central ion cyclotron emission in the DIII-D tokamak". *Nuclear Fusion* 59.8 (2019), p. 086011.
- [12] A I Zalzali et al. "Simulations of ion cyclotron emission from DIII-D tokamak plasmas". *47th EPS Conference on Plasma Physics* P4.1002 (2021).
- [13] G H DeGrandchamp et al. "Mode structure measurements of ion cyclotron emission and sub-cyclotron modes on DIII-D". *Nuclear Fusion* 62.10 (2022), p. 106033.
- [14] K E Thome et al. "Radio frequency measurements of energetic-particle-driven emission using the ion cyclotron emission diagnostic on the DIII-D tokamak". *Review of Scientific Instruments* 89.10 (2018), p. 10I102.
- [15] NA Crocker et al. "Novel internal measurements of ion cyclotron frequency range fast-ion driven modes". *Nuclear Fusion* 62.2 (2022), p. 026023.
- [16] GH DeGrandchamp et al. "Upgrades to the ion cyclotron emission diagnostic on the DIII-D tokamak". *Review of Scientific Instruments* 92.3 (2021).
- [17] Lunan Liu et al. "Explanation of core ion cyclotron emission from beam-ion heated plasmas in ASDEX Upgrade by the magnetoacoustic cyclotron instability". *Nuclear Fusion* 61.2 (2020), p. 026004.
- [18] R Ochoukov et al. "Core plasma ion cyclotron emission driven by fusion-born ions". *Nuclear Fusion* 59.1 (2018), p. 014001.
- [19] R Ochoukov et al. "Interpretation of core ion cyclotron emission driven by sub-Alfvénic beam-injected ions via magnetoacoustic cyclotron instability". *Nuclear Fusion* 59.8 (2019), p. 086032.
- [20] B Chapman et al. "Origin of ion cyclotron emission at the proton cyclotron frequency from the core of deuterium plasmas in the ASDEX-Upgrade tokamak". *Plasma Physics and Controlled Fusion* 62.9 (2020), p. 095022.
- [21] R Ochoukov et al. "Observations of core ion cyclotron emission on ASDEX Upgrade tokamak". *Review of Scientific Instruments* 89.10 (2018), 10J101.
- [22] Eric D Fredrickson et al. "Emission in the ion cyclotron range of frequencies (ICE) on NSTX and NSTX-U". *Physics of Plasmas* 26.3 (2019).
- [23] ED Fredrickson et al. "Chirping ion cyclotron emission (ICE) on NSTX-U". *Nuclear Fusion* 61.8 (2021), p. 086007.
- [24] M Ichimura et al. "Observation of spontaneously excited waves in the ion cyclotron frequency range on JT-60U". *Nuclear fusion* 48.3 (2008), p. 035012.

REFERENCES

16

- [25] Shoichi Sato et al. “Observation of ion cyclotron emission owing to DD fusion product H ions in JT-60U”. *Plasma and Fusion Research* 5 (2010), S2067–S2067.
- [26] Shuhei Sumida et al. “Characteristics of fast 3He ion velocity distribution exciting ion cyclotron emission on JT-60U”. *Plasma Physics and Controlled Fusion* 61.2 (2019), p. 025014.
- [27] L G Askinazi et al. “Ion cyclotron emission in NBI-heated plasmas in the TUMAN-3M tokamak”. *Nuclear Fusion* 58.8 (2018), p. 082003.
- [28] LG Askinazi et al. “The Spectrum of Ion Cyclotron Emission from Neutral Beam Injection Heated Plasma on the TUMAN-3M Tokamak”. *Technical Physics Letters* 44 (2018), pp. 1020–1023.
- [29] LG Askinazi et al. “Experimental Determination of the Dispersion Relation of Plasma Radiation at Frequencies of the Ion Cyclotron Resonance and Its Harmonics in Tokamak”. *Technical Physics Letters* 47.3 (2021), pp. 214–217.
- [30] L N Liu et al. “Ion cyclotron emission diagnostic system on the experimental advanced superconducting tokamak and first detection of energetic-particle-driven radiation”. *Review of Scientific Instruments* 90.6 (2019), p. 063504.
- [31] Lunan Liu et al. “Ion cyclotron emission driven by deuterium neutral beam injection and core fusion reaction ions in EAST”. *Nuclear Fusion* 60.4 (2020), p. 044002.
- [32] Lunan Liu et al. “Interpretation of ion cyclotron emission from sub-Alfvénic beam-injected ions heated plasmas soon after LH mode transition in EAST”. *Plasma Physics and Controlled Fusion* 66.1 (2023), p. 015007.
- [33] Lunan Liu et al. “High-frequency B-dot probes used to detect characteristics of ion cyclotron range of frequency waves in EAST”. *Journal of Plasma Physics* 85.2 (2019), p. 905850214.
- [34] Linzi Liu et al. “Identification of core ion cyclotron instabilities on HL-2A tokamak”. *Nuclear Fusion* 63.10 (2023), p. 104004.
- [35] RH Tong et al. “Development of the ion cyclotron emission diagnostic on the HL-2A tokamak”. *Journal of Instrumentation* 17.01 (2022), p. C01063.
- [36] Kenji Saito et al. “Measurement of ion cyclotron emissions by use of ICRF heating antennas in LHD”. *Fusion Engineering and Design* 84.7-11 (2009), pp. 1676–1679.
- [37] Kenji Saito et al. “Measurement of ion cyclotron emissions by using high-frequency magnetic probes in the LHD”. *Plasma Science and Technology* 15.3 (2013), p. 209.
- [38] B C G Reman et al. “Interpreting observations of ion cyclotron emission from large helical device plasmas with beam-injected ion populations”. *Nuclear Fusion* 59.9 (2019), p. 096013.
- [39] B C G Reman et al. “Density dependence of ion cyclotron emission from deuterium plasmas in the large helical device”. *Nuclear Fusion* 61.6 (2021), p. 066023.

REFERENCES

17

- [40] Bernard CG Reman et al. “First observation and interpretation of spontaneous collective radiation from fusion-born ions in a stellarator plasma”. *Plasma Physics and Controlled Fusion* 64.8 (2022), p. 085008.
- [41] Kenji Saito et al. “RF wave detection with high-frequency magnetic probes in LHD”. *Plasma and Fusion Research* 13 (2018), pp. 3402043–3402043.
- [42] R O Dendy et al. “The excitation of obliquely propagating fast Alfvén waves at fusion ion cyclotron harmonics”. *Physics of Plasmas* 1.6 (1994), pp. 1918–1928.
- [43] B Chapman et al. “Comparing theory and simulation of ion cyclotron emission from energetic ion populations with spherical shell and ring-beam distributions in velocity-space”. *Plasma Physics and Controlled Fusion* 62.5 (2020), p. 055003.
- [44] Kenneth G McClements et al. “Observations and modelling of ion cyclotron emission observed in JET plasmas using a sub-harmonic arc detection system during ion cyclotron resonance heating”. *Nuclear Fusion* 58.9 (2018), p. 096020.
- [45] J W S Cook, R O Dendy, and S C Chapman. “Particle-in-cell simulations of the magnetoacoustic cyclotron instability of fusion-born alpha-particles in tokamak plasmas”. *Plasma Physics and Controlled Fusion* 55.6 (2013), p. 065003.
- [46] L Carbajal et al. “Linear and nonlinear physics of the magnetoacoustic cyclotron instability of fusion-born ions in relation to ion cyclotron emission”. *Physics of Plasmas* 21.1 (2014), p. 012106.
- [47] M Ichimura et al. “Study of ion cyclotron emissions due to DD fusion product ions on JT-60U”. *22nd IAEA Fusion Energy Conf.(Geneva)*. 2008, EX–P8.
- [48] Rodolphe D’Inca. “Ion cyclotron emission on ASDEX Upgrade”. PhD thesis. Ludwig Maximilian University, 2014.
- [49] Shekar G Thatipamula et al. “Dynamic spectra of radio frequency bursts associated with edge-localized modes”. *Plasma Physics and Controlled Fusion* 58.6 (2016), p. 065003.
- [50] M H Kim et al. “Distinct stages of radio frequency emission at the onset of pedestal collapse in KSTAR H-mode plasmas”. *Nuclear Fusion* 58.9 (2018), p. 096034.
- [51] M H Kim et al. “Intense whistler-frequency emissions at the pedestal collapse in KSTAR H-mode plasmas”. *Nuclear Fusion* 60.12 (2020), p. 126021.
- [52] G A Cottrell. “Identification of minority ion-cyclotron emission during radio frequency heating in the JET tokamak”. *Physical Review Letters* 84.11 (2000), p. 2397.
- [53] P Jacquet et al. “Parasitic signals in the receiving band of the Sub-Harmonic Arc Detection system on JET ICRF Antennas”. *AIP Conference Proceedings*. Vol. 1406. 1. American Institute of Physics. 2011, pp. 17–20.
- [54] T D Arber et al. “Contemporary particle-in-cell approach to laser-plasma modelling”. *Plasma Physics and Controlled Fusion* 57.11 (2015), p. 113001.

REFERENCES

18

- [55] J W S Cook, S C Chapman, and R O Dendy. “Electron current drive by fusion-product-excited lower hybrid drift instability”. *Physical Review Letters* 105.25 (2010), p. 255003.
- [56] J W S Cook et al. “Self-consistent kinetic simulations of lower hybrid drift instability resulting in electron current driven by fusion products in tokamak plasmas”. *Plasma Physics and Controlled Fusion* 53.6 (2011), p. 065006.
- [57] AG Shalashov et al. “NBI-driven ion cyclotron instabilities at the W7-AS stellarator”. *Plasma physics and controlled fusion* 45.4 (2003), p. 395.
- [58] Mieko Toida et al. “Simulation Study of High-Frequency Magnetosonic Waves Excited by Energetic Ions in Association with Ion Cyclotron Emission”. *Plasma and Fusion Research* 13 (2018), pp. 3403015–3403015.
- [59] Mieko Toida et al. “Simulation study of energetic ion driven instabilities near the lower hybrid resonance frequency in a plasma with increasing density”. *Plasma and Fusion Research* 14 (2019), pp. 3401112–3401112.
- [60] RO Dendy and KG McClements. “Ion cyclotron emission from fusion-born ions in large tokamak plasmas: a brief review from JET and TFTR to ITER”. *Plasma Physics and controlled fusion* 57.4 (2015), p. 044002.
- [61] NN Gorelenkov. “Ion cyclotron emission studies: Retrospects and prospects”. *Plasma Physics Reports* 42.5 (2016), pp. 430–439.
- [62] VS Belikov and Ya I Kolesnichenko. “Magnetoacoustic cyclotron instability in a thermonuclear plasma”. *Sov. Phys.-Tech. Phys.(Engl. Transl.);(United States)* 20.9 (1975).
- [63] RO Dendy, Chris N Lashmore-Davies, and KF Kam. “A possible excitation mechanism for observed superthermal ion cyclotron emission from tokamak plasmas”. *Physics of Fluids B: Plasma Physics* 4.12 (1992), pp. 3996–4006.
- [64] RO Dendy, CN Lashmore-Davies, and KF Kam. “The magnetoacoustic cyclotron instability of an extended shell distribution of energetic ions”. *Physics of Fluids B: Plasma Physics* 5.7 (1993), pp. 1937–1944.
- [65] Kenneth George McClements et al. “Interpretation of ion cyclotron emission from sub-Alfvénic fusion products in the Tokamak Fusion Test Reactor”. *Physics of Plasmas* 3.2 (1996), pp. 543–553.
- [66] Tünde Fülöp and Mietek Lisak. “Ion cyclotron emission from fusion products and beam ions in the tokamak fusion test reactor”. *Nuclear fusion* 38.5 (1998), p. 761.
- [67] CK Birdsall and AB Langdon. “Plasma physics via computer simulation, Inst. of Phys”. *Publishing, Bristol/Philadelphia* 32 (1991).
- [68] Roger W Hockney and James W Eastwood. *Computer simulation using particles*. crc Press, 2021.
- [69] J W S Cook. “Doublet splitting of fusion alpha particle driven ion cyclotron emission”. *Plasma Physics and Controlled Fusion* 64.11 (2022), p. 115002.

REFERENCES

19

- [70] A L Verdon et al. “Warm electromagnetic lower hybrid wave dispersion relation”. *Physics of Plasmas* 16.5 (2009), p. 052105.
- [71] J W S Cook, R O Dendy, and S C Chapman. “Gyrobunching and wave–particle resonance in the lower hybrid drift instability”. *Plasma Physics and Controlled Fusion* 53.7 (2011), p. 074019.
- [72] Richard Fitzpatrick. *Plasma physics: an introduction*. Crc Press, 2022.
- [73] D Moseev et al. “Development of the ion cyclotron emission diagnostic for the W7-X stellarator”. *Review of Scientific Instruments* 92.3 (2021).
- [74] KG McClements and RO Dendy. “Ion cyclotron harmonic wave generation by ring protons in space plasmas”. *Journal of Geophysical Research: Space Physics* 98.A7 (1993), pp. 11689–11700.
- [75] RO Dendy and KG McClements. “Ion cyclotron wave emission at the quasi-perpendicular bow shock”. *Journal of Geophysical Research: Space Physics* 98.A9 (1993), pp. 15531–15539.
- [76] Kenneth George McClements, RO Dendy, and CN Lashmore-Davies. “A model for the generation of obliquely propagating ULF waves near the magnetic equator”. *Journal of Geophysical Research: Space Physics* 99.A12 (1994), pp. 23685–23693.
- [77] JL Posch et al. “Low-harmonic magnetosonic waves observed by the Van Allen Probes”. *Journal of Geophysical Research: Space Physics* 120.8 (2015), pp. 6230–6257.
- [78] VL Rekaa, Sandra C Chapman, and RO Dendy. “Ion pre-acceleration in fully self-consistent particle-in-cell simulations of supercritical perpendicular reforming shocks in multiple ion species plasmas”. *The Astrophysical Journal* 791.1 (2014), p. 26.
- [79] KG McClements et al. “Acceleration of cosmic ray electrons by ion-excited waves at quasiperpendicular shocks”. *Monthly Notices of the Royal Astronomical Society* 291.1 (1997), pp. 241–249.
- [80] Young C Kim and Edward J Powers. “Digital bispectral analysis and its applications to nonlinear wave interactions”. *IEEE Transactions on Plasma Science* 7.2 (1979), pp. 120–131.
- [81] T Dudok de Wit and Berlin Springer. “Spectral and statistical analysis of plasma turbulence: beyond linear techniques”. *Space Plasma Simulation* (2003), pp. 315–343.
- [82] V Kravtchenko-Berejnoi et al. “On the use of tricoherent analysis to detect nonlinear wave-wave interactions”. *Signal Processing* 42.3 (1995), pp. 291–309.
- [83] Ronald Davidson. *Methods in nonlinear plasma theory*. Elsevier, 2012.