



LJMU Research Online

Clarkson, GJ and Roesner, S

Synthesis of Benzofuopyridines and Dibenzofurans by a Metalation/Negishi Cross-Coupling/SNAr Reaction Sequence

<http://researchonline.ljmu.ac.uk/id/eprint/23110/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Clarkson, GJ and Roesner, S (2022) Synthesis of Benzofuopyridines and Dibenzofurans by a Metalation/Negishi Cross-Coupling/SNAr Reaction Sequence. The Journal of Organic Chemistry, 88 (1). pp. 684-689. ISSN 0022-3263

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Synthesis of Benzofuopyridines and Dibenzofurans by a Metalation/Negishi Cross-Coupling/ S_NAr Reaction Sequence

Guy J. Clarkson and Stefan Roesner*



Cite This: *J. Org. Chem.* 2023, 88, 684–689



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: An efficient methodology for the synthesis of benzofuopyridines and dibenzofurans from fluoropyridines or fluoroarenes and 2-bromophenyl acetates is reported. This streamlined one-pot procedure consists of a four-step directed *ortho*-lithiation, zincation, Negishi cross-coupling, and intramolecular nucleophilic aromatic substitution, allowing for the facile assembly of a diverse set of fused benzofuro heterocycles.



Nitrogen heterocycles are among the most significant structural motifs of pharmaceuticals with more than half of all FDA approved small-molecule drugs containing at least one N-heterocycle.¹ Among them, benzofuopyridines are tricyclic compounds containing an annulated pyridine, furan, and benzene ring. However, compared to the synthesis and biological evaluation of dibenzofurans,^{2,3} benzofuopyridines have been far less explored. This is surprising as this class of compounds shows diverse biological activity and interesting properties for potential applications in materials science. Examples include elblfluorene (I) and its derivatives that possess high activity as cyclin-dependent kinase (CDK) inhibitors⁴ and benzofuro[2,3-*c*]pyridine II with potential applications as an MDR modulator (Figure 1).⁵ In addition to




Figure 1. Benzofuopyridines and benzothienopyridines with interesting biological properties and applications in materials science.

their diverse biological activity, benzofuro- and benzothienopyridine derivatives possess interesting fluorescence properties suitable to be applied as green or blue OLED emitters.⁶ Moreover, dyes with a benzothieno[2,3-*c*]pyridine (III) anchoring group have received considerable attention in the development of dye-sensitized solar cells.⁷

Several synthetic strategies toward the preparation of benzofuopyridines have been reported. Intramolecular C–C bond formation in diaryl ethers^{8,9} and C–O bond formation in 2-biaryl phenols by intramolecular nucleophilic aromatic substitution (S_NAr)^{8,10} are the most common strategies. The

former approach was successfully applied by Yue and Li, who synthesized all four benzofuopyridine regioisomers.⁸ Here, a palladium catalyzed Stille coupling using toxic organotin reagents was a key step. Liu et al. reported a more environmental benign strategy generating biaryl phenols from dihalopyridines and 2-hydroxyphenylboronic acids via regioselective Suzuki cross-coupling followed by copper catalyzed intramolecular cyclization.^{9b} Alternative strategies for the synthesis of benzofuopyridines include the construction of the pyridine ring from benzofuran derivatives¹¹ or cascade reactions that generate two annulated rings in the same synthetic operation.¹² However, these approaches often rely on elaborated substrates. Consequently, the development of a concise and general synthesis of benzofuopyridines and its derivatives from readily available starting materials is still of considerable interest.


Recently, we reported an efficient synthesis of tricyclic carbolines employing a four-step *ortho*-lithiation/zincation/Negishi cross-coupling/ S_NAr reaction sequence (Scheme 1a).¹³ In this methodology a fluorine substituent serves both as directing group for the metalation¹⁴ as well as facile leaving group in the intramolecular cyclization. As a continuation of our efforts to develop new synthetic procedures for the construction of heterocyclic frameworks, we were curious if we could extend this procedure by using phenols and thiophenols as a route to tricyclic systems (Scheme 1b). Herein, we describe the first one-pot synthesis of benzofuro[2,3-*b*]- and benzofuro[2,3-*c*]pyridines 5 via a telescoped metalation/cross-coupling/ S_NAr reaction sequence from commercially available

Received: September 1, 2022

Published: December 9, 2022



Scheme 1. (a) Previously Reported Synthesis of Carbolines by a Lithiation/Zincation/Negishi Cross-Coupling/ S_NAr Reaction Sequence and (b) Extension of This Methodology to Benzofuro- and Benzothienopyridines as Well as Dibenzofurans



fluoropyridines and readily accessible 2-bromophenyl acetates.¹⁵ This strategy was expanded to benzothieno[2,3-*b*]pyridines. In addition, directed lithiation of fluoroarenes provided facile access to a range of functionalized dibenzofurans 6.

We initiated our studies by optimizing the reaction conditions for the one-pot Negishi cross-coupling/intramolecular S_NAr reaction sequence (Table 1). Using our previously reported conditions for the formation of organozinc species 2,¹³ 2-fluoropyridine (1a), and 2-bromophenyl acetate (3a) were converted to benzofuropyridine 5a in 96% yield for the four-step sequence using 2 mol % of an XPhos-based palladium precatalyst¹⁶ with additional 2 mol % XPhos ligand¹⁷ and 2.0 equiv of KOtBu after heating at 70 °C overnight (entry 1). Reducing either the catalyst loading to 1 mol % or omitting the additional XPhos ligand decreased the


yield of 5a (entry 2–3). Protection of the phenolic oxygen was essential as no product could be detected with 2-bromophenol, even in the presence of excess base (entry 4). We hypothesize that the free phenol is able to protonate 2, while the corresponding phenolate does not participate in the cross-coupling reaction. This is in stark contrast to 2-bromoanilines, which do not require protection under similar reaction conditions.¹³ Other phenolic protecting groups gave 5a either in lower yields or afforded the corresponding biaryls 4.¹⁸ Employing alternative palladium sources in the presence of 2 mol % XPhos ligand highlights the superior activity of the palladium precatalyst under the applied cross-coupling conditions (entries 5–7). It is worth noting that both the palladium catalyst and the XPhos ligand were essential for the reaction to take place (entries 8–9). In the absence of additional base, the cross-coupling was complete in 20 min as demonstrated by the isolation of biaryl acetate 4a in 84% yield (entry 10). Reducing either the amount of base or the reaction time led to incomplete conversion of the intermediate biaryl 4a (entries 11–13). Finally, while screening other bases to facilitate the deprotection of the acyl group and to promote the S_NAr reaction, we found NaHMDS as valuable alternative to KOtBu (entry 14), whereas Cs₂CO₃ did not fully deprotect the phenolic alcohol under the applied reaction conditions (entry 15).

The results from the optimization studies reveal the reaction order in the formation of 5a (Scheme 2). After Negishi cross-coupling between organozinc intermediate 2a and 2-bromophenyl acetate (3a) to form biaryl acetate 4a, deprotection of the acyl group occurs to provide probably biaryl phenol 7a, which was not observed in the optimization studies, indicating that it must undergo rapid intramolecular

Table 1. Optimization of the Reaction Conditions^a


entry	R	catalyst	base	time	yield 5a ^b (%)	yield 4a ^b (%)
1	Ac	Pd XPhos G3	KOtBu	o/n	96 ^c	–
2 ^d	Ac	Pd XPhos G3	KOtBu	o/n	81	–
3 ^e	Ac	Pd XPhos G3	KOtBu	o/n	74 ^c	–
4 ^f	H	Pd XPhos G3	KOtBu	o/n	0	–
5	Ac	Pd(PPh ₃) ₄	KOtBu	o/n	50	–
6	Ac	Pd(OAc) ₂	KOtBu	o/n	70	–
7 ^d	Ac	[PdCl(C ₃ H ₅) ₂] ₂	KOtBu	o/n	84	–
8	Ac	–	KOtBu	o/n	0	–
9 ^e	Ac	Pd(OAc) ₂	KOtBu	o/n	11	–
10	Ac	Pd XPhos G3	–	20 min	–	84 ^c
11	Ac	Pd XPhos G3	–	o/n	51	31
12 ^g	Ac	Pd XPhos G3	KOtBu	o/n	52	16
13	Ac	Pd XPhos G3	KOtBu	20 min	11	28
14	Ac	Pd XPhos G3	NaHMDS	o/n	86	–
15	Ac	Pd XPhos G3	Cs ₂ CO ₃	o/n	22	30

^a0.5 mmol scale; reaction conditions: (1) 1a (1.2 equiv), LDA (1.3 equiv), THF (0.25 M), 25 °C, 5 min; (2) ZnCl₂ (1.3 equiv), then –25 °C to rt; (3) 2-bromophenol derivative 3 (1.0 equiv), catalyst (2.0 mol %), and XPhos (2.0 mol %) in THF (0.5 M), base (2.0 equiv), 70 °C. ^bDetermined by ¹H NMR spectroscopy using 1,3,5-trimethoxybenzene as internal standard. ^cIsolated yield after column chromatography. ^d1 mol % catalyst loading. ^eNo additional XPhos ligand. ^f2.3 equiv of LDA used. ^g1.0 equiv of base.

Scheme 2. Reaction Sequence and Intermediates in the Formation of Benzofuropyridine 5a


cyclization under the basic conditions to generate the C–O bond and tricyclic **5a** via S_NAr .

With optimized reaction conditions in hand, we next examined the generality of the methodology for a variety of fluoropyridines (Table 2). With 2-fluoro-5-methylpyridine, 2,6- and 2,4-difluoropyridine, the corresponding benzofuro-

Table 2. Scope of Benzofuropyridines^a



^a0.5 mmol scale; reaction conditions: (1) **1** (1.2 equiv), LDA (1.3 equiv), THF (0.25 M), $-25\text{ }^\circ\text{C}$, 5 min; (2) ZnCl_2 (1.3 equiv), then $-25\text{ }^\circ\text{C}$ to rt; (3) **3** (1.0 equiv), Pd XPhos G3 (2.0 mol %), and XPhos (2.0 mol %) in THF (0.5 M), then KOtBu (2.0 equiv), $70\text{ }^\circ\text{C}$, o/n. ^bReaction conditions as shown for **a**, no KOtBu, $70\text{ }^\circ\text{C}$, 20 min, then solvent exchange to DMF (0.1 M), Cs_2CO_3 (5 equiv), $100\text{ }^\circ\text{C}$, 2 h. ^cReaction conditions as shown for **a**, isolation of **7**, then DMF (0.1 M), Cs_2CO_3 (5 equiv), $100\text{ }^\circ\text{C}$, 2 h. ^dFrom the diacetate of bromohydroquinone. ^e2.5 mmol scale. ^fX-ray structure of **5q** shown with 50% probability ellipsoids.

[2,3-*b*]pyridines **5b–d** were obtained as single regioisomers in 63–88% yield. However, for 3-fluoropyridine, 2,3- and 2,5-difluoropyridine no conversion to the corresponding benzofuro[2,3-*c*]pyridines was observed providing biaryl phenols instead. In other words, nucleophilic substitution of the less activated fluorine in 3-position of the pyridyl ring was not achieved. To promote the intramolecular S_NAr reaction, we conducted a solvent exchange from THF to DMF after the cross-coupling step and added additional base (Cs_2CO_3).^{10b} After heating to $100\text{ }^\circ\text{C}$ for 2 h, **5e–g** could be isolated in 39–88% yield. For **5g** we achieved better results conducting the nucleophilic substitution from the purified biaryl phenol.

We next explored the substrate scope of a variety of 2-bromophenyl acetates **3** (Table 2). Substitution in all positions of **3** with a number of electron-withdrawing and -donating groups was tolerated providing dibenzofuro[2,3-*b*]pyridines **5h–q** in 45–92% yield. The structure of tetracyclic **5q** was verified by single crystal X-ray diffraction.¹⁹ Additionally, we demonstrated the scalability of the procedure by the synthesis of **5n**, which could be obtained in excellent yield after simple recrystallization. Though, for substrates containing a strongly electron-withdrawing nitro group intramolecular nucleophilic substitution was not successful. Under standard conditions the intermediate biaryl phenols were isolated. Here, utilizing the reaction conditions for dibenzofuro[2,3-*c*]pyridines (solvent exchange to DMF, then Cs_2CO_3), **5r** and **5s** could be obtained in 46% and 66% yield, respectively. The procedure also furnished benzothienopyridine **5t**. It is not clear if the low isolated yield is due to poisoning of the palladium catalyst or too rapid deprotection of the acetyl group, which would make 2-bromothiophenol unreactive as coupling partner as previously shown for the related unprotected phenol.

Next, we extended our methodology to the synthesis of dibenzofurans. In contrast to fluoropyridines, the directed *ortho*-metalation of fluoroarenes requires a stronger base than LDA. Here, the use of the superbasic $n\text{BuLi}/\text{KOtBu}$ system has been well established.²⁰ Adapting our standard procedure using superbasic metalation conditions, the corresponding biaryl phenols were obtained, this means that the fluoroarenes were unreactive toward intramolecular S_NAr . Again, solvent exchange to DMF and addition of Cs_2CO_3 followed by heating ($120\text{ }^\circ\text{C}$, overnight) provided a solution converting fluoroarenes **1** and 2-bromophenyl acetates **3** to dibenzofurans **6** in a one-pot four-step procedure.

With fluorobenzene and 3,4-dimethylfluorobenzene, the corresponding dibenzofurans **6a** and **6b** were obtained in 78% and 50%, respectively (Table 3). In the case of fluoroanisoles, fluorine proved to be a stronger directing group under superbasic metalation conditions as demonstrated by the formation of **6c** and **6d**.^{20b} The structure of **6c** was confirmed by single crystal X-ray diffraction.¹⁹ Furthermore, 1,4- and 1,2-difluorobenzene provided the corresponding fluoro-substituted dibenzofurans **6e** and **6f** in reasonable yield. Finally, the 2-bromophenyl acetates **3** coupling partner was varied delivering **6d'–e'** and **6g–i** in 39–65% yield. Here, dibenzofurans **6e/e'** and **6i** have been reported as precursors for the synthesis of host materials for blue phosphorescent OLEDs.²¹ With dibenzofurans **6d** and **6d'** as well as **6e** and **6e'** being pairs of identical compounds, our methodology allows flexibility in terms of the choice of starting materials **1** and **3** (Scheme 3). Hereby, slightly higher yields were achieved utilizing fluorobenzene (**1b**) and substituted 2-bromophenyl acetates **3** via path b. Unfortunately, the synthesis of

Table 3. Scope of Dibenzofurans^a

^a0.5 mmol scale; reaction conditions: (1) **1** (1.2 equiv), KOtBu (1.3 equiv), LDA (1.3 equiv), THF (0.25 M), $-78\text{ }^{\circ}\text{C}$, 5 min; (2) ZnCl₂ (1.3 equiv), then $-78\text{ }^{\circ}\text{C}$ to rt; (3) **3** (1.0 equiv), Pd XPhos G3 (2.0 mol %), and XPhos (2.0 mol %) in THF (0.5 M), then, $70\text{ }^{\circ}\text{C}$, 20 min; and (4) solvent exchange to DMF (0.1 M), Cs₂CO₃ (5 equiv), $120\text{ }^{\circ}\text{C}$, o/n. ^bX-ray structure of **6c** shown with 50% probability ellipsoids.

dibenzothiophenes was not successful under these reaction conditions generating 2-bromothiophenol as main product from the corresponding acetate **3**.¹⁸

In conclusion, we have developed an efficient one-pot procedure for the facile assembly of fused benzofuro heterocycles. Fluoropyridines or fluoroarenes were subjected to a directed *ortho*-lithiation followed by zincation and Negishi cross-coupling with 2-bromophenyl acetates. In situ deprotection of the acyl group and subsequent intramolecular S_NAr facilitates the formation of benzofuro pyridines and dibenzofurans. This methodology takes advantage of readily available starting materials, mild reaction conditions and low catalyst loading to provide a diverse set of benzofuro pyridines and dibenzofurans.


EXPERIMENTAL SECTION

Typical Procedure for the Synthesis of 2-Bromophenyl Acetates 3. 2-Bromophenyl Acetate (**3a**). Following a procedure by Kónya,¹⁵ to a solution of 2-bromophenol (4.46 g, 40.0 mmol, 1.0 equiv) in anhydrous CH₂Cl₂ (40 mL) were added acetyl chloride (1.56 mL, 22.0 mmol, 1.1 equiv) and Et₃N (3.04 mL, 22.0 mmol, 1.1 equiv). The reaction mixture was stirred for 1 h at room temperature. Then, the organic layer was washed with saturated aqueous NaHCO₃ solution (20 mL) and the aqueous layer was extracted with CH₂Cl₂ (2 × 20 mL). The combined organic layers were dried over anhydrous MgSO₄, filtered, and the solvent was removed under reduced pressure

to give **3a** as a pale-yellow oil (4.26 g, 19.8 mmol, 99% yield), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ_H 7.61 (dd, *J* = 8.3, 0.9 Hz, 1H), 7.33 (t, *J* = 7.7 Hz, 1H), 7.16–7.09 (m, 2H), 2.36 (s, 3H); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ_C 168.7, 148.4, 133.5, 128.6, 127.5, 123.9, 116.4, 20.9. The analytical data match those reported in the literature.²²

Typical Procedure for the Synthesis of Benzofuro[2,3-*b*]pyridines 5. Benzofuro[2,3-*b*]pyridine (**5a**). To a solution of 2-fluoropyridine (52 μL, 0.60 mmol, 1.2 equiv) in anhydrous THF (2.4 mL), was added a solution of LDA (2.0 M, 325 μL, 0.65 mmol, 1.3 equiv) dropwise at $-25\text{ }^{\circ}\text{C}$. The reaction mixture was stirred at $-25\text{ }^{\circ}\text{C}$ for 5 min, followed by the addition of ZnCl₂ solution in THF (0.7 M, 930 μL, 0.65 mmol, 1.3 equiv). The cooling bath was removed, and the reaction mixture was allowed to warm to room temperature, after which a solution of **3a** (108 mg, 0.50 mmol, 1.0 equiv), precatalyst Pd XPhos G3 (8.5 mg, 10 μmol, 2.0 mol %), and XPhos (4.8 mg, 10 μmol, 2.0 mol %) in THF (1.0 mL) followed by a solution of KOtBu in THF (1.6 M, 625 μL, 1.00 mmol, 2.0 equiv) was added. The reaction mixture was stirred at $70\text{ }^{\circ}\text{C}$ in a heated oil bath overnight. Saturated NH₄Cl solution (20 mL) was added, and the mixture was extracted with EtOAc (3 × 20 mL). The combined organic layers were dried over anhydrous MgSO₄, filtered, and concentrated in vacuo. The crude product was purified by column chromatography on silica gel via 9:1 to 4:1 petroleum ether/EtOAc elution gradient to yield **5a** as an off-white solid (81 mg, 0.48 mmol, 96% yield). TLC(petroleum ether/EtOAc 9:1) *R*_f = 0.15; mp = 65–66 °C. Lit. 68–69 °C;⁸ ¹H NMR (500 MHz, CDCl₃) δ_H 8.44 (dd, *J* = 4.8, 1.3 Hz, 1H), 8.28 (dd, *J* = 7.5, 1.0 Hz, 1H), 7.95 (dd, *J* = 7.7, 0.6 Hz, 1H), 7.65 (d, *J* = 8.3 Hz, 1H), 7.53 (ddd, *J* = 8.4, 7.5, 1.3 Hz, 1H), 7.39 (td, *J* = 7.7, 0.8 Hz, 1H), 7.36 (dd, *J* = 6.8, 4.4 Hz, 1H); ¹³C{¹H} NMR (126 MHz, CDCl₃) δ_C 163.2, 154.7, 146.3, 129.9, 128.6, 123.5, 122.6, 121.5, 119.4, 117.3, 112.3. The analytical data match those reported in the literature.⁸

Typical Procedure for the Synthesis of Benzofuro[2,3-*c*]pyridines 5 and Benzofuro[2,3-*b*]pyridines with Strongly Electron-Withdrawing Substituents. Benzofuro[2,3-*c*]pyridine (**5e**). To a solution of 3-fluoropyridine (52 μL, 0.60 mmol, 1.2 equiv) in anhydrous THF (2.4 mL), was added a solution of LDA (2.0 M, 325 μL, 0.65 mmol, 1.3 equiv) dropwise at $-25\text{ }^{\circ}\text{C}$. The reaction mixture was stirred at $-25\text{ }^{\circ}\text{C}$ for 5 min, followed by the addition of ZnCl₂ solution in THF (0.7 M, 930 μL, 0.65 mmol, 1.3 equiv). The cooling bath was removed, and the reaction mixture was allowed to warm to room temperature, after which a solution of **3a** (108 mg, 0.50 mmol, 1.0 equiv), precatalyst Pd XPhos G3 (8.5 mg, 10 μmol, 2.0 mol %) and XPhos (4.8 mg, 10 μmol, 2.0 mol %) in THF (1.0 mL) was added. The reaction mixture was stirred at $70\text{ }^{\circ}\text{C}$ in a heated oil bath for 20 min. The solvent was removed under reduced pressure, and the residue was dissolved in anhydrous DMF (5.0 mL). Cs₂CO₃ (815 mg, 2.50 mmol, 5.0 equiv) was added, and the reaction mixture was stirred at $100\text{ }^{\circ}\text{C}$ in a heated oil bath for 2 h. After cooling to room temperature, saturated NH₄Cl solution (20 mL) was added, and the mixture was extracted with EtOAc (3 × 20 mL). The combined organic layers were washed with water (4 × 30 mL) and brine (30 mL), dried over anhydrous MgSO₄, filtered, and concentrated in vacuo. The crude product was purified by column chromatography on silica gel via 4:1 to 2:1 petroleum ether/EtOAc elution gradient to give **5e** as an off-white solid (33 mg, 0.20 mmol, 39% yield). TLC(petroleum ether/EtOAc 4:1) *R*_f = 0.14; mp = 90.5–91.5 °C. Lit. 93–95 °C;⁸ ¹H NMR (500 MHz, CDCl₃) δ_H 8.99 (s, 1H), 8.59 (d, *J* = 5.1 Hz, 1H), 8.03 (d, *J* = 7.8 Hz, 1H), 7.87 (d, *J* = 5.0 Hz, 1H), 7.65 (d, *J* = 8.2 Hz, 1H), 7.61 (t, *J* = 7.8 Hz, 1H), 7.42

Scheme 3. Convergent Synthesis of Dibenzofurans **6d/d'** and **6e/e'**

(t, $J = 7.4$ Hz, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ_{C} 156.9, 152.9, 143.1, 134.6, 131.1, 130.1, 123.6, 122.3, 122.2, 115.3, 112.6. The analytical data match those reported in the literature.⁸

Typical Procedure for the Synthesis of Dibenzofurans 6. *Dibenzo[b,d]furan (6a)*. To a solution of fluorobenzene (56 μL , 0.60 mmol, 1.2 equiv) and KOtBu (1.6 M, 403 μL , 0.65 mmol, 1.3 equiv) in anhydrous THF (2.4 mL), was added a solution of *n*BuLi (1.58 M, 411 μL , 0.65 mmol, 1.3 equiv) dropwise at -78 °C. The reaction mixture was stirred at -78 °C for 5 min, followed by the addition of ZnCl_2 solution in THF (0.7 M, 930 μL , 0.65 mmol, 1.3 equiv). The cooling bath was removed, and the reaction mixture was allowed to warm to room temperature, after which a solution of **3a** (108 mg, 0.50 mmol, 1.0 equiv), precatalyst Pd XPhos G3 (8.5 mg, 10 μmol , 2.0 mol %), and XPhos (4.8 mg, 10 μmol , 2.0 mol %) in THF (1.0 mL) was added. The reaction mixture was stirred at 70 °C in a heated oil bath for 20 min. The solvent was removed under reduced pressure, and the residue was dissolved in anhydrous DMF (5.0 mL). Cs_2CO_3 (815 mg, 2.50 mmol, 5.0 equiv) was added, and the reaction mixture was stirred at 120 °C in a heated oil bath overnight. After cooling to room temperature, saturated NH_4Cl solution (20 mL) was added, and the mixture was extracted with EtOAc (3 \times 20 mL). The combined organic layers were washed with water (4 \times 30 mL) and brine (30 mL), dried over anhydrous MgSO_4 , filtered, and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 95:5 petroleum ether/EtOAc to give **6a** as a waxy white solid (66 mg, 0.39 mmol, 78% yield). TLC (petroleum ether/EtOAc 9:1) $R_f = 0.59$; mp = 78–80 °C. Lit. 83–84 °C²³; ^1H NMR (400 MHz, CDCl_3) δ_{H} 7.97 (d, $J = 7.7$ Hz, 2H), 7.60 (d, $J = 8.2$ Hz, 2H), 7.48 (t, $J = 7.7$ Hz, 2H), 7.36 (t, $J = 7.5$ Hz, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ_{C} 156.3, 127.3, 124.4, 122.8, 120.8, 111.8. The analytical data match those reported in the literature.²³

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.2c02111>.

Experimental procedures and characterization data for all new compounds, copies of ^1H , ^{13}C , and ^{19}F NMR spectra, as well as XRD data for **5q** and **6c** (PDF)

Accession Codes

CCDC 2191069–2191070 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Stefan Roesner – Department of Chemistry, University of Warwick, Coventry CV4 7AL, United Kingdom;
 orcid.org/0000-0003-2143-4708; Email: s.roesner@warwick.ac.uk

Author

Guy J. Clarkson – Department of Chemistry, University of Warwick, Coventry CV4 7AL, United Kingdom

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.joc.2c02111>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the University of Warwick's Institutional Research Support Fund for financial support and Prof. Mike Shipman for helpful discussion.

REFERENCES

- Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274.
- Ram, V. J.; Sethi, A.; Nath, M.; Pratap, R. *The Chemistry of Heterocycles: Chemistry of Six-to-Eight-Membered N, O, S, P, and Se Heterocycles*; Elsevier: Amsterdam, 2019; pp 254–259.
- Love, B. E. Isolation and synthesis of polyoxygenated dibenzofurans possessing biological activity. *Eur. J. Med. Chem.* **2015**, *97*, 377–387.
- (a) Brachwitz, K.; Hilgeroth, A. Synthesis and First Biological Evaluation of 1-Aza-9-oxafluorenes as Novel Lead Structures for the Development of Small-Sized Cytostatics. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 411–413. (b) Voigt, B.; Meijer, L.; Lozach, O.; Schächtele, C.; Totzke, F.; Hilgeroth, A. Novel CDK inhibition profiles of structurally varied 1-aza-9-oxafluorenes. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 823–825.
- Krug, M.; Voigt, B.; Baumert, C.; Lüpken, R.; Molnár, J.; Hilgeroth, A. First biological evaluation of developed 3-benzyloxyfluorenes as novel class of MDR modulators. *Eur. J. Med. Chem.* **2010**, *45*, 2683–2688.
- (a) Lee, C. W.; Lee, J. Y. Benzo[4,5]thieno[2,3-*b*]pyridine derivatives as host materials for high efficiency green and blue phosphorescent organic light-emitting diodes. *Chem. Commun.* **2013**, *49*, 1446–1448. (b) Kim, H.; Lee, K. Y.; Kim, J. Y.; Noh, H. J.; Yoon, D.-W.; Yang, J. H.; Kim, J.; Kang, I. B.; Lee, S. J. Preparation of Acridine-modified Benzofuro-pyridine and Benzothienopyridine Derivatives and Their Thermally Activated Delayed Fluorescence Properties. *Bull. Korean Chem. Soc.* **2016**, *37*, 1870–1873. (c) Lee, G. H.; Kim, Y. S. Benzofuro-pyridine-based highly efficient thermally activated delayed fluorescence emitters. *J. Nanosci. Nanotechnol.* **2018**, *18*, 6635–6639.
- Ooyama, Y.; Sato, T.; Harima, Y.; Ohshita, J. Development of a D- π -A dye with benzothienopyridine as the electron-withdrawing anchoring group for dye-sensitized solar cells. *J. Mater. Chem. A* **2014**, *2*, 3293–3296.
- Yue, W. S.; Li, J. J. A concise synthesis of all four possible benzo[4,5]furo-pyridines via palladium-mediated reactions. *Org. Lett.* **2002**, *4*, 2201–2203.
- (a) Ames, D. E.; Opalko, A. Palladium-catalyzed cyclisation of 2-substituted halogenoarenes by dehydrohalogenation. *Tetrahedron* **1984**, *40*, 1919–1925. (b) Liu, J.; Fitzgerald, A. E.; Mani, N. S. Facile Assembly of Fused Benzo[4,5]furo Heterocycles. *J. Org. Chem.* **2008**, *73*, 2951–2954. (c) Yoon, W. S.; Lee, S. J.; Kang, S. K.; Ha, D.-C.; Ha, J. D. Facile synthesis of benzo[4,5]furo[3,2-*c*]pyridines via palladium-catalyzed intramolecular Heck reaction. *Tetrahedron Lett.* **2009**, *50*, 4492–4494.
- (a) Lai, L.-L.; Lin, P.-Y.; Huang, W.-H.; Shiao, M.-J.; Hwu, J. R. Sodium Trimethylsilanethiolate in Novel Cyclizations for Synthesis of Aromatic Heterocyclic Compounds. *Tetrahedron Lett.* **1994**, *35*, 3545–3546. (b) González, J. P.; Edgar, M.; Elsegood, M. R. J.; Weaver, G. W. Synthesis of fluorinated fused benzofurans and benzothiophenes: Smiles-type rearrangement and cyclisation of perfluoro(het)aryl ethers and sulfides. *Org. Biomol. Chem.* **2011**, *9*, 2294–2305. (c) Zhu, D.; Wu, Z.; Luo, B.; Du, Y.; Liu, P.; Chen, Y.; Hu, Y.; Huang, P.; Wen, S. Heterocyclic Iodoniums for the Assembly of Oxygen-Bridged Polycyclic Heteroarenes with Water as the Oxygen Source. *Org. Lett.* **2018**, *20*, 4815–4818.
- (a) Komine, Y.; Kamisawa, A.; Tanaka, K. Flexible Synthesis of Fused Benzofuran Derivatives by Rhodium-Catalyzed [2 + 2 + 2] Cycloaddition with Phenol-Linked 1,6-Diynes. *Org. Lett.* **2009**, *11*, 2361–2364. (b) Rao, Y.; Li, Z.; Yin, G. Clean and efficient assembly

of functionalized benzofuro[2,3-*c*]pyridines via metal-free one-pot domino reaction. *Green Chem.* **2014**, *16*, 2213–2218. (c) Hu, J.; Deng, Z.; Zhang, X.; Zhang, F.; Zheng, H. Synthesis of benzofuro[2,3-*c*]pyridines via a one-pot three-component reaction. *Org. Biomol. Chem.* **2014**, *12*, 4885–4889. (d) Kumar, S.; Cruz-Hernández, C.; Pal, S.; Saunthwal, R. K.; Patel, M.; Tiwari, R. K.; Juaristi, E.; Verma, A. K. Tandem Approach to Benzothieno- and Benzofuro-pyridines from *o*-Alkynyl Aldehydes via Silver-Catalyzed 6-*endo-dig* Ring Closure. *J. Org. Chem.* **2015**, *80*, 10548–10560. (e) Fu, X.; Yang, J.; Deng, K.; Shao, L.; Xia, C.; Ji, Y. Tandem C–C/C–N Formation via Palladium-Catalyzed C–H Activation/Styrenation and Sequential Annulation of *O*-Methylketoxime with Styrenes. *Org. Lett.* **2019**, *21*, 3505–3509.

(12) Xiong, W.; Hu, K.; Lei, Y.; Zhen, Q.; Zhao, Z.; Shao, Y.; Li, R.; Zhang, Y.; Chen, J. Palladium-Catalyzed Cascade Reactions of 2-(Cyanomethoxy)chalcones with Arylboronic Acids: Selective Synthesis of Emissive Benzofuro[2,3-*c*]pyridines. *Org. Lett.* **2020**, *22*, 1239–1243.

(13) Sathiyalingam, S.; Roesner, S. Synthesis of α - and β -Carbolines by a Metalation/Negishi Cross-Coupling/ S_NAr Reaction Sequence. *Adv. Synth. Catal.* **2022**, *364*, 1769–1774.

(14) Roesner, S.; Buchwald, S. L. Continuous-Flow Synthesis of Biaryls by Negishi Cross-Coupling of Fluoro- and Trifluoromethyl-Substituted (Hetero)arenes. *Angew. Chem. Int. Ed.* **2016**, *55*, 10463–10467.

(15) Pajtás, D.; Patonay, T.; Kónya, K. Synthesis of 8-Bromoflavone and Its Buchwald–Hartwig Reaction with Amines. *Synthesis* **2016**, *48*, 97–102.

(16) Bruno, N. C.; Tudge, M. T.; Buchwald, S. L. Design and preparation of new palladium precatalysts for C–C and C–N cross-coupling reactions. *Chem. Sci.* **2013**, *4*, 916–920.

(17) Note that later studies revealed that CPhos provided comparable results to XPhos. For CPhos in Negishi cross-coupling reactions, see: Zhang, H.; Buchwald, S. L. Palladium-Catalyzed Negishi Coupling of α -CF₃ Oxiranyl Zincate: Access to Chiral CF₃-Substituted Benzylic Tertiary Alcohols. *J. Am. Chem. Soc.* **2017**, *139*, 11590–11594.

(18) See the [Supporting Information](#) for a detailed description.

(19) CCDC 2191069 (**5q**) and 2191070 (**6c**) contain the supplementary crystallographic data for this paper.

(20) (a) Schlosser, M.; Takagishi, S. Fluorine- and Trifluoromethyl-Substituted Toluenes: Site Selective Metalation of Aromatic or Benzylic Positions. *Synlett* **1991**, *1991*, 119–121. (b) Katsoulos, G.; Takagishi, S.; Schlosser, M. The Metalation of Fluoroanisoles: Optional Regioselectivity Due to Metal Mediated Control. *Synlett* **1991**, *1991*, 731–732. (c) Mongin, F.; Schlosser, M. Regioselective *ortho*-Lithiation of Chloro and Bromo Substituted Fluoroarenes. *Tetrahedron Lett.* **1996**, *37*, 6551–6554.

(21) Yu, J. G.; Byeon, S. Y.; Han, S. H.; Lee, J. Y. Selective F or Br Functionalization of Dibenzofurans for Application as Host Materials of Phosphorescent Organic Light-Emitting Diodes. *Chem.-Eur. J.* **2017**, *23*, 16044–16050.

(22) Chyan, W.; Kilgore, H. R.; Gold, B.; Raines, R. T. Electronic and Steric Optimization of Fluorogenic Probes for Biomolecular Imaging. *J. Org. Chem.* **2017**, *82*, 4297–4304.

(23) Nervig, C. S.; Waller, P. J.; Kalyani, D. Palladium-Catalyzed Intramolecular C–H Arylation of Arenes Using Tosylates and Mesylates as Electrophiles. *Org. Lett.* **2012**, *14*, 4838–4841.