

# A general method to access underexplored ylideneamino sulfates as interrupted Beckmann-type rearrangement intermediates

Zhou, Yifei; Jones, Alan M

DOI:

[10.3390/molecules29071667](https://doi.org/10.3390/molecules29071667)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Zhou, Y & Jones, AM 2024, 'A general method to access underexplored ylideneamino sulfates as interrupted Beckmann-type rearrangement intermediates', *Molecules*, vol. 29, no. 7, 1667.  
<https://doi.org/10.3390/molecules29071667>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Article

# A General Method to Access Underexplored Ylideneamino Sulfates as Interrupted Beckmann-Type Rearrangement Intermediates

Yifei Zhou and Alan M. Jones \* 

School of Pharmacy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

\* Correspondence: a.m.jones.2@bham.ac.uk; Tel.: +44-(0)121-414-7288

**Abstract:** The Beckmann rearrangement of ketoximes to their corresponding amides, using a Brønsted acid-mediated fragmentation and migration sequence, has found wide-spread industrial application. We postulated that the development of a methodology to access ylideneamino sulfates using tributylsulfonium betaine (TBSAB) would afford isolable Beckmann-type intermediates and competent partners for subsequent rearrangement cascades. The ylideneamino sulfates generated, isolated as their tributylammonium salts, are sufficiently activated to undergo Beckmann rearrangement without additional reagent activation. The generation of sulfuric acid in situ from the ylideneamino sulfate giving rise to a routine Beckmann rearrangement and additional amide bond cleavage to the corresponding aniline was detrimental to reaction success. The screening of bases revealed inexpensive sodium bicarbonate to be an effective additive to prevent classic Brønsted acid-mediated fragmentation and achieve optimal conversions of up to 99%.

**Keywords:** Beckmann; rearrangement; sulfation; oxime; interrupted; ylideneamino sulfate



**Citation:** Zhou, Y.; Jones, A.M. A General Method to Access Underexplored Ylideneamino Sulfates as Interrupted Beckmann-Type Rearrangement Intermediates. *Molecules* **2024**, *29*, 1667. <https://doi.org/10.3390/molecules29071667>

Academic Editors: Florian F. Mulks and Renè Hommelsheim

Received: 18 March 2024

Revised: 26 March 2024

Accepted: 3 April 2024

Published: 8 April 2024




**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The amide functional group is one of the most important and widely encountered structural units in organic chemistry and biochemistry. The amide bond is commonly found in bioactive drug molecules including valsartan, an angiotensin receptor blocker (ARB) [1]; sitagliptin, an anti-diabetic medicine used to treat type-2 diabetes [2]; olaparib, a PARP inhibitor used to treat BRCA-mutated advanced ovarian cancer [3]; and apixaban, an anti-coagulant used to prevent stroke in nonvalvular atrial fibrillation via the inhibition of Factor Xa (Figure 1) [4].

The Beckmann rearrangement, which involves the conversion of an oxime to an amide [5,6], represents an attractive method for the synthesis of amides due to its simple operation, high selectivity, and atom economy. The Beckmann rearrangement is typically carried out by treating a ketoxime with a strong acid, such as sulfuric acid or phosphoric acid, at an elevated temperature (130 °C). The Beckmann rearrangement has been modified and updated using catalysts [7,8], acidic ionic liquids [9–11], radicals [12], photochemistry [13], reagent improvements [14,15], and recently, an example of an interrupted Beckmann rearrangement [16]. In turn, the amides generated are important substrates for electrosynthesis applications [17–20].


There are limited known methods for the use of ylideneamino sulfates as a Beckmann intermediate (Scheme 1). The rearrangement of potassium salts of ketoxime *O*-sulfates prepared via the reaction of the respective ketone with hydroxylamine-*O*-sulfonic acid gave amide products in very limited examples (5 exemplars, 7–47% yield) with multiple by-products formed using anhydrous hydrochloric acid mediated rearrangement conditions [21].




**Figure 1.** Amide bonds found in commonly encountered medicines (highlighted in red).

**Prior art**


**Smith, 1948 (5 examples reported)**




**Kelly, 1971**




**Fukui, 1973**



**Turbak, 1968**



**This work**



**Scheme 1.** Prior art and this work. R<sub>1</sub> and R<sub>2</sub> can be H, alkyl, aryl, or acyl. Dioxane cations omitted for clarity [21–24].

The use of reactive liquid sulfur trioxide in dioxane resulted in the isolation of the *O*-sulfate of cyclohexanone oxime [22–24]. Kelly observed that cyclohexanone oxime sulfate undergoes exothermic rearrangement under mild heating [22]. This contrasts with the work of Fukui who did not observe this phenomenon but instead explored a range of Lewis bases (alkylamines, arylamines), Lewis acids (sulfur trioxide, zinc chloride, tin tetrachloride), hydrochloric acid, water, and methanol-mediated effects on the cyclohexanone oxime sulfate intermediate [23]. It was found that sulfur trioxide was an optimal additive for the rearrangement step to caprolactam in this single example [23]. A series of complexing agents (pyridine, dioxane, triethyl phosphate, DMF) with sulfur trioxide have been explored to prepare caprolactam via the *O*-sulfate and subsequent treatment with aqueous sodium hydroxide [24].

Inspired by these findings in Scheme 1, we initially considered whether tributylsulfonium betaine (TBSAB) [25–30] would be an effective sulfation reagent for the generation of *O*-sulfated oximes (ylideneamino sulfates), a functional group with a paucity of methods to tractably access [31]. It was anticipated that these ylideneamino sulfates could act as intercepted Beckmann rearrangement intermediates [15].

## 2. Results and Discussion

Attempts to optimize the *O*-oxime sulfation step with TBSAB are reported in Table 1 using the model substrate 1-phenylethan-1-one oxime (**1a**). The optimal conditions identified (Table 1, entry 1) required 2.0 equivalents of TBSAB in acetonitrile at 90 °C for 3 h, consistent with the *O*-sulfation of alcohol functional groups [30].

**Table 1.** Identification of optimal *O*-oxime sulfation conditions on **1a**.


Entry	Solvent	Temperature [°C]	Time [h]	Conv. [%]
1	MeCN	90	3	>99
2	MeCN	60	3	90
3	MeCN	30	>24	<10

Due to a lack of commercially available oximes, 13 additional oximes were synthesized [32] from their respective ketones with hydroxylamine, in excellent yields of 90–99% (Chart 1).

With the optimal conditions in hand (Table 1) and a range of oxime substrates prepared (Chart 1), these were then in turn explored using TBSAB as the *O*-sulfating agent (Chart 2).

The *O*-sulfation of ketoximes (Chart 2) all showed good to excellent conversions as measured by <sup>1</sup>H NMR spectroscopy and isolated yields (typically 67–98%) after an extractive purification protocol (Supplementary materials). These results also showed that mono-substituted aromatic oximes were not influenced by the attached groups from highly electron-donating methoxy groups (**2e**) to highly electron-withdrawing nitro groups (**2f**). However, compound **2o**, starting from an aldoxime, gave a relatively low yield in comparison to the ketoximes. A potential rationale for this finding was due to the higher reactivity of the aldehyde functional group in comparison to the more hindered, and stabilized, ketone group found in the precursor to **2a** (*c.f.* **1a**). The instability of the aldehyde and resulting imine functionality may be one cause of the lower isolated yield in **2o**. For the poly-substituted examples, compounds **2i**, **2j**, and **2m** gave good yields whilst compounds **2g** and **2k** were affected by the two electron-withdrawing groups, giving a lower rate of conversion but acceptable yield of 67%. This contrasts with both the excellent

conversion in **2f** containing a *para*-nitro electron-withdrawing group and a *para*-chloro inductive withdrawing group and resonance electron-donating group found in **2b**. A plausible rationale is both the *meta*-positioning of the nitro group leading to a destabilizing delta-positive effect adjacent to the carbonyl group and reduced solubility of the halogen and nitro containing **2g** and **2k**.



**Chart 1.** Synthesis of oxime substrates. Percentage conversion is given in parentheses as measured by  $^1\text{H}$  NMR spectroscopy. An isolated yield is given thereafter.


With a range of differentially substituted ylideneamino sulfates isolated, our attention turned to whether these intermediates would undergo a Beckmann-type rearrangement. Optimization studies are shown in Table 2.

Entries 1–3 (Table 2) studied the effects of temperature on the rearrangement and identified that 100 °C gave the highest conversion to **3a** at 54% (entry 2). With increased temperatures, the formation of aniline (**4a**) became more pronounced (20%, entry 3) via presumably the hydrolysis of the amide motif in **3a**.

The analogous thermal conditions in entries 1–3 were compared with an additional equivalent of TBSAB (entries 4–6) and gave no improvement.


The possibility of the ylideneamino sulfate (**2a**) undergoing hydrolysis (entries 1–6, Table 2) to generate sulfuric acid and thus undergoing Brønsted-acid catalyzed rearrangement and/or further hydrolysis was initially overlooked. However, the low yields and multiple product formations made us consider two competing reaction pathways (Scheme 2). It should also be noted that the group *trans* to the oxime *O*-sulfate should migrate from a molecular orbital perspective [33]. However, under the reaction conditions, the *E/Z*

conversion of the oxime *O*-sulfate is rapid, such that the following order of migratory capacity is adhered to aryl, alkenyl > tertiary alkyl > secondary alkyl > primary alkyl based on electron density effects.




**Chart 2.** Screening of ylideneamino sulfate formation. \*Cation omitted for clarity ( $^+\text{NBu}_3\text{H}$ ). The percentage conversion is given in parentheses as measured by  $^1\text{H}$  NMR spectroscopy. An isolated yield is given thereafter.

Table 2. Optimization of the Beckmann rearrangement from 2a to 3a (and/or 1a and 4a).



Entry	T (°C)	Time (h)	TBSAB (1.0 Equiv.)	Base [6.0 Equiv.]	Conv. [%] <sup>a</sup> 1a	Conv. [%] <sup>a</sup> 3a	Conv. [%] <sup>a</sup> 4a
1	80	8	-	-	9	26	trace
2	100	8	-	-	6	54	14
3	120	8	-	-	7	37	20
4	80	8	Yes	-	5	40	trace
5	100	8	Yes	-	10	57	10
6	120	8	Yes	-	9	37	25
7 <sup>b</sup>	100	4	-	NaHCO <sub>3</sub>	trace	>99	0
8 <sup>b</sup>	100	4	Yes	NaHCO <sub>3</sub>	0	>99	0
9 <sup>b</sup>	100	4	Yes	KHCO <sub>3</sub>	0	90	0
10 <sup>b</sup>	100	4	Yes	pyridine	0	60	0

<sup>a</sup> Conversions were determined by <sup>1</sup>H NMR with internal standard. <sup>b</sup> Reaction time for entries 7–10 was reduced to 4 h.




**Scheme 2.** (a) Proposed mechanism of this work (blue) versus acid-mediated Beckmann mechanism (red); and (b) mechanism of the O-oxime sulfation step.

Entries 7–10 (Table 2) were designed to solve this issue through the addition of different bases to intercept any acid-mediated pathways in operation. Entry 8 (Table 2) was selected as the optimized condition with six equivalents of NaHCO<sub>3</sub> together with one equivalent of TBSAB at 100 °C for 4 h. Fortuitously, the increased yield and selectivity gained with the addition of sodium bicarbonate enabled a simple work-up to afford the amide products (Supplementary Materials).

The rationale for additional TBSAB being required is to return the formation of any trace oxime by-product (**1a**) to a competent reaction partner (**2a**) and increase the overall conversion to the desired amide product (**3a**) and concomitantly improve the purification protocol. The sulfur trioxide group in TBSAB may also activate the ylideneamino sulfate via coordination to the oxygen or the formation of pyrosulfate via analogy to the reported reactivity of sulfur trioxide-dioxane [23].

To consider the possibility of a counter ion effect of the base used, a controlled experiment was designed (Scheme 3). After sulfation of **1a**, the corresponding **2a** (as the HNBu<sub>3</sub> salt) was converted into the inorganic salts of sodium, **2a[Na]** and potassium **2a[K]**, respectively, in highly isolated yields for comparison. The lower conversions for the Beckmann-type rearrangement (40 and 15%, sodium and potassium salts, respectively) showed that the exchange of HNBu<sub>3</sub> in situ was unlikely with NaHCO<sub>3</sub> and the conclusion was that the base acted merely to intercept any hydrolytic acid generated. The lower conversions with the inorganic alkali metal salts may also be due to the tributylammonium cation acting as a phase transfer catalyst for the highly polar sulfate group.



**Scheme 3.** A control experiment to explore alternative sodium and potassium counter ion effects in the Beckmann-type rearrangement.


With the optimized conditions in place, the nineteen ylideneamino sulfates were screened for Beckmann-type rearrangement competency (Chart 3).

Electron-rich or moderately withdrawing mono-substituted aryl substrates gave excellent results, with 91–99% isolated yields (**3a**, **3c**, **3d**, **3e**, **3h** in Chart 3), and a trifluoromethyl example (**3l** in Chart 3) gave a lower 65% isolated yield under these conditions. More powerful electron withdrawing functional groups gave mixed results: **3p**, **3g**, and **3k** did not react and returned complex mixtures of products, whilst **3b** and **3f** returned the oxime starting material (**1b** and **1f**, respectively). A tentative reason for these unsuccessful examples can be elucidated from the electron withdrawing effect on the migratory capacity of the aryl ring system, which is clear in the starting materials **2p** and **2f**. Indeed, **2b** and **2f** gave clean (>99%) conversions back to **1b** and **1f** via inspection of <sup>1</sup>H NMR spectra and comparison to authentic samples.

A similar pattern with poly-substituted substrates with electron-rich **3i**, **3j**, **3n**, and **3m** (Chart 3) proceeded in modest to high yields. Electron-withdrawing examples **3g** and **3k** did not participate. A rationale for this effect derives from the electron-rich nature required for the migratory group. Compound **3o**, an example of a sulfated aldoxime, did not participate in the reaction despite the generally increased reactivity of aldoximes in the Beckmann rearrangement [34] or the possibility of benzonitrile formation via dehydration [23]. The



degradation of **2o** (precursor to **3o**) was observed, giving credence to the higher reactivity of the ketoxime example.



**Chart 3.** Screening for Beckmann-type rearrangement of the ylidenamino sulfates to amides. Percentage conversion is given in parentheses as measured by <sup>1</sup>H NMR spectroscopy. An isolated yield is given thereafter.

A cyclic example (**3q**) proved recalcitrant to these conditions. This may be attributed to the steric bulk around the reacting *O*-sulfate oxime and tributylammonium cation effect. This would be especially pronounced in a more rigid cyclic example, as was witnessed.

The deletion of one pyridyl group in **3s** vs. **3r** led to an excellent 88% yield vs. no reaction, respectively. Most likely, the deactivating nature of a bipyridyl example prevents the migratory step, as it is the phenyl group that migrates, not the pyridyl ring in **3s** which is not an available pathway to **3r**.

### 3. Conclusions

In summary, we have developed a general methodology to access nineteen isolable ylidenamino sulfates using tributylsulfoammonium betaine (TBSAB) in an up to 99% isolated yield, which was tolerant of a wide range of substituted and cyclic ket(al)oximes. In turn, these novel ylidenamino sulfates, isolated as their tributylammonium salts, are

sufficiently activated to undergo Beckmann-type rearrangement. The generation of sulfuric acid in situ from the ylideneamino sulfate giving rise to a classic Beckmann rearrangement and subsequent additional amide bond cleavage to the corresponding aniline was detrimental to reaction success. The screening of bases revealed inexpensive sodium bicarbonate to be an effective additive to prevent classic Brønsted acid-mediated fragmentation and achieve optimal conversions up to 99% in 11 examples with steric and electronic factors that influenced the outcome explored. A tentative mechanism is proposed based on the control experiments with alternative alkali metal cation intermediates. Taken together, these findings demonstrate an alternative protocol to activate and rearrange oximes to amides.

#### 4. Experimental

All reactions involving moisture-sensitive reagents were carried out using standard Schlenk techniques in a dry reaction vessel under argon. All solvents used under anhydrous conditions were decanted directly from an SPS dispensary or were stored over 4 Å molecular sieves 24 h prior to use.

Solvents used for workup procedures were of technical grade from Sigma-Aldrich, Honeywell, VWR, or Fisher Scientific. Unless stated otherwise, solvents were removed by rotary evaporation under reduced pressure between 30–50 °C. All chemical reagents were used as received unless stated otherwise. Reactions were monitored by TLC analysis on Merck silica gel 60 F254 using UV light (254 nm) and/or potassium permanganate.

$^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded either on a Bruker AVIII operating at 300 MHz for  $^1\text{H}$  and fitted with a 5mm BBFO probe or on a Bruker AVANCE NEO operating at 400 MHz for  $^1\text{H}$  fitted with a 5mm “smart” BBFO probe, respectively.  $^1\text{H}$ - $^1\text{H}$  COSY, DEPT-45,  $^1\text{H}$ - $^{13}\text{C}$  HSQC, and  $^1\text{H}$ - $^{13}\text{C}$  HMBC NMR spectra were recorded on a Bruker AVANCE NEO console operating at 400 MHz for  $^1\text{H}$  and fitted with a nitrogen-cooled BBFO probe. Chemical shift data for  $^1\text{H}$  are reported in parts per million (ppm,  $\delta$  scale) downfield from tetramethylsilane (TMS:  $\delta$  0.0) and referenced internally to the residual proton in the solvent. The deuterated solvents used for NMR analysis were chloroform ( $\text{CDCl}_3$ :  $\delta\text{H}$  7.26,  $\delta\text{C}$  77.2), methanol (MeOD:  $\delta\text{H}$  3.31,  $\delta\text{C}$  49.2), and dimethyl sulfoxide ( $\text{DMSO-d}_6$ :  $\delta\text{H}$  2.50,  $\delta\text{C}$  39.5). Coupling constants ( $J$ ) are given in hertz (Hz). The data are presented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, m = multiple, br = broad, app = apparent and combinations thereof), coupling constant, and integration and assignment.

Mass spectra were recorded on a Waters Xevo G2-XS ToF or Synap G2-S mass spectrometer using Zspray and Electro-spray ionization in negative (ESI-) and positive (ESI+) mode, respectively.

General procedure 1: Synthetic procedure for the preparation of oximes.

A magnetically stirred mixture of the corresponding acetophenone (5.0 mmol),  $\text{NH}_2\text{OH}\cdot\text{HCl}$  (0.520 g, 7.5 mmol), and sodium acetate (1.025 g, 12.5 mmol) in ethanol/water (20 mL of a 1:3 *v/v* mixture) was heated under reflux for 2 h. The precipitate was formed upon cooling and the reaction mixture was isolated by filtration and washed with water ( $3 \times 10$  mL) to give the desired oximes.

General procedure 2. Synthetic procedure for the preparation of sulfates using tributyl sulfoammonium betaine ( $\text{Bu}_3\text{NSO}_3$ , TBSAB).

A flask was charged with the respective oxime (1.0 mmol) and TBSAB (528 mg, 2.0 mmol, 2.0 eq) was added under an argon atmosphere. Anhydrous MeCN was added (giving a concentration of 0.50  $\text{Mol dm}^{-3}$  to the limiting oxime reagent). The reaction mixture was heated at 82 °C (reflux) for 3 h and monitored by TLC. After reaction completion, the flask was cooled to room temperature and the solvent was removed under reduced pressure. The reaction was quenched with cold water (10 mL) and filtered. The aqueous solution was extracted with EtOAc ( $4 \times 50$  mL). The organic layer was dried ( $\text{MgSO}_4$ ), filtered, and the filtrate solvent was removed in vacuo to afford the desired compound as a clear oil.

General procedure 3. Synthetic procedure for the preparation of amides.

A 25 mL round-bottom flask was charged with the respective *O*-sulfate (1.0 mmol) from General procedure 1, TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux (100 °C) for 4 h and monitored by TLC. The crude product was washed with cold water (3 × 10 mL), filtered, and freeze-dried to afford the desired product.

Selected compound characterization

*N*-phenylacetamide (**3a**)

Following general procedure 3: A round-bottom flask was charged with **2a** (400 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water (3 × 10 mL) and then filtered and freeze-dried to give the desired product as a white solid (163 mg, 99%). M.P. 112–114 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ<sub>H</sub> 9.90 (s, 1H), 7.56 (d, *J* = 7.8 Hz, 2H), 7.28 (t, *J* = 7.8 Hz, 2H), 7.01 (t, *J* = 7.8 Hz, 1H), 2.03 (s, 3H); <sup>13</sup>C NMR (101 MHz, DMSO-d<sub>6</sub>) δ<sub>C</sub> 168.2, 139.8, 128.6, 122.9, 119.0, 23.6; LRMS *m/z* (ESI+) 136.07 ([M+H]<sup>+</sup>, 100%); HRMS *m/z* (ESI+) C<sub>8</sub>H<sub>10</sub>NO requires 136.0711, found 136.0710 ([M+H]<sup>+</sup>).

*N*-(4-fluorophenyl)acetamide (**3c**)

Following general procedure 3: A round-bottom flask was charged with **2c** (419 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water (3 × 10 mL) and then filtered and freeze-dried to give the desired product as a white solid (279 mg, 91%). M.P. 152–153 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ<sub>H</sub> 9.96 (1 H, s), 7.61–7.55 (2 H, m), 7.15–7.09 (2 H, m), 2.02 (3 H, s); <sup>13</sup>C NMR (101 MHz, DMSO-d<sub>6</sub>) δ<sub>C</sub> 168.6, 159.5 (d, <sup>1</sup>*J*<sub>C-F</sub> = 239.4 Hz), 135.0 (d, <sup>4</sup>*J*<sub>C-F</sub> = 2.5 Hz), 121.2 (d, <sup>3</sup>*J*<sub>C-F</sub> = 7.5 Hz), 115.8 (d, <sup>2</sup>*J*<sub>C-F</sub> = 22.3 Hz), 25.0; <sup>19</sup>F NMR (377 MHz, MeOD) δ −120.75; LRMS *m/z* (ESI+) 154.06 ([M+H]<sup>+</sup>, 100%); HRMS *m/z* (ESI+) C<sub>8</sub>H<sub>9</sub>FNO requires 154.0598, found 154.0599 ([M+H]<sup>+</sup>).

*N*-(4-methylphenyl)acetamide (**3d**)

Following general procedure 3: A round-bottom flask was charged with **2d** (414 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water (3 × 10 mL) and then filtered and freeze-dried to give the desired product as a white solid (148 mg, 99%). M.P. 151–153 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ<sub>H</sub> 9.81 (1 H, s), 7.46–7.43 (2 H, m), 7.09–7.06 (2 H, m), 2.23 (3 H, s), 2.01 (3 H, s); <sup>13</sup>C NMR (101 MHz, DMSO-d<sub>6</sub>) δ<sub>C</sub> 168.5, 137.3, 132.6, 129.0, 121.2, 24.4, 19.5; LRMS *m/z* (ESI+) 150.08 ([M+H]<sup>+</sup>, 100%); HRMS *m/z* (ESI+) C<sub>9</sub>H<sub>12</sub>NO requires 150.0804, found 150.0802 ([M+H]<sup>+</sup>).

*N*-(4-methoxyphenyl)acetamide (**3e**)

Following general procedure 3: A round-bottom flask was charged with **2e** (411 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water (3 × 10 mL) and then filtered and freeze-dried to give the desired product as a white solid (163 mg, 99%). M.P. 125–128 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ<sub>H</sub> 9.77 (s, 1H), 7.50–7.43 (m, 2H), 6.88–6.82 (m, 2H), 3.70 (s, 3H), 2.00 (s, 3H); <sup>13</sup>C NMR (101 MHz, DMSO-d<sub>6</sub>) δ<sub>C</sub> 169.7, 154.1, 134.8, 119.7, 115.5, 55.0, 26.0; LRMS *m/z* (ESI+) 166.08 ([M+H]<sup>+</sup>, 100%); HRMS *m/z* (ESI+) C<sub>9</sub>H<sub>12</sub>NO<sub>2</sub> requires 166.0839, found 166.0842 ([M+H]<sup>+</sup>).

*N*-(4-methylthiophenyl)acetamide (**3h**)

Following general procedure 3: A 25 mL round-bottom flask was charged with **2h** (480 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h monitored by TLC. The crude product was washed with cold water (3 × 10 mL) and then filtered and freeze-dried to give the desired product as a white solid (179 mg, 99%). M.P. 125–128 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ<sub>H</sub> 9.92 (s, 1H), 7.57–7.49 (m, 2H), 7.25–7.17 (m, 2H), 2.43 (s, 3H), 2.02 (s, 3H);

$^{13}\text{C}$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_{\text{C}}$  170.6, 136.1, 131.8, 126.2, 118.4, 25.0, 17.1; LRMS  $m/z$  (ESI+) 182.06 ( $[\text{M}+\text{H}]^+$ , 100%); HRMS  $m/z$  (ESI+)  $\text{C}_9\text{H}_{12}\text{NOS}$  requires 182.0603, found 182.0607 ( $[\text{M}+\text{H}]^+$ ).

*N*-(3,4,5-trimethoxyphenyl)acetamide (**3i**)

Following general procedure 3: A round-bottom flask was charged with **2i** (490 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol),  $\text{NaHCO}_3$  (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water ( $3 \times 10$  mL) and then filtered and freeze-dried to give the desired product as a white solid (214 mg, 95%). M.P. 140–142 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  9.85 (s, 1H), 6.95 (s, 2H), 3.72 (s, 6H), 3.60 (s, 3H), 2.01 (s, 3H);  $^{13}\text{C}$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_{\text{C}}$  169.0, 155.2, 137.7, 134.3, 97.2, 59.5, 55.6, 24.5; LRMS  $m/z$  (ESI+) 226.10 ( $[\text{M}+\text{H}]^+$ , 100%); HRMS  $m/z$  (ESI+)  $\text{C}_{11}\text{H}_{16}\text{NO}_4$  requires 226.1027, found 226.1022 ( $[\text{M}+\text{H}]^+$ ).

*N*-(3-fluoro-4-methoxyphenyl)acetamide (**3j**)

Following general procedure 3: A round-bottom flask was charged with **2j** (468 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol),  $\text{NaHCO}_3$  (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water ( $3 \times 10$  mL) and then filtered and freeze-dried and then purified with ( $\text{SiO}_2$ ; Hexane/EtOAc, 1:1,  $R_f = 0.2$ ) to give the desired product as a white solid (119 mg, 65%). M.P. 165–168 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  9.93 (s, 1H), 7.56 (dd,  $J = 9.0, 2.5$  Hz, 1H), 7.20 (ddd,  $J = 9.0, 2.5, 1.4$  Hz, 1H), 7.08 (t,  $J = 9.0, 1.4$  Hz, 1H), 3.78 (s, 3H), 2.01 (s, 3H);  $^{13}\text{C}$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_{\text{C}}$  168.6 (d,  $J_{\text{C-F}} = 250.2$  Hz), 160.1 (d,  $J_{\text{C-F}} = 3.2$  Hz), 150.1 (d,  $J_{\text{C-F}} = 13.3$  Hz), 146.6 (d,  $J_{\text{C-F}} = 8.4$  Hz), 132.1 (d,  $J_{\text{C-F}} = 4.2$  Hz), 114.9 (d,  $J_{\text{C-F}} = 1.5$  Hz), 108.0 (d,  $J_{\text{C-F}} = 12.6$  Hz), 56.6, 24.3;  $^{19}\text{F}$  NMR (377 MHz, DMSO- $d_6$ )  $\delta_{\text{F}}$  −134.08; LRMS  $m/z$  (ESI+) 184.07 ( $[\text{M}+\text{H}]^+$ , 100%); HRMS  $m/z$  (ESI+)  $\text{C}_9\text{H}_{11}\text{FNO}_2$  requires 184.0695, found 184.0692 ( $[\text{M}+\text{H}]^+$ ).

*N*-(4-trimethylfluorophenyl)acetamide (**3l**)

Following general procedure 3: A round-bottom flask was charged with **2l** (468 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol),  $\text{NaHCO}_3$  (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water ( $3 \times 10$  mL) and then filtered, freeze-dried, and then purified with ( $\text{SiO}_2$ ; Hexane/EtOAc, 4:6,  $R_f = 0.2$ ) to give the desired product as a white solid (132 mg, 65%). M.P. 103–104 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  10.30 (s, 1H), 7.78 (d,  $J = 8.2$  Hz, 2H), 7.65 (d,  $J = 8.2$  Hz, 2H), 2.08 (s, 3H);  $^{13}\text{C}$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_{\text{C}}$  169.3, 150.3, 141.9 (d,  $^2J_{\text{C-F}} = 31$  Hz), 127.5 (q,  $^4J_{\text{C-F}} = 4.8$  Hz), 127.3 (q,  $^3J_{\text{C-F}} = 12.6$  Hz), 119.2 (d,  $^1J_{\text{C-F}} = 281.1$  Hz), 24.6;  $^{19}\text{F}$  NMR (377 MHz, DMSO- $d_6$ )  $\delta_{\text{F}}$  −59.25; LRMS  $m/z$  (ESI+) 204.06 ( $[\text{M}+\text{H}]^+$ , 100%); HRMS  $m/z$  (ESI+)  $\text{C}_9\text{H}_9\text{F}_3\text{NO}$  requires 204.0628, found 204.0627 ( $[\text{M}+\text{H}]^+$ ).

*N*-(3-chloro-4-methoxyphenyl)acetamide (**3m**)

Following general procedure 3: A round-bottom flask was charged with **2m** (465 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol),  $\text{NaHCO}_3$  (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water ( $3 \times 10$  mL) and then filtered, freeze-dried, and then purified with ( $\text{SiO}_2$ ; Hexane/EtOAc, 1:1,  $R_f = 0.2$ ) to give the desired product as a white solid (110 mg, 55%). M.P. 82–85 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  9.93 (s, 1H), 7.76 (d,  $J = 2.6$  Hz, 1H), 7.39 (dd,  $J = 8.9, 2.6$  Hz, 1H), 7.08 (d,  $J = 8.9$  Hz, 1H), 3.80 (s, 3H), 2.01 (s, 3H);  $^{13}\text{C}$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_{\text{C}}$  168.5, 150.7, 133.5, 121.0, 120.9, 119.2, 113.4, 56.6, 23.9; LRMS  $m/z$  (ESI+) 200.04 ( $[\text{M}^{35}\text{Cl}+\text{H}]^+$ , 100%), 202.04 ( $[\text{M}^{37}\text{Cl}+\text{H}]^+$ , 60%); HRMS  $m/z$  (ESI+)  $\text{C}_9\text{H}_{11}\text{ClNO}_2$  requires 200.0441, found 200.0439 ( $[\text{M}^{35}\text{Cl}+\text{H}]^+$ ).

*N*-(benzo[*d*][1,3]dioxol-5-yl)acetamide (**3n**)

Following general procedure 3: A round-bottom flask was charged with **2n** (414 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol),  $\text{NaHCO}_3$  (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h, monitored by TLC. The crude product was washed with cold water ( $3 \times 10$  mL) and then filtered and freeze-dried to give the desired product as a white solid (159 mg, 80%). M.P. 135–137 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  9.83 (s, 1H), 7.28 (d,  $J = 2.1$  Hz, 1H), 6.92 (dd,  $J = 8.4, 2.1$  Hz, 1H), 6.82 (d,  $J = 8.4$  Hz,

1H), 5.96 (s, 2H), 1.99 (s, 3H); <sup>13</sup>C NMR (101 MHz, DMSO-d<sub>6</sub>) δ<sub>C</sub> 168.1, 146.5, 142.4, 134.7, 114.9, 108.9, 101.7, 101.3, 22.9; LRMS *m/z* (ESI+) 180.06 ([M+H]<sup>+</sup>, 100%); HRMS *m/z* (ESI+) C<sub>9</sub>H<sub>10</sub>NO<sub>3</sub> requires 180.0622, found 180.0620 ([M+H]<sup>+</sup>).

*N*-(pyridin-2-yl)benzamide (**3s**)

Following general procedure 3: A round-bottom flask was charged with **2s** (445 mg, 1.0 mmol), TBSAB (256 mg, 1.0 mmol), NaHCO<sub>3</sub> (504 mg, 6.0 mmol), and water (10 mL). The reaction was heated under reflux for 4 h while being monitored by TLC. The crude product was washed with cold water (3 × 10 mL) and then filtered and freeze-dried to give the desired product as a white solid (160 mg, 89%). M.P. 78–79 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ<sub>H</sub> 10.62 (s, 1H), 8.74 (ddd, *J* = 4.8, 1.7, 1.1 Hz, 1H), 8.17 (dt, *J* = 7.9, 1.1 Hz, 1H), 8.07 (td, *J* = 7.9, 1.7 Hz, 1H), 7.93–7.89 (m, 2H), 7.68 (ddd, *J* = 7.9, 4.8, 1.1 Hz, 1H), 7.39–7.34 (m, 2H), 7.15–7.10 (m, 1H); <sup>13</sup>C NMR (101 MHz, DMSO-d<sub>6</sub>) δ<sub>C</sub> 162.9, 150.4, 148.9, 138.8, 138.6, 131.1, 129.2, 128.5, 127.4, 124.4, 122.8, 120.7; LRMS *m/z* (ESI+) 199.08 ([M+H]<sup>+</sup>, 100%); HRMS *m/z* (ESI+) C<sub>12</sub>H<sub>11</sub>N<sub>2</sub>O requires 199.0809, found 199.0813 ([M+H]<sup>+</sup>).

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules29071667/s1>, General procedures, compound characterization, and copies of <sup>1</sup>H, <sup>13</sup>C, and <sup>19</sup>F NMR spectra; References are cited in [35–45].

**Author Contributions:** Conceptualization, A.M.J.; methodology, Y.Z.; formal analysis, Y.Z. and A.M.J.; investigation, Y.Z.; resources, A.M.J.; data curation, Y.Z.; writing—original draft preparation, Y.Z. and A.M.J.; writing—review and editing, A.M.J.; supervision, A.M.J.; project administration, A.M.J. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Salim, H.; Jones, A.M. Angiotensin II receptor blockers (ARBs) and manufacturing contamination: A retrospective National Register Study into suspected associated adverse drug reactions. *Br. J. Clin. Pharmacol.* **2022**, *88*, 4812–4827. [CrossRef] [PubMed]
2. Jones, L.; Jones, A.M. Suspected adverse drug reactions of the type 2 antidiabetic drug class dipeptidyl-peptidase IV inhibitors (DPP4i): Can polypharmacology help explain? *Pharmacol. Res. Perspect.* **2022**, *10*, e01029. [CrossRef] [PubMed]
3. Sandhu, D.; Antolin, A.A.; Cox, A.R.; Jones, A.M. Identification of different side effects between PARP inhibitors and their polypharmacological multi-target rationale. *Br. J. Clin. Pharmacol.* **2022**, *88*, 742–752. [CrossRef] [PubMed]
4. Ferro, C.J.; Solkhon, F.; Jalal, Z.; Al-Hamid, A.M.; Jones, A.M. Relevance of physicochemical properties and functional pharmacology data to predict the clinical safety profile of direct oral anticoagulants. *Pharmacol. Res. Perspect.* **2020**, *8*, e00603. [CrossRef] [PubMed]
5. Beckmann, E. Zur Kenntniss der Isonitrosoverbindungen. *Ber. Dtsch. Chem. Ges.* **1886**, *19*, 988–993. [CrossRef]
6. Kaur, K.; Srivastava, S. Beckmann rearrangement catalysis: A review of recent advances. *New J. Chem.* **2020**, *44*, 18530–18572. [CrossRef]
7. Hashimoto, M.; Obora, Y.; Sakaguchi, S.; Ishii, Y. Beckmann Rearrangement of Ketoximes to Lactams by Triphosphazene Catalyst. *J. Org. Chem.* **2008**, *73*, 2894–2897. [CrossRef] [PubMed]
8. Mo, X.; Morgan, T.D.R.; Ang, H.T.; Hall, D.G. Scope and Mechanism of a True Organocatalytic Beckmann Rearrangement with a Boronic Acid/Perfluoropinacol System under Ambient Conditions. *J. Am. Chem. Soc.* **2018**, *140*, 5264–5271. [CrossRef] [PubMed]
9. Brzeczek-Szafran, A.; Erfurt, K.; Swadzba-Kwasny, M.; Piotrowski, T.; Chrobok, A. Beckmann Rearrangement with Improved Atom Economy, Catalyzed by Inexpensive, Reusable, Bronsted Acidic Ionic Liquid. *ACS Sustain. Chem. Eng.* **2022**, *10*, 13568–13575. [CrossRef]
10. Hu, H.; Cai, X.; Xu, Z.; Yan, Y.; Zhao, S. Beckmann Rearrangement of Ketoxime Catalyzed by *N*-methyl-imidazolium Hydrosulfate. *Molecules* **2018**, *23*, 1764. [CrossRef]
11. Munnuri, S.; Verma, S.; Chandra, D.; Anugu, R.R.; Falck, J.R.; Jat, J.L. Cu(OTf)<sub>2</sub>-Catalyzed Beckmann Rearrangement of Ketones Using Hydroxylamine-O-sulfonic Acid (HOSA). *Synthesis* **2019**, *51*, 3709–3714. [CrossRef] [PubMed]

12. Mahajan, P.S.; Humne, V.T.; Tanpure, S.D.; Mhaske, S.B. Radical Beckmann Rearrangement and Its Application in the Formal Total Synthesis of Antimalarial Natural Product Isocryptolepine via C–H Activation. *Org. Lett.* **2016**, *18*, 3450–3453. [CrossRef] [PubMed]
13. Chen, Y.; Cantillo, D.; Kappe, C.O. Visible Light-Promoted Beckmann Rearrangements: Separating Sequential Photochemical and Thermal Phenomena in a Continuous Flow Reactor. *Eur. J. Org. Chem.* **2019**, *2019*, 2163–2171. [CrossRef] [PubMed]
14. De Luca, L.; Giacomelli, G.; Porcheddu, A. Beckmann Rearrangement of Oximes under Very Mild Conditions. *J. Org. Chem.* **2002**, *67*, 6272–6274. [CrossRef] [PubMed]
15. Furuya, Y.; Ishihara, K.; Yamamoto, H. Cyanuric Chloride as a Mild and Active Beckmann Rearrangement Catalyst. *J. Am. Chem. Soc.* **2005**, *127*, 11240–11241. [CrossRef]
16. Tochette, S.J.; Dunkley, E.M.; Lowder, L.L.; Wu, J. Nucleophile-intercepted Beckmann fragmentation reactions. *Chem. Sci.* **2019**, *10*, 7812–7815. [CrossRef] [PubMed]
17. Alfonso-Suárez, P.; Kolliopoulos, A.V.; Smith, J.P.; Banks, C.E.; Jones, A.M. An experimentalist’s guide to electrocatalysis: The Shono oxidation. *Tetrahedron Lett.* **2015**, *56*, 6863–6867. [CrossRef]
18. Bal, M.K.; Banks, C.E.; Jones, A.M. Metabolism Mimicry: An Electrosynthetic Method for the Selective Deethylation of Tertiary Benzamides. *ChemElectroChem* **2019**, *6*, 4284–4291. [CrossRef]
19. Jones, A.M. Dialling-In New Reactivity into the Shono-Type Anodic Oxidation Reaction. *Chem. Rec.* **2021**, *21*, 2120–2129. [CrossRef]
20. Jones, A.M.; Banks, C.E. The Shono-type electroorganic oxidation of unfunctionalised amides. Carbon–carbon bond formation via electrogenerated N-acyliminium ions. *Beilstein J. Org. Chem.* **2014**, *10*, 3056–3072. [CrossRef]
21. Smith, P.A.S. Ketoxime-O-sulfonic Acids. *J. Am. Chem. Soc.* **1948**, *70*, 323–326. [CrossRef]
22. Fukui, K.; Uchida, M.; Masaki, M. Formation and Reaction of Cyclohexanone Oxime Hydrogen Sulfate. *Bull. Jpn Chem. Soc.* **1973**, *46*, 3168–3173. [CrossRef]
23. Kelly, K.K.; Matthews, J.S. Use of Lewis base-sulfur trioxide complexes as reagents for the Beckmann rearrangement of ketoximes. *J. Org. Chem.* **1971**, *36*, 2159–2161. [CrossRef]
24. Turbak, A.F. A Low Temperature Beckmann Rearrangement with SO<sub>3</sub> Including Separation of Caprimidyl Sulfate as a New Composition. *Ind. Eng. Chem. Prod. Res. Dev.* **1968**, *7*, 189–191. [CrossRef]
25. Zhou, Y.; Jones, A.M. Rearrangement of Arylsulfamates and Sulfates to Para-Sulfonyl Anilines and Phenols. *Molecules* **2024**, *29*, 1445. [CrossRef]
26. Alshehri, J.A.; Gill, D.M.; Jones, A.M. A Sulfuryl Group Transfer Strategy to Selectively Prepare Sulfated Steroids and Isotopically Labelled Derivatives. *Front. Mol. Biosci.* **2021**, *8*, 776900. [CrossRef] [PubMed]
27. Gill, D.M.; Povinelli, A.P.R.; Zazeri, G.; Mahmoud, A.M.; Shamir, S.A.; Wilkinson, F.L.; Alexander, M.Y.; Cornelio, M.L.; Jones, A.M. The modulatory role of sulfated and non-sulfated small molecule heparan sulfate-glycomimetics in endothelial dysfunction: Absolute structural clarification, molecular docking and simulated dynamics, SAR analyses and ADMET studies. *RSC Med. Chem.* **2021**, *12*, 779–790. [CrossRef]
28. Jones, A.M. Tributylsulfoammonium betaine. In *The Encyclopaedia of Reagents for Organic Synthesis (e-EROS)*; John Wiley & Sons Ltd.: Hoboken, NJ, USA, 2021; Available online: <https://onlinelibrary.wiley.com/doi/full/10.1002/047084289X.RN02393> (accessed on 27 March 2024).
29. Benedetti, A.M.; Gill, D.M.; Tsang, C.W.; Jones, A.M. Chemical Methods for N- and O-Sulfation of Small Molecules, Amino Acids and Peptides. *ChemBioChem* **2020**, *21*, 938–942. [CrossRef] [PubMed]
30. Gill, D.M.; Male, L.; Jones, A.M. Sulfation made simple: A strategy for synthesising sulfated molecules. *Chem. Commun.* **2019**, *55*, 4319–4322. [CrossRef]
31. Nishijima, K.; Nishida, H.; Yamashita, Y.; Ito, M.; Onuki, Y.; Mizota, M.; Miyano, S. Synthesis and diuretic activity of bicyclic fused heterocycles containing oxime-O-sulfonic acid moiety. *Eur. J. Med. Chem.* **2000**, *35*, 227–240. [CrossRef]
32. Chen, Y.; Gardiner, M.G.; Lan, P.; Banwell, M.G.  $\alpha$ -Iodo- $\alpha,\beta$ -Unsaturated Ketones as Vicinal Dielectrophiles: Their Reactions with Dinucleophiles Provide New Annulation Protocols for the Formation of Carbo- and Heterocyclic Ring Systems. *J. Org. Chem.* **2022**, *87*, 6146–6160. [CrossRef] [PubMed]
33. Fernández, A.B.; Boronat, M.; Blasco, T.; Corma, A. Establishing a Molecular Mechanism for the Beckmann Rearrangement of Oximes over Microporous Molecular Sieves. *Angew. Chem. Int. Ed.* **2005**, *44*, 2370–2373. [CrossRef] [PubMed]
34. Sharghi, H.; Sarvari, M.H. One-step Beckmann rearrangement from carbonyl compounds and hydroxylamine hydrochloride in Al<sub>2</sub>O<sub>3</sub>/CH<sub>3</sub>SO<sub>3</sub>H (AMA) as a new reagent. *J. Chem. Res.* **2001**, *2001*, 446–449. [CrossRef]
35. Han, Z.; Lv, J.; Zhang, J. One-pot synthesis of 2-amino-3,4-dicyanopyridines from ketoximes and tetracyanoethylene via Cu(I)-catalyzed cyclization. *Tetrahedron* **2019**, *75*, 2162–2168. [CrossRef]
36. Sonn, A. Über einige synthetische Versuche mit O-Trimethyl-gallusaldehyd. (Mitbearbeitet von Ernst Müller, Wolfgang Bülow und Walter Meyer). *Ber. Chem. Ges. A/B* **1925**, *58*, 1103–1110. [CrossRef]
37. Mahajan, P.S.; Mahajan, J.P.; Mhaske, S.B. Malonic ester amide synthesis. An efficient methodology for synthesis of amides. *Synth. Commun.* **2013**, *43*, 2508–2516. [CrossRef]
38. Tang, L.; Wang, Z.-L.; Wan, H.-L.; He, Y.-H.; Guan, Z. Visible-Light-Induced Beckmann Rearrangement by Organic Photoredox Catalysis. *Org. Lett.* **2020**, *22*, 6182–6186. [CrossRef] [PubMed]

39. Bhattacharya, S.; Ghosh, P.; Basu, B. Graphene oxide (GO) catalyzed transamidation of aliphatic amides: An efficient metal-free procedure. *Tetrahedron Lett.* **2018**, *59*, 899–903. [[CrossRef](#)]
40. Jat, J.L.; Kumar, P.; Verma, S.; Chandra, D.; Singh, V.; Tiwari, B. Metal-free synthesis of secondary amides using N-Boc-O-tosylhydroxylamine as nitrogen source via Beckmann rearrangement. *New J. Chem.* **2022**, *46*, 14782–14785. [[CrossRef](#)]
41. Ito, A.; Asami, Y.; Asato, M.; Fukuda, K.; Yamasaki, R.; Okamoto, I. Synthesis and conformational analysis of N-aryl-N-(3-thienyl)acetamides. *Tetrahedron Lett.* **2018**, *59*, 2454–2458. [[CrossRef](#)]
42. Ran, M.; He, J.; Yan, B.; Liu, W.; Li, Y.; Fu, Y.; Li, S.-J.; Yao, Q. Catalyst-free generation of acyl radicals induced by visible light in water to construct C–N bonds. *Org. Biomol. Chem.* **2021**, *19*, 1970–1975. [[CrossRef](#)] [[PubMed](#)]
43. Chip, G.K.; Grossert, J.S. Aromatic Halogenation with Titanium (IV) Chloride in Presence of Peroxytrifluoroacetic Acid. *Can. J. Chem.* **1972**, *50*, 1233–1240. [[CrossRef](#)]
44. Crisenza, G.E.M.; Sokolova, O.O.; Bower, J.F. Branch-Selective Alkene Hydroarylation by Cooperative Destabilization: Iridium-Catalyzed ortho-Alkylation of Acetanilides. *Angew. Chem. Int. Ed.* **2015**, *54*, 14866–14870. [[CrossRef](#)]
45. Fu, Z.; Wang, X.; Tao, S.; Bu, Q.; Wei, D.; Liu, N. Manganese Catalyzed Direct Amidation of Esters with Amines. *J. Org. Chem.* **2021**, *86*, 2339–2358. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.