

## Socioeconomic Deprivation and the Risk of Sight-Threatening Diabetic Retinopathy (STDR)

Tan, Luyuan; Wang, Jingya; Han, Jieun; Sainsbury, Chris; Denniston, Alastair; Crowe, Francesca; Toulis, Konstantinos; Nirantharakumar, Krishnarajah; Karamat, Muhammad Ali; Yao, Mi

DOI:  
[10.2337/dc23-1626](https://doi.org/10.2337/dc23-1626)

License:  
Other (please specify with Rights Statement)

*Document Version*  
Peer reviewed version

*Citation for published version (Harvard):*  
Tan, L, Wang, J, Han, J, Sainsbury, C, Denniston, A, Crowe, F, Toulis, K, Nirantharakumar, K, Karamat, MA & Yao, M 2024, 'Socioeconomic Deprivation and the Risk of Sight-Threatening Diabetic Retinopathy (STDR): A Population-Based Cohort Study in the U.K.', *Diabetes Care*. <https://doi.org/10.2337/dc23-1626>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

This is an accepted manuscript version of an article first published in *Diabetes Care*. The final version of record is available at <https://doi.org/10.2337/dc23-1626>

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Title:**  
2 Socioeconomic deprivation and the risk of sight-threatening diabetic retinopathy (STDR): a  
3 population-based cohort study in the UK

4  
5 **Authors**

6 Luyuan Tan MSc<sup>1\*</sup>, Jingya Wang PhD<sup>1\*†</sup>, Jieun Han MSc<sup>1</sup>, Christopher Sainsbury MD<sup>1</sup>,  
7 Alastair K. Denniston PhD<sup>2,3</sup>, Francesca L. Crowe PhD<sup>1</sup>, Konstantinos A Toulis PhD<sup>1</sup>,  
8 Muhammad Ali Karamat MD<sup>4</sup>, Mi Yao PhD<sup>5</sup>, Krishnarajah Nirantharakumar MD<sup>1†</sup>

9  
10 1 Institute of Applied Health Research, University of Birmingham, Birmingham, United  
11 Kingdom

12 2 Academic Unit of Ophthalmology, Institute of Inflammation and Ageing, University of  
13 Birmingham, Birmingham, United Kingdom

14 3 NIHR Birmingham Biomedical Research Centre, Birmingham, United Kingdom

15 4 Diabetes and Endocrinology Specialist Training Committee, Health Education West  
16 Midlands, Birmingham, United Kingdom

17 5 Department of general practice, Peking University First Hospital, Beijing, China

18

19 \* Joint First Authors.

20 † Joint Senior Authors

21

22 **Corresponding authors:** Jingya Wang

23 Current address: IOEM Building, University of Birmingham, Edgbaston, Birmingham,  
24 B15 2TT, UK

25 Telephone number: +44 (0)759 3945643

26 Email: [j.wang.6@bham.ac.uk](mailto:j.wang.6@bham.ac.uk)

27

28 and Krishnarajah Nirantharakumar

29 Current address: IOEM Building, University of Birmingham, Edgbaston, Birmingham,  
30 B15 2TT, UK

31 Telephone number: +44 (0)7737 773559

32 Email: [k.nirantharan@bham.ac.uk](mailto:k.nirantharan@bham.ac.uk)

33

1 **Twitter summary**

2 A UK study reveals an elevated risk of sight-threatening diabetic retinopathy in  
3 socioeconomically deprived individuals with diabetes, underscoring health equity issues.

4

5 Word count: 1479

6 Number of tables: 2

7 Number of figures: 1

8

1 **Objective**

2 To evaluate the associations between socioeconomic deprivation and the risk of sight-  
3 threatening diabetic retinopathy (STDR) in individuals with type 1 and type 2 diabetes.

4

5 **Research design and methods**

6 Using data derived from 175,628 individuals with diabetes in the Health Improvement Network,  
7 we assessed the risk of STDR across Townsend Deprivation Index quantiles using Cox  
8 proportional hazard regression models.

9

10 **Results**

11 Compared to the least deprived individuals, those in the most deprived quintile with type 1  
12 diabetes had a 2.85 times higher risk of developing STDR (95% CI 1.05-7.73), and those with  
13 type 2 diabetes had a 25% higher risk (1.13-1.40).

14

15 **Conclusions**

16 Increasing socioeconomic deprivation is associated with a higher risk of developing STDR in  
17 people with diabetes. This underscores persistent health disparities linked to poverty, even  
18 within a country offering free universal healthcare. Further research is needed to address health  
19 equity concerns in socioeconomically deprived regions.

20

1 **Article highlights**

2 • **Why did we undertake this study?**

3 Deprivation has been associated with various diabetes-related health issues, including  
4 morbidity, mortality, and blood glucose levels; however, its specific relationship with  
5 sight-threatening diabetic retinopathy (STDR) in people with diabetes remains  
6 relatively unclear.

7 • **What is the specific question(s) we wanted to answer?**

8 Whether deprivation played a significant role in the development of STDR in both  
9 people with type 1 and type 2 diabetes.

10 • **What did we find?**

11 A higher risk of developing STDR was observed in people with diabetes in more  
12 deprived areas.

13 • **What are the implications of our findings?**

14 Within the context of the UK's free universal healthcare system, addressing health  
15 equity in deprived areas has the potential to prevent STDR in people with diabetes.

## 1 **Introduction**

2 Diabetic retinopathy (DR), affecting over one-third of the 537 million adults with diabetes in  
3 2021, is a leading cause of preventable blindness (1). Sight-threatening diabetic retinopathy  
4 (STDR) is an advanced stage of DR affecting one in ten individuals with diabetes, which  
5 presents a significant health burden globally (2). While past studies linked low socioeconomic  
6 status with STDR in type 1 diabetes (T1DM), these studies often with small and  
7 nonrepresentative populations, lacked dedicated designs and were performed in countries  
8 without universal healthcare access (3, 4). In a country with universal healthcare, we aim to  
9 rigorously investigate the associations between socioeconomic deprivation and the risk of  
10 STDR in individuals with T1DM and type 2 diabetes (T2DM), respectively.

11

## 12 **Research Design and Methods**

13 Two open cohorts were performed in the Health Improvement Network (THIN), a large  
14 primary care-based electronic medical records database generalizable to the UK population for  
15 demographics, major condition prevalence, and death rates (5). We included individuals newly  
16 diagnosed with T1DM (aged below 40 and with insulin prescription) or those with newly  
17 diagnosed T2DM (aged over 16) between 1<sup>st</sup> January 2005 and 21<sup>st</sup> February 2020. To ensure  
18 only incident diabetes individuals were captured, the study entry began 12 months after  
19 registration. Townsend deprivation index quantile was used to measure socioeconomic  
20 deprivation, incorporating four components: unemployment, car ownership, homeownership,  
21 and household overcrowding (6). As the outcome was incident STDR, individuals with STDR  
22 at baseline were excluded. All conditions were identified using Read codes (7).

23

24 All participants were followed up from 15 months after the initial diagnosis date of diabetes (a  
25 latency period to minimize reverse causation bias) until the earliest occurrence of first diagnosis

1 of STDR, individuals left practice, practice ceased contributing to the database, death, or study  
2 end (21<sup>st</sup> February 2020).

3  
4 The incidence rates (IR) of STDR (per 1,000 person-years) across Townsend quintiles were  
5 estimated by Poisson regression. Cox proportional hazard regression models estimated the  
6 crude and adjusted hazard ratios (aHR) for STDR, adjusting for key confounding factors: age,  
7 sex, ethnicity (grouped and classified based on UK census) (8), weight and height (T1DM  
8 cohort), body mass index categories (T2DM cohort), HbA<sub>1c</sub> categories, smoking status,  
9 hypertension, peripheral vascular disease, ischemic heart disease, heart failure, chronic kidney  
10 disease, diabetic foot disease, antidiabetic drugs and lipid-lowering drugs. Missing data for  
11 ethnicity and smoking status were included in analyses as a missing category. A trend test was  
12 conducted using the multivariable-adjusted model. The risk of STDR across age groups and  
13 Townsend index missingness status were compared. All statistical tests were two-tailed and a  
14  $P < 0.05$  was considered statistically significant. Analyses were conducted using Stata 16.

15

## 16 **Results**

17 The T1DM cohort and the T2DM cohort comprised 4,406 participants and 171,222 participants,  
18 respectively (Figure 1 and Table 1).

19

### 20 **Type 1 Diabetes (T1DM) Cohort**

21 With a median follow-up of 3.7 years (interquartile range [IQR] 1.6-6.5), 89 of 4,406  
22 individuals in the T1DM Cohort developed STDR, reflecting a crude IR of 4.6 per 1,000  
23 person-years. Compared to individuals in the lowest two deprivation quintiles, the risk of  
24 STDR increased with deprivation levels (trend test:  $P < 0.001$ ) and reached its highest level in  
25 the highest deprivation quintile (aHR 2.85, 95%CI 1.05-7.73) (Table 2).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

**Type 2 Diabetes (T2DM) Cohort**

During a median follow-up of 3.8 years (IQR 1.7-6.6), 5,363 out of 171,222 individuals with T2DM developed STDR, representing a crude IR of 7.1 per 1,000 person-years. The risk of developing STDR was associated with escalating deprivation levels (trend test: P = 0.001) and reached its highest level in the highest deprivation quintile (aHR 1.25, 95%CI 1.13-1.40) compared to the lowest deprivation quintile (Table 2).

The results of age subgroup analyses were consistent with the main findings (Table 2). Unlike the unchanged STDR risk in the T1DM cohort, individuals without Townsend index exhibited a slightly higher STDR risk (aHR 1.20, 95%CI 1.12-1.28) compared to those with Townsend index in the T2DM cohort.

**Conclusions**

We found that the risk of developing STDR increases with elevated levels of deprivation in individuals with T1DM and those with T2DM. Compared with the least deprived group, individuals with T1DM and T2DM in the most deprived group had nearly three times and 25% higher risk of STDR, respectively. In a country with free universal healthcare, the observed difference in the risk of STDR among different socioeconomic deprivation levels could be largely attributed to underlying factors beyond healthcare access. Our findings highlight the importance of implementing comprehensive health equity strategies that engage diverse stakeholders and prioritize proactive management and care in socioeconomically disadvantaged areas.



1 Similar to our results, a recent US study using data derived from 1,116,361 adults with diabetes  
2 reported that the risk of hypoglycemic and hyperglycemic crises increases with the elevated  
3 level of socioeconomic deprivation (9). The results of a Scottish primary care-based study were  
4 partly similar to our findings. After adjusting for age and gender, a higher prevalence of DR in  
5 individuals with T1DM (n = 1861) was observed only in the most deprived quintile (aHR 2.40,  
6 95% CI 1.36-4.27), compared to the least deprived quintile. No association was observed  
7 between deprivation and the prevalence of DR in individuals with T2DM (n = 18,197) (10).  
8 Conversely, another large UK primary care-based study with 7.7 million participants reported  
9 an association of deprivation with the incidence of DR/severe retinopathy in individuals with  
10 T2DM (no trend was observed), but not in individuals with T1DM (11). While these studies  
11 provide valuable insights, their capacity to furnish precise estimates is hindered by flaws within  
12 their designs (e.g. failure to exclude individuals with the outcome of interests at baseline and  
13 to adjust for essential confounders [e.g. smoking, HbA<sub>1c</sub> levels, and comorbidities]) (10, 11).  
14  
15 The evident association of deprivation with the risk of STDR in individuals living with diabetes  
16 observed in a country with a universal healthcare system are possibly attributed to underlying  
17 factors beyond healthcare access. Firstly, the built environment plays a pivotal role, where  
18 inadequate housing conditions, deficient community infrastructure, and limited access to green  
19 spaces intertwine, impacting overall well-being and health conditions (12). Secondly, restricted  
20 educational opportunities and lower health literacy in these areas may lead to limited  
21 engagement in health awareness programs and social support networks, impacting preventive  
22 health behaviors, such as diabetes self-management and actively attending eye screening and  
23 treatment programs (13, 14). Thirdly, the racial composition and its related language barriers,  
24 even with free interpreter service and proactive investigations on race inequality, may still pose  
25 significant hurdles, impeding effective communication with healthcare providers and access to

1 healthcare services and essential health information (15, 16). Lastly, fragmented social  
2 structures and weakened community connections exacerbate the impact of the aforementioned  
3 factors, limiting resource access, hindering community engagement, and elevating stress levels  
4 within deprived areas, thus perpetuating health disparities (17-19). Addressing these  
5 multifaceted issues requires comprehensive interventions spanning socioeconomic,  
6 environmental, and educational domains to improve the overall well-being of deprived  
7 communities.

8

9 Given that the National Health Service (NHS) offers universal free healthcare services and  
10 medications to all individuals in the UK, the observed variations in the risk of STDR among  
11 different socioeconomic deprivation groups suggested that health equity becomes crucial in  
12 socioeconomically disadvantaged regions. Individuals with lower incomes were more likely to  
13 have greater healthcare requirements (20), suggesting that prioritized healthcare services,  
14 rather than equitable provision, should be addressed in deprived communities. When providing  
15 healthcare services, socioeconomic disadvantage and health requirements should be considered  
16 as critical determinants, since increased NHS resources invested in the most impoverished  
17 areas might generate a bigger absolute increase in health outcomes compared to the wealthier  
18 regions (21). Given that individuals with high levels of deprivation are less likely to utilize  
19 planned and preventative healthcare services (22), it is essential to implement targeted  
20 healthcare interventions, such as promoting participation in DR screening programs and  
21 providing educational activities to enhance self-glucose control skills.

22

23 The large sample size, longitudinal population-based study design, and long follow-up period  
24 have ensured that the data in this study are nationally generalizable. The adjustment of  
25 important potential confounders has improved the power of the associations between

1 socioeconomic deprivation and STDR. However, a limitation arises from the approximately  
2 16% missing data in Townsend deprivation indices. To address this, we conducted a sensitivity  
3 analysis comparing the risk of STDR between individuals with and without Townsend index  
4 to offer a comprehensive picture to readers. While ethnicity has been fully controlled in our  
5 analyses, if we hold a sufficient number of participants from minority groups, performing  
6 subgroup analyses in different ethnic groups would surely strengthen our conclusion. Lastly,  
7 our study had constraints in examining associations between individual components of the  
8 Townsend index and STDR due to data unavailability. Nonetheless, the Townsend index, a  
9 widely recognized composite measure, affords a more holistic assessment of deprivation than  
10 its individual elements.

11

12 In conclusion, we found socioeconomic deprivation was associated with the increased risk of  
13 STDR in people with T1DM and T2DM. This study highlights the importance of addressing  
14 health equity concerns and its relevance to the prevention of STDR.

15

16 **Acknowledgements.** The author wishes to thank Professor KK Cheng for his critical and  
17 insightful input in interpretation.

18 **Funding.** There is no funding support.

19 **Duality of Interest.** This is no potential conflict of interest relevant to this study.

20 **Author Contributions.** L.T. developed the idea, designed the study, analyzed the data, and  
21 led the writing of the manuscript under the supervision of J.W and K.N. J.W checked the  
22 analysis code parallelly and revised the paper. J.W and K.N. proposed the idea, critically  
23 appraised the paper, and made final suggestions. J.H., K.T., F.C., C.S., M.Y., M.K., and A.D.  
24 provided specialist input. All authors reviewed the manuscript and agreed to submit the final  
25 manuscript. L.T. and J.W. are the guarantors of this work. There is no funding support and no  
26 potential conflicts of interest in this study.

1 **Reference:**

- 2 1. Federation ID. Diabetes around the world in 2021: International Diabetes Federation;  
3 2022 [Available from:  
4 [https://diabetesatlas.org/#:~:text=1%20in%208%20adults%20\(206,caused%20by%20diabete](https://diabetesatlas.org/#:~:text=1%20in%208%20adults%20(206,caused%20by%20diabete)  
5 [s%20in%202021.](https://diabetesatlas.org/#:~:text=1%20in%208%20adults%20(206,caused%20by%20diabete)
- 6 2. Yau JWY, Rogers SL, Kawasaki R, Lamoureux EL, Kowalski JW, Bek T, et al. Global  
7 Prevalence and Major Risk Factors of Diabetic Retinopathy. *Diabetes Care*. 2012;35(3):556-  
8 64.
- 9 3. Bihan H, Laurent S, Sass C, Nguyen G, Huot C, Moulin JJ, et al. Association among  
10 individual deprivation, glycemic control, and diabetes complications: the EPICES score.  
11 *Diabetes care*. 2005;28(11):2680-5.
- 12 4. Alvarez-Ramos P, Jimenez-Carmona S, Alemany-Marquez P, Cordoba-Doña JA,  
13 Aguilar-Diosdado M. Socioeconomic deprivation and development of diabetic retinopathy in  
14 patients with type 1 diabetes mellitus. *BMJ Open Diabetes Research and Care*.  
15 2020;8(2):e001387.
- 16 5. Blak B, Thompson M, Dattani H, Bourke A. Generalisability of The Health  
17 Improvement Network (THIN) database: demographics, chronic disease prevalence and  
18 mortality rates. *Journal of Innovation in Health Informatics*. 2011;19(4):251-5.
- 19 6. Townsend P. Deprivation. *Journal of Social Policy*. 1987;16(2):125-46.
- 20 7. Luyuan Tan JH, Christopher Sainsbury, Alastair K. Denniston, Francesca L. Crowe,  
21 Konstantinos A Toulis, Muhammad Ali Karamat, Mi Yao, Krishnarajah Nirantharakumar,  
22 Jingya Wang. Appendix 1-3 and Appendix Table 1-2 Github: Github; 2023 [Available from:  
23 <https://github.com/Freyaaay/Deprivation/blob/85546e1c6f75aa3970421f439327515bccf2dcb>  
24 [3/Appendix.docx.](https://github.com/Freyaaay/Deprivation/blob/85546e1c6f75aa3970421f439327515bccf2dcb)
- 25 8. GOV.UK. List of ethnic groups: GOV.UK; 2021 [Available from:

- 1 <https://www.ethnicity-facts-figures.service.gov.uk/style-guide/ethnic-groups>.
- 2 9. Kurani SS, Heien HC, Sangaralingham LR, Inselman JW, Shah ND, Golden SH, et al.  
3 Association of area-level socioeconomic deprivation with hypoglycemic and hyperglycemic  
4 crises in US adults with diabetes. *JAMA Network Open*. 2022;5(1):e2143597-e.
- 5 10. Low L, Law JP, Hodson J, McAlpine R, O'Colmain U, MacEwen C. Impact of  
6 socioeconomic deprivation on the development of diabetic retinopathy: a population-based,  
7 cross-sectional and longitudinal study over 12 years. *BMJ open*. 2015;5(4).
- 8 11. Mathur R, Bhaskaran K, Edwards E, Lee H, Chaturvedi N, Smeeth L, et al. Population  
9 trends in the 10-year incidence and prevalence of diabetic retinopathy in the UK: a cohort study  
10 in the Clinical Practice Research Datalink 2004–2014. *BMJ open*. 2017;7(2):e014444.
- 11 12. Gibson M, Petticrew M, Bambra C, Sowden AJ, Wright KE, Whitehead M. Housing  
12 and health inequalities: a synthesis of systematic reviews of interventions aimed at different  
13 pathways linking housing and health. *Health & place*. 2011;17(1):175-84.
- 14 13. Tan ST, Quek RYC, Haldane V, Koh JJK, Han EKL, Ong SE, et al. The social  
15 determinants of chronic disease management: perspectives of elderly patients with  
16 hypertension from low socio-economic background in Singapore. *International journal for  
17 equity in health*. 2019;18(1):1-14.
- 18 14. Bachmann M, Eachus J, Hopper C, Davey Smith G, Propper C, Pearson N, et al. Socio-  
19 economic inequalities in diabetes complications, control, attitudes and health service use: a  
20 cross-sectional study. *Diabetic Medicine*. 2003;20(11):921-9.
- 21 15. Zheng Y, Lamoureux EL, Chiang P-CP, Anuar AR, Ding J, Wang JJ, et al. Language  
22 barrier and its relationship to diabetes and diabetic retinopathy. *BMC public health*. 2012;12:1-  
23 9.
- 24 16. Cheung N, Chee ML, Klein R, Klein BE, Shea S, Cotch MF, et al. Incidence and  
25 progression of diabetic retinopathy in a multi-ethnic US cohort: the Multi-Ethnic Study of

- 1 Atherosclerosis. *British Journal of Ophthalmology*. 2022;106(9):1264-8.
- 2 17. Whitley E, Gunnell D, Dorling D, Smith GD. Ecological study of social fragmentation,  
3 poverty, and suicide. *bmj*. 1999;319(7216):1034-7.
- 4 18. Allardyce J, Gilmour H, Atkinson J, Rapson T, Bishop J, McCreadie R. Social  
5 fragmentation, deprivation and urbanicity: relation to first-admission rates for psychoses. *The*  
6 *British Journal of Psychiatry*. 2005;187(5):401-6.
- 7 19. Grigoroglou C, Munford L, Webb RT, Kapur N, Ashcroft DM, Kontopantelis E. Spatial  
8 distribution and temporal trends in social fragmentation in England, 2001– 2011: a national  
9 study. *BMJ open*. 2019;9(1):e025881.
- 10 20. Marmot M. Fair society, healthy lives. *Fair society, healthy lives*. 2013:1-74.
- 11 21. Barr B, Bambra C, Whitehead M. The impact of NHS resource allocation policy on  
12 health inequalities in England 2001-11: longitudinal ecological study. *Bmj*. 2014;348.
- 13 22. Goddard M, Smith P. Equity of access to health care services:: Theory and evidence  
14 from the UK. *Social science & medicine*. 2001;53(9):1149-62.
- 15

**Table 1. Baseline characteristics of individuals with type 1 and type 2 diabetes across Townsend Deprivation Index quintiles**

Type 1 diabetes	Townsend Deprivation Index				
	1 (n = 811) (lowest deprivation)	2 (n = 752)	3 (n = 778)	4 (n = 687)	5 (n = 573) (highest deprivation)
<b>Age (years), mean <math>\pm</math> SD</b>	15.3 $\pm$ 8.5	17.1 $\pm$ 9.6	17.5 $\pm$ 9.5	18.0 $\pm$ 10.0	18.2 $\pm$ 10.5
<b>Male, n (%)</b>	479 (59.1)	431(57.3)	455 (58.5)	403 (58.7)	348 (60.7)
<b>Ethnicity, n (%)</b>					
White	347 (42.8)	323 (43.0)	361 (46.4)	321(46.7)	267 (46.6)
Mixed or Multiples	1 (0.1)	2 (0.3)	1 (0.1)	3 (0.4)	6 (1.1)
Other ethnic groups	4 (0.5)	1 (0.1)	2 (0.3)	5 (0.7)	3 (0.5)
Black, Black British, Black Welsh, Caribbean or African	0 (0.0)	5 (0.7)	4 (0.5)	4 (0.6)	9 (1.6)
Asian, Asian British, Asian Welsh	7 (0.9)	6 (0.8)	11 (1.4)	6 (0.9)	7 (1.2)
Missing	452(55.7)	415 (55.2)	399 (51.3)	348 (50.7)	281 (49.0)
<b>Smoking Status, n (%)</b>					
Smoker	43 (5.3)	70 (9.3)	86 (11.1)	93 (13.5)	125 (21.8)
Ex-smoker	34 (4.2)	40 (5.3)	59 (7.6)	46 (6.7)	31 (5.4)
Non-smoker	301 (37.1)	280 (37.2)	295 (37.9)	262 (38.1)	192 (33.5)
Missing	433 (53.4)	362 (48.1)	338 (43.4)	286 (41.6)	225 (39.3)
<b>HbA<sub>1c</sub> level, n (%)</b>					
<6.5%	113 (13.9)	118 (15.7)	100 (12.9)	102 (14.9)	73 (12.7)
6.5-7.5%	179 (22.1)	191 (25.4)	168 (21.6)	131 (19.1)	106 (18.5)
7.5-8.5%	177 (21.8)	152 (20.2)	160 (20.6)	143 (20.8)	120 (20.9)
$\geq$ 8.5%	179 (22.1)	172 (22.9)	217 (27.9)	187 (27.2)	176 (30.7)
Missing	163 (20.1)	119 (15.8)	133 (17.1)	124 (18.1)	98 (17.1)
<b>Hypertension, n (%)</b>	1 (0.1)	7 (0.9)	4 (0.5)	10 (1.5)	10 (1.8)
<b>Chronic kidney disease, n (%)</b>	5 (0.6)	9 (1.2)	7 (0.9)	10 (1.5)	12 (2.1)
<b>Diabetic Foot Disease, n (%)</b>	10 (1.2)	16 (2.1)	5 (0.6)	11 (1.6)	7 (1.2)
<b>Metformin, n (%)</b>	23 (2.8)	28 (3.7)	33 (4.2)	30 (4.4)	39 (6.8)

<b>Lipid-lowering drugs, <i>n</i> (%)</b>	14 (1.7)	14 (1.9)	19 (2.4)	21 (3.1)	21 (3.7)
---	----------	----------	----------	----------	----------

<b>Type 2 diabetes</b>	<b>Townsend Deprivation Index</b>				
	<b>1 (<i>n</i> = 30,909) (lowest deprivation)</b>	<b>2 (<i>n</i> = 29,765)</b>	<b>3 (<i>n</i> = 31,049)</b>	<b>4 (<i>n</i> = 29,087)</b>	<b>5 (<i>n</i> = 22,181) (highest deprivation)</b>
<b>Age (years), mean ± SD</b>	64.3 ± 12.3	64.4 ± 12.6	62.8 ± 13.1	61.9 ± 13.6	60.3 ± 13.6
<b>Male, <i>n</i> (%)</b>	18,314 (59.3)	17,044 (57.3)	17,326 (55.8)	15,647 (53.8)	11,679 (56.7)
<b>Ethnicity, <i>n</i> (%)</b>					
White	13,102 (42.4)	13,492 (45.3)	14,183 (45.7)	13,510 (46.5)	11,117 (50.1)
Mixed or Multiples	140 (0.5)	119 (0.4)	189 (0.6)	200 (0.7)	227 (1.0)
Other ethnic groups	45 (0.2)	49 (0.2)	61 (0.2)	85 (0.3)	78 (0.4)
Black, Black British, Black Welsh, Caribbean or African	157 (0.5)	167 (0.6)	352 (1.1)	484 (1.7)	681 (3.1)
Asian, Asian British, Asian Welsh	572 (1.9)	563 (1.9)	984 (3.2)	1,227 (4.2)	1,017 (4.6)
Missing	16,893 (54.7)	15,375 (51.7)	15,280 (49.2)	13,581 (46.7)	9,061 (40.9)
<b>Smoking Status, <i>n</i> (%)</b>					
Smoker	3,056 (9.9)	3,640 (12.2)	5,016 (16.2)	5,850 (20.1)	5,740 (25.9)
Ex-smoker	11,324 (36.6)	11,139 (37.4)	11,545 (37.2)	10,675 (36.7)	7,546 (34.0)
Non-smoker	16,474 (53.3)	14,954 (50.2)	14,443 (46.5)	12,533 (43.1)	8,865 (40.0)
Missing	55(0.2)	32 (0.1)	45 (0.1)	29 (0.1)	30 (0.1)
<b>BMI (kg/m<sup>2</sup>), mean ± SD</b>	30.6 ± 6.0	31.2 ± 6.3	31.8 ± 6.7	32.3 ± 6.9	32.6 ± 7.2
<b>HbA<sub>1c</sub> level, <i>n</i> (%)</b>					



≤6.5%	13,711 (44.4)	12,714 (42.7)	12,822 (41.3)	11,516 (39.6)	8,485 (38.3)
6.5-7.5%	11,975 (38.7)	11,637 (39.1)	11,918 (38.4)	11,233 (38.6)	8,531 (38.5)
7.5-8.5%	2,719 (8.8)	2,885 (9.7)	3,204 (10.3)	3,027 (10.4)	2,396 (10.8)
≥8.5%	1,844 (6.0)	1,989 (6.7)	2,446 (7.9)	2,624 (9.0)	2,269 (10.2)
Missing	660 (2.1)	540 (1.8)	659 (2.1)	687 (2.4)	500 (2.3)
<b>Hypertension, <i>n</i> (%)</b>	17,162(55.5)	16,668 (56.0)	17,060 (55.0)	15,535 (53.4)	11,546 (52.1)
<b>Peripheral vascular disease, <i>n</i> (%)</b>	730 (2.4)	832 (2.8)	947 (3.1)	1,025 (3.5)	928 (4.2)
<b>Stroke, <i>n</i> (%)</b>	2,088 (6.8)	2,227 (7.5)	2,233 (7.2)	2,137 (7.4)	1,714 (7.7)
<b>Ischemic heart disease, <i>n</i> (%)</b>	4,996 (16.2)	5,074 (17.1)	5,280 (17.0)	5,226 (18.0)	4,043 (18.2)
<b>Heart Failure, <i>n</i> (%)</b>	923 (3.0)	1,013 (3.4)	1,101 (3.6)	1,168 (4.0)	924 (4.2)
<b>Chronic kidney diseases, <i>n</i> (%)</b>	5,428 (17.6)	5,396 (18.1)	5,238 (16.9)	4,963 (17.1)	3,404 (15.4)
<b>Diabetic Foot Disease, <i>n</i> (%)</b>	4,226 (13.7)	4,309 (14.5)	46,97 (15.1)	4,462 (15.3)	3,635 (16.4)
<b>Metformin, <i>n</i> (%)</b>	15,560 (50.3)	14,995 (50.4)	16,831 (54.2)	16,467 (56.6)	13,125 (59.2)
<b>Insulin, <i>n</i> (%)</b>	1,756 (5.7)	1,775 (6.0)	2,154 (6.9)	2,213 (7.6)	1,873 (8.4)
<b>Other glucose-lowering medications, <i>n</i> (%)</b>	4,370 (15.9)	4,380 (16.0)	5,012 (18.3)	4,997 (18.2)	4,141 (15.1)
<b>Lipid-lowering drugs, <i>n</i> (%)</b>	22,600 (73.1)	21,720 (73.0)	22,716 (73.2)	21,220 (73.0)	16,347 (73.7)

SD, standard deviation; HbA<sub>1c</sub>, hemoglobin A1c.

**Table 2. The risk of developing sight-threatening diabetic retinopathy (STDR) across Townsend deprivation index quintiles in individuals with type 1 and type 2 diabetes**

Type 1 Diabetes	Townsend Deprivation Index				
	1 (lowest deprivation)	2	3	4	5 (highest deprivation)
N	811	752	778	687	573
STDR, <i>n</i> (%)	6 (0.7)	9 (1.2)	17 (2.2)	25 (3.6)	16 (2.8)
Person-years	3,582.1	3,377.7	3,454.1	2,913.9	2,568.2
Incidence rate (per 1,000 person-years)	1.7	2.7	4.9	8.6	6.2
Crude HR (95% CI)	Ref	1.69 (0.59-4.89)	2.67 (1.01-7.03)	5.08 (2.05-12.60)	4.15 (1.61-10.72)
Adjusted HR (95% CI)*	Ref	0.98 (0.32-3.08)	1.42 (0.50-4.03)	2.05 (0.76-5.55)	2.85 (1.05-7.73)
		<i>Trend test: P &lt; 0.001</i>			
<b>By age</b>					
<18 years (Adjusted HR, 95% CI)	Ref	0.38 (0.04-3.80)	1.03 (0.20-5.27)	2.56 (0.60-11.08)	2.88 (0.65-12.80)
≥18 years (Adjusted HR, 95% CI)	Ref	1.10 (0.29-5.32)	1.47 (0.36-6.01)	1.64 (0.43-6.35)	2.42 (0.61-9.59)

  

Type 2 Diabetes	Townsend Deprivation Index				
	1 (lowest deprivation)	2	3	4	5 (highest deprivation)
N	30,909	29,765	31,049	29,087	22,181
STDR, <i>n</i> (%)	881 (2.9)	858 (2.9)	942 (3.0)	930 (3.2)	701 (3.2)
Person-years	137,627.6	130,988.2	136,477.3	125,716.4	95,856.6
Incidence rate (per 1,000 person-years)	6.4	6.6	6.9	7.4	7.3
Crude HR (95% CI)	Ref	1.04 (0.94-1.15)	1.06 (0.96-1.17)	1.13 (1.02-1.24)	1.17 (1.05-1.30)
Adjusted HR (95% CI)†	Ref	1.06 (0.95-1.17)	1.08 (0.98-1.20)	1.18 (1.07-1.31)	1.25 (1.13-1.40)
		<i>Trend test: P = 0.001</i>			

**By age**

<b>&lt;65 years (Adjusted HR, 95% CI)</b>	Ref	1.02 (0.89-1.17)	1.12 (0.98-1.29)	1.20 (1.05-1.38)	1.21 (1.04-1.42)
<b>≥65 years (Adjusted HR, 95% CI)</b>	Ref	1.08 (0.93- 1.27)	1.01 (0.87-1.18)	1.11 (0.96-1.30)	1.24 (1.07-1.45)

---

STDR, sight-threatening diabetic retinopathy; HR, hazard ratio; Ref, reference group; CI, confidence interval.

\*Adjusted for age, sex, ethnicity, weight, height, HbA<sub>1c</sub>, smoking status, hypertension, chronic kidney disease (stage 3 to 5), diabetic foot disease, glucose-lowering medicine, and lipid-lowering medicine.

†Adjusted for age, sex, ethnicity, body mass index, HbA<sub>1c</sub>, smoking status, hypertension, chronic kidney disease (stage 3 to 5), diabetic foot disease, glucose-lowering medicine, lipid-lowering medicine, peripheral vascular disease, ischemic heart disease stroke, and heart failure.