



A global database of dissolved organic matter (DOM) concentration measurements in coastal waters (CoastDOM v1)

Christian Lønborg¹, Cátia Carreira², Gwenaël Abril³, Susana Agustí⁴, Valentina Amaral⁵,
Agneta Andersson⁶, Javier Arístegui⁷, Punyasloke Bhadury⁸, Mariana B. Bif⁹, Alberto V. Borges¹⁰,
Steven Bouillon¹¹, Maria Ll. Calleja¹², Luiz C. Cotoviz Jr.^{13,49}, Stefano Cozzi¹⁴, Maryló Doval¹⁵,
Carlos M. Duarte⁴, Bradley Eyre¹⁶, Cédric G. Fichot¹⁷, E. Elena García-Martín¹⁸,
Alexandra Garzon-Garcia¹⁹, Michele Giani^{20,21}, Rafael Gonçalves-Araujo²², Renee Gruber²³,
Dennis A. Hansell²⁴, Fuminori Hashihama²⁵, Ding He²⁶, Johnna M. Holding²⁷, William R. Hunter²⁸,
J. Severino P. Ibánhez²⁹, Valeria Ibello³⁰, Shan Jiang³¹, Guebuem Kim³², Katja Klun³³,
Piotr Kowalczyk³⁴, Atsushi Kubo³⁵, Choon-Weng Lee³⁶, Cláudia B. Lopes³⁷, Federica Maggioni³⁸,
Paolo Magni³⁹, Celia Marrase⁴⁰, Patrick Martin⁴¹, S. Leigh McCallister⁴², Roisin McCallum⁴³,
Patricia M. Medeiros⁴⁴, Xosé Anxelu G. Morán^{4,45}, Frank E. Muller-Karger⁴⁶, Allison Myers-Pigg⁴⁷,
Marit Norli⁴⁸, Joanne M. Oakes¹⁶, Helena Osterholz⁴⁹, Hyekyung Park³², Maria Lund Paulsen⁵⁰,
Judith A. Rosentreter^{16,51}, Jeff D. Ross⁵², Digna Rueda-Roa⁴⁶, Chiara Santinelli⁵³, Yuan Shen⁵⁴,
Eva Teira⁵⁵, Tinkara Tinta³³, Guenther Uher⁵⁶, Masahide Wakita⁵⁷, Nicholas Ward⁴⁷,
Kenta Watanabe⁵⁸, Yu Xin⁵⁹, Youhei Yamashita⁶⁰, Liyang Yang⁶¹, Jacob Yeo¹⁶, Huamao Yuan⁶²,
Qiang Zheng^{54,63}, and Xosé Antón Álvarez-Salgado²⁹

¹Section for Marine Diversity and Experimental Ecology, Department of Ecoscience,
Aarhus University, 4000 Roskilde, Denmark

²Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Portugal

³Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), CNRS,
Muséum National d'Histoire Naturelle, 61 Rue Buffon, 75005, Paris, France

⁴King Abdullah University of Science and Technology, Thuwal 23955-6900, Kingdom of Saudi Arabia

⁵Departamento Interdisciplinario de Sistemas Costero Marinos, Centro Universitario Regional Este,
Universidad de la República, Ruta 9 y 15, CP 27000, Rocha, Uruguay

⁶Umeå Marine Sciences Centre, Umeå University, Sweden

⁷Instituto de Oceanografía y Cambio Global (IOCAG), Universidad de Las Palmas de Gran Canaria,
Las Palmas, Spain

⁸Centre for Climate and Environmental Studies, Indian Institute of Science Education and Research Kolkata,
Mohanpur, Nadia, West Bengal, India

⁹Monterey Bay Aquarium Research Institute, Moss Landing, California, United States

¹⁰Chemical Oceanography Unit, University of Liège, Liège, Belgium

¹¹Department of Earth and Environmental Sciences, KU Leuven, Leuven, Belgium

¹²Marine Ecology and Systematics (MarES), Department of Biology, Universitat de les Illes Balears,
07122 Palma de Mallorca, Spain

¹³Departamento de Geoquímica, Universidade Federal Fluminense, Outeiro São João Batista s/n,
24020015 Niterói, Rio de Janeiro, Brazil

¹⁴Consiglio Nazionale delle Ricerche, Istituto di Scienze Marine (CNR-ISMAR),
Strada Statale 14, km 163.5, 34149 Trieste, Italy

¹⁵Instituto Tecnolóxico para o Control do Medio Mariño de Galicia, 36611 Vilagarcía de Arousa, Spain

¹⁶Centre for Coastal Biogeochemistry, Faculty of Science and Engineering,
Southern Cross University, Lismore, NSW, 2480, Australia

¹⁷Department of Earth and Environment, Boston University, Boston, Massachusetts, United States

- ¹⁸National Oceanography Centre, European Way, Southampton, SO14 3ZH, United Kingdom
- ¹⁹Department of Environment and Science, PO Box 5078, Brisbane, QLD, 4001, Australia
- ²⁰National Institute of Oceanography and Applied Geophysics (OGS), Trieste, Italy
- ²¹Istituto Centrale per la Ricerca scientifica e tecnologia Applicata al Mare, Chioggia, Italy
- ²²National Institute of Aquatic Resources, Technical University of Denmark, Kongens Lyngby, Denmark
- ²³Australian Institute of Marine Science, PMB 3, Townsville, QLD, 4810, Australia
- ²⁴Department of Ocean Sciences, Rosenstiel School of Marine, Atmospheric & Earth Science, University of Miami, Miami, Florida, United States
- ²⁵Department of Ocean Sciences, Tokyo University of Marine Science and Technology, Japan
- ²⁶Department of Ocean Science and Center for Ocean Research in Hong Kong and Macau, The Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong, China
- ²⁷Department of Ecoscience, Aarhus University, 8000 Aarhus, Denmark
- ²⁸Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute, Belfast, Northern Ireland, United Kingdom
- ²⁹CSIC, Instituto de Investigaci3n Mari3as, Eduardo Cabello 6, 36208 Vigo, Spain
- ³⁰Institute of Marine Sciences, Middle East Technical University, 33731 Erdemli, Mersin, T3rkiye
- ³¹State Key Laboratory of Estuarine and Coastal Research, East China Normal University, 200241, Shanghai, China
- ³²School of Earth and Environmental Sciences, Seoul National University, Seoul 08826, South Korea
- ³³Marine Biology Station, National Institute of Biology, Forna3e 41, 6330 Piran, Slovenia
- ³⁴Remote Sensing Laboratory, Institute of Oceanology, Polish Academy of Sciences, Sopot, Poland
- ³⁵Department of Geosciences, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka, 422-8529, Japan
- ³⁶Laboratory of Microbial Ecology, Institute of Biological Sciences, Institute of Ocean and Earth Sciences, Universiti Malaya, 50603 Kuala Lumpur, Malaysia
- ³⁷CICECO – Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal
- ³⁸ENTROPIE, IRD, Universit3 de la R3union, CNRS, IFREMER, Universit3 de Nouvelle-Cal3donie, Noum3a 98800, New Caledonia
- ³⁹Consiglio Nazionale delle Ricerche, Istituto per lo studio degli impatti Antropici e Sostenibilit3 in ambiente marino (CNR-IAS), Loc. Sa Mardini, Torregrande, 09170, Oristano, Italy
- ⁴⁰Institut de Ci3ncies del Mar (CSIC), Passeig Mar3tim de la Barceloneta, 37, 08003 Barcelona, Spain
- ⁴¹Asian School of the Environment, Nanyang Technological University, 639798, Singapore
- ⁴²Virginia Commonwealth University, Department of Biology, Richmond, Virginia, United States
- ⁴³Centre for Marine Ecosystems Research, School of Science, Edith Cowan University, 270 Joondalup Drive, Joondalup, WA, 6027, Australia
- ⁴⁴Department of Marine Sciences, University of Georgia, Athens, Georgia, United States
- ⁴⁵Centro Oceanogr3fico de Gij3n/Xix3n (IEO-CSIC), 33212 Gij3n/Xix3n, Spain
- ⁴⁶College of Marine Science, University of South Florida, Saint Petersburg, Florida 33701, United States
- ⁴⁷Pacific Northwest National Laboratory, Marine and Coastal Research Laboratory, Sequim, Washington, United States
- ⁴⁸Norwegian Institute for Water Research, Oslo, Norway
- ⁴⁹Leibniz Institute for Baltic Sea Research Warnem3nde, 18119 Rostock-Warnem3nde, Germany
- ⁵⁰Marine microbiology, University of Bergen, Bergen, Norway
- ⁵¹Yale School of the Environment, Yale University, New Haven, Connecticut, United States
- ⁵²Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 49, Hobart, TAS, 7001, Australia
- ⁵³Biophysics Institute, CNR, Pisa, Italy
- ⁵⁴State Key Laboratory of Marine Environmental Science (MEL) & College of Ocean and Earth Sciences, Xiamen University, China
- ⁵⁵Departamento de Ecolog3a y Biolog3a Animal, Universidade de Vigo, Centro de Investigaci3n Mari3a da Universidade de Vigo (CIM-UVigo), Vigo, Spain
- ⁵⁶School of Natural and Environmental Science, Newcastle University, Newcastle upon Tyne, United Kingdom
- ⁵⁷Mutsu Institute for Oceanography, Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology, 690 Kitasekine, Sekine, Mutsu, Aomori, Japan

⁵⁸Coastal and Estuarine Environment Research Group, Port and Airport Research Institute, Yokosuka 239-0826, Japan

⁵⁹Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education, Institute for Advanced Ocean Study, Ocean University of China, Qingdao, Shandong, China

⁶⁰Faculty of Environmental Earth Science, Hokkaido University, Hokkaido 060-0810, Japan

⁶¹College of Environment and Safety Engineering, Fuzhou University, China

⁶²Key Laboratory of Marine Ecology and Environmental Sciences, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China

⁶³Fujian Key Laboratory of Marine Carbon Sequestration, Xiamen University, Xiamen 361102, China

Correspondence: Christian Lønborg (c.lonborg@ecos.au.dk, clonborg@gmail.com)

Received: 1 September 2023 – Discussion started: 27 September 2023

Revised: 12 December 2023 – Accepted: 15 January 2024 – Published: 27 February 2024

Abstract. Measurements of dissolved organic carbon (DOC), nitrogen (DON), and phosphorus (DOP) concentrations are used to characterize the dissolved organic matter (DOM) pool and are important components of biogeochemical cycling in the coastal ocean. Here, we present the first edition of a global database (CoastDOM v1; available at <https://doi.org/10.1594/PANGAEA.964012>, Lønborg et al., 2023) compiling previously published and unpublished measurements of DOC, DON, and DOP in coastal waters. These data are complemented by hydrographic data such as temperature and salinity and, to the extent possible, other biogeochemical variables (e.g. chlorophyll *a*, inorganic nutrients) and the inorganic carbon system (e.g. dissolved inorganic carbon and total alkalinity). Overall, CoastDOM v1 includes observations of concentrations from all continents. However, most data were collected in the Northern Hemisphere, with a clear gap in DOM measurements from the Southern Hemisphere. The data included were collected from 1978 to 2022 and consist of 62 338 data points for DOC, 20 356 for DON, and 13 533 for DOP. The number of measurements decreases progressively in the sequence DOC > DON > DOP, reflecting both differences in the maturity of the analytical methods and the greater focus on carbon cycling by the aquatic science community. The global database shows that the average DOC concentration in coastal waters (average \pm standard deviation (SD): $182 \pm 314 \mu\text{mol CL}^{-1}$; median: $103 \mu\text{mol CL}^{-1}$) is 13-fold higher than the average coastal DON concentration ($13.6 \pm 30.4 \mu\text{mol NL}^{-1}$; median: $8.0 \mu\text{mol NL}^{-1}$), which is itself 39-fold higher than the average coastal DOP concentration ($0.34 \pm 1.11 \mu\text{mol PL}^{-1}$; median: $0.18 \mu\text{mol PL}^{-1}$). This dataset will be useful for identifying global spatial and temporal patterns in DOM and will help facilitate the reuse of DOC, DON, and DOP data in studies aimed at better characterizing local biogeochemical processes; closing nutrient budgets; estimating carbon, nitrogen, and phosphorus pools; and establishing a baseline for modelling future changes in coastal waters.

1 Introduction

Coastal waters are the most biogeochemically dynamic areas of the ocean, exhibiting the highest standing stocks, process rates, and transport fluxes of carbon (C), nitrogen (N), and phosphorus (P) per unit area (Bauer et al., 2013; Mackenzie et al., 2011). In these areas, organic matter plays a critical role in numerous biogeochemical processes, serving as both a C, N, and P reservoir and substrate (Carreira et al., 2021).

Organic material found in the marine environment is commonly distinguished by its size; material retained on a filter with a pore size typically between 0.2 and 0.7 μm is classified as particulate organic matter (POM), whereas organic matter that passes through the filter is referred to as dissolved organic matter (DOM). This partitioning is operational but has implications for biogeochemical cycling: POM can be suspended in the water column or sink to the sediments con-

trolled by its size, shape, and density (Laurenceau-Cornec et al., 2015), whereas DOM is a solute that mostly remains in the water column. In most coastal waters, DOM concentrations are higher than POM, with POM having a larger proportion of known biochemical classes (e.g. carbohydrates, proteins) than the dissolved fraction, suggesting that, generally, DOM is more reworked and recalcitrant (Boudreau and Rud-dick, 1991; Lønborg et al., 2018; Benner and Amon, 2015).

The DOM pool consists mainly of C (DOC), N (DON), and P (DOP), but it also includes other elements such as oxygen, sulfur, and trace elements (Lønborg et al., 2020). In coastal waters, DOM originates from multiple sources. Internal, or autochthonous, sources include planktonic organisms (Lønborg et al., 2009; Carlson and Hansell, 2015), benthic microalgae, macrophytes, and sediment porewater (Burdige and Komada, 2014; Wada et al., 2008). On the other hand, DOM from external, or allochthonous, sources, has

mainly terrestrial origins, including wetlands, river and surface runoff, groundwater discharges, and atmospheric deposition (Iavorivska et al., 2016; Raymond and Spencer, 2015; Taniguchi et al., 2019; Santos et al., 2021). The main sinks for DOM from the water column in coastal waters are the following: (1) bubble coagulation and abiotic flocculation (Kerner et al., 2003) or sorption to particles (Chin et al., 1998), (2) sunlight-mediated photodegradation (Mopper et al., 2015), and (3) microbial degradation by mainly heterotrophic prokaryotes (Lønborg and Álvarez-Salgado, 2012).

Given the importance of DOM as a source of nutrients and for coastal biogeochemical cycling in general, numerous studies have measured the C, N, and P contents of the DOM pool over the last few decades (e.g. García-Martín et al., 2021; Cauwet, 2002; Osterholz et al., 2021). Most data, however, are often unavailable or stored in an inaccessible manner, making it difficult to, for example, analyse global spatial and temporal patterns effectively. Global open ocean DOM data compilation for DOC, total dissolved nitrogen (TDN) (Hansell et al., 2021), and DOP (Liang et al., 2022; Karl and Björkman, 2015) already exists and contains a few coastal samples (< 200 m) (Hansell et al., 2021), but there are no compilations specifically focused on coastal waters. Hence, there is a clear need for a comprehensive global and integrated database of DOC, DON, and DOP measurements for coastal waters. To address this need, we have prepared the first edition of a coastal DOM database (named CoastDOM v1), by compiling both previously reported as well as unpublished data. These data have been obtained from authors of the original studies or extracted directly from the original studies. In order to allow the DOM measurements to be interpreted across larger scales, and to better understand their relationship with local environmental conditions, we have included concurrently collected ancillary data (such as physical and/or chemical seawater properties) whenever available. The objective of this database is multifaceted. Firstly, we aimed to compile all available coastal DOM data into a single repository. Secondly, our intention was to make these data easily accessible to the research community. And thirdly, we sought to achieve long-term consistency of the measurements to enable data intercomparison and establish a robust baseline for assessing, for example, the impacts of climate change and land use changes.

2 Methods

2.1 Data compilation

The measurements included in CoastDOM v1 were obtained directly from authors of previously published studies, from online databases, or from scientific papers. An extensive search of published reports, PhD theses, and peer-reviewed literature was performed to identify studies dealing with DOM in coastal waters. First, a formal search was

performed using Google Scholar in January 2022 using the search terms “dissolved organic carbon”, “dissolved organic nitrogen”, and “dissolved organic phosphorus” in connection with “marine” or “ocean”, which yielded a total of 897 articles (after filtering the query by searching content in the title and abstract and excluding non-coastal articles). When data could not be obtained directly from the corresponding authors, relevant data were extracted. Further searches for relevant datasets were conducted using the reference lists of the identified scientific papers as well as databases and repositories to capture as many datasets as possible. Additionally, research groups that were invited to participate in this effort were also encouraged to submit unpublished data to CoastDOM v1.

2.2 Dissolved organic matter analysis

The DOC concentrations included in CoastDOM v1 were commonly measured using a total organic carbon (TOC) high-temperature catalytic oxidation (HTCO) analyser (81 % of samples; Sharp et al., 1993). Some were measured by a combined wet chemical oxidation (WCO) step and/or UV digestion, after which the carbon dioxide generated was quantified (19 % of samples). Similarly, concentrations of total dissolved nitrogen (TDN; Sipler and Bronk, 2015) were determined using either a nitric oxide chemiluminescence detector connected in series with the HTCO analyser used for DOC analyses (31 % of the samples) or by employing a UV and/or chemical oxidation step (69 %). In the latter approach, both organic and inorganic N compounds were oxidized to nitrate, which was subsequently quantified through a colorimetric method to determine the concentration of inorganic N (Valderrama, 1981; Álvarez-Salgado et al., 2023; Halewood et al., 2022; Foreman et al., 2019). Another method used for DON determination is oxidizing the sample and measuring the resulting total nitrate by the nitric oxide chemiluminescence method (Knapp et al., 2005). However, none of the concentration measurements included in CoastDOM v1 applied this method. The reported DON concentrations were calculated as the difference between TDN and dissolved inorganic nitrogen (DIN: sum of ammonium (NH_4^+) and nitrate/nitrite ($\text{NO}_3^- + \text{NO}_2^-$); $\text{DON} = \text{TDN} - \text{DIN}$) (Álvarez-Salgado et al., 2023). Analyses of total dissolved phosphorus (TDP) were determined by UV (4 %), wet chemical oxidation (66 %), or a combination of these (30 %), and they were subsequently analysed for inorganic phosphorus by a colorimetric method (Álvarez-Salgado et al., 2023). Another method also previously used for TDP analysis is the ash/hydrolysis method (Solorzano and Sharp, 1980), even though none of the data included in CoastDOM v1 used this method. The DOP concentrations were calculated as the difference between TDP and soluble reactive phosphorus (SRP: HPO_4^{2-}) ($\text{DOP} = \text{TDP} - \text{SRP}$) (Álvarez-Salgado et al., 2023).

3 Description of the dataset

The data compiled in CoastDOM v1 were collected, analysed and processed by different laboratories; however, all data included have undergone quality control measures, either by using reference samples or internal quality assurance procedures. While many of the included DOC and TDN data have been systematically compared against consensus reference material (CRM), mainly provided by the University of Miami's CRM programme (Hansell, 2005), there is a limitation in CoastDOM v1 regarding the inter-calibration across different measurement systems used for both DOP and DON determination. While the CRM could be used for DOC, DON, and DOP measurements, this has not yet been attempted for DOP, and measurement uncertainties increase in the sequence $\text{DOC} > \text{DON} > \text{DOP}$. Although some of the reported measurements have quantified the DOP recovery based on commercially available DOP compounds such as adenosine triphosphate (ATP), it is not known if these were conducted systematically in all cases. Therefore, we strongly recommend undertaking further inter-calibration across laboratories for future measurements of TDP, as has been done for DOC and TDN measurements (e.g. Sharp et al., 2002). Since additional quality control is not possible in retrospect, we assessed the quality of CoastDOM v1 based on its internal consistency.

In CoastDOM v1, we defined “coastal water” as encompassing estuaries (salinity > 0.1) to the continental shelf break (water depth < 200 m). However, some locations, such as deep fjords which are close to the coast, cannot be classed as coastal due to bathymetry (deeper than > 200 m). Therefore, we evaluated the inclusion of some datasets on a case-by-case basis. For inclusion in the database, each DOM measurement needed, at a minimum, to contain the following information (if reported in the original publication or otherwise available):

- country where samples were collected,
- latitude of measurement (in decimal degree),
- longitude of measurement (in decimal degree),
- year of sampling,
- month of sampling,
- sampling day (when available),
- depth (m) at which the discrete samples were collected,
- temperature ($^{\circ}\text{C}$) of the sample,
- salinity of the sample,
- dissolved organic carbon (DOC) concentration ($\mu\text{mol L}^{-1}$),
- method used to measure DOC concentration,
- DOC – QA flag: quality flag for DOC measurement,
- dissolved organic nitrogen (DON) concentration ($\mu\text{mol L}^{-1}$),
- total dissolved nitrogen (TDN) concentration ($\mu\text{mol L}^{-1}$),
- method used to measure TDN concentration,
- TDN – QA flag: quality flag for TDN measurement,
- dissolved organic phosphorus (DOP) concentration ($\mu\text{mol L}^{-1}$),
- total dissolved phosphorus (TDP) concentration ($\mu\text{mol L}^{-1}$),
- method used to measure TDP concentration,
- TDP – QA flag: quality flag for TDP measurement,
- responsible person,
- originator institution,
- contact info of data originator.

It should be noted that in all entries at least DOC, DON, or DOP should have been measured. In addition, we also included other relevant data, when available, in the CoastDOM v1 dataset:

- depth at the station where the sample was collected (Bottom depth, metres),
- total suspended solid (TSS) concentration (mg L^{-1}),
- chlorophyll *a* (Chl *a*) concentration ($\mu\text{g L}^{-1}$),
- chl *a* – QA flag: quality flag for chlorophyll *a* measurement,
- sum of nitrate and nitrite ($\text{NO}_3^- + \text{NO}_2^-$) concentration ($\mu\text{mol L}^{-1}$),
- $\text{NO}_3^- + \text{NO}_2^-$ – QA flag: quality flag for $\text{NO}_3^- + \text{NO}_2^-$ measurement,
- ammonium (NH_4^+) concentration ($\mu\text{mol L}^{-1}$),
- NH_4^+ – QA flag: quality flag for NH_4^+ measurement,
- soluble reactive phosphorus (HPO_4^{2-}) concentration ($\mu\text{mol L}^{-1}$),
- HPO_4^{2-} – QA flag: quality flag for HPO_4^{2-} measurement,
- particulate organic carbon (POC) concentration ($\mu\text{mol L}^{-1}$),
- method used to measure POC concentration,

- POC – QA flag: quality flag for POC measurement,
- particulate nitrogen (PN) concentration ($\mu\text{mol L}^{-1}$),
- method used to measure PN concentration,
- PN – QA flag: quality flag for PN measurement,
- particulate phosphorus (PP) concentration ($\mu\text{mol L}^{-1}$),
- method used to measure PP concentration,
- PP – QA flag: quality flag for PP measurement,
- dissolved inorganic carbon (DIC) concentration ($\mu\text{mol kg}^{-1}$),
- DIC – QA flag: quality flag for DIC measurement,
- total alkalinity (TA) concentration ($\mu\text{mol kg}^{-1}$),
- TA – QA flag: quality flag for TA measurement.

Quality control of large datasets is crucial to ensure their reliability and usefulness. Thus, we have not included data that were deemed compromised, such as records that had not gone through quality control by the data originators. We also accepted a certain degree of measurement error since multiple groups have been involved in the collection, analysis, and/or compilation of the information. Some of these errors were corrected (e.g. when a value was placed in a wrong column or when clearly inaccurate locations were reallocated for consistency with the place of study), while others could not be rectified (e.g. values showing clear signs of contamination) and were consequently excluded from CoastDOM v1 (Fig. 1). It should also be noted that differences in analytical capabilities between laboratories and individual measurement campaigns likely caused additional uncertainty. Outliers, arising, for example, from contamination, were removed from the dataset. The data were moreover screened for zero values (i.e. concentrations below the detection limit or absence of data). In cases where concentrations were below the detection limit, the zero values were replaced with half the value of the limit of detection. Commonly reported detection limits are $4 \mu\text{mol L}^{-1}$ for DOC, $0.3 \mu\text{mol L}^{-1}$ for DON, and are $0.03 \mu\text{mol L}^{-1}$ for DOP.

To ensure the inclusion of only high-quality data, we only accepted entries with specific World Ocean Circulation Experiment (WOCE) quality codes: “2: Acceptable measurement” and “6: Mean of replicate measurements”. In our quality control assessments, we carefully avoided overly strict criteria, known as “data grooming”, which could potentially overlook genuine patterns and changes in the dataset that may be significant over longer temporal and/or wider spatial scales. Coastal waters are known to exhibit a wide range of environmental concentrations, influenced by factors such as seasonality and local anthropogenic activities. Consequently, these data points may encompass a wide concentration range.

However, obtaining consistent long-term datasets is important to enable data intercomparison and establish a robust baseline. Such long-term consistency can be achieved by using the CRM standards provided by the Hansell laboratory for DOC and TDN. Another helpful approach is comparing the DOM concentrations obtained by different laboratories in the same study area and time of year.

3.1 Summary of dissolved organic carbon (DOC) concentration observations

Measurements of DOC concentrations were conducted between 1978 to 2022, with a total of 62 338 individual data points (Table 1). The DOC concentrations ranged from 17 to $30\,327 \mu\text{mol CL}^{-1}$ (average \pm standard deviation (SD): $182 \pm 314 \mu\text{mol CL}^{-1}$; median: $103 \mu\text{mol CL}^{-1}$; Table 1). The majority (53 %) of the concentrations fell within the range of 60 to $120 \mu\text{mol CL}^{-1}$ (Fig. 2). A large number of DOC concentration observations (17 %) ranged between 300 and $600 \mu\text{mol CL}^{-1}$, which were predominantly collected in eutrophic and river-influenced coastal waters of the Northern Hemisphere, such as the Baltic Sea (Fig. 2). It was observed that 75 % of the DOC concentrations were higher than $77 \mu\text{mol CL}^{-1}$, while 25 % of the measurements surpassed $228 \mu\text{mol CL}^{-1}$ (Table 1).

Coastal environments that experience minimal continental runoff, such as Palmer Station in Antarctica, typically exhibit low DOC concentrations. On the other hand, coastal waters heavily influenced by humic-rich terrigenous inputs, such as the Sarawak region in Malaysia, tended to have high DOC concentrations. In addition, some extremely high DOC concentrations were measured in the River Derwent in Australia, which is impacted by paper mill effluents. There has been a large increase in the number of DOC concentration observations after 1992 (Fig. 3), and those measurements were from a wide range of locations. However, these concentration observations were not evenly distributed around the globe, with the Southern Hemisphere being under-sampled (10 % of observations), especially in the African, South American, and Antarctic continents (Figs. 3, 4).

3.2 Summary of dissolved organic nitrogen (DON) concentration observations

The DON concentration measurements were collected between 1990 and 2021, with a total of 20 356 data points (Table 1). Concentrations of DON ranged from < 0.1 to $2095.3 \mu\text{mol NL}^{-1}$ (average \pm SD: $13.6 \pm 30.4 \mu\text{mol NL}^{-1}$; median: $8.0 \mu\text{mol NL}^{-1}$; Table 1), with the most common range (42 %) for DON concentrations between 4 to $8 \mu\text{mol NL}^{-1}$ (Fig. 2). Overall, 75 % of DON concentrations were above $5.5 \mu\text{mol NL}^{-1}$, while 25 % were above $15.8 \mu\text{mol NL}^{-1}$ (Table 1).

The lowest DON concentrations were recorded in Young Sound, Greenland, which receives direct runoff from the




Figure 1. Flow diagram of data collation, quality control, and inclusion into the CoastDOM v1 database.

Table 1. Descriptive statistics for the dissolved organic carbon (DOC), dissolved organic nitrogen (DON), and dissolved organic phosphorus (DOP) concentration observations included in the CoastDOM v1 dataset. The DOC : DON, DOC : DOP, and DON : DOP for paired measurements are also reported. The minimum (Min), maximum (Max), average values (Avg.) with standard deviation (SD), median, coefficient of variation (CV %), 25th and 75th percentiles, and number of samples (N) for each variable are shown.

	DOC μmolL^{-1}	DON μmolL^{-1}	DOP μmolL^{-1}	DOC : DON	DOC : DOP	DON : DOP
Min	17	< 0.1	< 0.01	1	18	0.14
Max	30 327	2095.3	84.27	3046	248 024	8894
Avg. \pm SD	182 \pm 314	13.6 \pm 30.4	0.34 \pm 1.11	18 \pm 43	1171 \pm 4248	100 \pm 580
Median	103	8.0	0.18	14	583	47
CV	173	224	324	244	363	578
25th percentile	77	5.5	0.11	11	401	30
75th percentile	228	15.8	0.30	18	1034	78
N	62 338	20 356	13 533	12 632	7415	12 954

Greenland Ice Sheet, whereas the highest concentrations were detected during a flood event in the Richmond River estuary, Australia. Since 1995, there has been a large increase in the number of DON measurements conducted in coastal waters globally (Fig. 3); however, the majority of those measurements have been in the Northern Hemisphere (79 % of observations), mostly in Europe and the United States (Figs. 3, 4).

3.3 Summary of dissolved organic phosphorus (DOP) concentration observations

CoastDOM v1 includes a total of 13 533 DOP measurements, collected between 1990 and 2021 (Table 1). Overall, DOP concentrations ranged from < 0.10 to 84.27 μmolPL^{-1} (average \pm SD: 0.34 \pm 1.11 μmolPL^{-1} ; median: 0.18 μmolPL^{-1} ; Table 2). The majority (74 %) of DOP concentrations were below 0.30 μmolPL^{-1} (Fig. 2). Analysis of the DOP dataset revealed that 75 % of the con-

centrations were above 0.11 μmolPL^{-1} , while 25 % were above 0.30 μmolPL^{-1} (Table 1).

The lowest DOP concentrations were measured off the Kimberley Coast in Australia, while the highest concentrations were found in the Vasse-Wonnerup Estuary in the southwest region of Australia. Similar to DOC and DON, most of the DOP measurements have been conducted from the 1990s onwards, with a predominant focus in the Northern Hemisphere (70 % of observations), particularly in Europe and the United States (Figs. 3, 4).

3.4 Summary of dissolved organic matter (DOM) concentration observations

In CoastDOM v1, the number of measurements decreases progressively in the sequence DOC > DON > DOP (62 338, 20 356, and 13 533, respectively), reflecting both differences in the maturity of the analytical methods and the greater focus on carbon cycling by the aquatic science community. In




Figure 2. Histograms showing the distribution of observations for (a) dissolved organic carbon (DOC), (b) nitrogen (DON), and (c) phosphorus (DOP), within defined concentration ranges in the coastal ocean. Note that the concentration ranges are not uniform in all cases due to the large difference in concentrations.

addition, the average DOC concentration in coastal waters ($(182 \pm 314) \mu\text{mol C L}^{-1}$) was 13-fold higher than the average coastal DON concentrations ($(13.6 \pm 30.4) \mu\text{mol N L}^{-1}$), which was itself 39-fold higher than the average coastal DOP concentrations ($(0.34 \pm 1.11) \mu\text{mol P L}^{-1}$) (Table 1). Interestingly, the coefficient of variation (CV – dispersion of the data around the mean) increased from DOC (173 %) to

DON (224 %) and DOP (326 %), which is related to the fact that the percent contribution of refractory organic material decreases in the same sequence (Table 1). It should be noted that CoastDOM v1 only contains 7058 paired measurements of DOC, DON, and DOP; therefore, only a subset of observations reported all three element pools. The average C : N : P stoichiometry for these paired DOM measurements was $1171 (\pm 4248) : 100 (\pm 580) : 1$ (Table 1), which was very N and P depleted compared to the Redfield ratio (Redfield et al., 1963). However, the large variations in C : N, C : P, and N : P ratios reveals large variations in the composition of the DOM pool in coastal waters.

3.5 Potential use of the dataset

The use of the CoastDOM v1 dataset should be accompanied by the citation of this paper and the inclusion of the correct DOI reference. CoastDOM v1 is available in full open access on the PANGAEA web page as a *.csv file. The dataset includes a brief description of the metadata and methods employed, with emphasis on measurement techniques and data units. We chose the terminology most familiar to the ocean science community. It is important to note that all data included in CoastDOM v1, as well as this paper, are considered public domain; as such, a subset of this global dataset is also available in previous data compilations (e.g. Hansell et al., 2021). The list of citations and links referenced in CoastDOM v1 also provides users with information on how these data have been previously used in publications or databases.

4 Data availability

The dataset is available at the PANGAEA database (<https://doi.org/10.1594/PANGAEA.964012>; Lønborg et al., 2023). The file can be downloaded as a *.csv merged file and is available in full open access.

5 Recommendations and conclusions

In CoastDOM v1, we have compiled available coastal DOM data in a single repository, making it openly and freely available to the research community. This compilation has established a consistent global dataset, serving as a valuable information source to investigate a variety of environmental questions and to explore spatial and temporal trends. We suggest a set of recommendations for the future expansion of this global dataset. First, our analysis highlights a spatial bias, with a concentration of sampling efforts and/or data availability predominantly concentrated in the Northern Hemisphere. The data gap in coastal DOM measurements in the Southern Hemisphere needs to be addressed to provide a more representative global understanding of the role of DOM in coastal water biogeochemistry. Additionally, increased sampling efforts especially around Africa and South America as well as island nations are warranted due to the




Figure 3. (a) Cumulative number of concentration observations for dissolved organic carbon (DOC), nitrogen (DON), and phosphorus (DOP). Number of concentration observations shown as a function of (b) sampling month (“N.S.” represents samples for which the sampling month is not specified), (c) latitude, and (d) longitude, grouped into bins of 10° latitude or longitude.

vulnerability of many coastal areas to climate change and intensifying human activities, which will undoubtedly impact DOM biogeochemistry. Furthermore, there are comparatively few data from coastal waters affected by river discharge into the tropics, e.g. the Amazon or Indian and Indonesian rivers that together dominate freshwater inputs to the coastal ocean. Second, there is a need for more comprehensive temporal and spatial datasets to capture the variability of DOM concentrations in highly dynamic and productive coastal systems. Focused efforts should be made to resolve these temporal and spatial changes. Third, only a fraction of data entries report paired DOC, DON, and DOP measurements; we encourage that these be measured and reported together in order to better determine changes in stoichiometry and composition. Fourth, collecting and reporting ancillary data, such as temperature, salinity, nutrient measurements, and particulate components, are important to provide context and to better understand the underlying processes driving the observed DOM concentrations. Fifth, studies need to collect

a minimum of metadata and report them in a standardized manner. Lastly, we recommend regular inter-calibration exercises to establish standardized and interoperable methods and data, particularly for DON and DOP measurements. This will ensure the comparability and reliability of data across different studies and enhance our understanding of DON and DOP dynamics in coastal waters.

In light of ongoing global environmental changes, the mobilization and open sharing of existing data for important biogeochemical variables, such as the DOM pool, are crucial for establishing baselines and determining global trends and changes in coastal waters. The aim is to publish an updated version of the database periodically to determine global trends of DOM levels in coastal waters, and we therefore encourage researchers to submit new data to the corresponding author. The CoastDOM v1 dataset was developed according to the FAIR principles regarding findability, accessibility, interoperability, and reusability of data. Thus, CoastDOM v1 will serve as a reliable open-source information resource, en-




Figure 4. Global distribution of concentration observations included in CoastDOM v1 for (a) dissolved organic carbon (DOC), (b) nitrogen (DON), and (c) phosphorus (DOP). The black dots on the map represent the reported data that are included in the CoastDOM v1 database. Histograms show the distribution of observations for DOC, DON, and DOP within defined concentration ranges in the continents where measurements are available. Maps were created using the GIS shapefile obtained from Laruelle et al. (2013).

abling in-depth analyses and providing quality-controlled input data for large-scale ecosystem models.

Author contributions. CL, CC, and XAÁS started the initiative and finalized the data compilation. All other co-authors contributed data. CL wrote the manuscript with input from all co-authors.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. Qi Chen is acknowledged for his very skilful help with Fig. 4. Cláudia B. Lopes thanks FCT for her researcher contract, 2021.03739.CEECIND/CP1659/CT0025 (<https://doi.org/10.54499/2021.03739.CEECIND/CP1659/CT0025>, PTCRIS, 2022–2024a), and project CICECO-Aveiro Institute of Materials, UIDB/50011/2020 (<https://doi.org/10.54499/UIDB/50011/2020>, PTCRIS, 2020–2023) & LA/P/0006/2020 (<https://doi.org/10.54499/LA/P/0006/2020>, PTCRIS, 2022–2024b), financed by national funds through the FCT/MCTES (PIDDAC).

Financial support. During the drafting of the manuscript, Christian Lønborg received funding from the Independent Research Fund Denmark (grant no. 1127-00033B). The monitoring data obtained from Bermuda received funding from the Bermuda Government Department of Environment and Natural Resources. A subset of the data obtained from UK coastal estuaries received funding from the Natural Environment Research Council (grant no. NE/N018087/1). Data retrieved from the Palmer LTER data were collected with support from the Office of Polar Programs, US National Science Foundation. Data obtained from the Great Barrier Reef Marine Monitoring Program for Inshore Water Quality, which is a partnership between the Great Barrier Reef Marine Park Authority, the Australian Institute of Marine Science, James Cook University, and the Cape York Water Partnership. The contribution by Piotr Kowalczyk was supported by DiSeDOM (project contract no. UMO-2019/33/B/ST10/01232) funded by the NCN – National Science Centre, Poland. Nicholas Ward and Allison Myers-Pigg participated in this synthesis effort with funding provided by the U.S. Department of Energy funded COMPASS-FME project; the provided data was collected with funding from the PREMIS Initiative, conducted under the Laboratory Directed Research and Development Program at Pacific Northwest National Laboratory. The data obtained from the Levantine Sea (Med Sea) received funding from the Scientific and Technological Research Council of Türkiye (TÜBİTAK, 1001 programme, grant no. 115Y629). The

data obtained from Gulf of Trieste (Slovenian waters) were financed by Research Program No. P1-0237 (Slovenian Research and Innovation Agency). Data obtained from the northern Baltic Sea were financed by the research programme EcoChange (Swedish research council FORMAS). The contribution by Ding He was supported by the National Natural Science Foundation of China (grant no. 42222061) and funding support from the Center for Ocean Research in Hong Kong and Macau (CORE). CORE is a joint research centre for ocean research between Laoshan Laboratory and HKUST. The contribution by Yuan Shen was supported by National Natural Science Foundation of China (grant no. 42106040), the Fundamental Research Funds for the Central Universities of China (grant no. 20720210076), and Fujian Provincial Central Guided Local Science and Technology Development Special Project (grant no. 2022L3078). The data provided by Luiz C. Cotovicz Jr. were supported by the Brazilian National Council of Research and Development (CNPq-Pve No. 401.726/2012-6) and by the Carlos Chagas Foundation for Research Support of the State of Rio de Janeiro (FAPERJ; No. E-26202.785/2016). Data provided by Stefano Cozzi were collected in the framework of Italian (PRISMA 1 and 2, ANOCSIA and VECTOR) and European (OCEANCERTAIN) research projects. Data provided by MG were collected within the Project “Mucilages in the Adriatic and Tyrrhenian Seas (MAT)”, coordinated by the Istituto Centrale per la Ricerca Scientifica e Tecnologica Applicata al Mare and financially supported by the Italian Ministry of the Environment. Data obtained from the Georgia coast (USA) were supported by the National Science Foundation through grants OCE-1832178 (GCE-LTER Program) and OCE-1902131. Observations in the southern Caribbean Sea including the Cariaco Basin were collected as part of the CARIACO Ocean Time Series programme (supported by the Consejo Nacional de Ciencia y Tecnología of Venezuela, the Ley de Ciencia, Tecnología e Innovación de Venezuela, the Estación de Investigaciones Marinas de Venezuela; the National Science Foundation (grant nos. OCE-0752139, OCE-9216626, OCE-9729284, OCE-491 9401537, OCE-9729697, OCE-9415790, OCE-9711318, OCE-0326268, OCE-0963028, OCE-0326313, and OCE-0326268, NASA grants NAG5-6448, NAS5-97128, and NNX14AP62A); the Inter-American Institute for Global Change Research grant IAI-CRN3094), and the Marine Biodiversity Observation Network/M-BON of the Group on Earth Observations Biodiversity Observation Network). Data provided by Digna Rueda-Roa and Bradley Eyre was supported by ARC Linkage (LP0770222), with Norske-Skog Boyer and the Derwent Estuary Program providing financial and in-kind assistance.

Review statement. This paper was edited by Xingchen Wang and reviewed by two anonymous referees.

References

Álvarez-Salgado, X. A., Nieto-Cid, M., and Rossel, P. E.: Dissolved Organic Matter, in: *Marine Analytical Chemistry*, edited by: Blasco, J., and Tovar-Sánchez, A., Springer International Publishing, Cham, 39–102, https://doi.org/10.1007/978-3-031-14486-8_2, 2023.

- Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. A.: The changing carbon cycle of the coastal ocean, *Nature*, 504, 61–70, <https://doi.org/10.1038/nature12857>, 2013.
- Benner, R. and Amon, R. M.: The size-reactivity continuum of major bioelements in the ocean, *Annu. Rev. Marine Sci.*, 7, 185–205, <https://doi.org/10.1146/annurev-marine-010213-135126>, 2015.
- Boudreau, B. P. and Ruddick, B. R.: On a reactive continuum representation of organic matter diagenesis, *Am. J. Sci.*, 291, 507–538, 1991.
- Burdige, D. J. and Komada, T.: Sediment pore waters, in: *Biogeochemistry of marine dissolved organic matter*, edited by: Hansen, D. A., and Carlson, C. A., Elsevier, 535–577, 2014.
- Carlson, C. A. and Hansell, D. A.: DOM Sources, Sinks, Reactivity, and Budgets, in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Carlson, C. A., and Hansell, D. A., Elsevier Science & Technology, 65–126, <https://doi.org/10.1016/b978-0-12-405940-5.00003-0>, 2015.
- Carreira, C., Talbot, S., and Lønborg, C.: Bacterial consumption of total and dissolved organic carbon in the Great Barrier Reef, *Biogeochemistry*, 489–508, <https://doi.org/10.1007/s10533-021-00802-x>, 2021.
- Cauwet, G.: DOM in coastal areas, in: *Biogeochemistry of Dissolved organic matter*, edited by: Hansell, D., and Carlson, C. A., Academic Press, London, 579–609, 2002.
- Chin, W. C., Orellana, M. V., and Verdugo, P.: Spontaneous assembly of marine dissolved organic matter into polymer gels, *Nature*, 391, 568–572, 1998.
- Foreman, R. K., Björkman, K. M., Carlson, C. A., Opalk, K., and Karl, D. M.: Improved ultraviolet photo-oxidation system yields estimates for deep-sea dissolved organic nitrogen and phosphorus, *Limnol. Oceanogr.-Methods*, 17, 277–291, <https://doi.org/10.1002/lom3.10312>, 2019.
- García-Martín, E. E., Sanders, R., Evans, C. D., Kitidis, V., Lapworth, D. J., Rees, A. P., Spears, B. M., Tye, A., Williamson, J. L., Balfour, C., Best, M., Bowes, M., Breimann, S., Brown, I. J., Burden, A., Callaghan, N., Felgate, S. L., Fishwick, J., Fraser, M., Gibb, S. W., Gilbert, P. J., Godsell, N., Gomez-Castillo, A. P., Hargreaves, G., Jones, O., Kennedy, P., Lichtschlag, A., Martin, A., May, R., Mawji, E., Mounteny, I., Nightingale, P. D., Olszewska, J. P., Painter, S. C., Pearce, C. R., Pereira, M. G., Peel, K., Pickard, A., Stephens, J. A., Stinchcombe, M., Williams, P., Woodward, E. M. S., Yarrow, D., and Mayor, D. J.: Contrasting Estuarine Processing of Dissolved Organic Matter Derived From Natural and Human-Impacted Landscapes, *Global Biogeochem. Cycles*, 35, e2021GB007023, <https://doi.org/10.1029/2021GB007023>, 2021.
- Halewood, E., Opalk, K., Custals, L., Carey, M., Hansell, D. A., and Carlson, C. A.: Determination of dissolved organic carbon and total dissolved nitrogen in seawater using High Temperature Combustion Analysis, *Front. Marine Sci.*, 9, 1061646, <https://doi.org/10.3389/fmars.2022.1061646>, 2022.
- Hansell, D. A.: Dissolved Organic Carbon Reference Material Program, *Eos, Transactions American Geophysical Union*, 86, 318–318, <https://doi.org/10.1029/2005EO350003>, 2005.
- Hansell, D. A., Carlson, C. A., Amon, R. M. W., Álvarez-Salgado, X. A., Yamashita, Y., Romera-Castillo, C., and Bif, M. B.: Compilation of dissolved organic matter (DOM) data obtained from global ocean observations from 1994 to 2021. Version 2, NCEI Accession 0227166 [data set], <https://doi.org/10.25921/s4f4-ye35>, 2021.
- Iavorivska, L., Boyer, E. W., and DeWalle, D. R.: Atmospheric deposition of organic carbon via precipitation, *Atmos. Environ.*, 146, 153–163, <https://doi.org/10.1016/j.atmosenv.2016.06.006>, 2016.
- Karl, D. M. and Björkman, K. M.: Dynamics of Dissolved Organic Phosphorus, in: *Biogeochemistry of marine dissolved organic matter*, edited by: Hansell, D. A., and Carlson, C. A., 233–334, <https://doi.org/10.1016/b978-0-12-405940-5.00005-4>, 2015.
- Kerner, M., Hohenberg, H., Ertl, S., Reckermann, M., and Spitz, A.: Self-organization of dissolved organic matter micelle-like microparticles in river water, *Nature*, 422, 150–154, 2003.
- Knapp, A. N., Sigman, D. M., and Lipschultz, F.: N isotopic composition of dissolved organic nitrogen and nitrate at the Bermuda Atlantic Time-series Study site, *Global Biogeochem. Cycles*, 19, 1–15, <https://doi.org/10.1029/2004GB002320>, 2005.
- Laruelle, G. G., Dürr, H. H., Lauerwald, R., Hartmann, J., Slomp, C. P., Goossens, N., and Regnier, P. A. G.: Global multi-scale segmentation of continental and coastal waters from the watersheds to the continental margins, *Hydrol. Earth Syst. Sci.*, 17, 2029–2051, <https://doi.org/10.5194/hess-17-2029-2013>, 2013.
- Laurenceau-Cornec, E. C., Trull, T. W., Davies, D. M., De La Rocha, C. L., and Blain, S.: Phytoplankton morphology controls on marine snow sinking velocity, *Marine Ecol. Prog. Ser.*, 520, 35–56, 2015.
- Liang, Z., McCabe, K., Fawcett, S. E., Forrer, H. J., Hashihama, F., Jeandel, C., Marconi, D., Planquette, H., Saito, M. A., Sohm, J. A., Thomas, R. K., Letscher, R. T., and Knapp, A. N.: A global ocean dissolved organic phosphorus concentration database (DOPv2021), *Sci. Data*, 9, 772, <https://doi.org/10.1038/s41597-022-01873-7>, 2022.
- Lønborg, C. and Álvarez-Salgado, X. A.: Recycling versus export of bioavailable dissolved organic matter in the coastal ocean and efficiency of the continental shelf pump, *Global Biogeochem. Cycles* 26, GB3018, <https://doi.org/10.1029/2012GB004353>, 2012.
- Lønborg, C., Álvarez-Salgado, X. A., Davidson, K., and Miller, A. E. J.: Production of bioavailable and refractory dissolved organic matter by coastal heterotrophic microbial populations, *Estuar. Coast. Shelf Sci.*, 82, 682–688, <https://doi.org/10.1016/j.ecss.2009.02.026>, 2009.
- Lønborg, C., Álvarez-Salgado, X. A., Duggan, S., and Carreira, C.: Organic matter bioavailability in tropical coastal waters: The Great Barrier Reef, *Limnol. Oceanogr.*, 63, 1015–1035, <https://doi.org/10.1002/lno.10717>, 2018.
- Lønborg, C., Carreira, C., Jickells, T., and Álvarez-Salgado, X. A.: Impacts of Global Change on Ocean Dissolved Organic Carbon (DOC) Cycling, *Front. Marine Sci.*, 7, 466, <https://doi.org/10.3389/fmars.2020.00466>, 2020.
- Lønborg, C., Carreira, C., Abril, G., Agustí, S., Amaral, V., Andersson, A., Aristegui, J., Bhadury, P., Bif, M. B., Borges, A. V., Bouillon, S., Calleja, M. L., Cotovicz Jr., L. C., Cozzi, S., Doval, M. D., Duarte, C. M., Eyre, B. D., Fichot, C. G., García-Martín, E. E., Garzon-Garcia, A., Giani, M., Gonçalves-Araujo, R., Gruber, R., Hansell, D. A., Hashihama, F., He, D., Holding, J. M., Hunter, W. R., Ibáñez, J. S., Ibello, V., Jiang, S., Kim, G., Klun, K., Kowalczyk, P., Kubo, A., Weng Lee, C., Lopes, C., Maggioni, F., Magni, P., Marrase, C., Martin, P., McCallister, S. L.,

- McCallum, R., Medeiros, P. M., Morán, X. A. G., Muller-Karger, F., Myers-Pigg, A., Norli, M., Oakes, J. M., Osterholz, H., Park, H., Paulsen, M. L., Rosentreter, J. Ross, J. A., Rueda-Roa, D., Santinelli, C., Shen, Y., Teira, E., Tinta, T., Uher, G., Wakita, M., Ward, N., Watanabe, K., Xin, Y., Yamashita, Y., Yang, L., Yeo, J., Yuan, H., Zheng, Q., and Álvarez-Salgado, X. A.: A global database of dissolved organic matter (DOM) concentration measurements in coastal waters (CoastDOM v1), PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.964012>, 2023.
- Mackenzie, F. T., De Carlo, E. H., and Lerman, A.: 5.10 – Coupled C, N, P, and O Biogeochemical Cycling at the Land–Ocean Interface, in: *Treatise on Estuarine and Coastal Science*, edited by: Wolanski, E., and McLusky, D., Academic Press, Waltham, 317–342, <https://doi.org/10.1016/B978-0-12-374711-2.00512-X>, 2011.
- Mopper, K., Kieber, D. J., and Stubbins, A.: Marine Photochemistry of Organic Matter, in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Carlson, C. A., and Hansell, D. A., 389–450, <https://doi.org/10.1016/b978-0-12-405940-5.00008-x>, 2015.
- Osterholz, H., Burmeister, C., Busch, S., Dierken, M., Frazão, H. C., Hansen, R., Jeschek, J., Kremp, A., Kreuzer, L., and Sadkowiak, B.: Nearshore dissolved and particulate organic matter dynamics in the southwestern Baltic Sea: environmental drivers and time series analysis (2010–2020), *Front. Marine Sci.*, 8, 795028, <https://doi.org/10.3389/fmars.2021.795028>, 2021.
- PTCRIS: CICECO-Instituto de Materiais de Aveiro, Universidade de Aveiro, <https://doi.org/10.54499/UIDB/50011/2020>, 2020–2023.
- PTCRIS: GraphChem – Providing solutions for clean water production and recycling of technological critical elements: from ‘the wonder material’ to chemobionics systems, Universidade de Aveiro, <https://doi.org/10.54499/2021.03739.CEECIND/CP1659/CT0025>, 2022–2024a.
- PTCRIS: CICECO – Instituto de Materiais de Aveiro, Universidade de Aveiro, <https://doi.org/10.54499/LA/P/0006/2020>, 2022–2024b.
- Raymond, P. A. and Spencer, R. G. M.: Riverine DOM, in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Hansell, D. A., and Carlson, C. A., Elsevier, Amsterdam, 509–533, <https://doi.org/10.1016/b978-0-12-405940-5.00011-x>, 2015.
- Redfield, A. C., Ketchum, B. K., and Richards, F. A.: The influence of organisms on the composition of sea-water, in: *The sea*, vol. 2, *The composition of sea water: Comparative and descriptive oceanography*, edited by: Hill, M. N., Wiley-Interscience, 26–77, 1963.
- Santos, I. R., Burdige, D. J., Jennerjahn, T. C., Bouillon, S., Cabral, A., Serrano, O., Wernberg, T., Filbee-Dexter, K., Guimond, J. A., and Tamborski, J. J.: The renaissance of Odum’s outwelling hypothesis in “Blue Carbon” science, *Estuar. Coast. Shelf Sci.*, 255, 107361, <https://doi.org/10.1016/j.ecss.2021.107361>, 2021.
- Sharp, J. H., Benner, R., Bennett, L., Carlson, C. A., Dow, R., and Fitzwater, S. E.: Re-evaluation of high temperature combustion and chemical oxidation measurements of dissolved organic carbon in seawater, *Limnol. Oceanogr.*, 38, 1774–1782, <https://doi.org/10.4319/lo.1993.38.8.1774>, 1993.
- Sharp, J. H., Rinker, K. R., Savidge, K. B., Abell, J., Benaim, J. Y., Bronk, D., Burdige, D. J., Cauwet, G., Chen, W., Doval, M. D., Hansell, D., Hopkinson, C., Kattner, G., Kaumeyer, N., McGlathery, K. J., Merriam, J., Morley, N., Nagel, K., Ogawa, H., Pollard, C., Pujo-Pay, M., Raimbault, P., Sambrotto, R., Seitzinger, S., Spyres, G., Tirendi, F., Walsh, T. W., and Wong, C. S.: A preliminary methods comparison for measurement of dissolved organic nitrogen in seawater, *Marine Chem.*, 78, 171–184, 2002.
- Sipler, R. E. and Bronk, D. A.: Dynamics of Dissolved Organic Nitrogen, in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Hansell, D. A., and Carlson, C. A., 127–232, <https://doi.org/10.1016/b978-0-12-405940-5.00004-2>, 2015.
- Solorzano, L. and Sharp, J. H.: Determination of total dissolved phosphorus and particulate phosphorus in natural waters, *Limnol. Oceanogr.*, 25, 754–758, 1980.
- Taniguchi, M., Dulai, H., Burnett, K. M., Santos, I. R., Sugimoto, R., Stieglitz, T., Kim, G., Moosdorf, N., and Burnett, W. C.: Submarine Groundwater Discharge: Updates on Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects, *Front. Environ. Sci.*, 7, 141, <https://doi.org/10.3389/fenvs.2019.00141>, 2019.
- Valderrama, J. C.: The simultaneous analysis of total nitrogen and total phosphorus in natural waters, *Marine Chem.*, 10, 109–122, 1981.
- Wada, S., Aoki, M. N., Mikami, A., Komatsu, T., Tsuchiya, Y., Sato, T., Shinagawa, H., and Hama, T.: Bioavailability of macroalgal dissolved organic matter in seawater, *Marine Ecol. Prog. Ser.*, 370, 33–44, 2008.