



University of Dundee

Diffusion Weighted Imaging and Grading of Brain Tumours

Changmai, Manah; Reza, Mohammed Faruque; Gohain, Kastury

DOI:
[10.16965/ijar.2023.142](https://doi.org/10.16965/ijar.2023.142)

Publication date:
2024

Licence:
CC BY-NC-SA

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Changmai, M., Reza, M. F., & Gohain, K. (2024). Diffusion Weighted Imaging and Grading of Brain Tumours. *International Journal of Anatomy and Research*, 12(1), 8800-8807. <https://doi.org/10.16965/ijar.2023.142>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Diffusion Weighted Imaging and Grading of Brain Tumours

Manah Chandra Changmai^{1a,b}, Mohammed Faruque Reza², Kastury Gohain³.

*^{1a}Lecturer, Centre for Anatomy and Human Identification, University of Dundee, Dundee, United Kingdom, ^b Universiti Sains, Malaysia.

² Senior Lecturer, Department of Neuroscience, School of Medicine, Universiti Sains Malaysia, Kubang kerian, Kelantan, Malaysia.

³ Lecturer, School of Business, University of Dundee, Dundee, United Kingdom.

ABSTRACT

Background: Most prevalent primary cerebral tumours are meningiomas. The other frequent intracranial tumours are pituitary adenomas, which are benign, and gliomas, which are intra-axial brain tumours. The objective of this study is to understand the importance of DW MRI imaging with standard b-value in differentiating presurgical grading of the brain tumour.

Design: A total of 24 DWI patients, including 12 meningiomas, 8 gliomas, and 4 pituitary adenomas, were included in this retrospective analysis.

Method: The Stejskal-Tanner equation is used to analyse the ADC_{mean} , ADC_{min} , and ADC_{max} values from the healthy and tumour core that are obtained out from area of interest (ROI).

Result: The ADC_{mean} value of Gliomas ranges from $0.09 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ to $0.99 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ with a median value of $0.25 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. ADC_{mean} value $1.82 \times 10^{-3} \text{ mm}^2/\text{s}$ (sensitivity: 67%. Specificity: 81.8%) and $0.94 \times 10^{-3} \text{ mm}^2/\text{s}$ (sensitivity: 75%. specificity: 81.3%) can discriminate grade II –IV meningioma from grade II-IV glioma.

Conclusion: The ADC and its threshold levels offer crucial details on the grades, consistency, and characterization of tumour, aiding accurate diagnosis and therapy.

KEYWORDS: Brain tumour, Imaging, MRI, DWI, ADC, b-value.

Corresponding Author: Dr Manah Chandra Changmai, PhD Graduate Universiti Sains Malaysia, Malaysia. Lecturer, Centre for Anatomy and Human Identification, University of Dundee, Dundee, United Kingdom. **E-Mail:** manahchangmai@gmail.com

Access this Article online	Journal Information
Quick Response code  DOI: 10.16965/ijar.2023.142	International Journal of Anatomy and Research ISSN (E) 2321-4287 ISSN (P) 2321-8967 https://www.ijmhr.org/ijar.htm DOI-Prefix: https://dx.doi.org/10.16965/ijar 
	Article Information
	Received: 30 Aug 2023 Peer Review: 02 Sep 2023 Revised: 12 Dec 2023
	Accepted: 20 Jan 2024 Published (O): 05 Mar 2024 Published (P): 05 Mar 2024

INTRODUCTION

The meningiomas are the most common primary intracranial tumours and represents 36% of central nervous system (CNS) tumour [1]. The gliomas are intra-axial brain tumour located frequently in the supratentorial region and accounts for 3.4% of all intracranial tumours [2]. The pituitary adenomas are mainly benign slow growing tumour arising from the pituitary gland. They are intensifying

accompanies with disorder of hormones [3].

These tumours are frequently identified by histopathological examination. However, an improper selection of area for biopsy sample from brain tumours leads to error in microscopic diagnosis. Magnetic resonance imaging (MRI) is the noninvasive procedure suitable in resolving, management and forecasting prognosis of brain tumours. The Diffusion weighted imaging (DWI) is a

technique in magnetic resonance imaging which has brought distinctive change in imaging. It catches small motions of water molecules augmenting expression of intrinsic directionality in the brain [4]. Our body spaces contain water which provides a considerable element in body weight.

The diffusion of water molecules in tissues follows a definite pattern that captures the physiology by and reflects the difference in rate of diffusion in between tissues. The DWI is advised to trace the distinction between a benign and malignant brain tumour. Diffusion imaging predicated on ADC has indeed been tested for tumour assessment throughout the past two decades [5,6]. It has been proposed that analyzing the ADC of the lesion data could be of predictive as well as diagnostic significance [7,8]. Furthermore, preoperative diffusion values appear to be useful for differentiating radiation-induced brain injury from advanced disease and prognostic of clinical response to radiotherapy [8-11]. The Apparent diffusion coefficient (ADC) is the quantification of proportions of water molecules in the tissue analysed with DWI. The malignant tumour exhibit lower ADC value compared to benign tumours [12].

The values of ADC lower than $1 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ considered to be a malignant tumour [13]. However, some benign tumours demonstrate low ADC values and identified as malignant tumour. The value of ADC is considerably inversely proportional to cellularity of the brain tumour [14]. An improper selection of area for biopsy sample leads to error in microscopic diagnosis. The ADC values are utilized as a guide to forecast the outcome of the treatment in malignant tumours. It is necessary to identify the grade of the brain tumour and its expression possibilities with the aid of diffusion weighted imaging (DWI). The objective of this study is to analyze the apparent diffusion coefficient (ADC) value in distinguishing three common tumours that includes Meningioma, Gliomas and Pituitary adenomas.

Aim: The aim of this study is to understand the importance of DW MRI imaging with standard b-value in differentiating presurgical

grading of the brain tumour.

METHOD

Study Design: It is a retrospective investigation of ADC values of three tumours. The cases of meningioma and glioblastoma were identified to persuade the research. A patient age limit of 5 years to 70 years was chosen comprising both genders and different races of Malaysia. A total of 33 patients (n=33) were selected for the study. These patients were diagnosed with brain tumour. Out of these 33 patients, 12 (n=12) meningiomas, 8 (n=8) glioma and 4 (n=4) pituitary adenoma patients underwent DWI. Patients with head injury, demyelinating diseases and meningitis were excluded from the study. The patient's data was collected from their case file (ARCHIVE) from Department of Neuroscience and the images of DWI were recovered from a public University hospital in Malaysia's picture archiving and communication system (PACS).

DWI images: The MRI of the patients were performed in Radiology in a public university hospital in Malaysia with Philips ACHEVA 3.0 Tesla MRI machine. A spin echo pulse sequence of [TR/TE] of 3433/93.8, flip angle 90° and 5mm slice thickness was stipulated in an axial plane. In DWI, magnitude of diffusion gradient (b value) of $b = 0 \text{ s/mm}^2$ and b-value of 1000 s/mm^2 is applied in X, Y and Z axis maintaining a space of 6ms separating two slices.

Image Evaluation: A skilled radiologist and a well-trained radiographer retrospectively evaluated the DWI images A region of interest ROI were placed in the tumour core and another ROI was placed on the contralateral healthy area of the cerebral hemisphere to determine the apparent diffusion coefficient (ADC) which is estimated with Stejskal-Tanner equation (15). $ADC = -(1/b)\ln(S/S_0)$ where S_0 is intensity of the signal with gradient factor $b=0$; S is intensity of the signal with gradient factor $b=1000 \text{ mm}^2/\text{s}$; \ln is natural algorithm and b in $1/b$ is 1000; The Values of ADC were expressed were illustrated with $10^{-3} \text{ mm}^2/\text{s}$

Statistical evaluation: The analysis of the data was carried out with SPSS version 23. The accumulated information of ADC values was estimated by descriptive analysis. Box plot and




Fig. 1: A 56-year-old female with features of Grade I meningioma A) DWI at b_0 showing a hyperintense mass in the left anterior parasagittal region B) DWI at b_{1000} showing hypointense mass in left frontal region.




Fig. 2: A 48-year-old male with features of Grade I Glioma A) DWI at b_0 showing a isointense to hyperintense mass in the right posterior parietal lobe B) DWI at b_{1000} showing hypointense mass in right posterior parietal lobe.




Fig. 3: A 46-year-old female with features of Grade I pituitary adenoma A) DWI at b_0 showing a hyperintense mass in the suprasellar region B) DWI at b_{1000} showing hypointense mass in the suprasellar region.

ROC curve analysis was performed to compare the tumour core ADC values of meningioma, glioma, and pituitary adenoma. The 2-tailed t test values of ADC of Meningiomas, Gliomas and Pituitary adenomas. The recorded variability was considered statistically significant if $P < 0.05$. Normalized ADC (NADC) was calculated in every case as a ratio $ADC_{mean} \text{ meningioma} / ADC_{mean} \text{ white matter}$.

RESULTS

Meningioma: The ADC_{mean} value of meningioma ranges from $0.15 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ to $2.55 \times$

$10^{-3} \text{ mm}^2\text{s}^{-1}$. Meningioma grade I has an ADC_{mean} value of $0.79 \pm 0.76 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$, while meningioma grades II to IV have an ADC_{mean} value of $0.16 \pm 0.50 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. In high grade compared to grade I meningioma, the ADC_{mean} is lower. Meningioma grade I has an ADC_{min} value of $0.72 \pm 0.86 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$, and meningioma grades II through IV have an ADC_{min} value of $0.15 \pm 0.07 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. Meningioma grade I has an ADC_{max} value of $0.61 \pm 0.67 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$, and meningioma grades II through IV have an ADC_{max} value of $0.17 \pm 0.10 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. Meningioma grade I has a

Table 1: ADC_{mean}, ADC_{min}, ADC_{max}, and NADC values in three different brain tumours.

	Meningioma		Glioma		Pituitary adenoma
	Grade I	Grade II-IV	Grade I	Grade II-IV	Grade I
ADC _{mean}	0.79 ± 0.76	0.16 ± 0.50	0.40 ± 0.32	0.20 ± 0.14	0.74 ± 0.33
ADC _{min}	0.72 ± 0.86	0.15 ± 0.07	0.50 ± 0.56	0.13 ± 0.12	0.82 ± 1.53
ADC _{max}	0.61 ± 0.67	0.17 ± 0.10	0.33 ± 0.18	0.28 ± 0.24	0.51 ± 0.84
NADC	0.60 ± 0.47	0.07 ± 0.03	0.19 ± 0.15	0.14 ± 0.05	0.57 ± 1.03

Table 2: The mean ADC values of different grade of brain tumours using standard b-values by diffusion weighted imaging.

	Area under curve(AUC)(95%CI)	P value	Cut off ADC value	Sensitivity (%) (95%CI)	Specificity (%) (95%CI)	Younde-n Index
ADC _{mean} Meningioma G1	0.8(65.2-97.8)	0.02	0.88x10 ⁻³	72.0 (64.0-100.0)	81.4(64.3-8.7)	0.54
ADC _{mean} Meningioma G2, G4	0.73 (56.0-82.1)	0.25	1.82x10 ⁻³	67.0 (50.0-100.0)	81.8(54.5-90.9)	0.56
ADC _{min} Meningioma G1	0.71 (52.1-87.2)	0.29	0.85x10 ⁻³	80.0(60.0-90.0)	78.6(57.1-85.7)	0.64
ADC _{min} Meningioma G2,G4	0.69.5(51.0-72.5)	0.52	0.74x10 ⁻³	75.1(57.3-86.1)	86.4(63.3-95.5)	0.62
ADC _{max} Meningioma G1	0.93(73.5-95.3)	0.34	0.90x10 ⁻³	78.3(56.9-90.0)	72.4(57.1-85.7)	0.67
ADC _{max} Meningioma G2, G4	73.5(53.0-85.0)	0.45	0.85x10 ⁻³	70.0(55.4-84.8)	86.4(59.1-100.0)	0.66
ADC _{mean} Glioma G1,G4	0.82(72.0-94.0)	0.62	0.94x10 ⁻³	75.0(64.3-85.5)	81.3(56.3-100.0)	0.68
ADC _{min} Glioma G1,G4	0.80(65.1-90.0)	0.36	0.87x10 ⁻³	76.0(62.5-87.5)	75.0(62.8-100.0)	0.63
ADC _{max} Glioma G1,G4	0.85(68.0-92.8)	0.54	0.90x10 ⁻³	87.5(52.3-1.00)	87.5(55.4-100.0)	0.62

NADC value of $0.60 \pm 0.47 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$, and meningioma grades II through IV have a value of $0.07 \pm 0.03 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$.

Gliomas

The ADC_{mean} value of Gliomas ranges from $0.09 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ to $0.99 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ with a median value of $0.25 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. The ADC_{mean} value for grades I-IV glioma is $0.40 \pm 0.32 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ and $0.20 \pm 0.14 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ respectively. In high grade compared to grade I glioma, the ADC_{mean} is lower. The ADC_{min} value for gliomas in grades I through IV is $0.50 \pm 0.56 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ and $0.13 \pm 0.12 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ respectively. The ADC_{max} value for gliomas in grades I through IV is $0.33 \pm 0.18 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ and $0.28 \pm 0.24 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ respectively. Meningioma grade I has a NADC value of $0.19 \pm 0.15 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$, and meningioma grades II through IV have a value of $0.14 \pm 0.05 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$.

Pituitary adenoma: The ADC_{mean} value of Pituitary adenoma ranges from $0.60 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ to $2.77 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$ with a median value of $0.19 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. Grade I pituitary adenomas are recognized. The value of its ADC_{mean} is $0.74 \pm 0.33 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$.

The tumours ADC_{min} mean value is $0.82 \pm 1.53 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. These tumours ADC_{max} value is calculated to be $0.51 \pm 0.84 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$. The tumor’s NADC value is $0.57 \pm 1.03 \times 10^{-3} \text{ mm}^2\text{s}^{-1}$.

According to Table I, the ADC_{mean}, ADC_{min}, ADC_{max}, and NADC values were higher in Grade I meningioma compared to Grade I pituitary adenomas. Grade II-IV meningiomas have lower values than Grade II-IV gliomas.

Only ADC_{mean} for Grade I meningioma from the ROC curve analysis were statistically significant, despite the fact that there were some statistically significant disparities across the grades.

DISCUSSION

DWI is crucial for imaging brain tumour, particularly for finding their grade. This MRI modality is also essential to determine the outcome of the management of these tumours. There are few investigations on meningiomas, gliomas and pituitary adenomas on quality of their diffusion and differentiation between their grades in accordance with DWI and ADC values [16]. ADC offers critical extra information beyond what MRI can. This study has

demonstrated that there are differences in ADC grades between meningioma, glioblastoma, and pituitary adenoma, but not enough to differentiate between the different primary cancer types [17,18].

This could be because of the ADC expressing more universal traits like cellular proliferation. ADC levels often are less in primary brain tumours with more cellular proliferation in contrast to normal brain parenchyma [19,20].

In present study the ADC_{mean} of grade I meningioma is slightly higher than glioblastoma. In a study in 39 meningioma patients, the ADC_{mean} was higher in lower grade compared to the higher grade. However, when compared to benign lesions, atypical and malignant meningiomas exhibits lower ADC values [21,22]. ADC_{mean} is less relevant to tumour cellularity than ADC_{min} [23]. ADC_{mean} value may be used to identify cancers with a high differentiation activity using the specified threshold. Many explanations, including greater tumour cellularity, tumour pattern, fibrous or gliotic tissues, or a combination of these characteristics, have been put up to explain the lower ADC in high-grade tumours [24]. Analyses of ADC values would make a big difference in the identification and differential diagnosis. Main benefit include not requiring contrast material and quick and easy imaging capture within seconds [25]. When employing a typical b-value DW-MRI, a minimum ADC threshold of $1.070 \times 10^{-3} \text{ mm}^2/\text{s}$ gave 79.7% sensitivity and 60.0% specificity is pertinent for identifying high-grade gliomas(26). The results of this study indicated an ADC_{min} threshold of $0.74 \times 10^{-3} \text{ mm}^2/\text{s}$ (sensitivity: 75%. Specificity: 86.4%) and $0.87 \times 10^{-3} \text{ mm}^2/\text{s}$ (sensitivity:76%. specificity:75%) to discriminate grade II –IV meningioma from grade II-IV glioma. Additionally, ADC_{mean} value $1.82 \times 10^{-3} \text{ mm}^2/\text{s}$ (sensitivity: 67%. Specificity: 81.8%) and $0.94 \times 10^{-3} \text{ mm}^2/\text{s}$ (sensitivity:75%. specificity:81.3%) to discriminate grade II –IV meningioma from grade II-IV glioma. A similar study with standard b-value ADC images and a histogram estimation of the whole tumour volume was conducted to distinguish different glioma grading. According to their findings, grades II, III, and IV gliomas had distinct ADCmin

values for standard b-value scans. They also found significant difference of ADC_{mean} for standard b-value between Grade II-IV tumour (27). The present study has shown a significant difference in ADC_{mean} that can discriminate Grade II-IV meningioma from grade II-IV glioma. The minimal level ADC value from $b = 1000 \text{ DW-MRI}$ inside that solid tumour region was substantially lower in high-grade (III and IV) gliomas than low-grade (II) gliomas at 3T, and a cutoff value of $0.90 \times 10^{-3} \text{ mm}^2/\text{s}$ was recommended to distinguish high and low-grade gliomas with 85% specificity and 71% specificity. The ADC_{max} is also higher in Grade I meningioma than Grade I pituitary adenoma. The ADC_{min} has been thoroughly investigated for glioma grading with 1.5T, high-grade (III and IV) and low-grade (I and II) tumours could be distinguished with 95% sensitivity and 80.6% specificity using a threshold of 1.48 for the lowest ADC within the tumour adjusted with the contralateral side collected using a standard b-value [28].

Strong tissue mechanisms and substantial water flow across capillaries hinder the tumor's definition. The ADC values are impacted by myelin, which has a stronger effect on the diffusing pattern. Despite the fact that various factors influence ADC values in brain tissues [29]. As an indicator for tumour cellularity, DWI is useful to discriminate between benign and malignant tumours by evaluating the ADC. It is also used to investigate heterogeneity [30].

The box plot Figure (A) depicts that NADC decreased from 0.25 units in Meningioma grade I to about 0.10 units. There is no significant increase for Glioma grade I – grade IV but potential increase is observed for Pituitary adenoma grade1 which ranged more than 2.0 units. There also appears to be major decrease in median NADC for Pituitary adenoma grade1 compared to Meningioma grade I- grade IV, and Glioma grade I – grade IV which maintained consistent spread out. However, there is one outlier with high NADC in Glioma grade I – grade IV. This demonstrates that NADC values are significantly varies between Meningiomas, Gliomas and pituitary adenomas. In Figure (B) the boxplot exhibits




Fig. 4: Boxplots of ADC values of different grades of (A) Meningioma (B) Glioma (C) pituitary adenoma.

that ADC_{mean} for Meningioma grade I – grade IV and Glioma grade I- grade IV having same median unit. Consequently, Pituitary adenoma grade I showing low median unit. There is one high ADC_{mean} for Glioma grade I-Grade IV whereas there are no extreme indicators for the rest of the tumours. In Figure (C) Similarly, for ADC_{min} , Pituitary adenoma grade I has significant rise of patients compared to Meningioma and Glioma. There are high outliers for both Meningioma grade I and Glioma grade I- grade IV while the same spread for Meningioma grade I – grade IV and Glioma grade I – grade IV is noticed. Pituitary adenoma grade I specify uneven distribution with median lying at the bottom of the box.

This study has few limitations. The patient group is wide and there are only a limited

number of patients. The use of ROI analysis to examine the performance of ADC derived parameters in tumor tissue areas for tumor grading was another limitation. Other limitation is that it is a retrospective study. The study only included one centre and had a smaller batch size of brain tumours. It is advised to do more research using a larger tumour collection and a multicenter strategy.

CONCLUSION

The study revealed numerous correlations between various DWI data. The Grade I meningioma had statistically significant higher ADC_{mean} than Grade I glioma. Overall, the findings of the present investigation point to the possibility that the ADC parameters obtained from conventional b-value DWI MRI

can be used to predict the grade of a tumour. The effectiveness of b-value of DW-MRI in the pre-operative screening of brain tumours needs to be confirmed in future investigations that should include more comparable patient populations.

ACKNOWLEDGEMENTS

The authors want to thank Department of Radiology and Department of neurosciences of public university hospital to support the research.

Source of funding: The present research is self-funded, and no grant has been received from financing organizations in the public, private, or not-for-profit sectors.

Ethical Statement: The University has given ethical consent to perform the research with code JEPeM/20080420.

ORCID

Manah Chandra Changmai: 0000-0001-6440-2752

Mohammed Faruque Reza: 0000-0002-0302-6623

Kastury Gohain: 0000-0002-5190-8620

Author Contributions

The authors affirm that everyone listed is eligible to be an author and have checked the article for plagiarism. If plagiarism is found, all authors will be held accountable on an equal basis. The study was created and prepared by **MC, MFR and KG**, who also carried out the research, supplied the research tools, and gathered and organised the data. Data were examined and interpreted by **MC**. The article's first and last drafts were written by **MC**, who also helped with the logistics. The content and similarity score of the paper are the responsibility of all authors, who also gave the final text a critical assessment and approval.

Conflicts of Interests: None

REFERENCES

- [1]. Ostrom QT, Ph D, Gittleman H, Truitt G, Boscia A, Kruchko C, et al. Neuro-Oncology CBTRUS Statistical Report/ : Primary Brain and Other Central Nervous System Tumors Diagnosed in the. 2018;20:1-86.
<https://doi.org/10.1093/neuonc/noy131>
PMid:30445539 PMCID:PMC6129949
- [2]. Grossman R, Ram Z. Literature Review Posterior Fossa Intra-Axial Tumors in Adults. World Neurosurg [Internet]. 2015;88:140-5.
<https://doi.org/10.1016/j.wneu.2015.12.066>
PMid:26743385
- [3]. Benitez-rosario MA, Mcdarby G, Doyle R, Fabby C. Chronic Cluster-Like Headache Secondary to Prolactinoma/ : Uncommon Cephalalgia in Association with Brain Tumors. J Pain Symptom Manage [Internet]. 2009;37(2):271-6.
<https://doi.org/10.1016/j.jpainsymman.2008.02.013>
PMid:18694630
- [4]. Changmai MC, Reza MF, Idris Z, Kandasamy RA, Gohain K, others. The Diffusion Weighted Imaging and Its Contribution to the Distinction of Brain Tumour. Int Med J. 2021;28(4).
- [5]. Kang Y, Choi SH, Kim Y-J, Kim KG, Sohn C-H, Kim J-H, et al. Gliomas: histogram analysis of apparent diffusion coefficient maps with standard-or high-b-value diffusion-weighted MR imaging-correlation with tumor grade. Radiology. 2011;261(3):882-90.
<https://doi.org/10.1148/radiol.11110686>
PMid:21969667
- [6]. Raab P, Hattingen E, Franz K, Zanella FE, Lanfermann H. Cerebral gliomas: diffusional kurtosis imaging analysis of microstructural differences. Radiology. 2010;254(3):876-81.
<https://doi.org/10.1148/radiol.09090819>
PMid:20089718
- [7]. Rumboldt Z, Camacho DLA, Lake D, Welsh CT, Castillo M. Apparent diffusion coefficients for differentiation of cerebellar tumors in children. Am J Neuroradiol. 2006;27(6):1362-9.
- [8]. Berghoff AS, Spanberger T, Ilhan-Mutlu A, Magerle M, Hutterer M, Woehrer A, et al. Preoperative Diffusion-Weighted Imaging of Single Brain Metastases Correlates with Patient Survival Times. PLoS One. 2013;8(2):1-8.
<https://doi.org/10.1371/journal.pone.0055464>
PMid:23393579 PMCID:PMC3564927
- [9]. Mardor Y, Roth Y, Ocherashvili A, Spiegelmann R, Tichler T, Daniels D, et al. Pretreatment Prediction of Brain Tumors Response to Radiation Therapy Using High b -Value Diffusion-Weighted MRI 1. Neoplasia [Internet]. 2004;6(2):136-42.
<https://doi.org/10.1593/neo.03349>
PMid:15140402 PMCID:PMC1502089
- [10]. Hein PA, Eskey CJ, Dunn JF, Hug EB. Diffusion-Weighted Imaging in the Follow-up of Treated High-Grade Gliomas/ : Tumor Recurrence versus Radiation Injury. 2004;2(February):201-9.
- [11]. Asao C, Korogi Y, Kitajima M, Hirai T, Baba Y, Makino K. Diffusion-Weighted Imaging of Radiation- Induced Brain Injury for Differentiation from Tumor Recurrence. 2005;(July):1455-60.
- [12]. Subhawong TK, Durand DJ, Thawait GK, Jacobs MA, Fayad LM. Characterization of soft tissue masses/ : can quantitative diffusion weighted imaging reliably distinguish cysts from solid masses/ ? 2013;1583-92.
<https://doi.org/10.1007/s00256-013-1703-7>

- [13]. Wang J, Takayama F, Kawakami S, Saito A, Matsushita T, Momose M. Head and Neck Imaging Head and Neck Lesions/ : Characterization with. :621-30.
- [14]. Caldas-magalhaes J, Kooij N. Imaging in Laryngeal and Hypopharyngeal Carcinoma/ : Association between Apparent Diffusion Coefficient and Histologic. 2014;272(2).
<https://doi.org/10.1148/radiol.14131173>
PMid:24749712
- [15]. Sener RN. Diffusion MRI/ : apparent diffusion coefficient (ADC) values in the normal brain and a classification of brain disorders based on ADC values. 2001;25.
[https://doi.org/10.1016/S0895-6111\(00\)00083-5](https://doi.org/10.1016/S0895-6111(00)00083-5)
PMid:11356324
- [16]. Sener RN. Diffusion magnetic resonance imaging of solid vestibular schwannomas. J Comput Assist Tomogr. 2003;27(2):249-52.
<https://doi.org/10.1097/00004728-200303000-00024>
PMid:12703020
- [17]. Duygulu G, Ovali GY, Calli C, Kitis Ö, Yünter N, Akalin T, et al. Intracerebral metastasis showing restricted diffusion: correlation with histopathologic findings. Eur J Radiol. 2010;74(1):117-20.
<https://doi.org/10.1016/j.ejrad.2009.03.004>
PMid:19359117
- [18]. Hirai T, Morishita S, Kitajima M, Murakami R, Korogi Y, Makino K, et al. Diffusion-weighted Imaging of Metastatic Brain Tumors/ : Comparison with Histologic Type and. 2006,
- [19]. Holodny AI, Ollenschlager M. Diffusion imaging in brain tumors. Neuroimaging Clin N Am [Internet]. 2002;12(1):107-24.
[https://doi.org/10.1016/S1052-5149\(03\)00072-8](https://doi.org/10.1016/S1052-5149(03)00072-8)
PMid:11998248
- [20]. Hu R, Hoch MJ. Application of Diffusion Weighted Imaging and Diffusion Tensor Imaging in the Pre-treatment and Post-treatment of Brain Tumor. Radiol Clin North Am [Internet]. 2021;59(3):335-47.
<https://doi.org/10.1016/j.rcl.2021.01.003>
PMid:33926681
- [21]. Hakyemez B, Gokalp G, Erdogan C, Parlak M. The contribution of diffusion-weighted MR imaging to distinguishing typical from atypical meningiomas. 2006;513-20.
<https://doi.org/10.1007/s00234-006-0094-z>
PMid:16786348
- [22]. Ng WH, Chan YH, Hui F, Lee CK, Lim CCT. Diffusion-Weighted MR Imaging/ : Diagnosing Atypical or Malignant Meningiomas and. 2008;(January 1999):1147-52.
<https://doi.org/10.3174/ajnr.A0996>
PMid:18356472 PMCID:PMC8118818
- [23]. Chen L, Liu M, Bao J, Xia Y, Zhang J, Zhang L, et al. The correlation between apparent diffusion coefficient and tumor cellularity in patients: A meta-analysis. PLoS One. 2013;8(11).
<https://doi.org/10.1371/journal.pone.0079008>
PMid:24244402 PMCID:PMC3823989
- [24]. Sugahara T, Korogi Y, Kochi M, Ikushima I, Shigematu Y, Hirai T, et al. Usefulness of diffusion-weighted MRI with echo-planar technique in the evaluation of cellularity in gliomas. J Magn Reson Imaging An Off J Int Soc Magn Reson Med. 1999;9(1):53-60.
- [25]. Tan FC, Gokoglu A, Tucer B, Caner Y. The Apparent Diffusion Coefficient in Glial Tumors. Omi J Radiol. 2017;6(256):2.
- [26]. Server A, Kulle B, Gadmar ØB, Josefsen R, Kumar T, Nakstad PH. Measurements of diagnostic examination performance using quantitative apparent diffusion coefficient and proton MR spectroscopic imaging in the preoperative evaluation of tumor grade in cerebral gliomas. 2011;80:462-70.
<https://doi.org/10.1016/j.ejrad.2010.07.017>
PMid:20708868
- [27]. Chen Z, Ma L, Lou X, Zhou Z. Diagnostic value of minimum apparent diffusion coefficient values in prediction of neuroepithelial tumor grading. J Magn Reson Imaging. 2010;31(6):1331-8.
<https://doi.org/10.1002/jmri.22175>
PMid:20512884
- [28]. Arvinda HR, Kesavadas C, Sarma PS, Thomas B, Radhakrishnan V V, Gupta AK, et al. Glioma grading: sensitivity, specificity, positive and negative predictive values of diffusion and perfusion imaging. J Neurooncol. 2009;94(1):87-96.
<https://doi.org/10.1007/s11060-009-9807-6>
PMid:19229590
- [29]. Brunberg JA, Chenevert TL, McKeever PE, Ross DA, Junck LR, Muraszko KM, et al. In vivo MR determination of water diffusion coefficients and diffusion anisotropy: Correlation with structural alteration in gliomas of the cerebral hemispheres. Am J Neuroradiol. 1995;16(2):361-71.
- [30]. Fornasa F. Diffusion-weighted magnetic resonance imaging: What makes water run fast or slow? J Clin Imaging Sci. 2011;1(1):1-7.
<https://doi.org/10.4103/2156-7514.81294>
PMid:21966624 PMCID:PMC3177415

How to cite this article: Manah Chandra Changmai, Mohammed Faruque Reza, Kastury Gohain. Diffusion Weighted Imaging and Grading of Brain Tumours. *Int J Anat Res* 2024;12(1):8800-8807. DOI: 10.16965/ijar.2023.142