

A systematic review of the genus *Bolbelasmus* Boucomont (Coleoptera, Geotrupidae, Bolboceratinae) from Indochina and surrounding areas

Chun-Lin Li¹, Chuan-Chan Wang²

¹ The Experimental Forest, National Taiwan University, Chushan, Nantou County 557, Taiwan

² Department of Life Science, Fu Jen Catholic University, Hsinchuang, New Taipei City 24205, Taiwan

Corresponding author: Chuan-Chan Wang (gwennywang@gmail.com)

Abstract

Indochinese species of the genus *Bolbelasmus* (Coleoptera, Geotrupidae, Bolboceratinae) are reviewed. Three new species, *Bolbelasmus chifengi* Wang & Li, **sp. nov.**, *Bolbelasmus concavisuturalis* Li & Wang, **sp. nov.** and *Bolbelasmus yutangi* Li & Wang, **sp. nov.**, are described and illustrated. An annotated checklist and modified key to species of the genus are provided. Information for each species in the checklist includes literature review, synonymy, distribution and type locality.

Key words: Checklist, earth-boring beetles, key, natural history, new species, taxonomy



Academic editor: Andrey Frolov

Received: 11 October 2023

Accepted: 16 January 2024

Published: 14 February 2024

ZooBank: <https://zoobank.org/F412786C-54D1-408D-8BC7-D630A05F086F>

Citation: Li C-L, Wang C-C (2024)

A systematic review of the genus *Bolbelasmus* Boucomont (Coleoptera, Geotrupidae, Bolboceratinae) from Indochina and surrounding areas. ZooKeys 1191: 287–305. <https://doi.org/10.3897/zookeys.1191.114021>

Copyright: © Chun-Lin Li & Chuan-Chan Wang. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Bolbelasmus Boucomont, 1910 (Coleoptera: Geotrupidae: Bolboceratinae: Bolboceratini) is one of the largest bolboceratine genera, currently consisting of 29 species and two subspecies (Schoolmeesters 2023) (including subgenus *Kolbeus*). Among bolboceratine genera, *Bolbelasmus* has the widest distribution, occurring throughout the Holarctic and Oriental regions (Li et al. 2008; Hillert et al. 2016). Krikken (1977) published the most recent review dealing with eastern and southeastern Asian species and provided a complete checklist and summary of generic affinities and notes of species from the Middle East and North Africa. In that paper, two new species were described including one, *Bolbelasmus meridionalis* Krikken, 1977, from Southeast Asia. Additional distribution data were provided for *B. coreanus* (Kolbe, 1886) in southwestern China, Thailand and India. The Indochinese *Bolbelasmus* species have received little attention since Krikken's (1977) work. Zinchenko (2016) described a new species, *B. korshunovi* Zinchenko, from northern Thailand, the only species described during the past four decades. Currently, ten species are recorded in eastern and southeastern Asia, including the three new species described here

based on specimens recently obtained from the Indochinese Peninsula and neighboring areas, including the first record from Myanmar. These collections constitute the basis for further detailed knowledge of the Indochinese bolboceratine fauna.

The natural histories of eastern and southeastern Asian *Bolbelasmus* species are poorly documented. Tsukamoto et al. (2017) reported that two of the Japanese species, *B. nativus ishigakiensis* Masumoto and *B. shibatai* Masumoto inhabit densely forested montane areas, where adults are nocturnal and fly about 30 cm over the forest floor from late dusk to midnight. Adults of both species can be sporadically attracted to light, as observed with *B. coreanus* (Kolbe) in Taiwan (Li et al. 2008). *Bolbelasmus* specimens can occasionally be collected using baited traps (e.g., Kawai et al. 2005; Li et al. 2008).

In this paper, we review the species of *Bolbelasmus* occurring in the Indochinese Peninsula and neighboring areas, with descriptions of three new species. We also provide an annotated checklist, distributional data for known species in eastern and southeastern Asia, and a modified key.

Material and methods

The type specimens of the two new species, *B. chifengi* and *B. yutangi*, were collected by flight interception traps (FIT). The depository of the type specimens is in the following institutions: National Museum of Nature and Science (NSMT, Tsukuba, Japan); Zoological Museum, University of Copenhagen (ZMUC, Copenhagen, Denmark); Taiwan Agriculture Research Institute (TARI, Taichung, Taiwan); and the private collection of Chun-lin Li (CCLI, Nantou, Taiwan). Habitus images of *Bolbelasmus* specimens were taken using a Canon 7D digital camera with a Canon EF 100 mm macro lens and a Canon Macro Twin Lite MT-24EX Flash. Detailed images of specimens, body parts and male genitalia were captured using a Leica M205C stereo microscope equipped with a Leica MC190HD microscope camera or by a Hitachi TM3030 Plus tabletop scanning electron microscope. Color images were processed using Helicon Focus v.8.2.0 to increase depth of field, and all images were edited in Adobe Photoshop v.24.0.0 (background removed, images integrated, numbered and scale bars added). Measurements, treatments, and preservation of specimens and genitalia are based on methods described by Li et al. (2008).

Taxonomy

Bolbelasmus Boucomont, 1910

Diagnosis. Species of *Bolbelasmus* are small to medium-sized (5.6–15.2 mm in body length) and can be recognized by the glossy, unicolored, reddish-brown to black dorsal surface; presence of a conical frontal tubercle with a rounded or bifurcated tip in males; transverse frontal carina present in females; eyes protruding, divided by a canthus anteriorly, canthus with anterior margin smooth; antennal club with first segment glabrous on inner side; pronotum usually quadrituberculate in major males (vestigial or reduced to bituberculate in minor males), females with transverse carina only; first elytral stria terminated by scutellum; parameres usually weakly sclerotized.

Key to eastern and southeastern Asian *Bolbelasmus* species based on males

(excluding *Bolbelasmus orientalis*)

- 1 Frontal tubercle located at center of frons (Fig. 15) **2**
 - Frontal tubercle located in junction of clypeofrontal suture (Figs 11–14, 16) **4**
- 2 Sutural intervals of elytra more convex than other intervals (Figs 25, 26) **3**
 - Sutural intervals (Fig. 23, 24, 27, 28) of elytra equally convex as other intervals; pronotal median tubercles well developed, primary punctures moderately distributed throughout disc (Fig. 21) except for a small area near posterior margin impunctate; posterior margin punctate; parameres small, narrowed apically, curved in lateral view ***Bolbelasmus yutangi* sp. nov.**
- 3 Pronotal median tubercles weakly developed, center of disc typically impunctate, primary punctures moderately distributed at sides of center, posterior margin sparsely punctate at center; parameres small, narrowed apically, flat in lateral view ***Bolbelasmus nativus* Krikken, 1979**
 - Pronotal median tubercles well developed, disc sparsely punctate, primary punctures sparsely distributed at sides of center (Fig. 20), posterior margin impunctate at center; parameres (Figs 35, 36, 42) moderate in size, trapezoidal, bases swollen in lateral view ***Bolbelasmus korshunovi* Zinchenko, 2016**
- 4 Elytral sutural intervals (Fig. 25) distinctly convex **5**
 - Elytral sutural intervals concave (Fig. 23), flat or partially convex (Figs 24, 27, 28) **6**
- 5 Lateral margins of pronotum widely explanate; parameres with tips angulate at anterolateral angles ***Bolbelasmus meridionalis* Krikken, 1979**
 - Lateral margins of pronotum narrowly explanate; parameres with tips narrowed anteriorly ***Bolbelasmus minutus* Li & Masumoto, 2008**
- 6 Elytral sutural intervals completely concave; parameres (Figs 29, 30, 39) small, anteriorly 1/2 curved ventrally in lateral view ***Bolbelasmus concavisuturalis* sp. nov.**
 - Elytral sutural intervals flat or partially, moderately convex **7**
- 7 Elytral sutural intervals partially, moderately convex; parameres with bases contracted in dorsal view, length longer than one-half of basal piece **8**
 - Elytral sutural intervals flat; parameres straight, small, length shorter than one-half of basal piece ***Bolbelasmus krikkeni* Nikolajev, 1979**
- 8 Pronotal disc sparsely punctate **9**
 - Pronotal disc with many primary punctures (Fig. 18); parameres (Figs 31, 32) large with tips acute and vertically curved inward ***Bolbelasmus chifengi* sp. nov.**
- 9 Pronotal disc with primary punctures finer (Fig. 22); elytral intervals (Fig. 28) slightly convex; parameres with inner margins straight and separate, moderately evenly sclerotized ***Bolbelasmus coreanus* (Kolbe, 1886)**
 - Pronotal disc with primary punctures coarser; elytral intervals convex; parameres with inner margins broadened basally and overlapping, distinctly sclerotized, partly with median lobe ***Bolbelasmus shibatai* Masumoto, 1984**

Checklist of species of *Bolbelasmus* Boucomont from eastern and southeastern Asia

Bolbelasmus chifengi Wang & Li, sp. nov.

<https://zoobank.org/624857C3-4224-46EF-80FA-FDD3E4B1F169>

Figs 3, 4, 12, 18, 24, 31, 32, 40, 46, 47

Type materials. **Holotype male.** “China: Yunnan, Bangdashan (邦達山), 16.IX.2015. leg. Y.-T. Wang.” (glued on label, TARI), Taichung, Taiwan. **Paratypes.** 3♂♂, 1♀ (TARI). same collecting data as the holotype. 1♀ (TARI). “China: Yunnan, Wudian (武甸), 17.IX.2014. leg. Y.-T. Wang”. 1♂, 2♀♀ (TARI). “China: Yunnan, CCCC, Nabang (那邦), 21.VI.2017. leg. Y.-T. Wang”. 1♀ (CCLI). “China: Yunnan, Banggunjianshan (邦棍尖山), 19.IX.2015. leg. Y.-T. Wang”. 1♂ (CCLI). “China: Yunnan, Bangdashan (邦達山), 01.IX.2015. leg. Y.-T. Wang”. 2♂♂, 1♀ (CCLI). “China: Yunnan, Ruili (瑞麗), 01.IX.2014. leg. Y.-T. Wang”.

Description. **Holotype male** (Figs 3, 4). Body length 9.7 mm; width across humeri 6.0 mm. Dorsum moderately shiny. Head, pronotum and scutellum dark brown with elytra reddish brown. **Head** (Fig. 12): labrum with anterior margin crenulate, disc transversally rugose. Clypeus trapezoidal, surface densely rugopunctate; protrusion at basal angle moderately developed. Clypeofrontal suture well defined, distinctly curved in front of frontal tubercle. Frons with surface sparsely punctate, punctures fine, frontal tubercle vertically located in junction of suture, right-triangle in shape in lateral view. Eye prominent, canthus simple, not exceeding eye. **Thorax:** pronotum (Fig. 18) quadrituberculate, tubercles placed in a line, lateral tubercle greatly reduced in size; anterior face of median tubercles almost perpendicular to plane of pronotum; primary punctures coarse, moderately distributed on disc and intermixed with impunctate area, punctures between lateral margins of pronotum and fovea bigger and denser, posterior area between elytral humeri and suture impunctate except for four coarse punctures in front of scutellum, secondary punctures tiny, evenly scattered throughout surface of pronotum; frontal and lateral margins beaded, posterior margin beaded only in front of elytral humeri. Scutellum elongate, secondary punctures sparse throughout surface with a coarse puncture at center. **Elytron** (Fig. 24): elytral striae shallow, punctures mostly spaced 2–3 times diameters of punctures. Intervals slightly convex including sutural one, surface with scattered secondary punctures. Male genitalia (Figs 31, 32, 40).

Female (Figs 46, 47). Body length 7.7–10.0 mm; width across humeri 5.4–6.5 mm. Similar to male with minor differences in the form of strongly wrinkled surface of clypeus, transverse frontal carina trilobed, central lobe more prominent than lateral lobes, punctures on frons and vertex rugose, transverse pronotal carina feebly bilobed with lobes broadly developed, punctures on pronotal disc coarser and denser than males.

Variation in male. Dorsum brown, smaller body size, 6.6 mm in length and 5.2 mm in width across humeri, frontal tubercle less developed and not in junction of clypeofrontal suture, pronotal tubercles feebly convex, and number of coarse punctures arranged at pronotal posterior margin variable.

Diagnosis. *Bolbelasmus chifengi* is morphologically similar to *B. concavisuturalis*, but can be distinguished from the latter by the coarser and denser primary punctures on the pronotal disc (finer and scattered in *B. concavisuturalis*), elytral




Figures 1–6. Dorsal and lateral views of male *Bolbelasmus* spp. **1, 2** *B. concavisuturalis* sp. nov.sp. nov., holotype **3, 4** *B. chifengi* sp. nov., holotype **5, 6** *B. meridionalis*.

intervals evenly convex (elytral intervals flat with sutural interval concave in *B. concavisuturalis*) and by the longer parameres (shorter in *B. concavisuturalis*).

Distribution. Southern Yunnan, China (Fig. 52).

Etymology. *Bolbelasmus chifengi* sp. nov. is named after Dr Chi-feng Lee, the curator of the Department of Applied Zoology, Taiwan Agriculture Research Institute, Taichung, Taiwan, who kindly provided materials used in this study.



Figures 7–10. Dorsal and lateral views of male *Bolbelasmus* spp. **7**, **8** *B. korshunovi* **9**, **10** *B. yutangi* sp. nov., holotype.

***Bolbelasmus concavisuturalis* Li & Wang, sp. nov.**

<https://zoobank.org/B3BDDA3C-0CE3-4ACF-9470-E2750D968B93>

Figs 1, 2, 11, 17, 23, 29, 30, 39, 44, 45

Type materials. Holotype male. “Mon-Angget, near Chiangmai, North Thailand, 31-V-1990, K. Masumoto leg.” (glued on label, NSMT). **Paratypes.** 1♂ (ZMUC) “Northern Thailand, Doi Sutep, 21.6.1958, B. Degerbøl leg., Pr. 548 (1.7.59)”. 1♀ (NSMT) “Doi Suthep, Chiang Mai, THAILAND, 15-VIII-1989, Y. MANIT leg.”

Description. Holotype male (Figs 1, 2). Body length 10.2 mm; width across humeri 6.3 mm. Dorsum distinctly shiny. Head, pronotum and scutellum reddish brown with elytra brown in color. **Head** (Fig. 11): labrum with anterior margin crenulate, disc transversally rugose. Clypeus trapezoidal, surface densely rugopunctate; protrusion at basal angle reduced. Clypeofrontal suture well defined, distinctly curved in front of frontal tubercle. Frons with surface sparsely punctate, punctures fine, frontal tubercle vertically located in junction of suture, triangular in lateral view. Eye prominent, canthus simple, not exceeding eye. **Thorax:** pronotum (Fig. 17) quadrituberculate, tubercles situated in a line, lateral tubercle smaller; anterior face of median tubercles almost perpendicular to surface of pronotum; primary punctures weakly defined, sparse on disc except between lateral margin of pronotum and fovea, these coarser and denser, line in front of scutellum with a coarse puncture, secondary punctures tiny, evenly

scattered on surface of pronotum; frontal and lateral margins beaded, posterior margin beaded only anterior to humeri of elytra. Scutellum elongate, secondary punctures sparsely distributed. **Elytron** (Fig. 23): elytral striae shallowly impressed, punctures mostly spaced 2–3 times diameters of punctures. Intervals flat, with sutural interval weakly concave, surface scattered with secondary punctures. Male genitalia (Figs 29, 30, 39).

Female paratype (Figs 44, 45). Body length 9.8 mm; width across humeri 5.8 mm. Similar to male with minor differences in the form of strongly wrinkled surface of clypeus, transverse frontal carina trilobed, central lobe more prominent than lateral lobes, punctures on frons and vertex rugose, transverse pronotal carina feebly bilobed, lobes broad, punctures on pronotal disc coarser and denser than those of males.

Male paratype. The single male paratype is smaller in body size, 9.4 mm in length and 5.1 mm in width across humeri, frontal tubercle less developed and with three coarse punctures along pronotal posterior margin in front of scutellum.

Diagnosis. *Bolbelasmus concavisuturalis* sp. nov. is morphologically similar to *B. coreanus*, but can be distinguished from the latter by having denser punctures along the midline of the pronotum (Fig. 17) (sparser punctures in *B. coreanus* (Fig. 22)), punctures in elytral striae moderately developed (Fig. 23) (punctures weakly developed in *B. coreanus* (Fig. 28)) and ventrally curved parameres (straight in *B. coreanus*).

Distribution. Northern Thailand (Fig. 52).

Etymology. *Concavi-* (L.) = concave, *-suturalis* (L.) = suture. In reference to the concave sutural intervals of the elytra.

***Bolbelasmus coreanus* (Kolbe, 1886)**

Bolboceras coreanus Kolbe, 1886: 188. Original combination (type locality: Seoul, Korea, female type in the Museum für Naturkunde, Berlin, Germany).


Bolbelasmus coreanus (Kolbe, 1886): Cartwright 1953: 97 (generic combination); Krikken 1977: 288 (notes; diagnosis; illustration); Kim 2000: 45 (diagnosis; collecting records); Li et al. 2008: 480 (redescription, illustrations, collecting records, distribution, remarks); Král, Löbl and Nikolajev 2006: 83 (catalog, subgenus *Kolbeus*); Nikolajev, Král and Bezděk 2016: 33 (catalog, subgenus *Kolbeus*).

Kolbeus coreanus (Kolbe, 1886): Boucomont 1911: 335 (generic combination); Boucomont 1912: 17 (catalog); Boucomont and Gillet 1921: 72 (record to Taiwan; diagnosis); Miwa 1930: 164 (catalog); Miwa 1931: 276 (catalog); Miwa and Chûjô 1939: 30 (catalog); Paulian 1945: 42 (diagnosis; figures; distribution).

Bolbelasmus kurosawai Masumoto, 1984: 76; Li et al. 2008: 481 (junior synonym).

Bolboceras conicifrons Fairmaire, 1896: 82; Boucomont and Gillet 1921: 72 (junior synonym).

Distribution. Korean Peninsula; China (Anhui, Zhejiang, Kweichow, Szechuan, Yunnan); Taiwan.



Figures 11–16. Scanning electron micrographs of heads of male *Bolbelasmus* spp. **11** *B. concavisuturalis* sp. nov. **12** *B. chifengi* sp. nov. **13** *B. meridionalis* **14** *B. korshunovi* **15** *B. yutangi* sp. nov. **16** *B. coreanus*. Scale bar: 1 mm.

Remarks. The detailed distribution of *B. coreanus* in China requires further investigation, particularly those from southwestern areas. Based on a large number of *Bolbelasmus* specimens recently collected from Yunnan and neighboring areas, we found no representatives of *B. coreanus* among them. Therefore, we reserve a decision about whether *B. coreanus* occurs in Yunnan, Thailand and India, as recorded by Krikken (1977). Voucher specimens from the areas mentioned above are required.

***Bolbelasmus korshunovi* Zinchenko, 2016**

Figs 7, 8, 14, 20, 26, 35, 36, 42

Bolbelasmus korshunovi Zinchenko, 2016: 328. Original combination (type locality: Nong Bun Nak, Nakhon Prov., Thailand).

Material examined. (5♂♂). 2♂♂ (ZMUC). THAILAND: Loei Province, Phu Luang Wildlife Sanctuary, 8.–14.x.1984, 700–900 m, Karsholt, Lomboldt & Nielsen leg., Pral Siaw, 1923-9-33, Paül Fogh/ Coll. Roseberg. 3♂♂ (NSMT). Sansai, Chiang Mai, THAILAND, 17. VI. 1993.

Diagnosis. Body length, males, 6.8–9.0 mm, greatest width at pronotal base 4.4–5.7 mm; females, 7.1–8.7 mm in length, 4.3–5.6 mm in width (Zinchenko 2016). *Bolbelasmus korshunovi* is distinguished from the other Oriental *Bolbelasmus* species by elytral sutural intervals that are moderately convex, primary punctures sparsely distributed either side of the center of the pronotum, and shapes of the parameres.

Notes. Thirteen type specimens were designated in the original description of the species (Zinchenko 2016), 12 of them collected from June to August, and the holotype during November. Accordingly, the temporal activity of adults is likely at least half the year during both rainy and dry seasons. This is identical to the sympatric species, *B. meridionalis*.

Distribution. Northern Thailand.

Remarks. *Bolbelasmus korshunovi* inhabits plains to low-elevation montane areas and occurs sympatrically with *B. meridionalis* in northern Thailand.

***Bolbelasmus krikkeni* Nikolajev, 1979**

Bolbelasmus krikkeni Nikolajev, 1979: 225. Original combination (type locality: Gopaldhara, Sikkim, India); Král, Löbl and Nikolajev 2006: 83 (catalog, in subgenus *Kolbeus*); Nikolajev, Král and Bezděk 2016: 33 (catalog, subgenus *Kolbeus*).

Distribution. Northern India.


Remarks. Based on the collecting data from the monotypic specimen, *B. krikkeni* occurs in mid-elevation forests above 1000 m and is unique compared to its congeners that usually inhabit plains to low-elevation montane areas in the region. No additional specimens have been recorded since the publication of the original description.

***Bolbelasmus meridionalis* Krikken, 1977**

Figs 5, 6, 13, 19, 25, 33, 34, 41, 48, 49

Bolbelasmus meridionalis Krikken, 1977: 285. Original combination (type locality: Java, Indonesia); Král, Löbl and Nikolajev 2006: 83 (catalog, subgenus *Kolbeus*); Nikolajev, Král and Bezděk 2016: 33 (catalog, subgenus *Kolbeus*).

Material examined. (11♂♂, 8♀♀). 1♂ (NSMT). THAILAND: Sansai, Chiang Mai, 17. VI. 1993 (1 male at NSMT). 9♂♂, 7♀♀ (NSMT); same locality, 12-V-1996.



Figures 17–22. Scanning electron micrographs of pronota of male *Bolbelasmus* spp. 17 *B. concavisuturalis* sp. nov. 18 *B. chifengi* sp. nov. 19 *B. meridionalis* 20 *B. korshunovi* 21 *B. yutangi* sp. nov. 22 *B. coreanus*. Scale bar: 1 mm.

1♀ (NSMT); near Chiang Mai, N Thailand, VII-1996, native collector. 1♂ (NSMT). TAIWAN: Formosa, Heito, 10-VII-1941, H. Kondo/Sizumu Nomura Bequest, 1981.

Diagnosis. Body length, males, 6.1–8.2 mm, greatest width at pronotal base, 3.7–5.0 mm; females, 5.6–8.2 mm in length, 3.3–5.2 mm in width. Both *B. meridionalis* and *B. minutus* constitute a distinctive group among southeastern Asian congeners based on sharing the distinctly convex elytral sutural intervals and the tips of the parameres acute and curved ventrally in lateral view. Due to a lack of further material being available of the later species, *B. meridionalis* and *B. minutus* can

only be separated by the shape of male genitalia and the lateral margin of the pronotum in *B. meridionalis*, which is more widely explanate than that of *B. minutus*.

Chinese name. 脊背厚角金龜

Notes. Li et al. (2008) excluded *B. meridionalis* from the registered Taiwan fauna due to the lack of verified records. During the present study, we examined a male *B. meridionalis* specimen housed in NSMT bearing identical labels as the paratype of the species collected in Heito (now Pingtung) by the late Japanese coleopterist, Yushiro Miwa. We therefore confirm the record of *B. meridionalis* in Taiwan, though it has been lacking in reports of the genus for 90 years. Consequently, the conservation status of *B. meridionalis* in Taiwan is in urgent need of study, along with that of *B. minutus* Li & Masumoto, 2008 and *Bolbotrypes davidis* (Fairmaire, 1891). These species are restricted to habitats in highly urbanized areas and/or intensively farmed plains of Taiwan.

Distribution. Indonesia (Java); eastern China; Thailand; Vietnam; Taiwan.

Remarks. *Bolbelasmus meridionalis* has the widest known distribution among congeners in the region in eastern and southeastern Asia. Also, the records from Java for the holotype and paratypes indicated that it is the only member from the Sunda Islands of the genus.

***Bolbelasmus minutus* Li & Masumoto, 2008**

Bolbelasmus minutus Li & Masumoto, 2008: 482. Original combination (type locality: Heito (presently Pingtung), Taiwan); Nikolajev, Král and Bezděk 2016: 33 (catalog, subgenus *Kolbeus*).

Distribution. Taiwan.

Remarks. *Bolbelasmus minutus* was described from a pair of specimens collected during 1931, and no further records of the species have been recorded. This species occurs sympatrically with *B. meridionalis* and *B. nativus* in the plains of southern Taiwan.

***Bolbelasmus nativus nativus* Krikken, 1977**


Bolbelasmus nativus Krikken, 1977: 287. Original combination (type locality: Heito (presently Pingtung), Taiwan); Nikolajev, Král and Bezděk 2016: 33 (catalog, in nominate subgenus *Bolbelasmus*).

B. n. ishigakiensis ssp. Masumoto, 1984.

Bolbelasmus ishigakiensis Masumoto, 1984: 73. Original combination (type locality: Ishigaki island, Okinawa, Japan); Král, Löbl and Nikolajev 2006: 83 (catalog, subgenus *Kolbeus*); Ochi and Masumoto 2005: 244 (as subspecies of *B. nativus*); Nikolajev, Král and Bezděk 2016: 33 (catalog, subgenus *Kolbeus*).

Distribution. Taiwan (southern areas and Lanyu island); Japan (Iriomote, Ishigaki and Tarama islands, Okinawa Prefecture).

Remarks. *Bolbelasmus nativus* was originally described based on a single male from Taiwan. Ochi and Masumoto (2005) treated the population distributed on a few small islands near Taiwan as a subspecies, *B. nativus ishigakiensis*.



Figures 23–28. Scanning electron micrographs of elytra of male *Bolbelasmus* spp. **23** *B. concavisuturalis* sp. nov. **24** *B. chifengi* sp. nov. **25** *B. meridionalis* **26** *B. korshunovi* **27** *B. yutangi* sp. nov. **28** *B. coreanus*. Scale bar: 1 mm.

***Bolbelasmus shibatai* Masumoto, 1984**

Bolbelasmus shibatai Masumoto, 1984: 75. Original combination (type locality: Amami Oshima Island, Japan); Nikolajev, Král and Bezděk 2016: 33 (catalog).

Distribution. Japan (Amami oshima and Okinawa island).

Remarks. Populations of *B. shibatai* are restricted to a few small islands in the southwestern archipelagos of Japan. Males possess strongly sclerotized parts of the parameres that can be distinguished from the similar species, *B. coreanus*.

***Bolbelasmus yutangi* Li & Wang, sp. nov.**

<https://zoobank.org/A57879F9-F8A6-4D40-B922-DF3429C8BA4D>

Figs 9, 10, 15, 21, 27, 37, 38, 43, 50, 51


Type materials. Holotype male. “Myanmar: Bago Region, Moe Yin Gyi, CCCC, 21.V.2017. leg. Y.-T. Wang.” (glued on label, TARI). **Paratypes.** 1♀ (TARI). data as the holotype. 5♂♂ (TARI). “China: Yunnan, Wudian (武甸), 17.IX.2014. leg. Y.-T. Wang”. 1♀ (CCLI) “China: Yunnan, Banggunjianshan (邦棍尖山), 18.IX.2015. leg. Y.-T. Wang”. 6♂♂ (CCLI). “China: Yunnan, Bangdashan (邦達山), 01.IX.2015. leg. Y.-T. Wang”. 1♂ (CCLI). “China: Yunnan, Ruili (瑞麗), 15.IX.2014. leg. Y.-T. Wang”. 1♂ (NSMT). “Doi Saket, Chiang Mai, THAILAND, 12-X-1989, Y. MANIT leg”. 1♂ (NSMT). “Doi Mon Unggate, Samoeng Distr., Chiang Mai Prov., THAILAND, 18-VII-1989, Y. MANIT leg”.

Description. Holotype male (Figs 9, 10). Body length 9.3 mm; width across humeri 5.9 mm. Dorsum moderately shiny. Head, pronotum and scutellum dark brown with elytra reddish brown. **Head** (Fig. 15): labrum with anterior margin crenulate, disc transversally rugose. Clypeus trapezoidal, surface densely rugopunctate; protrusion at basal angle moderately developed. Clypeofrontal suture well defined, slightly curved in front of frontal tubercle. Frons with surface moderately punctate, punctures coarse, frontal tubercle vertically located at center of disc, triangular when viewed laterally. Eye prominent, canthus simple, not exceeding eye. **Thorax:** pronotum (Fig. 21) quadrituberculate, tubercles placed in a line, lateral tubercle greatly reduced in size; anterior face of median tubercles almost perpendicular to plane of pronotum; primary punctures coarse, dense on disc with small impunctate area in front of middle of posterior margin, punctures between lateral margins of pronotum and fovea bigger and denser, scattered coarse punctures distributed along posterior margin with seven punctures in front of scutellum, secondary punctures tiny, evenly scattered on surface of pronotum; frontal and lateral margins beaded, posterior margin beaded only in front of elytral humeri. Scutellum elongate, fine punctures sparsely distributed on surface. **Elytron** (Fig. 27): elytral striae shallowly impressed, punctures mostly spaced by 1–3 times diameters of punctures. Intervals slightly convex, including sutural interval, surface with scattered secondary punctures. Male genitalia. Figs 37, 38, 43.

Female (Figs 50, 51). Body length 8.4–8.7 mm; width across humeri 5.1–5.3 mm. Similar to male with minor differences in the form of strongly wrinkled surface of clypeus, transverse frontal carina trilobed, central lobe more prominent than lateral lobes, punctures on frons and vertex rugose, transverse pronotal carina feebly bilobed, with lobes broadly developed to reduced, bigger punctures on pronotal disc denser than those of males and scutellum with 1 or 2 bigger punctures.

Variation in male. Male paratypes differ from the holotype in the following respects: smaller body size, 6.6 mm in body length and 5.2 mm in width across humeri, frontal tubercle less developed and not in junction of clypeofrontal suture, pronotal tubercle feebly convex, reduced, and number of coarse punctures along pronotal posterior margin variable.

Diagnosis. *Bolbelasmus yutangi* is similar to *B. nativus* in sharing the frontal tubercle location at the center of the frons as well as having smaller parameres. It can be distinguished from the latter by the weakly convex basal angle of the




Figures 29–38. Male genitalia of *Bolbelasmus* spp. (29, 31, 33, 35, 37, dorsal views; 30, 32, 34, 36, 38, ventral views) 29, 30 *B. concavisuturalis* sp. nov. 31, 32 *B. chifengi* sp. nov. 33, 34 *B. meridionalis* 35, 36 *B. korshunovi* 37, 38 *B. yutangi* sp. nov. Scale bar: 0.3 mm.

clypeus (distinctly bulging in *B. nativus*), primary punctures densely distributed on the disc (almost impunctate along the middle of disc in *B. nativus*), sutural interval convex, similar to discal intervals (distinctly more convex than discal intervals in *B. nativus*) and parameres with the tips tapered (parameres broader at tips in *B. nativus*).

Distribution. Eastern Myanmar, northern Thailand and western Yunnan, China (Fig. 52).

Etymology. *Bolbelasmus yutangi* sp. nov. is named after Mr Yu-tang Wang, a beetle enthusiast of Taiwan, who collected most of the material used in this study.




Figures 39–43. Male genitalia of *Bolbelasmus* spp., lateral views 39 *B. concavisuturalis* sp. nov. 40 *B. chifengi* sp. nov. 41 *B. meridionalis* 42 *B. korshunovi* 43 *B. yutangi* sp. nov. Scale bar: 0.2 mm.

Species with doubtful locality record

***Bolbelasmus orientalis* Petrovitz, 1968**

Bolbelasmus orientalis Petrovitz, 1968: 185. Original combination (type locality: Wladiwostok [Vladivostok], Primorskii Territory, Russia); Krikken 1977: 289 (notes; diagnosis; illustration); Nikolajev, Král and Bezděk 2016: 33 (catalog).

Distribution. Described from Vladivostok, Russian Far East.



Figures 44–51. Dorsal and lateral views of female *Bolbelasmus* spp. 44, 45 *B. concavisuturalis* sp. nov., paratype 46, 47 *B. chifengi* sp. nov., paratype 48, 49 *B. meridionalis* 50, 51 *B. yutangi* sp. nov., paratype.

Remarks. *Bolbelasmus orientalis* was described from one male and one female. Krikken (1977) examined both type specimens and found that the female allotype to be a minor male. Also, he noted that the species has no direct




Figure 52. Distribution map of the eastern and southeastern Asian *Bolbelasmus* species. Question marks indicate doubtful distributional records.

affinity with three congeners, *B. coreanus*, *B. meridionalis* and *B. nativus*, which are geographically close to *B. orientalis* but are allied to the western Palearctic *B. unicornis*. Bezborodov and Koshkin (2014) doubted the locality label attached to the type specimens because there were no additional records of the species documented in Russia or nearby territories other than that of type specimens. We, therefore, exclude *B. orientalis* from the *Bolbelasmus* fauna in the eastern Palearctic and the Oriental regions.

Acknowledgements

We thank Chi-Feng Lee (TARI) and Shûhei Nomura (National Museum of Nature and Science, Tsukuba, Japan) for lending valuable specimens. The first author acknowledges Paul Schoolmeesters (Belgium) for his help in providing literature. The authors thank Michele Rossini for valuable comments on the earlier version of the manuscript. Chris Carlton is thanked for reading the draft and editing the English.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was supported by the Experimental Forest, National Taiwan University, Taiwan (grant no. 110 RG2 to C.-L. Li).

Author contributions

Funding acquisition: CLL. Investigation: CLL. Methodology: CLL. Supervision: CCW. Writing – original draft: CLL. Writing – review and editing: CCW.

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bezborodov VG, Koshkin ES (2014) A review of Bolboceratidae (Coleoptera, Scarabaeoidea) species from the Russian Far East. *Entomological Review* 93(8): 953–959. <https://doi.org/10.1134/S0013873814090127>
- Boucomont A ([1911] 1910). Contribution à la classification des Geotrupidae. *Annales de la Société entomologique de France*. Paris (1911) 79: 333–350. <https://doi.org/10.5962/bhl.part.4697>
- Boucomont A (1912) Scarabaeidae: Taurocerastinae, Geotrupinae. In: Junk W, Schenkling S (Eds) *Coleopterorum catalogus*, pars 46. W. Junk, Berlin, 1–47.
- Boucomont A, Gillet J (1921) Faune entomologique de l'Indo-chine française. Fam. Scarabaeidae Laparosticti. Imprimerie Nouvelle Albert Portail. Saigon 4: 1–76.
- Cartwright OL (1953) Scarabaeid beetles of the genus *Bradycinetulus* and closely related genera in the United States. *Proceedings of the United States National Museum* 103(3318): 95–120. <https://doi.org/10.5479/si.00963801.103-3318.95>
- Fairmaire L (1896) Coléoptères de l'Inde boréale, Chine et Malaisie. *Notes of the Leyden Museum* 18: 81–129.
- Hillert O, Arnone M, Král D, Massa B (2016) The genus *Bolbelasmus* in the western and southern regions of the Mediterranean Basin. *Acta Entomologica Musei Nationalis Pragae* 56(1): 211–254.
- Kawai S, Hori S, Kawahara M, Inagaki M (2005) Atlas of Japanese Scarabaeoidea, volume 1 Coprophagous group. Roppon-Ashi Entomological Books, Tokyo, 189 pp.
- Kim JI (2000) Scarabaeoidea (I). *Economic Insects of Korea*, 4. Ins. Koreana Suppl. 11. National Institute of Agricultural Science and Technology, Suwon, 149 pp.
- Kolbe HJ (1886) Beiträge zur Kenntniss der Coleopteren Fauna Koreas. *Archiv für Naturgeschichte* 52: 139–240. <https://doi.org/10.5962/bhl.part.28437>
- Král D, Löbl I, Nikolajev GV (2006) Bolboceratidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera*, volume 3. Apollo Books, Stenstrup, Denmark, 82–84. https://doi.org/10.1163/9789004260917_004

- Krikken J (1977) The genus *Bolbelasmus* Boucomont in Asia, with notes on species occurring in other regions (Coleoptera: Geotrupidae). *Zoölogische Mededeelingen* 51: 277–292. <https://repository.naturalis.nl/pub/319326>
- Li CL, Wang CC, Masumoto K, Ochi T, Yang PS (2008) Review of the tribe Bolboceratini s.l. from Taiwan (Coleoptera: Scarabaeoidea: Geotrupidae) with a key to the Eurasian genera. *Annals of the Entomological Society of America* 101(3): 474–490. [https://doi.org/10.1603/0013-8746\(2008\)101\[474:ROTTBS\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2008)101[474:ROTTBS]2.0.CO;2)
- Masumoto K (1984) New coprophagous lamellicornia from Japan and Formosa (I). *Entomological Review of Japan* 39: 73–83. <https://archive.org/details/entomologicalreviewofjapan19840039001/page/n75/mode/2up>
- Miwa Y (1930) An enumeration of the coprophagid-coleoptera from Formosa, with a table of the geographical distribution. *Insecta Matsumurana* IV(4): 163–180.
- Miwa Y (1931) A systematic Catalogue of Formosan Coleoptera. Report of the Department of Agriculture of the Government Research Institute 55: 1–359.
- Miwa Y, Chûjô M (1939) Scarabaeidae. *Catalogus Coleopterorum Japonicorum*, Pars 5. Noda-Syobo, Taihoku, 94 pp.
- Nikolajev GV (1979) Eine neue *Bolbelasmus*-Art aus Asien (Coleoptera, Scarabaeidae, Bolboceratinae). *Reichenbachia* 17: 225–227.
- Nikolajev GV, Král D, Bezděk B (2016) Bolboceratinae. In: Löbl I, Löbl D (Eds) *Catalogue of Palaearctic Coleoptera*, volume 3. Koninklijke Brill NV, Leiden, The Netherlands, 33–36. <https://doi.org/10.1163/9789004309142>
- Ochi T, Masumoto K (2005) Systematic position of *Bolbelasmus ishigakiensis* Masumoto (Coleoptera, Scarabaeidae). *Elytra* 3: 244. <https://archive.org/details/elytra20050033001/page/n245/mode/1up>
- Paulian R (1945) *Faune de l'empire français III coléoptères Scarabéides de l'Indochine*. Librairie Larose, Paris, 227 pp.
- Petrovitz R (1968) Bekannte und unbekante Scarabaeidae (Hybosorinae, Troginae, Orphninae, Dynamopinae, Geotrupinae, Aegialinae). *Entomologische Arbeiten aus dem Museum G. Frey* 19: 179–187. https://www.zobodat.at/pdf/Entomologische-Arbeiten-Museum-Frey_19_0179-0187.pdf
- Schoolmeesters P (2023) World Scarabaeidae Database. In: Bánki O, Roskov Y, Döring M, Ower G, Vandepitte L, Hobern D, Remsen D, Schalk P, DeWalt RE, Keping M, Miller J, Orrell T, Aalbu R, Abbott J, Adlard R, Adriaenssens EM, Aedo C, Aescht E, Akkari N, et al. (Eds) *Catalogue of Life Checklist (Version 2023-02-06)*.
- Tsukamoto K, Inagaki M, Kawahara M, Mori M (2017) *Geotrupes* and related species of Japan, Handbook Series of Insects 9. Mushi-sha, Tokyo, 116 pp.
- Zinchenko VK (2016) A new species of the genus *Bolbelasmus* (Coleoptera, Bolboceratidae) from Thailand. *Evraziatskii Entomologicheskii Zhurnal* 15(4): 328–329. https://kmkjournals.com/upload/PDF/EEJ/15/EEJ15_4_06.pdf