

Chapman University

## Chapman University Digital Commons

---

Physical Therapy Faculty Articles and Research

Physical Therapy

---

11-17-2023

### **Multi-Site Identification and Generalization of Clusters of Walking Behaviors in Individuals With Chronic Stroke and Neurotypical Controls**

Natalia Sánchez

Nicolas Schweighofer

Sara J. Mulroy

Ryan T. Roemmich

Trisha M. Keshar

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.chapman.edu/pt\\_articles](https://digitalcommons.chapman.edu/pt_articles)



Part of the [Nervous System Diseases Commons](#), [Other Rehabilitation and Therapy Commons](#), and the [Physical Therapy Commons](#)

---

---

# Multi-Site Identification and Generalization of Clusters of Walking Behaviors in Individuals With Chronic Stroke and Neurotypical Controls

## Comments

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Neurorehabilitation and Neural Repair*, volume 37, issue 11-12, in 2023 following peer review. This article may not exactly replicate the final published version. The definitive publisher-authenticated version is available online at <https://doi.org/10.1177/15459683231212864>.

## Copyright

The authors

## Authors

Natalia Sánchez, Nicolas Schweighofer, Sara J. Mulroy, Ryan T. Roemmich, Trisha M. Keshar, Gelsy Torres-Oviedo, Beth E. Fisher, James M. Finley, and Carolee J. Winstein

Clusters of walking impairment post-stroke

**Multi-site identification and generalization of clusters of walking behaviors in individuals with chronic stroke and neurotypical controls**

Natalia Sánchez, PhD<sup>1,2,3</sup>, Nicolas Schweighofer, PhD<sup>3,4,5</sup>, Sara J. Mulroy PT, PhD<sup>3,7\*</sup>, Ryan T. Roemmich, PhD<sup>8</sup>, Trisha M. Kesar PT, PhD<sup>9</sup>, Gelsy Torres-Oviedo, PhD<sup>10</sup>, Beth E. Fisher PT, PhD<sup>3,6</sup>, James M. Finley, PhD<sup>3,4,5+</sup>, Carolee J. Winstein, PT, PhD<sup>3,6+</sup>

<sup>1</sup> Department of Physical Therapy, Chapman University, Irvine, CA

<sup>2</sup> Fowler School of Engineering, Chapman University, Orange, CA

<sup>3</sup> Division of Biokinesiology and Physical Therapy, <sup>4</sup> Department of Biomedical Engineering, <sup>5</sup> Neuroscience Graduate Program, <sup>6</sup> Department of Neurology Keck School of Medicine, University of Southern California, Los Angeles, CA

<sup>7</sup> Pathokinesiology Lab, Rancho Los Amigos National Rehabilitation Center, Downey, CA

<sup>8</sup> Center for Movement Studies, Kennedy Krieger Institute, Baltimore, Department of Physical Medicine and Rehabilitation, Johns Hopkins University School of Medicine, Baltimore, MD

<sup>9</sup> Department of Rehabilitation Medicine, Emory University School of Medicine. Atlanta GA

<sup>10</sup> Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA

\*Additional Information: Coauthor Sara J. Mulroy, PT, PhD, passed away on May 3<sup>rd</sup>, 2022.

+ Shared senior authorship

**Corresponding Author**

Natalia Sánchez

Department of Physical Therapy, Chapman University

9401 Jeronimo Road, Irvine CA 92618

[sanchezaldana@chapman.edu](mailto:sanchezaldana@chapman.edu)

Tel: 714-516-5503

**Word Counts:**

Abstract: 250/250 words

Main Text: 5333/4500

Figures: 4

Tables: 2

References: 60

Supplementary Tables: 1

Supplementary Figures: 5

## Clusters of walking impairment post-stroke

### Abstract

**Background:** Walking patterns in stroke survivors are highly heterogeneous, which poses a challenge in systematizing treatment prescriptions for walking rehabilitation interventions.

**Objective:** We used bilateral spatiotemporal and force data during walking to create a multi-site research sample to: 1) identify clusters of walking behaviors in people post-stroke and neurotypical controls, and 2) determine the generalizability of these walking clusters across different research sites. We hypothesized that participants post-stroke will have different walking impairments resulting in different clusters of walking behaviors, which are also different from control participants.

**Methods:** We gathered data from 81 post-stroke participants across four research sites and collected data from 31 control participants. Using sparse K-means clustering, we identified walking clusters based on 17 spatiotemporal and force variables. We analyzed the biomechanical features within each cluster to characterize cluster-specific walking behaviors. We also assessed the generalizability of the clusters using a leave-one-out approach.

**Results:** We identified four stroke clusters: a fast and asymmetric cluster, a moderate speed and asymmetric cluster, a slow cluster with frontal plane force asymmetries, and a slow and symmetric cluster. We also identified a moderate speed and symmetric gait cluster composed of controls and participants post-stroke. The moderate speed and asymmetric stroke cluster did not generalize across sites.

**Conclusions:** Although post-stroke walking patterns are heterogeneous, these patterns can be systematically classified into distinct clusters based on spatiotemporal and force data. Future interventions could target the key features that characterize each cluster to increase the efficacy of interventions to improve mobility in people post-stroke.

**Clusters of walking impairment post-stroke**

**Key Words: Stroke, clusters, walking impairment, ground reaction forces, spatiotemporal variables.**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

Walking patterns differ between individuals to the point that we can recognize people by how they walk<sup>1</sup>. These individual differences are more marked after a neurologic injury, such as stroke<sup>2</sup>, due to heterogeneity in stroke lesion type, size, location, and differences in recovery<sup>3-6</sup>. These individual differences in walking patterns make systematizing treatment prescription for walking rehabilitation interventions a difficult clinical endeavor. The heterogeneity in walking patterns has been acknowledged by Knutsson and Richards as early as 1979, stating that “therapy and training in the hemiparetic patient should preferably be adapted to the disturbance in each individual case”<sup>4</sup>. In this study<sup>4</sup>, researchers used paretic electromyography (EMG) and walking kinematics to qualitatively identify three subgroups of abnormal muscle activation during walking post-stroke – early triceps surae activation, decreased activation of paretic musculature, and paretic muscle coactivation. Similarly, Olney and Richards qualitatively identified different subgroups of walking impairments in the spatiotemporal, kinematic, and kinetic domains<sup>7</sup>. Also, a seminal study by our co-authors\* investigated paretic leg kinematics and used hierarchical clustering to identify four clusters of walking behaviors, distinguished by walking speed and paretic knee flexion angles at different points during the gait cycle<sup>5</sup>. While these studies identified individual differences in paretic leg function during walking, walking patterns post-stroke are defined not only by paretic leg function but also by non-paretic leg function<sup>8</sup>.

To characterize functional differences between the paretic and non-paretic legs during walking, recent studies have used asymmetry measures in kinematics<sup>9-14</sup> or forces<sup>11,15-18</sup>. These asymmetry measures are compared to asymmetries observed in neurotypical controls to classify individuals post-stroke into subgroups with higher, comparable, or lower asymmetries than

**Clusters of walking impairment post-stroke**

1  
2  
3 control participants<sup>9-19</sup>. However, individuals might be classified as symmetric in one feature,  
4  
5 for example step lengths, while remaining asymmetric in other features, for example joint  
6  
7 kinematics<sup>19</sup>. Therefore, we reasoned that using an approach that allowed simultaneously  
8  
9 assessing multiple measures of walking, we can quantitatively identify distinct subgroups of  
10  
11 walking behaviors beyond those identified via a single asymmetry measure, which could inform  
12  
13 more specific clinical rehabilitation targets.  
14  
15  
16  
17  
18

19           Identifying subgroups of walking behaviors requires a heterogeneous sample that  
20  
21 encompasses the different combinations of gait deviations observed in survivors of stroke.  
22  
23 Achieving this heterogeneity necessitates a large sample size. However, studies in walking post-  
24  
25 stroke typically collect small, single site samples: for example, a systematic review of 46 studies  
26  
27 in walking post-stroke reported sample sizes between 8 and 39 participants<sup>20</sup> that lack the  
28  
29 heterogeneity of walking behaviors observed post-stroke. Additionally, activity levels,  
30  
31 socioeconomic, and ethnic disparities in post-stroke care across geographical locations influence  
32  
33 the walking patterns of the samples engaged in research at different sites<sup>21-25</sup>. Therefore,  
34  
35 combining data across different sites increases sample size and heterogeneity of behaviors  
36  
37 measured in research studies, which ultimately can improve the generalizability of research  
38  
39 findings to the overall post-stroke population.  
40  
41  
42  
43  
44  
45  
46

47           Here, we obtained measures derived from ground reaction forces (GRFs), which is the  
48  
49 simplest data collected across research labs that use instrumented treadmills, to generate a multi-  
50  
51 site sample and used a data-driven approach to identify subgroups of walking behaviors in  
52  
53 people post-stroke and controls. We used sparse K-means clustering to obtain a subset of  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

1  
2  
3 features that define walking clusters, and determined whether different levels of function and  
4  
5 impairment distinguish these walking clusters, such that they are indicative of different walking  
6  
7 subgroups. We determined whether the observed walking clusters are generalizable across  
8  
9 research sites. We hypothesized that participants post-stroke will have different walking features  
10  
11 resulting in different clusters of walking behaviors, which are also different from control  
12  
13 participants<sup>5,14,26</sup>. Our results could provide the basis for designing and testing targeted  
14  
15 interventions aimed at improving walking quality in people post-stroke.  
16  
17  
18  
19  
20  
21  
22

## Methods:

### Data Curation

23  
24  
25  
26  
27  
28  
29  
30 The lack of standardized protocols for collecting, processing, and analyzing walking data  
31  
32 can limit researchers' ability to combine walking data<sup>27</sup>. However, bilateral ground reaction  
33  
34 forces (GRFs) measured using instrumented treadmills are commonly collected across research  
35  
36 sites. These GRFs result from muscles generating forces at each segment which then is applied to  
37  
38 the ground during walking<sup>28,29</sup>, providing insight into how each extremity contributes to the  
39  
40 main objectives of walking, defined as shock absorption, stance stability, and forward  
41  
42 progression<sup>29</sup>. Additionally, these GRFs can be used to derive spatiotemporal walking metrics  
43  
44 such as step lengths, step times, and speed<sup>30</sup>. Thus, we gathered GRF data from individuals with  
45  
46 chronic hemiparetic stroke walking at their self-selected speed. We gathered data from: Rancho  
47  
48 Los Amigos (Rancho, N=7<sup>31</sup>), collected from an overground force plate at 2500 Hz (Kistler  
49  
50 Instrument Corp., Amherst, NY) for the paretic extremity only. University of Southern California  
51  
52 (USC2018 N=22<sup>9</sup> and USC2021 N=23<sup>12</sup>, the first author reviewed all data to ensure no  
53  
54  
55  
56  
57  
58  
59  
60



**Clusters of walking impairment post-stroke**

1  
2  
3 duplicates between studies): participants walked for three minutes on a Bertec instrumented  
4 treadmill (Columbus, OH, USA) that measured ground reaction forces at 1000 Hz. Kennedy  
5 Krieger Institute/Johns Hopkins University School of Medicine (JHU, N=10<sup>19</sup>): participants  
6 walked for five minutes on a Motek instrumented treadmill (Amsterdam, NL) that recorded  
7 forces at 1000 Hz. Emory University (Em, N=9<sup>32</sup>): participants walked for one minute on a  
8 Bertec instrumented treadmill (Columbus, OH, USA) that measured ground reaction forces at  
9 1000 Hz. University of Pittsburgh (Pitt, N=21<sup>33-35</sup>) participants walked between 150 and 320  
10 strides on a Bertec instrumented treadmill (Columbus, OH, USA) that measured ground reaction  
11 forces at 1000 Hz (Fig. 1A). We received GRF data for each gait cycle, normalized to 100 points  
12 per gait cycle, filtered with a fourth order lowpass Butterworth filter with a 15 Hz cut-off  
13 frequency for JHU and with a 20 Hz cutoff for Pitt. Data from Emory were shared as raw data.  
14 We filtered data from Em and USC using a 20 Hz cut-off low-pass zero-lag digital Butterworth  
15 filter.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

We also collected data for 32 age and gender-matched neurotypical control participants using a Bertec instrumented treadmill (Columbus, OH, USA). We used the control data in the definition of the different walking clusters, to determine whether individuals post-stroke could be comparable to neurotypical adults walking at the reduced speed of people post-stroke, such that controls and participants post-stroke would belong to the same clusters. Control participants were age and sex matched to the participants in the USC2021 study<sup>12</sup>.

Participants post-stroke held onto a front handrail during walking in all studies except USC2021<sup>12</sup> and Pitt<sup>33-35</sup>. Control participants did not hold onto a handrail. The respective IRB

## Clusters of walking impairment post-stroke

1  
2  
3 approved all studies, and all participants provided written informed consent before testing. Data  
4 collection for neurotypical control participants was approved by the USC IRB number HS-19-  
5  
6 00430. Data gathering was approved by the USC IRB number HS-19-00075. IRB-approved Data  
7  
8 Use Agreements (DUA) were established between USC and each research institution. Data from  
9  
10 Rancho were excluded as they were collected overground; all DUAs and data sharing manuals  
11  
12 were first implemented with Rancho, which was vital in setting up this project. We accumulated  
13  
14 GRF data for 92 participants post-stroke (Fig. 1A). We compiled information about participant  
15  
16 age, sex, time post-stroke in months, paresis, mass, treadmill walking speed, and lower extremity  
17  
18 Fugl-Meyer score <sup>36</sup> (out of 34 points for the motor scale) (Table 1).  
19  
20  
21  
22  
23  
24  
25  
26

### Data processing

27  
28  
29  
30 From the GRF data collected across labs, we used custom code and derived 17 walking  
31  
32 variables common across all labs. Variable definitions are presented in Table 2. We obtained  
33  
34 averages across strides for all variables for each participant. Data processing was done using  
35  
36 custom code written in Matlab 2021b (The MathWorks, Natick, MA).  
37  
38  
39  
40  
41

42 For data collected in neurotypical adults, leg dominance was defined as the leg they  
43  
44 would use to kick a ball, which was the right leg for all participants. We compared the non-  
45  
46 dominant leg of control participants to the paretic leg and the dominant leg to the non-paretic leg.  
47  
48  
49  
50

51 Stride-by-stride data in a subset of participants post-stroke and controls are available for  
52  
53 download from the Stroke Initiative for Gait Data Evaluation (STRIDE) database<sup>37</sup> hosted by the  
54  
55 Archive of Data on Disability to Enable Policy and Research (ADDEP).  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke**

## Statistical analyses

Statistical analyses were done in RStudio, R version 4.1.2. Since multiple researchers collected data, identifying potential acquisition issues and quality assessment was done post-hoc. To remove noisy data and outliers, we quantified the mean and standard deviation of all variables in our sample, including both participants post-stroke and controls and removed participants with datapoints outside of the mean  $\pm 3$ \*standard deviation range (Supplementary Table 1). The final sample comprised 81 participants post-stroke and 31 controls for 112 participants (Fig. 1A). For all analyses comparing participants post-stroke with neurotypical controls, we used data with neurotypical control participants walking at a matched speed to participants post-stroke.

Given that many walking variables are correlated due to the inherent coordination found in the walking pattern, the clusters found in the data can be accounted for by a small subset of features. Thus, we used sparse K-means clustering analyses<sup>38</sup>, which uses a Lasso penalty<sup>39</sup> to derive a sparse subset of features with non-zero predictors. We z-scored all variables before analyses, and set K=5 clusters in agreement with prior work identifying four subgroups of participants post-stroke<sup>5</sup>, and an additional control group. We verified that K=5 provided clusters that maximized the between clusters distance via the Krzanowski and Lai index<sup>40</sup>. Using the sparcl package in R, we chose the tuning parameter for the Lasso penalty<sup>39</sup>, which determines the number of non-zero predictors to use in our analyses that maximizes between cluster variance and minimizes within cluster variance. We ran 100 permutations in a search space between 1 and the square root of the number of candidate variables included in the sparse

**Clusters of walking impairment post-stroke**

1  
2  
3 clustering analyses, i.e.,  $\sqrt{17}$  and set the number of random starts to 100 to avoid finding a  
4  
5 local minimum.  
6  
7  
8  
9

10 We assessed stability of the clusters identified in this study via the Jaccard similarity  
11 index<sup>41-43</sup>. We resampled the 112 participants with replacement via bootstrap to obtain 10,000  
12  
13 new samples, and identified  $K=5$  clusters for each bootstrap iteration. We then measured the  
14  
15 proportion of observations consistently assigned to the same cluster over each iteration; this  
16  
17 proportion is the Jaccard index. A Jaccard below 0.6 indicates that the clusters are unstable, a  
18  
19 Jaccard between 0.6 and 0.75 shows moderately stable clusters, between 0.75 and 0.85 stable  
20  
21 clusters, and a Jaccard above 0.85 indicates highly stable clusters.  
22  
23  
24  
25  
26  
27

28 We used linear models with *cluster* as a categorical fixed effect to determine whether  
29  
30 there were significant differences across clusters in participants' biomechanical features, as well  
31  
32 as participant demographics and Fugl-Meyer scores. We performed multiple comparisons using  
33  
34 Tukey-Kramer adjusted critical values. We used the results from these multiple comparisons to  
35  
36 characterize the cluster-specific walking behaviors, by identifying which features were  
37  
38 significantly different across clusters. Within each cluster, we used t-tests with false discovery  
39  
40 rate (FDR) corrected p-values for all spatiotemporal and force variables, to compare paretic  
41  
42 values relative to non-paretic values to determine asymmetries in each cluster.  
43  
44  
45  
46  
47  
48

49 We assessed generalizability of each research site by determining the number of  
50  
51 participants from each site that were included in each cluster. To assess whether the observed  
52  
53 clusters can be generalized, we performed a leave-one-out sparse K-means clustering approach,  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

1  
2  
3 leaving out data from each site simultaneously (Em, Pitt, JHU, USC2018, USC2021). We  
4 mapped the clusters obtained in each iteration of the leave-one-out approach to the clusters using  
5 the full dataset. We interpret cluster generalizability based on its Jaccard<sup>43</sup> and the similarity of  
6 the descriptive walking and impairment measures within each cluster.  
7  
8  
9  
10  
11  
12  
13  
14

### Results

15  
16  
17  
18  
19 Our final sample was composed of 112 participants (Fig. 1, Table 1 and 2,  
20 Supplementary Fig. 1). At a group level, participants post-stroke showed significant differences  
21 in paretic swing ( $p=0.007$ ) and stance times ( $p=0.005$ ), medial and lateral ground reaction forces  
22 ( $p<0.001$ ), braking ( $p=0.032$ ), and propulsion ( $p<0.001$ ) compared to the non-paretic extremity.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

35 Sparse K-means clustering identified 8/17 features with non-zero weights from which  
36 five distinct clusters could be obtained (Fig. 2, Supplementary Figs. 2 – 5). The Krzanowski and  
37 Lai index (Supplementary Fig. 2) was maximal for  $K=5$  clusters. Analyses of the clusters  
38 obtained showed different walking behaviors from those observed at a group level. The variables  
39 that maximized between cluster variance and minimized within cluster variance, as shown by the  
40 non-zero weights returned by the Lasso analyses, were paretic and non-paretic stance times, non-  
41 paretic propulsion, speed, paretic step length, paretic and non-paretic braking, and non-paretic  
42 step length (Fig. 2). The data for the paretic extremity was combined with the non-dominant  
43 extremity in controls, and the data for the non-paretic extremity was combined with the  
44 dominant extremity in controls. We describe the obtained clusters next:  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke**

Cluster 1 – Fast speed and asymmetric stroke cluster: N=10 stroke participants.

Composed of four participants from JHU, three from Pitt, two from USC2018 and one from USC2021. Participants had mild impairment (FM  $29\pm 3$ , Fig. 2D).

Walking characteristics: Participants in the fast and asymmetric cluster had the fastest speed ( $0.97\pm 0.17$  m/s,  $p<0.001$ , Fig. 2C, Fig. 3), and were classified as community ambulators<sup>6,44</sup>. Step lengths were  $0.49\pm 0.05$  and  $0.51\pm 0.07$  m for the paretic and non-paretic extremities, the longest of all post-stroke participants. Participants had the largest braking and propulsive GRFs bilaterally ( $p<0.001$ ), and showed marked asymmetries: non-paretic propulsion was 72% greater than paretic propulsion ( $p<0.001$ ), and non-paretic stance ( $p=0.015$ ) and paretic swing times ( $p=0.017$ ) were longer than the contralateral extremity.

Cluster stability and generalizability: The Jaccard index for the fast and asymmetric cluster was 0.72 (Fig. 4), and during the leave-one-out validation varied between 0.63 and 0.75. Leaving out the data from JHU, which was the sample with the highest walking speed, had the greatest effect on cluster stability and reduced the Jaccard to 0.63. Given the absence of the JHU participants, many participants changed cluster membership and the number of participants in the fast and asymmetric cluster increased to 32, and the average speed decreased to 0.78m/s as the speed requirement for this cluster became less stringent. Despite the change in membership, the general characteristics of the fast and asymmetric cluster remained consistent, thus, the cluster is generalizable.

Cluster 2– Moderate speed, symmetric and short stance times in controls and stroke:

N=22 composed of 11 control and 11 post-stroke participants. Participants post-stroke were four

**Clusters of walking impairment post-stroke**

1  
2  
3 from Pitt, one from USC2018 and six from USC2021. Participants post-stroke in the moderate  
4 speed, symmetric cluster had mild impairment (FM 29±4, Fig. 2D).

5  
6 Walking characteristics: Participants in the moderate speed, symmetric cluster walked at  
7  
8 0.61±0.14 m/s (greater than the slow clusters,  $p<0.005$  and less than the fast cluster,  $p<0.001$ ),  
9  
10 classified as least limited community ambulators<sup>6,44</sup>. Step lengths were 0.35±0.06 m for both the  
11  
12 paretic and non-paretic extremities. Additional analyses also showed that stance times were  
13  
14 shorter in the moderate speed, symmetric cluster compared to all other clusters (Fig. 3). Post-  
15  
16 stroke participants in the moderate speed, symmetric cluster were symmetric across all variables  
17  
18 measured, making them comparable to controls walking at this slow speed.

19  
20 Cluster stability and generalizability: The Jaccard index for the moderate speed, symmetric  
21  
22 cluster was 0.63 and varied between 0.6 and 0.87 during validation, bordering on stable to highly  
23  
24 stable. Leaving out the data from Em, JHU, USC2018, and USC2021 made the moderate speed,  
25  
26 symmetric cluster exclusively a control cluster with Jaccard indices of 0.87, 0.63, 0.80, and 0.85.  
27  
28 Leaving out the data from Pitt reduced the Jaccard to 0.6 and reduced the number of control  
29  
30 participants in the moderate speed, symmetric cluster to eight, with one participant from Em,  
31  
32 four from JHU and four from USC. Given that this cluster is primarily a control cluster during  
33  
34 validation and that the Jaccard remains in the moderately stable to the highly stable range, the  
35  
36 moderate speed, symmetric cluster is generalizable and may indicate changes in the walking  
37  
38 pattern that are adopted when people walk at a slower speed.

39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49 Cluster 3 – Moderate speed, asymmetric cluster of participants post-stroke and controls:

50  
51 N=34. 11 controls and 23 participants post-stroke from all sites. Post-stroke participants had mild  
52  
53 impairment (FM 27±3, Fig. 2D).

**Clusters of walking impairment post-stroke**

1  
2  
3 Walking characteristics: Participants in the moderate, asymmetric cluster walked at  $0.69 \pm 0.16$   
4 m/s, classified as least limited community ambulators<sup>6,44</sup> (faster than Cluster 4 and Cluster 5,  
5  $p < 0.005$  and slower than Cluster 1,  $p < 0.001$ ). This was the only cluster that showed cluster-level  
6  
7 step length asymmetry, with longer paretic steps ( $p = 0.032$ ).  
8  
9

10  
11 Stance times were a shorter percentage of the stride sample compared to all other clusters (Fig.  
12  
13 3). Forces were highly asymmetric both for participants post-stroke and control participants in  
14  
15 this cluster. The non-paretic peak lateral GRF was 62% greater than the paretic lateral GRF  
16  
17 ( $p < 0.001$ ), non-paretic propulsion was 62% greater than paretic propulsion ( $p < 0.001$ ), and  
18  
19 paretic braking was 28% greater than non-paretic braking ( $p = 0.001$ ).  
20  
21  
22

23 Cluster stability and generalizability: The Jaccard index for the moderate, asymmetric cluster  
24  
25 using the entire dataset was 0.75 and during the leave-one-out validation varied between 0.51  
26  
27 and 0.81, from unstable to stable. Leaving out the data from Pitt or USC2018 reduced Jaccard  
28  
29 indices to 0.51 (Fig. 4A). Leaving out the data from Em, JHU, USC2018, or USC2021 made the  
30  
31 moderate, asymmetric cluster a primarily control cluster composed of three participants post-  
32  
33 stroke from Pitt and nine controls. In contrast, leaving out the data from Pitt made the moderate,  
34  
35 asymmetric cluster a primarily stroke cluster with three control participants and 13 participants  
36  
37 post-stroke. Given the change in cluster from a control cluster to a stroke cluster during  
38  
39 validation, this cluster is not generalizable across sites.  
40  
41  
42  
43  
44  
45  
46

47 Cluster 4 – Slow speed and frontal plane force asymmetries:  $N = 16$  primarily stroke  
48  
49 participants, three from Em, five from Pitt, three from USC2018, two from USC2021 with  
50  
51 moderate to mild impairment ( $FM 20 \pm 8$ , Fig. 2D). This cluster included three control  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Clusters of walking impairment post-stroke**

1  
2  
3 participants who walked with a frontal plane waddling pattern to account for the slower speeds to  
4  
5 match participants post-stroke.  
6

7  
8 Walking characteristics: Participants in the slow cluster with frontal plane asymmetry walked at  
9  
10 0.35±0.13 m/s (slower than the fast and the two moderate speed clusters,  $p<0.001$ ), classified as  
11  
12 unlimited household ambulators<sup>6,44</sup>. Step lengths were 0.29±0.08 and 0.31±0.09m for the paretic  
13  
14 and non-paretic extremities, respectively. Participants within the slow cluster with frontal plane  
15  
16 asymmetry had stance times that were a longer percentage of the gait cycle compared to all other  
17  
18 participants ( $p<0.001$ ), non-paretic stance was markedly longer than paretic stance ( $p<0.001$ ),  
19  
20 and paretic swing time was also longer than non-paretic swing time ( $p=0.038$ ). The paretic  
21  
22 medial GRF was 25% greater than the non-paretic counterpart ( $p=0.013$ ), and a paretic lateral  
23  
24 GRF that was half of its non-paretic counterpart ( $p=0.005$ , Fig. 3). Non-paretic propulsion was  
25  
26 61% greater than paretic propulsion ( $p=0.006$ ). Participants in the slow cluster with frontal plane  
27  
28 asymmetry generated non-paretic propulsion at around 31% of the gait cycle, significantly later  
29  
30 than all other clusters ( $p<0.0001$ ), and had significantly longer double support times than all  
31  
32 other clusters ( $p<0.001$ ).  
33  
34  
35  
36

37  
38 Cluster stability and generalizability: The Jaccard index for the slow cluster with frontal plane  
39  
40 asymmetry using the entire dataset was 0.87, and during the leave-one-out validation varied  
41  
42 between 0.73 and 0.80, indicating a stable cluster. Given the high stability with the entire dataset  
43  
44 and maintained stability during the leave-one-out approach, this cluster is generalizable across  
45  
46 sites.  
47  
48  
49

50  
51 Cluster 5 – Slow speed, symmetric: N=30, primarily stroke cluster of participants from all  
52  
53 sites with moderate impairment (FM 21±4, Fig. 2D).  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

1  
2  
3 Walking characteristics: Participants in the slow speed symmetric cluster walked at  $0.45 \pm 0.13$   
4 m/s (slower than the fast and moderate speed clusters,  $p < 0.001$ ), classified as the most limited  
5  
6 community ambulators<sup>6,44</sup>. Step lengths were  $0.31 \pm 0.08$  and  $0.31 \pm 0.09$ m for the paretic and non-  
7  
8 paretic extremities, respectively. The percent of the gait cycle in the swing phase was  
9  
10 significantly shorter for participants in the slow speed, symmetric cluster. Participants within the  
11  
12 slow speed symmetric cluster were symmetric except with regards to propulsion, which was 30%  
13  
14 greater in the non-paretic extremity ( $p = 0.005$ ), and the medial GRF, which was greater in the  
15  
16 paretic extremity ( $p = 0.003$ ).

17  
18  
19 Cluster stability and generalizability: The Jaccard index in the slow speed, symmetric cluster  
20  
21 using the entire dataset was 0.73 and during the leave-one-out validation varied between 0.57  
22  
23 and 0.72, from unstable to moderately stable. Leaving out the data from JHU, which contributed  
24  
25 two participants to this cluster, had the greatest effect on cluster stability, increasing the Jaccard  
26  
27 to 0.57 and walking speed to 0.60 m/s. Leaving out the USC2018 data decreased the Jaccard to  
28  
29 0.64 and increased walking speed to 0.63 m/s. Despite the lower Jaccard, the characteristics of  
30  
31 the slow speed symmetric cluster, mainly low propulsion and short swing times remained  
32  
33 consistent, making this cluster generalizable.  
34  
35  
36  
37  
38  
39  
40

## Discussion

41  
42  
43  
44  
45 We identified five clusters of walking behaviors in a combined sample of people post-  
46  
47 stroke and controls. Contrary to our hypothesis, we did not identify a cluster composed  
48  
49 exclusively of control individuals: we identified a cluster with an equal number of control  
50  
51 participants and participants post-stroke, the latter of whom seemed to have reduced walking  
52  
53 impairment. We also identified clusters with mostly participants post-stroke and a few controls,  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

1  
2  
3 which points to the fact that slow walking speeds can lead to aberrant gait patterns even in  
4 healthy controls <sup>45</sup>. The clusters obtained in our study point at different levels of function and  
5  
6 impairment defining each cluster, which correspond to different walking subgroups post-stroke.  
7  
8 We assessed the generalizability of clusters and observed that 4/5 clusters were generalizable  
9  
10 across research sites. These clusters were: Cluster 1: a cluster of post-stroke participants with fast  
11  
12 walking speed and asymmetric propulsion; Cluster 2: a cluster of controls and stroke participants  
13  
14 walking with moderate speed and symmetric steps, with apparent impairments due to reduced  
15  
16 speed, such as short stance times, low propulsion; Cluster 4: a cluster of participants post-stroke  
17  
18 walking with a slow speed and asymmetric medial and lateral ground reaction forces; and Cluster  
19  
20 5: a cluster of post-stroke participants who walked with a slow speed, short swing times and slow  
21  
22 but symmetric steps. It may be possible to develop more personalized intervention targets by  
23  
24 considering the cluster to which a given patient is assigned. For example, a post-stroke  
25  
26 participant in the fast cluster can benefit from an intervention to increase paretic propulsion <sup>17,46–</sup>  
27  
28 <sup>48</sup>; a post-stroke participant in the moderate symmetric cluster can benefit from an intervention to  
29  
30 increase speed <sup>13,49–51</sup>; a post-stroke participant in the slow, frontal asymmetry cluster can benefit  
31  
32 from interventions to reduce asymmetries in medial and lateral GRFs <sup>52</sup>; and a post-stroke  
33  
34 participant in the slow symmetric cluster can benefit from increasing speed <sup>53</sup>, swing times, and  
35  
36 step lengths. The effects of these targeted interventions are still unknown, and future work will  
37  
38 determine whether addressing cluster specific impairments will lead to a shift towards a control  
39  
40 cluster, or a shift towards a different stroke cluster. Taken together, our results provide a  
41  
42 preliminary cross-sectional analyses to estimate which subject-specific walking-related variables  
43  
44 could be targeted in interventions to improve mobility and promote neuroplasticity in the long  
45  
46 term <sup>54</sup>.

**Clusters of walking impairment post-stroke**

1  
2  
3  
4  
5  
6 Our findings complement those of our co-authors who showed speed to be the greatest  
7  
8 determinant in allocating participants to specific clusters <sup>5</sup>, and previous work using speed alone  
9  
10 to classify participants to ambulation categories <sup>6</sup>. Similar to Mulroy, 2003, we found one fast  
11  
12 cluster of participants post-stroke, one moderate speed cluster and two slow clusters. A recent  
13  
14 study used paretic and non-paretic kinematic data in 36 stroke survivors and identified six  
15  
16 distinct walking clusters <sup>55</sup> based on range of motion for the paretic and non-paretic side. Like  
17  
18 our results, Kim et al. observed different types of impairment associated with the different  
19  
20 clusters. A potential limitation of these two studies is that they comprise single-site datasets,  
21  
22 which might be biased by geographical affordances. We complement the clusters described by  
23  
24 Mulroy by providing insights into both paretic and non-paretic spatiotemporal characteristics and  
25  
26 forces generated by each limb. One of our slow clusters had asymmetric mediolateral forces  
27  
28 (Cluster 4), which corresponds to the slow extended cluster in Mulroy 2003: this cluster in  
29  
30 Mulroy 2003 showed knee hyperextension in mid-stance, which limits pre-swing and swing knee  
31  
32 flexion and toe clearance, leading to frontal plane compensations and asymmetric frontal plane  
33  
34 forces, as observed in our study. Our other slow cluster, Cluster 5 also corresponds to the very  
35  
36 slow velocity and excessive knee flexion cluster from Mulroy 2003. Thus, comparison of our  
37  
38 clusters to those previously reported show consistency of subgroups of walking patterns, and  
39  
40 provide additional insights into non-paretic function in these subgroups.  
41  
42  
43  
44  
45  
46  
47  
48

49 The variables that had the largest effect on the between cluster variance, were paretic and  
50  
51 non-paretic stance times, non-paretic propulsion, speed, paretic step length, paretic and non-  
52  
53 paretic braking, and non-paretic step length. It is worth noting that none of these variables on  
54  
55  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke**

1  
2  
3 their own are sufficient to identify the clusters that we observed, similar to what was concluded  
4 by Mulroy, 2003. For example, speed ranges overlapped for participants in Clusters 2 and 3 and  
5  
6 for participants in Clusters 4 and 5, yet different stance times and forces were observed between  
7  
8 clusters despite overlapping speeds, indicating that speed alone was not the only factor driving  
9  
10 the between cluster differences. Thus, here we show that while speed influences many gait  
11  
12 features, speed alone is not enough to classify post-stroke individuals in more specific subgroups  
13  
14 of impairment.  
15  
16  
17  
18  
19  
20

21 Cluster 3, the moderate speed and asymmetric cluster was not stable or generalizable, consistent  
22  
23 with the fact that the common impairments of stroke participants, and how they compare to  
24  
25 control participants within Cluster 3 was less clear. For example, participants in Cluster 3 had  
26  
27 highly asymmetric forces between the paretic and non-paretic extremity whereas control  
28  
29 participants within the cluster were not asymmetric. Similarly, at a group level, participants  
30  
31 within Cluster 3 were the only ones to show marked step length asymmetry with longer paretic  
32  
33 steps. It could be the case that Cluster 3 is composed of participants for whom the measured  
34  
35 spatiotemporal and kinetic variables included in analyses does not account for variability in the  
36  
37 data. Future work will aim to use joint level kinematic, kinetic or EMG measures to determine  
38  
39 whether these more specific measures can detect evident differences in walking patterns in these  
40  
41 individuals.  
42  
43  
44  
45  
46  
47  
48

49 We observed differences in cluster specific speed and FM scores compared to group level  
50  
51 averages in these variables. For example, the average FM score for all participants was  $24.6 \pm$   
52  
53  $5.6$  points, yet as seen in Figure 2D, this average value does not align with any of the cluster-  
54  
55  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke**

1  
2  
3 specific averages. We also observed that similar levels of impairment measured via overlapping  
4  
5 FM scores between clusters could be associated with vastly different walking speeds:  
6  
7 participants in Cluster 1 walked significantly faster than those in Clusters 2 and 3, despite no  
8  
9 differences in FM scores. Clinically this might imply that participants in Clusters 2 and 3 have  
10  
11 the capacity to walk at faster speeds, given their impairment. Note that these differences in  
12  
13 the capacity to walk at faster speeds, given their impairment. Note that these differences in  
14  
15 behavior and capacity between clusters could inform clinical practice beyond group level  
16  
17 averages.

18  
19 Interestingly, our findings show that some commonly reported spatiotemporal impairments  
20  
21 post-stroke are observed in only a subset of post-stroke participants <sup>8,28,61,66</sup>. For example, we  
22  
23 observed asymmetric stance times only for participants in the fast cluster, and asymmetries in  
24  
25 step lengths only in Cluster 3 <sup>11,61,66,67</sup>, as well as varying degrees of paretic propulsion that were  
26  
27 not uniformly associated with walking speed or impairment <sup>56</sup>. Finally, and surprisingly, we did  
28  
29 not observe any asymmetries in the peak vertical GRF among our participants. Thus, our results  
30  
31 show in a sample of 81 participants post-stroke that spatiotemporal asymmetries are less  
32  
33 pronounced than what has been shown in the seminal literature using smaller samples.  
34  
35  
36  
37  
38  
39

40 We observed significant differences in participant impairment and function across research  
41  
42 sites. Consistent with prior literature, control participants' self-selected walking speed was faster  
43  
44 than post-stroke participants<sup>7</sup>. Similarly, participants from Pitt had lower impairment compared  
45  
46 to Emory and USC<sup>2018</sup>, while participants from JHU walked at faster speeds than those from  
47  
48 Em, and USC. Additional information about activity levels and access to post-stroke care across  
49  
50 geographical locations could provide information about the causes of the differences in  
51  
52 impairment and function between sites. Given these differences, we assessed generalizability by  
53  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

1  
2  
3 assessing cluster stability when removing each experimental sample from our dataset. This  
4  
5 approach assessed both generalizability of the cluster such that its definition did not depend on  
6  
7 the experimental samples, as well as generalizability of the experimental samples such that if a  
8  
9 sample needed its own cluster, it would imply that participants in that sample are distinct from  
10  
11 all other participants. We did not find a cluster of participants post-stroke from a single research  
12  
13 site. Additionally, 4/5 clusters were generalizable across research sites. However, we did find  
14  
15 that some research sites did not contribute to specific clusters. For example, participants from  
16  
17 JHU were not part of Cluster 4, and participants from Em were not part of the fast cluster,  
18  
19 consistent with the significant differences in walking speeds observed between both samples.  
20  
21 These results confirm that some samples may not encompass all different walking behaviors  
22  
23 observed after stroke.  
24  
25  
26  
27  
28  
29  
30

### 31 Limitations

32  
33  
34  
35 There are multiple limitations in our study. First, this study used variables derived from  
36  
37 gait analyses to identify clusters indicative of subgroups of walking impairment. The variables  
38  
39 used in this study are easy to capture via GRFs and represent global measures of walking that  
40  
41 relate to the coordinated action of multiple segments and joints. However, joint-level kinematic  
42  
43 and kinetic measures might capture impairments that cannot be inferred from GRF data alone.  
44  
45 Future research efforts should aim to establish standards for data collection that allow combining  
46  
47 more complex data across sites, such as joint-level kinematics, kinetics and EMG patterns.  
48  
49 Second, we extracted peak force as the feature representing GRF data due to the ease to capture  
50  
51 these measures. Future work can assess whether other GRF features, such as impulses, or GRFs  
52  
53 during more specific points during the gait cycle provide more insight into the range of walking  
54  
55  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke**

1  
2  
3 impairments in people post-stroke. Third, we were also limited in the sample size of control  
4 participants which was unbalanced and matched in speed to the USC2021 sample only. The  
5 sample size across study sites was also unbalanced, with samples from Em and JHU consisting  
6 of ~10 participants and the samples from USC and Pitt consisting of ~20 participants, providing  
7 different heterogeneity within each sample and different contribution of each site to the overall  
8 sample. Stability analyses show however, that changes in stability when removing each of the  
9 samples were not just due to sample size: removing the JHU sample which consisted of 10  
10 participants with the highest speeds decreased cluster stability for most clusters, while removing  
11 the USC2018 sample did not change cluster stability uniformly. We interpret this to indicate that  
12 the characteristics of the samples had a greater influence on cluster stability than did sample size.  
13 Future multi-site studies can address these points. Fourth, participants in this study were all in  
14 the chronic phase of stroke recovery; future work should assess if these clusters are consistent or  
15 change across recovery phases. Fifth, data were collected with participants walking on a  
16 treadmill which may induce changes in walking patterns compared to overground walking <sup>57</sup>.  
17 However, some of our clusters were similar to those reported by Mulroy 2003 <sup>5</sup>, indicating  
18 consistency of clusters on the treadmill compared to overground. In addition, the use of a  
19 handrail in some of the experimental protocols could have influenced participant's walking  
20 patterns <sup>58</sup>. Future work could systematically assess whether participants are assigned to the same  
21 clusters measured during treadmill walking as to during overground walking. Finally, we used  
22 average metrics within participants, despite different patterns of stride-to-stride variance during  
23 post-stroke walking <sup>13</sup>. Future work in larger samples can include stride-to-stride variance as  
24 additional features to characterize post-stroke walking patterns.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Clusters of walking impairment post-stroke****Conclusions**

We compiled and curated GRF data across multiple research sites in people post-stroke and controls. Using simple measures derived from GRFs, we identified five clusters of different walking behaviors. Four of these clusters captured walking subgroups that were generalizable across study sites. Our findings provide new information about how to classify the heterogeneity of gait patterns post-stroke. Identifying more specific types of walking impairment and different intervention targets for each subgroup can move the field of neurorehabilitation toward a precision medicine approach<sup>59</sup>, and improve the effectiveness of rehabilitation interventions.

**Acknowledgments**

We want to thank Chang Liu, PhD, Sungwoo Park, PhD, Tara Cornwell, Ryan Novotny, Catherine Yunis, Carly Sombric, PhD, Digna DeKam, PhD, Thu Nguyen, PhD, Justin Liu, Purnima Padmadaban and all trainees who contributed to data collection at all study sites.

This work was funded by:

NIH NCATS grants nos. KL2TR001854 (N Sánchez), R03TR004248 (N Sánchez), and NIH

NCMRR grant no. P2CHD065702 (N Sánchez);

NIH NICHD grant no. R01-HD091184 (J.M. Finley),

NIH NINDS grant no. R21 NS120274 (N. Schweighofer)

NIH NIA grant no. R21 AG059184 (R. Roemmich)

NIH NINCHD grant nos. R01 HD095975 and R21 HD095138 (T. Kesar)

**Clusters of walking impairment post-stroke**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES

1. Cutting JE, Kozlowski LT. Recognizing friends by their walk: Gait perception without familiarity cues. *Bull. Psychon. Soc.* 1977;9:353–356.
2. Winner TS, Rosenberg MC, Kesar TM, Ting LH, Berman GJ. *Discovering individual-specific gait signatures from data-driven models of neuromechanical dynamics*. Neuroscience; 2022. Available at: <http://biorxiv.org/lookup/doi/10.1101/2022.12.22.521665>. Accessed April 26, 2023.
3. Shiavi R, Bugle HJ, Limbird T. Electromyographic gait assessment, part 2: Preliminary assessment of hemiparetic synergy patterns. *Journal of rehabilitation research and development*. 1987;24:24–30.
4. Knutsson E, Richards CL. Different types of disturbed motor control in gait of hemiparetic patients. *Brain*. 1979;102:405–430.
5. Mulroy S, Gronley JA, Weiss W, Newsam C, Perry J. Use of cluster analysis for gait pattern classification of patients in the early and late recovery phases following stroke. *Gait & Posture*. 2003;18:114–125.
6. Perry J, Garrett M, Gronley JK, Mulroy SJ. Classification of walking handicap in the stroke population. *Stroke*. 1995;26:982–989.
7. Olney SJ, Richards C. Hemiparetic gait following stroke. Part I: Characteristics. *Gait and Posture*. 1996;4:136–148.
8. Roerdink M, Beek PJ. Understanding Inconsistent Step-Length Asymmetries Across Hemiplegic Stroke Patients: Impairments and Compensatory Gait. *Neurorehabil Neural Repair*. 2011;25:253–258.
9. Sanchez N, Finley JM. Individual differences in locomotor function predict the capacity to reduce asymmetry and modify the energetic cost of walking post-stroke. *Neurorehabilitation and Neural Repair*. 2018;25:1–13.
10. Balasubramanian CK, Neptune RR, Kautz SA, Bowden MG, Neptune RR, Kautz SA. Relationship Between Step Length Asymmetry and Walking Performance in Subjects With Chronic Hemiparesis. *Archives of Physical Medicine and Rehabilitation*. 2007;88:43–49.
11. Allen JL, Kautz SA, Neptune RR. Step length asymmetry is representative of compensatory mechanisms used in post-stroke hemiparetic walking. *Gait and Posture*. 2011;33:538–543.
12. Park S, Liu C, Sánchez N, Tilson JK, Mulroy SJ, Finley JM. Using Biofeedback to Reduce Step Length Asymmetry Impairs Dynamic Balance in People Poststroke. *Neurorehabilitation and Neural Repair*. 2021;35:738–749.
13. Sanchez N, Schweighofer N, Finley JM. Different biomechanical variables explain within-subjects versus between-subjects variance in step length asymmetry post-stroke. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*. 2021;29:1–1.
14. Kettlety SA, Finley JM, Reisman DS, Schweighofer N, Leech KA. Speed-dependent biomechanical changes vary across individual gait metrics post-stroke relative to neurotypical adults. *bioRxiv*. 2022.

## Clusters of walking impairment post-stroke

15. Kim CM, Eng JJ. Symmetry in vertical ground reaction force is accompanied by symmetry in temporal but not distance variables of gait in persons with stroke. *Gait and Posture*. 2003;18:23–28.
16. Roelker SA, Bowden MG, Kautz SA, Neptune RR. Paretic propulsion as a measure of walking performance and functional motor recovery post-stroke: A review. *Gait and Posture*. 2019;68:6–14.
17. Hsiao H, Knarr BA, Pohlig RT, Higginson JS, Binder-Macleod SA. Mechanisms used to increase peak propulsive force following 12-weeks of gait training in individuals poststroke. *Journal of Biomechanics*. 2016;49:388–395.
18. Allen JL, Kautz SA, Neptune RR. Forward propulsion asymmetry is indicative of changes in plantarflexor coordination during walking in individuals with post-stroke hemiparesis. *Clinical Biomechanics*. 2014;29:780–786.
19. Padmanabhan P, Rao KS, Gulhar S, Cherry-allen KM, Leech KA, Roemmich RT. Persons post-stroke improve step length symmetry by walking asymmetrically. *Journal of NeuroEngineering and Rehabilitation*. 2020;17:1–14.
20. Wonsetler EC, Bowden MG. A systematic review of mechanisms of gait speed change post-stroke. part 1: Spatiotemporal parameters and asymmetry ratios. *Topics in Stroke Rehabilitation*. 2017;24:435–446.
21. Althoff T, Sosič R, Hicks JL, King AC, Delp SL, Leskovec J. Large-scale physical activity data reveal worldwide activity inequality. *Nature*. 2017;547:336–339.
22. Reistetter TA, Kuo YF, Karmarkar AM, et al. Geographic and Facility Variation in Inpatient Stroke Rehabilitation: Multilevel Analysis of Functional Status. *Archives of Physical Medicine and Rehabilitation*. 2015;96:1248–1254.
23. Freburger JK, Holmes GM, Ku LJE, Cutchin MP, Heatwole-Shank K, Edwards LJ. Disparities in postacute rehabilitation care for stroke: An analysis of the state inpatient databases. *Archives of Physical Medicine and Rehabilitation*. 2011;92:1220–1229.
24. Morgenstern LB, Sais E, Fuentes M, et al. Mexican Americans Receive Less Intensive Stroke Rehabilitation Than Non-Hispanic Whites. *Stroke*. 2017;48:1685–1687.
25. Morgenstern LB, Hollenhorst CN, Gallo LC, et al. Ethnic Differences in Informal Caregiving After Stroke. *Stroke*. 2021;52:3938–3943.
26. Kramers De Quervain IA, Simon SR, Leurgans S, Pease WS, McAllister D. Gait pattern in the early recovery period after stroke. *Journal of Bone and Joint Surgery - Series A*. 1996;78:1506–1514.
27. Ferrari A, Grazia M, Pavan E, et al. Quantitative comparison of five current protocols in gait analysis. 2008;28:207–216.
28. John CT, Seth A, Schwartz MH, Delp SL. Contributions of muscles to mediolateral ground reaction force over a range of walking speeds. *Journal of Biomechanics*. 2012;45:2438–2443.
29. Perry J, Burnfield JM. *Gait analysis : normal and pathological function*. 2nd ed. Thorofare, NJ: SLACK; 2010.

## Clusters of walking impairment post-stroke

- 1  
2  
3 30. Zeni JA, Richards JG, Higginson JS. Two simple methods for determining gait events during  
4 treadmill and overground walking using kinematic data. *Gait and Posture*. 2008;27:710–714.  
5
- 6 31. Sullivan KJ, Brown DA, Klassen T, et al. Effects of Task-Specific Locomotor and Strength Training  
7 in Adults Who Were Ambulatory After Stroke: Results of the STEPS Randomized Clinical Trial.  
8 *Physical Therapy*. 2007;87:1580–1602.  
9
- 10 32. Liu J, Kim H Bin, Wolf SL, Kesar TM. Comparison of the Immediate Effects of Audio, Visual, or  
11 Audiovisual Gait Biofeedback on Propulsive Force Generation in Able-Bodied and Post-stroke  
12 Individuals. *Applied Psychophysiology Biofeedback*. 2020;45:211–220.  
13
- 14 33. Nguyen TM, Jackson RW, Aucie Y, et al. Self-selected step length asymmetry is not explained by  
15 energy cost minimization in individuals with chronic stroke. *Journal of NeuroEngineering and  
16 Rehabilitation*. 2020;17:1–27.  
17
- 18 34. De Kam D, Iturralde PA, Torres-Oviedo G. Cerebral Contribution to the Execution , But Not  
19 Recalibration , of Motor Commands in a Novel Walking Environment. *eNeuro*. 2020;7:1–14.  
20
- 21 35. Sombric CJ, Torres-Oviedo G. Augmenting propulsion demands during split-belt walking increases  
22 locomotor adaptation of asymmetric step lengths. *Journal of NeuroEngineering and Rehabilitation*.  
23 2020;17:1–15.  
24
- 25 36. Fugl-Meyer AR. Assessment of motor function in hemiplegic patients. *Neuropsychological aspects of  
26 rehabilitation medicine*. 1976.  
27
- 28 37. Sanchez N. Stroke Initiative for Gait Data Evaluation (STRIDE). *Inter-university Consortium for  
29 Political and Social Research*. 2021. Available at:  
30 <https://www.icpsr.umich.edu/web/ADDEP/studies/38002>.  
31  
32
- 33 38. Witten DM, Tibshirani R, Itten DMW, Ibsirani RT. A Framework for Feature Selection in Clustering  
34 A Framework for Feature Selection in Clustering. 2012;1459.  
35
- 36 39. Tibshirani R. Regression Shrinkage and Selection via the Lasso. *Journal of the Royal Statistical  
37 Society*. 1996;58:267–288.  
38
- 39 40. Krzanowski WJ, Lai YT. A Criterion for Determining the Number of Groups in a Data Set Using  
40 Sum-of-Squares Clustering. *Biometrics*. 1988;44:23.  
41
- 42 41. Hennig C. Cluster-wise assessment of cluster stability. *Computational Statistics and Data Analysis*.  
43 2007;52:258–271.  
44
- 45 42. Yin Y, Yasuda K. Similarity coefficient methods applied to the cell formation problem: A taxonomy  
46 and review. *International Journal of Production Economics*. 2006;101:329–352.  
47
- 48 43. Yu H, Chapman B, Di Florio A, et al. Bootstrapping estimates of stability for clusters, observations  
49 and model selection. *Computational Statistics*. 2019;34:349–372.  
50
- 51 44. Fulk GD, He Y, Boyne P, Dunning K. Predicting Home and Community Walking Activity Poststroke.  
52 *Stroke*. 2017;48:406–411.  
53  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

- 1  
2  
3 45. Chen G, Patten C, Kothari DH, Zajac FE. Gait differences between individuals with post-stroke  
4 hemiparesis and non-disabled controls at matched speeds. *Gait & Posture*. 2005;22:51–56.  
5  
6 46. Hsiao H, Knarr BA, Higginson JS, Binder-Macleod SA. The relative contribution of trailing limb  
7 angle and ankle moment to propulsive force during speed modulation for individuals poststroke. *Journal*  
8 *of NeuroEngineering and Rehabilitation*. 2015;39:212–221.  
9  
10 47. Schenck C, Kesar TM. Effects of unilateral real-time biofeedback on propulsive forces during gait.  
11 *Journal of NeuroEngineering and Rehabilitation*. 2017;14:1–10.  
12  
13 48. Hsiao H, Higginson JS, Binder-Macleod SA. Baseline predictors of treatment gains in peak  
14 propulsive force in individuals poststroke. *Journal of NeuroEngineering and Rehabilitation*. 2016;13:2.  
15  
16 49. Helm EE, Reisman DS. The Split-Belt Walking Paradigm. Exploring Motor Learning and  
17 Spatiotemporal Asymmetry Poststroke. *Physical Medicine and Rehabilitation Clinics of North America*.  
18 2015;26:703–713.  
19  
20 50. Reisman DS, Wityk R, Silver K, Bastian AJ. Locomotor adaptation on a split-belt treadmill can  
21 improve walking symmetry post-stroke. *Brain*. 2007;130:1861–1872.  
22  
23 51. Reisman DS, McLean H, Keller J, Danks KA, Bastian AJ. Repeated split-belt treadmill training  
24 improves poststroke step length asymmetry. *Neurorehabilitation and neural repair*. 2013;27:460–8.  
25  
26 52. Awad LN, Bae J, Kudzia P, et al. Reducing Circumduction and Hip Hiking During Hemiparetic  
27 Walking Through Targeted Assistance of the Paretic Limb Using a Soft Robotic Exosuit. *American*  
28 *Journal of Physical Medicine & Rehabilitation*. 2017;96:1.  
29  
30 53. Moore JL, Nordvik JE, Erichsen A, Rosseland I, Bø E, Hornby TG. Implementation of High-Intensity  
31 Stepping Training during Inpatient Stroke Rehabilitation Improves Functional Outcomes. *Stroke*.  
32 2020;56:563–570.  
33  
34 54. Lim SB, Louie DR, Peters S, Liu-Ambrose T, Boyd LA, Eng JJ. Brain activity during real-time  
35 walking and with walking interventions after stroke: a systematic review. *J NeuroEngineering Rehabil*.  
36 2021;18:8.  
37  
38 55. Kim H, Kim YH, Kim SJ, Choi MT. Pathological gait clustering in post-stroke patients using motion  
39 capture data. *Gait and Posture*. 2022;94:210–216.  
40  
41 56. Roerdink M, Geurts ACH, de Haart M, Beek PJ. On the relative contribution of the paretic leg to the  
42 control of posture after stroke. *Neurorehabilitation and neural repair*. 2010;23:267–274.  
43  
44 57. Kautz SA, Bowden MG, Clark DJ, Neptune RR. Comparison of Motor Control Deficits During  
45 Treadmill and Overground Walking Poststroke. *Neurorehabil Neural Repair*. 2011;25:756–765.  
46  
47 58. IJmker T, Lamoth CJ, Houdijk H, et al. Effects of handrail hold and light touch on energetics, step  
48 parameters, and neuromuscular activity during walking after stroke. *J NeuroEngineering Rehabil*.  
49 2015;12:70.  
50  
51 59. NIH. *National Institutes of Health Research Plan on Rehabilitation: Analysis and Progress.*; 2021.  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke**

60. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*. 1995;57:289–300.

## Clusters of walking impairment post-stroke

## Figure Captions

**Figure 1. Data consolidation and quality assessments and demographics for the final**

**sample.** A) Pipeline for participant inclusion. N indicates the sample size at each point for the total sample (bold) and the sample from each research site. The final sample size was N=112 participants. B) Time post-stroke onset in months across all research sites. Participants from Pitt were more chronic than those from USC2021 ( $p=0.033$ ). C) Post-stroke participant lower extremity Fugl-Meyer score. The maximum score is 34 points. Participants from Pitt were less impaired than those from Em and USC. D) Treadmill self-selected walking speed for all participants. Control self-selected speed was significantly higher than the average speed in the stroke group ( $p<0.001$ ). Control participants also walked at speed matched to that of a stroke participant of the same age and sex (Control Matched). Participants from JHU walked at a significantly greater speed than those from Em and both samples from USC ( $p<0.050$ ).

**Figure 2. The sparse variables make up K=5 clusters with distinct walking speeds and**

**impairments.** A) Weights for each of the 17 candidate features for clustering. 8/17 features had non-zero weights and were used in clustering analyses. B) Individual observations within each cluster, plotted in discriminant component space, colored by cluster. Cluster 1 comprised 10 participants; four from JHU, three from Pitt, two from USC2018, and one from USC2021. Cluster 2, 22 participants; 11 controls, four from Pitt, one from USC2018, and six from USC2021. Cluster 3, 34 participants; 11 control participants, one from Em, four from JHU, four from Pitt, seven from USC2018, and seven from USC2021. Cluster 4 had 16 participants; three controls, three from Em, five from Pitt, three from USC2018, and three from USC2021. Cluster



**Clusters of walking impairment post-stroke**

1  
2  
3 5 had 30 participants; six controls, four from Em, two from JHU, three from Pitt, seven from  
4 USC2018, and seven from USC2021. C) Participant speed (for participants post-stroke and  
5 controls) and impairment (for stroke participants) were measured via FM score across the  
6 different clusters. Only speed was used in the cluster definition. Solid horizontal lines indicate  
7 post-hoc significant differences between clusters ( $p < 0.050$ ). The dashed horizontal line indicates  
8 group-level average speed across all participants and average Fugl-Meyer (FM) scores for all  
9 stroke participants compared with the cluster-level speed and FM. P: Paretic NP: Non-Paretic.  
10 St: Stance. Sw: Swing. Lat: Lateral. Prop: Propulsion. GRF: Ground reaction force.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 **Figure 3. Gait features for participants post-stroke and controls within each cluster**  
25 **compared to all other participants outside each cluster for a subset of sparse variables used**  
26 **in k-means clustering.** C1, ... C5 indicates participants included in the respective cluster  
27 number, and ~C1, ... ~C5 indicates all participants not included in the respective cluster. Stance  
28 and swing times are expressed as a percentage of stride duration to account for differences in  
29 walking speed. Cluster 1: participants post-stroke with a fast-walking speed and asymmetric  
30 propulsion; Cluster 2: participants post-stroke and controls with moderate speed, short stance  
31 times, low propulsion, and symmetric steps. Cluster 3: participants post-stroke and controls with  
32 moderate speed, short stance times, and asymmetric forces; Cluster 4: participants post-stroke  
33 with a slow speed and frontal plane force asymmetries; and Cluster 5: post-stroke participants  
34 who walked slowly and symmetrically, with short swing times. Color conventions as in Figure 3.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49 \* FDR corrected  $p < 0.010$  for all variables indicated as significant. Abbreviations as in Figure 2.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Clusters of walking impairment post-stroke****Figure 4. Stability and generalizability of clusters assessed via leave one out approach. A)**

Jaccard stability index for each cluster for the entire dataset (horizontal black lines) and leaving out each sample. Colors indicate the sample left out to assess the Jaccard. The dashed gray line is the line above which clusters are considered stable. The Jaccard Index is the proportion of observations consistently assigned to the same cluster over the bootstrap iterations. A Jaccard below 0.6 indicates that the clusters are unstable, a Jaccard between 0.6 and 0.75 shows moderately stable clusters, 0.75 and 0.85 stable clusters, and a Jaccard above 0.85 indicates highly stable clusters. B) Average and standard deviation of the speed for each cluster for the entire dataset in black and the leave one out approach with colors indicating the sample left out from analyses.

## Clusters of walking impairment post-stroke

Table 1: Participant Demographics.

	Stroke	Control
<b>N</b>	81	31
<b>Sex</b>	28F/53M <sup>†</sup>	19F/12M
<b>Age</b>	58.9 ± 10.5 [32 – 77] years	63.3 ± 13.6 [24 – 81]
<b>Mass</b>	85 ± 18.6* [47 – 131] kg	73 ± 15.7 [46 – 110]
<b>Speed</b>	0.58 ± 0.24 [0.13 – 1.25] m/s	SS: 0.84 ± 0.18* [0.48 – 1.43] m/s Matched: 0.60 ± 0.25 [0.3 – 1.0] m/s
<b>Paresis</b>	42R/39L	
<b>Time post-stroke</b>	90 ± 83 [6 – 467] months	
<b>Fugl-Meyer Score</b>	24.6 ± 5.6 [7 – 33]	

Descriptive statistics are presented as average ± standard deviation with the range in brackets.

F: Female

M: Males

L: Left

R: Right

SS: Self-selected

\*Significant differences between participants post-stroke and controls (p<0.050)

<sup>†</sup>p<0.050 significant difference in the frequency of males compared to females in the study sample

## Clusters of walking impairment post-stroke

**Table 2: Group-level averages for gait variables in participants post-stroke and control participants walking at matched speeds.**

Variable	Definition	Stroke		Control	
		Paretic Mean $\pm$ SD	Non-Paretic Mean $\pm$ SD	Non-dominant Mean $\pm$ SD	Dominant Mean $\pm$ SD
<b>Speed</b>	Treadmill speed	0.58 $\pm$ 0.24		0.60 $\pm$ 0.18	
<b>Step Length (m)</b>	Distance between limbs at leading limb initial contact	0.37 $\pm$ 0.10	0.36 $\pm$ 0.11	0.39 $\pm$ 0.09	0.39 $\pm$ 0.10
<b>Stance Time (s)</b>	Time between initial contact and lift-off on the same side	0.97 $\pm$ 0.25 <sup>*,+</sup>	1.00 $\pm$ 0.28 <sup>+</sup>	0.86 $\pm$ 0.23	0.87 $\pm$ 0.23
<b>Swing Time (s)</b>	Time as the time between lift-off to initial contact on the same side	0.47 $\pm$ 0.09 <sup>*</sup>	0.43 $\pm$ 0.07	0.45 $\pm$ 0.06	0.45 $\pm$ 0.06
Double Support Time (s)	Time from contralateral initial contact to ipsilateral foot-off	0.29 $\pm$ 0.13 <sup>+</sup>	0.24 $\pm$ 0.19	0.20 $\pm$ 0.08	0.21 $\pm$ 0.09
<b>Peak Medial GRF (N/kg)</b>	medially directed force during the contralateral toe-off <sup>28</sup>	0.93 $\pm$ 0.22 <sup>*</sup>	0.82 $\pm$ 0.20	0.85 $\pm$ 0.28	0.82 $\pm$ 0.26
<b>Peak Lateral GRF (N/kg)</b>	Laterally directed GRF during loading response	-0.24 $\pm$ 0.21 <sup>*</sup>	-0.39 $\pm$ 0.24 <sup>+</sup>	-0.20 $\pm$ 0.18	-0.21 $\pm$ 0.17
<b>Peak Propulsive GRF (N/kg)</b>	Force during ipsilateral push-off	0.67 $\pm$ 0.34 <sup>*,+</sup>	1.03 $\pm$ 0.46 <sup>+</sup>	0.87 $\pm$ 0.34	0.87 $\pm$ 0.34
<b>Peak Braking GRF (N/kg)</b>	Force during weight acceptance/weight transfer	-0.86 $\pm$ 0.44 <sup>*</sup>	-0.75 $\pm$ 0.44	-0.80 $\pm$ 0.33	-0.84 $\pm$ 0.31
<b>Vertical GRF (N/kg)</b>	Force due to gravity	10.18 $\pm$ 1.05	10.15 $\pm$ 0.99	10.2 $\pm$ 2.50	10.7 $\pm$ 2.68
% Gait Cycle of Peak Propulsive GRF	Relative to paretic limb initial contact	47 $\pm$ 12%	19 $\pm$ 22%	57 $\pm$ 9%	11 $\pm$ 12%
% Gait Cycle of Peak Braking GRF	Relative to paretic limb initial contact	16 $\pm$ 16%	52 $\pm$ 17%	16 $\pm$ 3%	66 $\pm$ 4%

t-tests corrected for multiple comparisons using the Benjamini and Hochberg FDR<sup>60</sup>.

\*Significant differences between the paretic and non-paretic extremity of post-stroke participants

<sup>+</sup>Significant differences between participants post-stroke and controls

Variables in bold were included in sparse clustering analyses.




Figure 1. Data consolidation and quality assessments and demographics for the final sample. A) Pipeline for participant inclusion. N indicates the sample size at each point for the total sample (bold) and the sample from each research site. The final sample size was N=112 participants. B) Time post-stroke onset in months across all research sites. Participants from Pitt were more chronic than those from USC2021 ( $p=0.033$ ). C) Post-stroke participant lower extremity Fugl-Meyer score. The maximum score is 34 points. Participants from Pitt were less impaired than those from Em and USC. D) Treadmill self-selected walking speed for all participants. Control self-selected speed was significantly higher than the average speed in the stroke group ( $p<0.001$ ). Control participants also walked at speed matched to that of a stroke participant of the same age and sex (Control Matched). Participants from JHU walked at a significantly greater speed than those from Em and both samples from USC ( $p<0.050$ ).

177x124mm (300 x 300 DPI)




Figure 2. The sparse variables make up K=5 clusters with distinct walking speeds and impairments. A) Weights for each of the 17 candidate features for clustering. 8/17 features had non-zero weights and were used in clustering analyses. B) Individual observations within each cluster, plotted in discriminant component space, colored by cluster. Cluster 1 comprised 10 participants; four from JHU, three from Pitt, two from USC2018, and one from USC2021. Cluster 2, 22 participants; 11 controls, four from Pitt, one from USC2018, and six from USC2021. Cluster 3, 34 participants; 11 control participants, one from Em, four from JHU, four from Pitt, seven from USC2018, and seven from USC2021. Cluster 4 had 16 participants; three controls, three from Em, five from Pitt, three from USC2018, and three from USC2021. Cluster 5 had 30 participants; six controls, four from Em, two from JHU, three from Pitt, seven from USC2018, and seven from USC2021. C) Participant speed (for participants post-stroke and controls) and impairment (for stroke participants) were measured via FM score across the different clusters. Only speed was used in the cluster definition. Solid horizontal lines indicate post-hoc significant differences between clusters ( $p < 0.050$ ). The dashed horizontal line indicates group-level average speed across all participants and average Fugl-Meyer (FM) scores for all stroke participants compared with the cluster-level speed and FM. P: Paretic NP: Non-Paretic. St: Stance. Sw: Swing. Lat: Lateral. Prop: Propulsion. GRF: Ground reaction force.

165x165mm (300 x 300 DPI)




Figure 3. Gait features for participants post-stroke and controls within each cluster compared to all other participants outside each cluster for a subset of sparse variables used in k-means clustering. C1, ... C5 indicates participants included in the respective cluster number, and ~C1, ... ~C5 indicates all participants not included in the respective cluster. Stance and swing times are expressed as a percentage of stride duration to account for differences in walking speed. Cluster 1: participants post-stroke with a fast-walking speed and asymmetric propulsion; Cluster 2: participants post-stroke and controls with moderate speed, short stance times, low propulsion, and symmetric steps. Cluster 3: participants post-stroke and controls with moderate speed, short stance times, and asymmetric forces; Cluster 4: participants post-stroke with a slow speed and frontal plane force asymmetries; and Cluster 5: post-stroke participants who walked slowly and symmetrically, with short swing times. Color conventions as in Figure 3. \* FDR corrected p < 0.010 for all variables indicated as significant. Abbreviations as in Figure 2.

126x185mm (300 x 300 DPI)




Figure 4. Stability and generalizability of clusters assessed via leave one out approach. A) Jaccard stability index for each cluster for the entire dataset (horizontal black lines) and leaving out each sample. Colors indicate the sample left out to assess the Jaccard. The dashed gray line is the line above which clusters are considered stable. The Jaccard Index is the proportion of observations consistently assigned to the same cluster over the bootstrap iterations. A Jaccard below 0.6 indicates that the clusters are unstable, a Jaccard between 0.6 and 0.75 shows moderately stable clusters, 0.75 and 0.85 stable clusters, and a Jaccard above 0.85 indicates highly stable clusters. B) Average and standard deviation of the speed for each cluster for the entire dataset in black and the leave one out approach with colors indicating the sample left out from analyses.

168x68mm (300 x 300 DPI)



## Clusters of walking impairment post-stroke

## Supplementary Materials


Supplementary Table 1. Demographics for participants excluded from analyses.

Group	Speed (m/s)	Mass* (kg)	Age (years)	Sex	FM Score	Affected Side	Months Post-Stroke
Emory	0.45	71.4	74	F	26	L	24
Pitt	1.11	74.5	75	M	32	R	370
Pitt	0.53	53.9	66	F	26	R	117
USC	0.13	67.0	28	F	19	L	25
Control	1.05 (matched 0.13)	51.4	28	F			

\*p<0.05 significant differences for the independent samples t-test comparing the sample of participants excluded vs. the sample of participants included in analyses


Clusters of walking impairment post-stroke

Supplementary Figure 1



**Supplementary Figure 1. Demographics.** A) Participant age in years across all sites tested. B) Participant mass in kilograms. Mass in control participants was significantly lower than in post-stroke participants ( $p=0.002$ ). C) Self-reported sex across different sites. D) Paretic side for participants post-stroke across different sites.


## Supplementary Figure 2



**Supplementary Figure 2.** A) Krzanowski and Lai index for D-Index plots for the dataset in our study. The optimal number of clusters is the one with the highest index. B) Mean and 95% confidence intervals for the Krzanowski and Lai index obtained via 1000 bootstrap iterations. Bootstrap analyses indicate overlapping confidence intervals for the Krzanowski and Lai index for 2, 5, 6 and 13 clusters, which are also higher than all other number of clusters. Given previous work identifying 4 stroke clusters plus our group of control participants we maintain K=5 clusters.

## Clusters of walking impairment post-stroke


## Supplementary Figure 3



**Supplementary Figure 3. Participant demographics across the different clusters.** A) Age in years for participants post-stroke and controls in each cluster. Participants in C1 were marginally younger than C2 ( $p=0.051$ ). B) Mass in kg for participants post-stroke and controls across clusters. C) Time post-stroke in months for participants post-stroke. No differences were observed across clusters ( $p=0.925$ ). Horizontal lines indicate post-hoc differences between clusters. D) Sex for participants post-stroke and controls across clusters for all participants. E) Paresis across clusters for participants post-stroke.

Clusters of walking impairment post-stroke

Supplementary Figure 4




**Clusters of walking impairment post-stroke**

1  
2  
3  
4  
5 **Supplementary Figure 4. Spatiotemporal and peak forces across the different clusters for**  
6 **participants post-stroke and controls.** Post-hoc significant differences between clusters are  
7 indicated by the solid horizontal lines ( $p < 0.050$ ). The dashed horizontal line indicates the  
8 average value of the variable across all 112 participants to allow comparisons of group-level  
9 averages with cluster-level averages. DST refers to double support time, and SLS refers to single  
10 limb support time, which was available for all participants except those from Pitt and therefore  
11 were not used as candidate variables for clustering. All other variables were used as candidate  
12 variables in sparse analyses.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Clusters of walking impairment post-stroke

Supplementary Figure 5



**Supplementary Figure 5. Temporal variables are expressed as the percent duration of the overall stride.** Post-hoc significant differences between clusters are indicated by the solid horizontal lines ( $p < 0.050$ ). The dashed horizontal line indicates the average value of the variable across all 112 participants for stance and swing times and 93 participants for double and single support times to allow comparisons of group-level averages with cluster-level averages. DST refers to double support time, and SLS refers to single limb support time, which was available for all participants except those from Pitt.