




Article

Accumulation of Heavy Metal Ions from Urban Soil in Spontaneous Flora

Maria Harja ¹, Ramona Carla Ciocinta ², Gabrijel Ondrasek ³, Daniel Bucur ^{4,*} and Marcel Dirja ⁵

¹ Faculty of Chemical Engineering and Environmental Protection, “Gheorghe Asachi” Technical University of Iasi, 700050 Iasi, Romania

² Regional Directorate of Road and Bridge from Iasi, 700481 Iasi, Romania

³ Department of Soil Amelioration, Faculty of Agriculture, The University of Zagreb, 10000 Zagreb, Croatia

⁴ Department of Pedotechnics, Faculty of Agriculture, Iasi University of Life Sciences, 700490 Iasi, Romania

⁵ Department of Land Surveys and Exact Sciences, Faculty of Forestry and Cadastre, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, 400372 Cluj-Napoca, Romania

* Correspondence: dbucur@uaiasi.ro

Abstract: This study aimed to analyse the bioaccumulation of heavy metal ions in plants of spontaneous flora. An urban area was established from which soil samples were taken and analyses were performed on the content of heavy metal ions. The soil samples were collected from Iasi’s urban area to determine the heavy metal concentrations, obtain maps of sensitive land, and determine the content of heavy metals in spontaneous flora and transfer these into an aqueous extract. The investigation of heavy metal ion levels in certain plants revealed the following: (i) all plants from the *Brassicaceae* accumulate heavy metals, and *B. juncea* has a great ability to accumulate and transfer Cu, Cr(VI), Cd, Ni, Pb, and Zn towards the shoots; (ii) heavy metals (Fe, Cu, Cr, Mn, Zn) were present in variable concentrations, with mint and nettle being notable for their increased level of iron and thyme, and rattle for its zinc levels; (iii) toxic metals (Pb, Cd, Ni) are present in low concentrations in plants as well as in infusions, except for in primula and plantain, which do not have high levels of Pb and Cd. The results showed that values exceeded the maximum recommended values in areas with industrial pollution. Taking into account the potential for the bioaccumulation of heavy metal ions by plants from spontaneous flora and their use as medicinal plants, it is recommended, based on the studies conducted, to harvest and use plants from soils that do not contain heavy metal ions.

Keywords: soil contamination; heavy metals; medicinal plants; herbs; chemical composition



Citation: Harja, M.; Ciocinta, R.C.; Ondrasek, G.; Bucur, D.; Dirja, M. Accumulation of Heavy Metal Ions from Urban Soil in Spontaneous Flora. *Water* **2023**, *15*, 768. <https://doi.org/10.3390/w15040768>

Academic Editor: Saglara S. Mandzhieva

Received: 31 December 2022

Revised: 6 February 2023

Accepted: 13 February 2023

Published: 15 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The chemical composition of soil is extremely diverse, and soils contain practically all known chemicals [1]. Organic-mineral soil constituents are characterised by organic complexes with metal ions (Fe^{2+} , Cu^{2+} , Mn^{2+}) and associations formed between clays and humic and non-humic organic substances [2–4]. The pollutants released into all soils reach a distance of between 20 and 50 km. Soils support large amounts of pollutants from sludge from wastewater treatment plants, sludge from ports, waste from farms, etc. Waste from different industrial activities causes soil pollution with organic and inorganic compounds [5–7]. Due to their high toxicity and long-term stability in the environment, the accumulation of heavy metals in soil, and consequently in spontaneous flora, is a significant environmental problem [8]. Through ingesting food, breathing in particles, inhaling particles, and absorption through the skin, the bioaccumulation of heavy metals may enhance the dangers to human health [9,10]. Heavy metals are highly toxic to living organisms [7]. Above a specific point, toxic consequences occur, whereas below that point, some elements (Co, Cu, Fe, Ni, and Zn) may even be necessary building blocks for proteins involved in different metabolic pathways [11]. Heavy metal pollution is a global problem that is now of great interest for both industrial sites and for protected natural areas [12–14].

Heavy metal pollution is caused primarily by the metallurgical and ferrous industries, which release large amounts of dust into the atmosphere, and also by solid-fuel boilers, cement factories, road transport, waste dumps, and tailing deposits [15–17].

In Romania, heavy metals represented by oxides and gases pollute industrial areas around cities such as Baia Mare [18], Zlatna [19], and Copșa Mică [20,21]. In Baia Mare and Zlatna, pollution is mainly caused by lead [18], and it is caused by cadmium, zinc, lead, and copper in Copșa Mică [20]. In Copșa Mică, economic activities with a metallurgical and chemical profile determined soil pollution with heavy metals on more than 8000 ha, of which 1500 ha were found with cadmium, 422 ha were found with zinc, 335 ha were found with lead, and 32 ha were found with copper [21]. After 1990, metallurgical activities were drastically reduced and even stopped. However, the locals cultivate these lands, a fact that may affect their health [21].

In the last two decades, the concentration of lead in the human body in some areas has increased 200-fold, even exceeding the recommended concentration. Regarding the studied area, the main sites contaminated with heavy metals and hydrocarbons in the Iasi area are represented by the lands on which the Arcelor Mittal plant is located, as well as in the location of the former non-compliant municipal waste deposit at Tomesti (15.3 ha), which is located 8 km from the city of Iasi [22,23]. The deposit was closed in 2015 as part of the “Integrated Waste Management System” project, in which measures were adopted to ensure good soil quality and protect the health of the population and the environment.

Heavy metals can enter the soil or plants from fertilisers, supplements, and pesticides used in food production, or from gases released into the atmosphere from industry and by combustion [24]. Worldwide, a large number of plants that can bioaccumulate heavy metal ions from the soil have been tested both in the laboratory and in field conditions [25–29]. Currently, there are over 400 recognized species of plants that hyperaccumulate metals. Some species and genotypes can grow on lands that contain increased concentrations of heavy metals, and they hyperaccumulate metals that either occur there naturally or are there due to human activity [30,31]. Heavy metal pollution has particularly strong effects on land [32]; therefore, it is useful to know the content of heavy metals in urban lands to accurately establish their pollution levels. Several studies have reported soil contamination by heavy metals released due to industrial activities, especially in urban areas [33,34], and these raise dangers to human health [35,36]. Negative health effects on the population living in an area with contaminated soils have been observed.


Different studies have been conducted on the heavy metals in urban soils [23,37], but there is an absence of a link between soil pollution and heavy metal content in different plants. In this regard, soil samples were collected from the Iasi urban area for the following reasons: (1) to explore the characteristics of soil and heavy metal concentrations, (2) to generate maps of heavy metals in the studied areas, and (3) to determine the content of heavy metals in spontaneous flora. This study analysed, for the first time in an urban area, the possibility of bioaccumulation of heavy metal ions in plants from spontaneous flora. The findings demonstrated that plants grown in industrially polluted soils contain significant amounts of heavy metals that are transferred in aqueous media.

The water-soluble fractions of bioaccumulated heavy metals present toxicity risks, because the studied plants are used for herbal and medicinal teas and in different modern and traditional foods as aromatic herbs. In the studied area, Cr, Cu, Ni, Pb, and Zn are enriched in the soil samples; however, their content is below the potential human exposure limits. If levels of heavy metal cations are low risk for individual health, the presence of multiple cations exceeded the acceptable limit.


2. Materials and Methods

The study area was 16 km² and included the built-up area of Iasi and its surroundings (Figure 1a). The population numbers almost 400,000 inhabitants, with a density of about 150 inhabitants/km². Geochemical distribution maps of potentially polluting heavy elements were constructed using GIS methods for interpolating discrete data. For graphical

representation, the ESRI ArcGIS Desktop platform was used [38]. A map of land use is presented in Figure 1b.



(a)



(b)

Figure 1. (a) Sample collection grid from the study area (16 km²) including the built-up area of Iasi and its surroundings (1030 soil samples were taken at a depth of 0–25 cm from a network of 500 m); (b) land use.

2.1. Soil Sample Preparation

The level of contamination was assessed based on the collection and analysis of 1030 soil samples, collected at a depth of 0–25 cm, from the city of Iasi and the neighbouring areas, within a radius of 500 m. The collected soil samples were naturally dried at room temperature (20 °C). After this, the soil samples were ground with an agate mortar, sieved, and stored in polypropylene containers as samples for analysis.

The method described in the literature to determine the net concentration of heavy metals in soil was used to undertake acidic extraction [39]. A 250 cm³ glass vessel containing 10.0 g of soil sample was mixed with 50 cm³ of diluted nitric acid (1:1), and the mixture was then homogenised. The glass was covered and boiled for 10 min on low heat. The mixture was cooled to room temperature, then 10 cm³ of hydrogen peroxide was added dropwise under stirring. The glass content was heated and boiled under weak heating for 10 min, then cooled and filtered on a G4 filter funnel into a volumetric flask with a capacity of 100 cm³. The precipitate was mineralised with nitric acid following the described procedure, and finally, the volume of the filtrate was diluted to 100 cm³ with bidistilled water. Sample analysis was performed via atomic absorption spectroscopy using an iCE 3400 AAS Spectrometer (Thermo Fisher Scientific, Waltham, MA, USA), and fluorescence spectrometry RX was carried out with an EDXRF (Thermo Fisher Scientific™ Niton™ XL3t XRF-Analyzer, Waltham, MA, USA). Soil samples were stored in glass containers for further study and during further processing. Specifically, the plants were rinsed with deionised water and dried first in the shade and then in an oven at 110 °C for 2–4 h. The dried and ground samples were subsequently stored in bottles for further study. During all of these steps, necessary measures were taken to avoid any loss or contamination of the samples with heavy metals.

2.2. Spontaneous Flora Sample Preparation

A furnace was used to heat 1 g of ground plant material (shoots) for 5 h at 550 °C. After cooling under desiccating conditions, the crucible contents were added, and 2.5 mL 6M HNO₃ was added to dissolve the samples. The necessary glassware was thoroughly cleaned using regular soap and tap water before being submerged in an acid bath (30% nitric acid) and dried in a fume hood. Glassware was washed with deionised water. The sample solution was filtered, diluted to a total volume of 20 mL, and analysed with an atomic absorption spectrophotometer. In the laboratory, the soil content of the heavy metals Cd, Cr, Cu, Hg, Ni, Pb, and Zn was determined. For some of these heavy metals, maps of their accumulation distribution in soil were made. The determination of heavy metals in plants was carried out via UV/VIS and atomic absorption spectrophotometry [40]. After harvesting, the plants were washed with distilled water, dried at room temperature, and then processed by wet digestion (conc. HNO₃).

2.3. Methods for Heavy Metals' Determination from Aqueous Extract

The aqueous extracts were obtained by infusing the dried plants (1.00 g) with 100 mL of drinking water (Timisesti-Iasi source). The resulting infusions using all 11 plants from spontaneous flora were analysed by atomic absorption spectrophotometry to evaluate their content of the following elements: Cu, Cr, Cd, Pb, Fe, Mn, Zn, and Ni.

Principle of operation: With the specific reagent and analysis purity, the heavy metals form a substance of a certain colour through which, depending on the nature of the lamp, a light beam is transmitted. The device measures the absorbance at a certain wavelength and determines the concentration of the desired substance in the solution [41,42]. The absorption of light by a coloured substance is crossed by a beam of light. It is translated with the formula:

$$A = \lg \frac{I_0}{I} = -\lg T = \varepsilon \cdot b \cdot C \quad (1)$$

where I and I_0 represent light intensity, A —absorbance, C —concentration (mg L⁻¹),

T —transmittance, b —the thickness of the absorbent layer traversed (centimetres), and ε —extension coefficient.

Reagent and method blanks, duplicates, and standard reference materials were used in the quality assurance and control of chemical analyses. The reference materials were from CPA Chem. Ltd., Stara Zagora, Bulgaria. Standard solutions were used for calibration; when necessary, digested soil solutions were diluted by a factor of 10 to fit the calibration range.

Each element underwent calibration using a linear regression result (R^2) of 0.999. The least amount of random and systematic error was possible. To have trust in the precision, repeatability, and dependability of the results, after calibration curves were developed, all of the tests were run twice.

To find the relevance of the analytical results, standard statistical procedures were applied: the arithmetic mean of the measurements, absolute error, mean deviation, and standard deviation were applied; a propagation technique was applied to eliminate unreliable data error [43,44].

The legislation that established the permissive value of heavy metals content is Council Directive 86/278/EEC for Protection of the Environment (European Communities Council 1986) and the Romanian Regulation of Allowable Quantities of Hazardous and Harmful Substance in Soil (Order of the Ministry of Waters, Forests, and Environmental Protection No. 756/3 November 1997). According to the Romanian legislation, the intervention thresholds for soils with less sensitive use are 10 mg kg⁻¹ Cd, 600 mg kg⁻¹ Cr, 1500 mg kg⁻¹ Zn, 500 mg kg⁻¹ Cu, 4000 mg kg⁻¹ Mn, 1000 mg kg⁻¹ Pb, 500 mg kg⁻¹ Ni, and 250 mg kg⁻¹ Co [45]. As for the content of heavy metals in plants, it is regulated in the documents of the European Commission. For example, the maximum cadmium content allowed in non-leafy Brassica is 0.04 mg/kg wet weight, and in leafy Brassica, it is 0.1 mg/kg wet weight [46].

3. Results and Discussion

3.1. Heavy Metal Concentrations in Soils

Soils from the north part of the studied area were formed in excessive temperate climate conditions through an intense bioaccumulation process, and chernozems were dominant in the area. The chemical characteristics of the soil, such as organic carbon content, pH, oxide forms, carbonates, and some physical properties such as clay content, might influence the concentration of chemical elements [47]. The soil content of the heavy metals Cr, Cu, Ni, Pb, and Zn was analysed, determined, and interpreted [48–52].

Chromium is recognized to have lithophile geochemical affinity [53]; the maximum recommended value for total Cr content in soil is 30 mg kg⁻¹. The alert threshold (AT) for sensitive land (SL) use is 100 mg kg⁻¹, and for less sensitive land (LSL), it is 300 mg kg⁻¹. The distribution of Cr in the investigated soils appeared to be very close to the natural state, because 99.32% of the samples showed values within the normal range, whereas those that exceeded the alert threshold represented only 0.68% of samples, and those that exceeded the intervention threshold (IT) represented 0.48% of samples (Figure 2).

Copper has a chalcophile affinity that leads to its association in some minerals and ores with other heavy metals such as Pb, Zn, Mo, Ag, and Ni [54]. In the topsoil, a good correlation exists between Cu and Al, Ni, Cr, Mn, and Zn. The maximum recommended value for the total content of Cu in soil is 20 mg kg⁻¹. The AT for SL is 100 mg kg⁻¹, and for LSL, it is 250 mg kg⁻¹. The IT is at 200 mg kg⁻¹ for SL and 500 mg kg⁻¹ for LSL (Figure 3).

In the study area, the natural distribution of Cu contained values that not only exceeded the threshold value (5.73% of samples) but also the IT (1.94% of samples). These values determine the geochemical halos in chernozem soils located outside the southern part of Iasi city.




Figure 2. Map of Cr distribution in the analysed soil (mg kg^{-1}).




Figure 3. Map of Cu distribution in the analysed soil (mg kg^{-1}).

Nickel has a chalcophile geochemical affinity, the latter explaining the association of Ni with Co, Cu, and Pt in some sulphide ores [55]. The maximum limit for total Ni content in soil is 20 mg kg^{-1} . The AT for SL use is 75 mg kg^{-1} , and for LSL use, it is 200 mg kg^{-1} . The IT is 150 mg kg^{-1} for SL and 500 mg kg^{-1} for LSL. Throughout the studied soil profiles, the Ni content is presented in Figure 4. Samples for which the Ni content exceeded the alert threshold have a punctuated arrangement in the investigated area, an arrangement that is spatially related to the occurrence of chernozems.




Figure 4. Map of Ni distribution in the analysed soil (mg kg^{-1}).

Lead has a geochemical affinity for chalcophiles [56], which explains why it is found in sulphide ores with Fe, Zn, Cu, Sb, and Ag. The maximum limit for total Pb content in soil is 20 mg kg^{-1} . The AT for SL is 50 mg kg^{-1} , and for LSL land, it is 250 mg kg^{-1} . The IT is 100 mg kg^{-1} for sensitive land and 1000 mg kg^{-1} for less sensitive land. The Pb content determined in the study area varied within a fairly wide range, from 4.5 to $1995.4 \text{ mg kg}^{-1}$ (Figure 5).

Furthermore, 92.52% of the analysed samples had a Pb content ranging within normal limits; only 5.73% of samples had Pb content higher than the alert limit, and 1.75% of samples exceeded the IT.

Zinc interacts with several heavy metals, including Cu, Pb, Ag, Au, Sb, As, and Se, in certain sulphide ores [57]. The maximum recommended limit for the total Zn content in soil is 100 mg kg^{-1} . The AT for SL is 300 mg kg^{-1} , and for LSL, it is 700 mg kg^{-1} . The IT is 600 mg kg^{-1} for SL and 1500 mg kg^{-1} for LSL. Since zinc is a necessary nutrient for most plants, algae in rivers and lakes may absorb most of the dissolved zinc in water. However, excessive zinc in soils can be hazardous to many plant species. The Zn content determined for soils from Iasi city and its surroundings varied widely ($10.1\text{--}5624 \text{ mg kg}^{-1}$), with a majority of samples (96.5%) in which the content did not exceed normal values for soils. Samples with a Zn content that exceeded the AT (2.04%) and were higher than the IT (1.46%) belonged to soils in the urban areas with a southern exposure. Differences in the Zn content spatially overlapped mostly with those of Pb, suggesting that they were generated by the same common pollutant sources (Figure 6). In the study area, soil

pollution was characterised by the presence of three heavy metals, Pb, Zn and Cu, above the recommended limits.




Figure 5. Map of Pb distribution in the analysed soil (mg kg^{-1}).




Figure 6. Map of Zn distribution in the analysed soil (mg kg^{-1}).

We can conclude that heavy-metal pollution in Iasi was caused primarily by solid fuel boilers and road transport. For Zn, three genera and 20 species of plants are known to show hyperaccumulation. This group also includes the *Thlaspi* genus, which is known for hyperaccumulating different types of metals. For example, *T. caerulescens* accumulates Cd, Ni, Pb, and Zn; *T. geosingense* accumulates Ni and Zn; *T. ochroleucum* accumulates Ni and Zn; and *T. rotundifolium* accumulates Ni, Pb, and Zn.

3.2. Concentrations of Heavy Metals in Plants from the Spontaneous Flora

Many hyperaccumulators belong to Brassicaceae and are capable of accumulating large amounts of heavy metal ions [58]. Kumar et al. [59] analysed several Brassicaceae species with rapid growth, such as Indian mustard, black mustard, turnip, rape, and kale, for their ability to tolerate and accumulate metals. All plants from Brassicaceae accumulate metals, but *B. juncea* specifically has a great capacity to accumulate and transfer Cu, Cr(VI), Cd, Ni, Pb, and Zn to the shoots. The main families of metal-hyperaccumulator plants are Asteraceae, Brassicaceae, Caryophyllaceae, Cyperaceae, Fabaceae, Lamiaceae, Poaceae, Violaceae, and Euphorbiaceae [60]. The plants from Brassicaceae are the most numerous, comprising 11 genera and 87 species. A hyperaccumulator is defined as a plant that phytoaccumulates a minimum of 0.01% Cd and As, or a minimum of 0.1% Co, Cu, Cr, Zn, or Pb, or 1% Mn and Ni. The investigation of heavy metal ion levels in certain plants revealed the following: (i) All species from the Brassicaceae accumulate metals. From these, *B. juncea* has a great ability to accumulate and transfer Cu, Cr(VI), Cd, Ni, Pb, and Zn. (ii) Heavy metals (Fe, Cu, Cr, Mn, Zn) are present in variable concentrations, with mint and nettle having a notably increased level of iron and thyme, and rattle a higher level of zinc. (iii) Toxic metals (Pb, Cd, Ni) are present at low concentrations in plants as well as in infusions, except in primula and plantain, which do not have high levels of Pb and Cd.

In this study, 11 plants of spontaneous flora were collected from 2018–2019 from areas in which the existence of heavy metal ions was determined: mint (*Folium menthae*), chamomile (*Flores chamomillae*), yarrow (*Flores miellefolii*), field thyme (*Herba cichorii*), nettle (*Folium urticae*), dandelion (*Folium taraxaci*), celandine (*Herba chelidonii*), plantain (*Folium plantaginis*), primula (*Flores primulae*), shepherd's purse (*Herba bursae pastoris*), and rattle (*Herba hyperici*). Analytical data revealed that of the five herbs used in nutrition, mint and nettle contained the highest amounts of iron of $80.52 \pm 5.06 \mu\text{g g}^{-1}$ dry matter (DM) and $49.62 \pm 3.82 \mu\text{g g}^{-1}$, respectively. Zinc was present in higher concentrations in field thyme ($967.71 \pm 6.28 \mu\text{g g}^{-1}$) and rattle ($811.09 \pm 9.69 \mu\text{g g}^{-1}$), and the maximum manganese content was found in rattle ($55.66 \pm 5.15 \mu\text{g g}^{-1}$). Chromium and copper were detected at lower concentrations (Table 1).

Table 1. Mineral elements in the studied plants ($\mu\text{g g}^{-1}$).

Heavy Metals Ions	Iron	Chromium	Copper	Manganese	Zinc
<i>Folium urticae</i>	49.62 ± 3.82	2.19 ± 0.13	6.74 ± 0.45	14.98 ± 0.49	30.40 ± 4.12
<i>Folium taraxaci</i>	31.11 ± 1.41	5.21 ± 0.42	2.78 ± 0.11	14.33 ± 0.34	49.35 ± 2.01
<i>Herba cichorii</i>	22.08 ± 1.63	10.44 ± 0.80	2.09 ± 0.24	20.54 ± 1.78	967.61 ± 6.28
<i>Herba hyperici</i>	10.23 ± 1.27	1.84 ± 0.82	3.31 ± 0.61	55.66 ± 5.15	811.09 ± 9.69
<i>Folium menthae</i>	80.52 ± 5.06	2.67 ± 0.84	4.15 ± 0.31	25.67 ± 0.96	309.63 ± 45.1
Min	8.96	1.02	1.85	13.99	26.28
Max	85.58	11.25	4.46	60.81	973.28
Mean	38.712	4.47	3.814	26.236	433.616
Standard deviation	27.44327	3.5897	1.80067	17.0806	26.35

3.3. Concentrations of Heavy Metals in Aqueous Extract

In the aqueous extracts obtained from plantain and primula, no Pb or Cd were found. The rest of the extracts contained Pb, Cd, and Ni in concentrations between 0.0014 – 0.0283 mg L^{-1} . The water used for the preparation of plant extracts contained only iron in low concentrations (0.0073 mg L^{-1}).

The content of heavy metals in aqueous extract is presented in Figure 7.




Figure 7. The content of heavy metals in aqueous extract of analysed herbs.

From Figure 7, it can be seen that each herb accumulates and can solubilise other heavy metals. For example, in aqueous extract of nettle, dandelion, celandine, and plantain, high concentrations of lead (between 3.38 and 8.89 mg L⁻¹), were found, and in aqueous extract of yarrow and primula, nickel was found. The results are in accordance with the literature [61].

4. Conclusions

In urban residential areas, the soil can be polluted by a wide range of pollutants, resulting from industrial sources and human activities.

Beyond the potential toxicity of contaminated soil to human health through direct contact, contact through inhalation or through the plants from spontaneous flora is very important.

The results of the present investigation revealed that the concentrations of heavy metals (Cu, Cr(VI), Cd, Ni, Pb, and Zn) found in soil samples taken from Iasi, Romania suggest high levels of Pb and Zn, above the alert thresholds in limited areas, as required by Romanian legislation.

The investigation of heavy metal levels in some plants revealed the following:

- The presence of heavy metals (Fe, Cu, Cr, Mn, Zn) was detected in variable concentrations, with mint and nettle being notable for their increased level of iron and thyme and rattle for an increased level of zinc;
- Potentially toxic metals (Pb, Cd, Ni) are present at low concentrations in plants as well as in infusions, except in primula and plantain, which do not contain Pb and Cd.

In order to avoid contamination and potentially negative effects on the environment and on human health, it is crucial to analyse and monitor the amounts of heavy metals in soil and in plants from spontaneous flora suitable for consumption. The obtained information should be helpful in recommending the implementation of remediation strategies and in getting heavy metal contamination in soil and plants from spontaneous flora under control.

Taking into account the potential for the accumulation of heavy metal ions by plants from the spontaneous flora in the area and these same plants' use in medicine and based on the studies conducted, it is recommended to harvest and use the plants from soils that do not contain heavy metal ions, considering the expansion of inhabited areas in the studied metropolitan area.

Author Contributions: Conceptualisation, M.H. and D.B.; methodology, G.O.; software, R.C.C.; validation, M.H.; formal analysis, R.C.C.; investigation, M.D.; data curation, M.H.; writing—original draft preparation, M.H. and D.B.; writing—review and editing, M.D.; visualisation, G.O.; supervision, M.H.; project administration, M.H.; funding acquisition, G.O. All authors have read and agreed to the published version of the manuscript.

Funding: Subsidiary contract no. 8.D.119611/05.07.2022-S.C. DICOR LAND S.R.L. This research constitutes a partial phase of the contract entitled “Research on the Efficiency of the Use of Different Types of Irrigation Facilities in Agriculture” (partial phase). The realisation of the study is part of Type D collaborative activities—industrial research and/or experimental development activities carried out by the research organisation in effective collaboration with the enterprise, as well as Subactivity D.1—industrial research activities in effective collaboration within the project “Constitution and implementation of partnerships for knowledge transfer between the Research Institute for Agriculture and Environment Iasi and the agricultural economic environment”—acronym AGRIECOTEC, SMIS code 119611, financing contract no. 4/1.2.3G/5 June 2018. Project co-financed by the European Fund for Regional Development through the Competitiveness Operational Program 2014–2020.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Wang, Z.; Qin, H.; Wang, J. Accumulation of uranium and heavy metals in the soil–plant system in Xiazhuang uranium ore field, Guangdong Province, China. *Environ. Geochem. Health* **2019**, *41*, 2413–2423. [[CrossRef](#)] [[PubMed](#)]
2. Adimalla, N. Heavy metals contamination in urban surface soils of Medak province, India, and its risk assessment and spatial distribution. *Environ. Geochem. Health* **2019**, *42*, 59–75. [[CrossRef](#)] [[PubMed](#)]
3. Vinceković, M.; Šegota, S.; Jurić, S.; Harja, M.; Ondrasek, G. Development and Characterization of a Novel Soil Amendment Based on Biomass Fly Ash Encapsulated in Calcium Alginate Microspheres. *Int. J. Mol. Sci.* **2022**, *23*, 9984. [[CrossRef](#)]
4. Harja, M.; Cimpeanu, S.M.; Dirja, M.; Bucur, D. Synthesis of zeolites from fly ash and their use as soil amendment. In *Zeolites—Useful Minerals*; IntechOpen: London, UK, 2016.
5. Meng, C.; Wang, P.; Hao, Z.; Gao, Z.; Li, Q.; Gao, H.; Feng, F. Ecological and health risk assessment of heavy metals in soil and Chinese herbal medicines. *Environ. Geochem. Health* **2022**, *44*, 817–828. [[CrossRef](#)]
6. Buema, G.; Harja, M.; Lupu, N.; Chiriac, H.; Forminte, L.; Ciobanu, G.; Bucur, R.D. Adsorption performance of modified fly ash for copper ion removal from aqueous solution. *Water* **2021**, *13*, 207. [[CrossRef](#)]
7. Harja, M.; Buema, G.; Bucur, D. Recent advances in removal of Congo Red dye by adsorption using an industrial waste. *Sci. Rep.* **2022**, *12*, 6087. [[CrossRef](#)]
8. Anaman, R.; Peng, C.; Jiang, Z.; Liu, X.; Zhou, Z.; Guo, Z.; Xiao, X. Identifying sources and transport routes of heavy metals in soil with different land uses around a smelting site by GIS based PCA and PMF. *Sci. Total Environ.* **2022**, *823*, 153759. [[CrossRef](#)]
9. Hagh Nazar, H.; Belmont, P.; Johannesson, K.H.; Aghayani, E.; Mehraein, M. Human-induced pollution and toxicity of river sediment by potentially toxic elements (PTEs) and accumulation in a paddy soil-rice system: A comprehensive watershed-scale assessment. *Chemosphere* **2023**, *311*, 136842. [[CrossRef](#)]
10. Wu, J.; Li, J.; Teng, Y.; Chen, H.; Wang, Y. A partition computing-based positive matrix factorization (PC-PMF) approach for the source apportionment of agricultural soil heavy metal contents and associated health risks. *J. Hazard. Mater.* **2020**, *388*, 121766. [[CrossRef](#)]
11. Bisht, V.K.; Uniyal, R.C.; Sharma, S.M. Assessment of heavy metals content in herbal raw materials traded in India. *S. Afr. J. Bot.* **2022**, *148*, 154–161. [[CrossRef](#)]
12. Harja, M.; Bucur, D.; Cimpeanu, S.M.; Ciocinta, R.C. Conversion of ash on zeolites for soil application. *J. Food Agric. Environ.* **2012**, *10*, 1056–1059.
13. Jeong, H.I.; Han, J.E.; Shin, B.C.; Jang, S.B.; Won, J.H.; Kim, K.H.; Sung, S.H. Monitoring of Heavy Metals and Pesticide Residues of Herbal Decoctions in Traditional Korean Medicine Clinics in Korea. *Int. J. Environ. Res. Public Health* **2022**, *19*, 8523. [[CrossRef](#)] [[PubMed](#)]
14. Savić, R.; Maksimović, L.; Cimpeanu, S.; Bucur, D.; Ondrašek, G.; Vasin, J.; Harja, M.; Milić, S. Hazardous and harmful substances in sediments of the Jegricka stream. *J. Food Agric. Environ.* **2013**, *11*, 1152–1156.

15. Esfandbod, M.; Forghani, A.; Adhami, E.; Rezaei Rashti, M. The Role of CEC and pH in Cd Retention from Soils of North of Iran. *Soil Sediment Contam.* **2011**, *20*, 908–920. [[CrossRef](#)]
16. Harja, M.; Rusu, L.; Bucur, D.; Ciocinta, R.C. Fly ash-derived zeolites as adsorbents for Ni removal from wastewater. *Rev. Roum. Chim.* **2012**, *56*, 84–90.
17. Zhou, J.M.; Dang, Z.; Cai, M.F.; Liu, C.Q. Soil heavy metal pollution around the Dabaoshan Mine, Guangdong Province, China. *Pedosphere* **2007**, *17*, 588–594. [[CrossRef](#)]
18. Sur, I.M.; Micle, V.; Polyak, E.T.; Gabor, T. Assessment of soil quality status and the ecological risk in the Baia Mare, Romania area. *Sustainability* **2022**, *14*, 3739. [[CrossRef](#)]
19. Iosob, G.A.; Bute, A.; Calara, M.; Tremurici, A.A.; Benchea, C.M.; Bouruc, D.; Muscalu, S.P. Heavy metals contamination of soil and vegetables in three regions from Romania: A review. *Sci. Study Res. Biol.* **2020**, *29*, 79–87.
20. Hoaghia, M.A.; Cadar, O.; Moisa, C.; Roman, C.; Kovacs, E. Heavy metals and health risk assessment in vegetables grown in the vicinity of a former non-metallic facility located in Romania. *Environ. Sci. Pollut. Res.* **2022**, *29*, 40079–40093. [[CrossRef](#)]
21. Oprea, B.Ş.; Motelică, D.-M.; Vrînceanu, N.O.; Costea, M.; Plopeanu, G.I.; Carabulea, V. Research on the heavy metal content in onion bulbs correlated with soil from private households located in the Copşa Mică area, Central Romania. *J. Appl. Life Sci. Environ.* **2022**, *55*, 92–99. [[CrossRef](#)]
22. Carabulea, V.; Motelică, D.-M.; Vrînceanu, N.O.; Costea, M.; Plopeanu, G.I.; Oprea, B.Ş. Bioaccumulation of heavy metals in garlic bulbs (*Allium sativum* L.) in correlation with soil from private household in the Copşa Mică area, Romania. *J. Appl. Life Sci. Environ.* **2022**, *55*, 245–255. [[CrossRef](#)]
23. Secu, C.V.; Juravle, D.T.; Paiu, M.; Lesenciuc, D.C.; Costica, M.; Negru, D.; Breaban, I.G. Potentially toxic elements in urban soils of Iasi (Romania). *Environ. Eng. Manag. J.* **2016**, *15*, 687–698.
24. Onache, P.A.; Ionete, R.E.; Iordache, A.M.; Florea, A.; Sumedrea, D.I.; Iliescu, M.; Tanase, A. Monitoring the Content of Heavy Metals in The Soil and Wines from Different Wine-Growing Areas of Romania. *Int. Multidiscip. Res. Anal.* **2022**, *5*, 343–354.
25. Chhotu, J.; Fulekar, M. Phytoremediation of heavy metals: Recent techniques. *Afr. J. Biotechnol.* **2009**, *8*, 921–928.
26. Grispen, V.M.J.; Nelissen, H.J.M.; Verkleij, J.A.C. Phytoextraction with *Brassica napus* L.: A tool for sustainable management of heavy metal contaminated soils. *Environ. Pollut.* **2006**, *144*, 77–83. [[CrossRef](#)] [[PubMed](#)]
27. Liu, H.B.; Guo, P.T.; Wu, W.; Wang, Z.Y. Assessment of Soil Arsenic, Chromium, Mercury, and Lead at an Agricultural Landscape Scale. *Soil Sediment Contam.* **2011**, *20*, 995–1007. [[CrossRef](#)]
28. Nadgorska-Socha, A.; Ptasiński, B.; Kita, A. Heavy metal bioaccumulation and antioxidative responses in *Cardaminopsis arenosa* and *Plantago lanceolata* leaves from metalliferous and non-metalliferous sites: A field study. *Ecotoxicology* **2013**, *22*, 1422–1434. [[CrossRef](#)]
29. Schmidt, W. Biodiversity and plant productivity in a grassland succession: Effects of nutrient levels and disturbance regimes. *Pol. Bot. Stud.* **2006**, *22*, 437–448.
30. Wang, G.; Yan, X.; Zhang, F.; Zeng, C.; Gao, D. Traffic-Related Trace Element Accumulation in Roadside Soils and Wild Grasses in the Qinghai-Tibet Plateau, China. *Int. J. Environ. Res. Public Health* **2013**, *11*, 456–472. [[CrossRef](#)]
31. Yan, X.; Zhang, F.; Zeng, C.; Zhang, M.; Prasad Devkota, L.; Yao, T. Relationship between Heavy Metal Concentrations in Soils and Grasses of Roadside Farmland in Nepal. *Int. J. Environ. Res. Public Health* **2012**, *9*, 3029–3226. [[CrossRef](#)]
32. Sarim, M.; Jan, T.; Khattak, S.A.; Mihoub, A.; Jamal, A.; Saeed, M.F.; Radicetti, E. Assessment of the Ecological and Health Risks of Potentially Toxic Metals in Agricultural Soils from the Drosh-Shishi Valley, Pakistan. *Land* **2022**, *11*, 1663. [[CrossRef](#)]
33. Zhang, K.; Chai, F.; Zheng, Z.; Yang, Q.; Zhong, X.; Fomba, K.W. Size distribution and source of heavy metals in particulate matter on the lead and zinc smelting affected area. *J. Environ. Sci.* **2018**, *71*, 188–196. [[CrossRef](#)] [[PubMed](#)]
34. Liu, Z.; Du, Q.; Guan, Q.; Luo, H.; Shan, Y.; Shao, W. A Monte Carlo simulation-based health risk assessment of heavy metals in soils of an oasis agricultural region in northwest China. *Sci. Total Environ.* **2023**, *857*, 159543. [[CrossRef](#)] [[PubMed](#)]
35. Jurowski, K.; Krośniak, M. The Toxicological Assessment of Content and Exposure of Heavy Metals (Pb and Cd) in Traditional Herbal Medicinal Products with Marshmallow Root (*Althaea officinalis* L. radix) from Polish Pharmacies. *Toxics* **2022**, *10*, 188. [[CrossRef](#)]
36. Wu, L.; Yue, W.; Wu, J.; Cao, C.; Liu, H.; Teng, Y. Metal-mining-induced sediment pollution presents a potential ecological risk and threat to human health across China: A meta-analysis. *J. Environ. Manag.* **2023**, *329*, 117058. [[CrossRef](#)]
37. Xue, S.; Korna, R.; Fan, J.; Ke, W.; Lou, W.; Wang, J.; Zhu, F. Spatial distribution, environmental risks, and sources of potentially toxic elements in soils from a typical abandoned antimony smelting site. *J. Environ. Sci.* **2023**, *127*, 780–790. [[CrossRef](#)]
38. Zhao, Y.; Zhao, X.; Huang, X.; Guo, J.; Chen, G. Identifying a Period of Spatial Land Use Conflicts and Their Driving Forces in the Pearl River Delta. *Sustainability* **2023**, *15*, 392. [[CrossRef](#)]
39. Safarova, V.I.; Shaidullina, G.F.; Mikheeva, T.N.; Kudasheva, F.K.; Nizamutdinova, N.R. Methods of sample preparation of soil, bottom sediments, and solid wastes for atomic absorption determination of heavy metals. *Inorg. Mater.* **2011**, *47*, 1512–1517. [[CrossRef](#)]
40. Sadee, B.A.; Ali, R.J. Determination of heavy metals in edible vegetables and a human health risk assessment. *Environ. Nanotechnol. Monit. Manag.* **2023**, *19*, 100761.
41. Sevik, H.; Ozel, H.B.; Cetin, M.; Özel, H.U.; Erdem, T. Determination of changes in heavy metal accumulation depending on plant species, plant organism, and traffic density in some landscape plants. *Air Qual. Atmos. Health* **2019**, *12*, 189–195. [[CrossRef](#)]

42. Casasanta, G.; Falcini, F.; Garra, R. Beer–Lambert law in photochemistry: A new approach. *J. Photochem. Photobiol. A* **2022**, *432*, 114086. [[CrossRef](#)]
43. Santos-Francés, F.; Martínez-Graña, A.; Ávila Zarza, C.; García Sánchez, A.; Alonso Rojo, P. Spatial distribution of heavy metals and the environmental quality of soil in the Northern Plateau of Spain by geostatistical methods. *Int. J. Environ. Res. Public Health* **2017**, *14*, 568. [[CrossRef](#)] [[PubMed](#)]
44. Agyeman, P.C.; Kebonye, N.M.; John, K.; Haghaziar, H.; Borůvka, L.; Vašát, R. Compositional mapping, uncertainty assessment, and source apportionment via pollution assessment-based receptor models in urban and peri-urban agricultural soils. *J. Soils Sediments* **2022**, 1–22. [[CrossRef](#)]
45. Petrean, I.A.; Micle, V.; Sur, I.M.; Şenilă, M. Characterization of Sterile Mining Dumps by the ICP-OES Analytical Method: A Case Study from Baia Mare Mining Area (Maramures, Romania). *Sustainability* **2023**, *15*, 1158. [[CrossRef](#)]
46. Commission Regulation (EU) 2021/1323 of 10 August 2021 Amending Regulation (EC) No 1881/2006 as Regards Maximum Levels of Cadmium in Certain Foodstuff. Official Journal of the European. 11.08.2021 L228/13. Available online: <http://data.europa.eu/eli/reg/2021/1323/oj> (accessed on 10 December 2022).
47. Shi, S.; Hou, M.; Gu, Z.; Jiang, C.; Zhang, W.; Hou, M.; Xi, Z. Estimation of Heavy Metal Content in Soil Based on Machine Learning Models. *Land* **2022**, *11*, 1037. [[CrossRef](#)]
48. Cabrini, T.M.; Barboza, C.A.; Skinner, V.B.; Hauser-Davis, R.A.; Rocha, R.C.; Saint' Pierre, T.D.; Cardoso, R.S. Investigating heavy metal bioaccumulation by macrofauna species from different feeding guilds from sandy beaches in Rio de Janeiro, Brazil. *Ecotoxicol. Environ. Saf.* **2018**, *162*, 655–662. [[CrossRef](#)] [[PubMed](#)]
49. Chen, Y.; Jiang, X.; Wang, Y.; Zhuang, D. Spatial characteristics of heavy metal pollution and the potential ecological risk of a typical mining area: A case study in China. *Process Saf. Environ. Prot.* **2018**, *113*, 204–219. [[CrossRef](#)]
50. Mortensen, L.H.; Rønn, R.; Vestergård, M. Bioaccumulation of cadmium in soil organisms—With focus on wood ash application. *Ecotoxicol. Environ. Saf.* **2018**, *156*, 452–462. [[CrossRef](#)]
51. Wang, K.; Qiao, Y.; Zhang, H.; Yue, S.; Li, H.; Ji, X.; Liu, L. Bioaccumulation of heavy metals in earthworms from field contaminated soil in a subtropical area of China. *Ecotoxicol. Environ. Saf.* **2018**, *148*, 876–883. [[CrossRef](#)]
52. Shi, H.; Wang, P.; Zheng, J.; Deng, Y.; Zhuang, C.; Huang, F.; Xiao, R. A comprehensive framework for identifying contributing factors of soil trace metal pollution using Geodetector and spatial bivariate analysis. *Sci. Total Environ.* **2023**, *857*, 159636. [[CrossRef](#)]
53. Tang, Q.; Zhang, H.; Zhao, X.; Zheng, L.; Miao, C.; Liu, Y.; Fu, B. Chromium in Chinese coals: Geochemistry and environmental impacts associated with coal-fired power plants. *Environ. Geochem. Health* **2022**, 1–22. [[CrossRef](#)] [[PubMed](#)]
54. Osei, L.B.; Fosu, S.; Ndur, S.A.; Nyarko, S.Y. Assessing heavy metal contamination and ecological risk of urban topsoils in Tarkwa, Ghana. *Environ. Monit. Assess.* **2022**, *194*, 710. [[CrossRef](#)] [[PubMed](#)]
55. Adhami, E.; Salmanpour, A.; Omidi, A.; Khosravi, N.; Ghasemi-Fasaei, R.; Maftoun, M. Nickel adsorption characteristics of selected soils as related to some soil properties. *Soil Sediment Contam.* **2008**, *17*, 643–653. [[CrossRef](#)]
56. Yang, X.; Wen, E.; Ge, C.; El-Naggar, A.; Yu, H.; Wang, S.; Rinklebe, J. Iron-modified phosphorus-and silicon-based biochars exhibited various influences on arsenic, cadmium, and lead accumulation in rice and enzyme activities in a paddy soil. *J. Hazard. Mater.* **2023**, *443*, 130203. [[CrossRef](#)]
57. Xu, J.; Li, Y.; Wang, S.; Long, S.; Wu, Y.; Chen, Z. Sources, transfers and the fate of heavy metals in soil-wheat systems: The case of lead (Pb)/zinc (Zn) smelting region. *J. Hazard. Mater.* **2023**, *441*, 129863. [[CrossRef](#)] [[PubMed](#)]
58. Dwivedi, S.K. Medicinal herbs: A potential source of toxic metal exposure for man and animals in India. *Arch. Environ. Health* **2002**, *57*, 229–231. [[CrossRef](#)]
59. Kumar, P.B.A.N.; Dushenkov, V.; Motto, H.; Raskin, I. Phytoextraction: The use of plants to remove heavy metals from soils. *Environ. Sci. Technol.* **1995**, *29*, 1232–1238. [[CrossRef](#)]
60. Mrak, T.; Grašič, B.; Prislán, P.; Gričar, J.; Laznik, Ž.; Voglar, G.E. Soil contamination with potentially toxic elements and root herbivory: Effects on root surface area and stem secondary xylem of young common beech (*Fagus sylvatica* L.). *Acta Physiol. Plant.* **2023**, *45*, 23. [[CrossRef](#)]
61. Gao, L.; Di, D.; Liu, X.; Teng, F. Comparative Study of Heavy Metals Analysis in Mongolian Medicines Based on High Sensitivity X-ray Fluorescence Spectroscopy and ICP-MS. *Spectroscopy* **2022**, *37*, 20–27. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.