



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

How to Assess Degradation of Freshwater Communities due to Hydromorphological Changes

Friberg, Nikolai; Sandin, Leonard; Pedersen, Morten Lauge

Published in:

Risk Assessment in European River Basins - State of the Art and Future Challenges

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Friberg, N., Sandin, L., & Pedersen, M. L. (2008). How to Assess Degradation of Freshwater Communities due to Hydromorphological Changes. In Risk Assessment in European River Basins - State of the Art and Future Challenges: 12-14 November 2007, Leipzig, Germany (pp. 36-39). Helmholtz, Centre for Environmental Research, UFZ.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

How to assess degradation of freshwater communities due to hydromorphological changes

Nikolai Friberg¹, Leonard Sandin², Morten L. Pedersen³

¹Macaulay Institute, Catchment Management Group, Aberdeen, Scotland, UK, n.friberg@macaulay.ac.uk

²Swedish Agricultural University, Uppsala, Sweden

³Aalborg University, Aalborg, Denmark

Background

Man-made modifications to in-stream channel morphology are one of the main reasons that rivers do not achieve good ecological quality. Channelisation, dredging and weed cutting have seriously degraded habitat diversity thereby excluding some species and reduce numbers of others. In a Water Framework Directive context these are known as hydromorphological pressures and to mitigate these pressure are identified as being of key importance in fulfilling the Directive.

The current physical structure of streams and diversity of biological communities are closely linked past and present human activities in the entire catchment. Human activities influence stream ecosystems on multiple scales, ranging from direct manipulation of the in-stream environment on the stream reach to altering the landscape and land use in the catchment thereby influencing the hydrological pathways and morphological structure (Vannote et al., 1980; Frissell et al., 1986; Fitzpatrick et al., 2001; Allan, 2004). Past and present disturbances act simultaneously with different intensity on the stream ecosystem elements and it can thus be difficult to distinguish the exact disturbance from the individual stressors on the biotic communities (Lane & Richards, 1997; Harding et al., 1998; Allan, 2004).

Thesis 1: Physical-biological coupling are important in determining freshwater communities

The scientific literature reports many linkages between individual parameters (substratum, current velocity etc.) describing the in-stream physical environment and different attributes of the macroinvertebrate community. On a fine scale using high resolution measurements without other confounding factors, the importance of physical-biological coupling in streams have been shown albeit many aspects are still not fully understood (Hart & Finelli 1999). One reason is that numerous physical factors interact across different temporal and spatial scales, resulting in very different biotopes (riffles, pools etc.) as well as inducing variation within these biotopes. One example of the latter is shown in figure 1 as the response of two macroinvertebrate metrics (total abundance and EPT taxa abundance) to mean particle size varied considerably between to adjacent riffle biotypes in the same stream. The majority of studies have shown that diversity and abundance of macroinvertebrates increase with particle size as this increase habitat complexity and volume of interstitial spaces. In one riffle (B and D) there was the expected positive relationship between particle size and macroinvertebrate metrics whereas there was no relationship in the other riffle (A and C), the reason being that it was consolidated by fine sediments reducing habitat availability. This study illustrates that even within the same biotope that would appear completely similar when visually assessed there can be considerable differences in the physical-biological coupling.




Fig. 1: Relationship between median particle size and total macroinvertebrate abundance/EPT abundance in two riffles spaced 100 m apart in the same 3. order stream. A,C is the upstream riffle whereas B,D is the downstream riffle. From Pedersen & Friberg (2007).

Thesis 2: Multiple stressors interact and this will influence impacts of hydromorphological degradation

One key issue when assessing the influence of hydromorphology is that interpretation of results is often confounded by multiple stressors influencing freshwater communities. Typically, streams that have undergone a high degree of habitat degradation or alterations of flow regimes will be situated in areas with multiple anthropogenic pressures. A recent analysis of 1127 stream sites in three European countries showed that the strongest relationship to a biological metric (ASPT) was found when including both land use, chemistry and habitat modification in a multiple regression (Table 1). In addition, stressors can interact in a synergistically manner and increased concentrations of easy degradable organic matter will have a more detrimental impact in a habitat degraded stream because number of reactive surfaces and re-aeration capacity are reduced (Andersen, 1994).

Table 1: Relationship between the macroinvertebrate based metric ASPT and explanatory environmental variables (PTOT=total phosphorous concentration; ARABLE=percentage agriculture in the catchment; MD=morphological degradation). Data from the EU REBECCA project (SSPI-CT-2003-502158).

Metric	Countries	Relationship	R ²
ASPT	Sweden Slovakia Denmark	$y = 5.26 - 3.33PTOT - 0.014ARABLE + 2.32MD$	0.61

Thesis 3: Hydromorphology can be assessed

The linkage of reach scale physical parameters and biotic samples on sites only disturbed by physical alterations are scarce. There has been moderate success in linking biota with the currently used habitat surveys which is exemplified by the generally weak relationships found when investigation the relationship between morphological degradation and number of EPT families in 1541 stream sites from four countries (Figure 2). This is partly due to the mixed nature of pressures acting of a river reach and the habitat surveys using parameters of at least two spatial scales. It furthermore reflects that methods for sampling macroinvertebrates often are on a different scale to that of the hydromorphological assessment and that the sampling strategy was developed to target mainly organic pollution. At present, a few assessment systems that target impacts of low flow (Extence et al. 1999) and degraded hydromorphology (Barbour et al. 1996, Lorenz et al. 2004) do exist but they are not generally applicable and could be further developed. Therefore, the development of more indicator systems sensitive to hydromorphological degradation, using appropriate sampling techniques, will be a key issue in future.




Fig. 2: The number of EPT families along a gradient in hydromorphological quality in 4 countries, (a) Sweden, (b) Slovakia, (c) UK, (d) Denmark. Data from the EU REBECCA project (SSPI-CT-2003-502158).

Recommendations & perspectives

Hydromorphological degradation is one the most important pressures on European and imposes a serious risk to freshwater communities either when acting alone or in combination with other pressures. Impairment of hydromorphology in river ecosystems will significant reduce its resistance towards other pressures on the ecosystem. Hydromorphology are currently assessed using a range of techniques that are suboptimal as they lack appropriate sensitivity as well as the ability to quantify the importance of individual pressures. There is an urgent need to develop refined and updated assessment systems targeting hydromorphology.

References

- Allan, J.D. 2004. Landscapes and riverscapes: The influence of land use on stream ecosystems. *Annu. Rev. Ecol. Evol. Syst.* 35:257-284.
- Andersen J.M. 1994. Water quality management in the River Gudena, a Danish lake-stream-estuary system. *Hydrobiol.*, 275/276: 499-507.
- Barbour, M.T., Diamond, J. & Yoder, C. 1996. Biological assessment strategies : applications and limitations. In: Grothe, D., Dickson, K. Reed, D. (Eds.), Whole-effluent toxicity testing: An evaluation of methods and predictability of receiving systems responses. SETAC publications, Pensacola, Florida, pp. 245-270.
- Extence CA, Balbi DM & Chadd RP. 1999. River flow indexing using benthic macroinvertebrates: a framework for setting hydrobiological objectives. *Regul Rivers: Res Manag*, 15: 543-574.
- Fitzpatrick, F.A., Scudder, B.C., Lenz, B.N. & Sullivan, D.J. 2001. Effects of multi-scale environmental characteristics on agricultural stream biota in eastern Wisconsin. *J. Am. Water Resour. Assoc.* 37:1489-507.
- Frissell CA, Liss WJ, Warren CE, Hurley MD. 1986. A hierarchical framework for stream habitat classification: viewing streams in a watershed context. *Environ. Manag.* 12:199-214.
- Harding, J.S., Benfield, E.F., Bolstad, P.V., Helfman, G.S. & Jones, E.B.D. 1998. Stream biodiversity: the ghost of land use past. *Proc. Natl. Acad. Sci.* 95:14843-14847.
- Hart D.D. & Finelli C.M. 1999. Physical-biological coupling in streams: the pervasive effects of flow on benthic organisms. *Ann Rev Ecol System*, 30, 363-395.
- Lane, S.N. & Richards, K.S. 1997. Linking river channel form and process: time space and causality revisited. *Earth Surface Processes and Landforms*, 22: 249-260.
- Lorenz, A., Hering, D., Feld, C.K. & Rolauuffs, P. 2004: A new method for assessing the impact of hydromorphological degradation on the macroinvertebrate fauna of five German stream types. *Hydrobiologia*, 516: 107-127.
- Pedersen, M.L. & Friberg, N. 2007. Two lowland stream riffles – linkages between physical habitats and macroinvertebrates across multiple spatial scales. *Aquat Ecol*, 41: 475-490.
- Vannote RL, Minshall GW, Cummins KW, Sedell JR & Cushing CE 1980. The river continuum concept. *Can J Fish Aqua Sci* 37: 130-137.



Terrestrial-aquatic linkages and how recovery of riparian zones can improve ecological quality, assessment of pressures on stream ecosystems with specific emphasis on hydromorphology, interactions between organism groups and ecosystems function as indicator of impairment, global changes impacts on aquatic ecosystems and biodiversity