



**AALBORG UNIVERSITY**  
DENMARK

**Aalborg Universitet**

## **On directed coverings**

Fajstrup, Lisbeth

*Publication date:*  
2008

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Fajstrup, L. (2008). On directed coverings. Aalborg: Department of Mathematical Sciences, Aalborg University. (Research Report Series; No. R-2008-21).

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

AALBORG UNIVERSITY

**On directed coverings**

by

Lisbeth Fajstrup

R-2008-21

December 2008

DEPARTMENT OF MATHEMATICAL SCIENCES  
AALBORG UNIVERSITY

Fredrik Bajers Vej 7 G ■ DK-9220 Aalborg Øst ■ Denmark

Phone: +45 99 40 80 80 ■ Telefax: +45 98 15 81 29

URL: <http://www.math.aau.dk>



# ON DIRECTED COVERINGS

LISBETH FAJSTRUP

ABSTRACT. In [1], we study coverings in the setting of directed topology. Unfortunately, there is a condition missing in the definition of a directed covering. Some of the results in [1] require this extra condition and in fact it was claimed to follow from the original definition. It is the purpose of this note to give the right definition and point out how this affects the statements in that paper. Moreover, we give an example of a dicovering in the sense of [1], which does not satisfy the extra condition. Fortunately, with the extra condition, the subsequent results are now correct.

## 1. INTRODUCTION

In [1], we give a construction  $\pi : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$  which is denoted a universal dicovering. Moreover, we define dicoverings as maps  $p : (Y, y_0) \rightarrow (X, x_0)$  with certain lifting properties. We claim the existence of a map  $\phi : (\tilde{X}, \tilde{x}_0) \rightarrow (Y, y_0)$  s.t.  $\pi = p \circ \phi$ ; and it is in this sense, that the universal dicovering is universal. However, for the map  $\phi$  to be well defined, we need an extra lifting condition in the definition 2.4 of a dicovering. The purpose of this note is to make this clear.

## 2. THE PROBLEM AND THE SOLUTION

The directed spaces in [1] are locally partially ordered spaces. We will work in **d-Top**, the category of d-spaces, here. The approach through locally partially ordered spaces as in [1] requires extra conditions on the spaces, i.e., longer definitions, and hence would make this note longer; moreover, **d-Top** seems by now to be the right category for directed topology.

**Definition 2.1.** *A d-space is a topological space  $X$  with a set of paths  $\vec{P}(X) \subset X^I$  such that*

- $\vec{P}(X)$  contains all constant paths.
- $\gamma, \mu \in \vec{P}(X)$  implies  $\gamma \star \mu \in \vec{P}(X)$ , where  $\star$  is concatenation.
- If  $\phi : I \rightarrow I$  is monotone,  $t \leq s \Rightarrow \phi(t) \leq \phi(s)$ , and  $\gamma \in \vec{P}(X)$ , then  $\gamma \circ \phi \in \vec{P}(X)$

*A d-map or dimap  $f : X \rightarrow Y$  is a continuous map, such that if  $\alpha \in \vec{P}(X)$  then  $f \circ \alpha \in \vec{P}(Y)$ .*

---

*Date:* December 16, 2008.

The set of distinguished paths,  $\vec{P}(X)$  are called the dipaths. They are  $d$ -maps from the ordered interval  $\vec{I}$  to  $X$ . The category of  $d$ -spaces is denoted **d-Top**

**Definition 2.2.** For a  $d$ -space  $(X, \vec{P}(X))$  and  $A, B$  subsets of  $X$ , let  $\vec{P}(X, A, B)$  be the set of dipaths  $\gamma : \vec{I} \rightarrow X$ ,  $\gamma(0) \in A$ ,  $\gamma(1) \in B$ .

Let  $I \times \vec{I}$  be the unit square with discrete order (equality) in one coordinate and the standard order along the other. Dipaths are increasing in the second and constant in the first coordinate. Let  $x, y$  be points in  $X$  and  $\gamma, \mu \in \vec{P}(X, x, y)$ . Then  $\gamma$  is dihomotopic to  $\mu$  if there is a  $d$ -map  $H : I \times \vec{I} \rightarrow X$  with  $H(0, t) = \gamma(t)$ ,  $H(1, t) = \mu(t)$  and  $H(s_0, t) \in \vec{P}(X, x, y)$  for all  $s_0$ .

$H$  is a dihomotopy with fixed endpoints and the equivalence classes are called dihomotopy classes.

$\vec{\pi}_1(X, x, y)$  is the set of equivalence classes.

We define a universal dcovering as in [1] disregarding the extra conditions on the topology given in [1] on  $X$ , since these are not relevant for this note. See Rem. 2.8

**Definition 2.3.** [Following [1] Def. 3.1] Let  $((X, x_0), \vec{P}(X))$  be a pointed  $d$ -space and let  $\mathcal{U}$  be a basis for the topology on  $X$ . The universal dcovering space  $(\tilde{X}, \tilde{x}_0)$  of  $X$  with respect to  $x_0$  is the set

$$\{[\gamma] \mid \gamma : (\vec{I}, 0) \rightarrow (X, x_0)\}$$

where  $[\gamma]$  is the dihomotopy class of  $\gamma$  with fixed endpoints and  $\tilde{x}_0$  is the dihomotopy class of the constant dipath  $[x_0]$ . The topology on  $(\tilde{X}, \tilde{x}_0)$  is given by the following subbasis:

For  $\gamma : (\vec{I}, 0) \rightarrow (X, x_0)$  such that  $\gamma(1) \in U$ , where  $U \in \mathcal{U}$ , let

$$U_{[\gamma]} = \{[\eta] \in \cup_{y \in U} \vec{\pi}_1(X, x_0, y) \mid [\eta] \sim_U [\gamma]\}$$

where  $[\gamma] \sim_U [\eta]$  if there is a  $d$ map  $H : I \times \vec{I} \rightarrow X$  such that  $H(0, t) = \gamma(t)$ ,  $H(1, t) = \eta(t)$ ,  $H(s, 0) = x_0$  and  $H(s, 1) \in U$  for all  $s \in I$ .

The set of dipaths  $\vec{P}(\tilde{X})$  is the closure of  $\{\Gamma(t) = [\gamma|_{[0, t]}\}$  where  $\gamma : ((\vec{I}, 0) \rightarrow (X, x_0))$  under composition, reparametrization and subpath.

Let  $\pi : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$  be the endpoint map,  $\pi([\gamma]) = \gamma(1)$ .

A directed covering is defined in terms of lifting properties in [1]. We did not require condition 3, i.e., that dihomotopies with fixed endpoints and initial point  $x_0$  lift to dihomotopies with fixed endpoints. Condition 2 is a lifting property for dihomotopies with fixed initial point.

**Definition 2.4.** [Following [1] Def. 4.1] Let  $\Pi : \hat{X} \rightarrow X$  be a  $d$ -map. Then  $\Pi$  is a dcovering with respect to  $x_0 \in X$  if for every  $y_0 \in \Pi^{-1}(x_0)$ :

- (1) For every dipath  $\gamma : \vec{I} \rightarrow X$  such that  $\gamma(0) = x_0$ , there is a unique lift  $\hat{\gamma} : \vec{I} \rightarrow \hat{X}$ , such that  $\Pi \circ \hat{\gamma} = \gamma$  and  $\hat{\gamma}(0) = y_0$ .

- (2) For every  $d$ -map  $H : I \times \vec{I} \rightarrow X$  with  $H(s, 0) = x_0$  there is a unique lift  $\hat{H} : I \times \vec{I} \rightarrow \hat{X}$  s.t.  $\Pi \circ \hat{H} = H$  and  $\hat{H}(s, 0) = y_0$ .
- (3) For every  $d$ -map  $H : I \times \vec{I} \rightarrow X$  with  $H(s, 0) = x_0$  and  $H(s, 1) = x_1$  there is a unique lift  $\hat{H} : I \times \vec{I} \rightarrow \hat{X}$  s.t.  $\Pi \circ \hat{H} = H$ ,  $\hat{H}(s, 0) = y_0$  and  $\hat{H}(s, 1)$  is constant.

When  $X = \uparrow_X x_0 := \{x \in X \mid \exists \gamma : (\vec{I}, 0) \rightarrow (X, x_0) : \gamma(1) = x\}$ ,  $\Pi^{-1}(x_0) = \hat{x}_0$  and  $\hat{X} = \uparrow_{\hat{X}} \hat{x}_0$ , the dicovering is a simple dicovering, and all dipaths lift uniquely, not only the ones initiating in  $x_0$ .

Our universal dicovering is universal in the sense of

**Proposition 2.5.** [[1] 4.6] Let  $\Pi : (\hat{X}, \hat{x}_0) \rightarrow (X, x_0)$  be a dicovering w.r.t.  $x_0 \in X$ , such that  $\Pi^{-1}(x_0) = \hat{x}_0$ . Then there is a map  $\phi : (\tilde{X}, \tilde{x}_0) \rightarrow (\hat{X}, \hat{x}_0)$  covering the identity, i.e.,  $\Pi \circ \phi = \pi$ .

*Proof.* Let  $\phi([\gamma]) = \hat{\gamma}(1)$ , where  $\hat{\gamma}$  is the unique lift of  $\gamma$  with initial point  $\hat{x}_0$ . This is well defined, since if  $[\lambda] = [\gamma] \in \tilde{X}_{x_0}$ ,  $\lambda$  is dihomotopic to  $\gamma$ . Since dihomotopies with fixed endpoints lift to dihomotopies with fixed endpoints, it follows that  $\hat{\lambda}(1) = \hat{\gamma}(1)$ .  $\square$

The map  $\phi$  is not well defined if we remove condition 3 in Def. 2.4, and hence  $\tilde{X}$  is not “universal” in that setting. In [1] we claimed that condition 3 follows from condition 2 and continuity of dihomotopies. This claim is made, and condition 3 is needed, in the proof of [1] Prop. 4.5 as well:

**Proposition 2.6.** [[1] 4.5] Let  $\Pi : (\hat{X}, \hat{x}_0) \rightarrow (X, x_0)$  be a simple dicovering. Then for  $x \in X$ ,  $|\Pi^{-1}(x)| \leq |\tilde{\pi}_1(X, x_0, x)|$

However, as the following example shows, condition 3 does not follow from the original definition. In particular, Prop. 2.6 does not hold in this example.

**Example 2.7.** (See Fig.1) Let  $\hat{X}$  be the quotient of  $I \times \vec{I}$  under the relation  $(s, 0) \sim (0, 0)$  and let  $X$  be the quotient of  $\hat{X}$  under the relation  $(s, 1) \sim (0, 1)$ . Let  $p : (\hat{X}, (0, 0)) \rightarrow (X, (0, 0))$  be the quotient map. Then Def. 2.4 1 and 2 certainly hold with  $x_0 = (0, 0)$ , i.e., dipaths and of dihomotopies initiating in  $(0, 0)$  lift uniquely. But the dihomotopy (with fixed endpoints)  $H : I \times \vec{I} \rightarrow X$  given by projection to the quotient  $X$  lifts to projection to the quotient  $\hat{X}$ , which does not have fixed endpoint  $H(s, 1)$ . Hence this example is a dicover with regard to the original definition but not to the one given here.

All examples of dicoverings in [1] in fact do satisfy the extra lifting condition, hence it is still true that the map  $\phi$  in 2.5 is not necessarily continuous; the “Hawaiian star” example ([1] 4.7) proves that.

*Remark 2.8.* In [1], we study universal dicoverings with base space  $X$  a locally partially ordered space satisfying some technical requirements on the interplay

$$\begin{array}{ccc}
(0,0) \hookrightarrow & I \times \vec{I} /_{(s,0) \sim (0,0)} & \\
\downarrow & \nearrow & \downarrow p \\
I \times \vec{I} & \xrightarrow{H} & I \times \vec{I} /_{(s,0) \sim (0,0)}^{(s,1) \sim (0,1)}
\end{array}$$

FIGURE 1. All maps are projections or inclusions

between the topology and the directed paths. These cases can be studied in the above framework of d-spaces. Since the universal d-covering construction satisfies the present definition of a d-covering, the results, examples and counterexamples are still relevant: For instance, our construction 2.3 will not always provide a Hausdorff universal d-covering even if the base space is Hausdorff, and the fibers are not discrete (by [1] Ex. 3.7) but extra requirements on the topology of  $X$  will ensure that.

*Remark 2.9.* In [2] we prove that in certain subcategories of  $\mathbf{d-Top}$ , there exists a universal d-covering w.r.t. the original definition. The methods in [2] provide existence of a universal d-covering  $(\bar{X}, \bar{x}_0) \rightarrow (X, x_0)$  w.r.t. the definition here as well, and the resulting d-map  $f : \bar{X} \rightarrow \tilde{X}$  is a bijection of sets. We study this in a subsequent paper.

#### REFERENCES

1. L. Fajstrup, *Dicovering spaces*, Homology Homotopy Appl. **5** (2003), no. 2, 1–17.
2. J. Rosicky L. Fajstrup, *A convenient category for directed homotopy*, Theory Appl. Categ. **21** (2001), no. 1, 7–20.