



AALBORG UNIVERSITY
DENMARK

Aalborg Universitet

Fate of xenobiotic compounds and plants activity in reed bed sludge treatment

Chen, Xijuan; Pauli, Udo; Rehfus, Stefan; Bester, Kai

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Chen, X., Pauli, U., Rehfus, S., & Bester, K. (2009). ***Fate of xenobiotic compounds and plants activity in reed bed sludge treatment***. Poster presented at SETAC 2009, Göteborg, Sweden.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Fate of xenobiotic compounds and plants activity in reed bed sludge treatment

Xijuan Chen ^{1,3}, Udo Pauly ², Stefan Rehfus ², Kai Bester ^{1,3}

1) Section of Environmental Engineering, Aalborg University, Aalborg, Denmark

2) EKO-PLANT GmbH, Neu-Eichenberg, Germany

3) Institute of Environmental Analytical Chemistry, University Duisburg-Essen, Germany

Email: xc@bio.aau.dk, kb@bio.aau.dk

INTRODUCTION


Sludge reed beds have been used for dewatering (draining and evapotranspiration) and mineralisation of sludge in Europe since 1988. Although reed beds are considered as a low cost and low contamination method in reducing volume, breaking

down organic matter and increasing the density of sludge, it is not yet clear whether this enhanced biological treatment is suitable for degradation xenobiotic compounds. The first of two aims of this project is to assess the role of macrophytes in the sludge reed bed treatment technologies concerning different effects on removal of persistent organic compounds. The second aim of this project is to investigate the fate of xenobiotic compounds during the sludge reed bed treatment process.



METHODS

16 containers with a size of 1 m X 1 m X 1 m were built from stainless steel and filled with a 20 cm layer of gravel (16-32 mm) and 50 cm pre-treated sludge, from which plants and roots have been removed to prevent the cross-impact of reeds. The containers were





placed outdoors. Four containers were planted with reed canary grass (*Phalaris arundinacea*), four with bulrush (*Typha latifolia*), another four with reed (*Phragmites australis*) at a density of 12 plants m⁻² to study the plants effect on the degradation process of organic-pollutants, four containers were left unplanted in order to distinguish the impact of the root system on the performance of the containers for the degradation of the target compounds.

RESULTS AND DISCUSSIONS

During the twelve months experiment the concentration of xenobiotic compounds were decreased. There was no significant difference between samples from different containers with different macrophyte species. However, reed (*Phragmites australis*) grows better in sludge than the other species.

	OTNE	HHCB	AHTN	Triclosan	DEHP
reed canary grass	204	1155	866	433	408
bulrush	187	990	770	330	315
reed	198	1386	990	462	365
unplanted	187	1155	770	385	533



CONCLUSIONS

The sludge reed bed container study showed that the reed bed sludge treatment technology is able to reduce persistent organic pollutant significantly. The different macrophyte species did not have a significant effect on the dewatering process as well as degradation of the respective compounds in this experiment, this is possibly due to the small size of container (1 m³) in comparison with the large scale of reed bed.

REFERENCES

- [1] Bester K. Personal Care Compounds in the Environment. VCH-Wiley, Weinheim, Germany, 2007, 9-13 pp.
- [2] Chen X, Pauly U, Rehfus S, Bester K. Personal care compounds in advanced biological sludge treatment. Chemosphere 2009;accepted.