



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Encapsulation of lutein in amphiphilic -cyclodextrin nanoparticles for the application in drug delivery.

Løjkner, Lars Damgaard; Mazzaglia, Antonino; Larsen, Kim Lambertsen; Micali, Norberto; Stancanelli, Rosanna; Guardo, Marta; Cannavá, Carmela; Ficarra, Paola

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Løjkner, L. D., Mazzaglia, A., Larsen, K. L., Micali, N., Stancanelli, R., Guardo, M., Cannavá, C., & Ficarra, P. (2009). *Encapsulation of lutein in amphiphilic -cyclodextrin nanoparticles for the application in drug delivery..* Abstract from 2^o Congresso Nazionale Chimica E Tecnologia delle Ciclodestrine, Asti, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Incapsulation of lutein in amphiphilic β -cyclodextrin nanoparticles for the application in drug delivery.

Lars Damgaard Løjknær^a, Antonino Mazzaglia^b, Kim Lambertsen Larsen^a, Norberto Micali^c,
Rosanna Stancanelli^d, Marta Guardo^d, Carmela Cannavà^d, Paola Ficarra^d.

^a Department of chemistry, biotechnology and environmental engineering, Aalborg University, Sohngaardsholmsvej 49 & 57, 9000 Aalborg, Denmark.

^b ISMN-CNR c/o Dipartimento di Chimica Inorganica, Chimica Analitica e Chimica Fisica, Università di Messina, Salita Sperone 31, 98166 Messina, Italy.

^c IPCF-CNR, Salita Sperone, Contrada Papardo 98158, Messina, Italy.

^d Dipartimento Farmaco-Chimico, Università di Messina, Viale Annunziata 98168, Italy

Amphiphilic cyclodextrins (ACDs) have been studied and utilised in many different applications, as ACDs have the possibility to form micelles, liposomes and inclusion complexes with hydrophobic substances [1]. In the pharmaceutical area, the ACDs are used e.g. carrier systems for lipophilic drugs, to enhance bioavailability, solubility and stability [2].

In this work, primary face butyrate modified β -CDs (C47) [3] were used to encapsulate lutein in water soluble nanoparticles. Lutein is a hydrophobic molecule acting as a strong dye when solubilised in proper solvent and is commonly used in pharmaceutical, nutraceutical and food industries. The characteristic absorption peaks of lutein are measured to be at 430, 455 and 484 nm in dichloromethane. These characteristics make lutein an excellent test-molecule for evaluation of the encapsulation properties of the ACDs. The encapsulation of lutein was measured using UV-Vis spectroscopy and DLS for size determination of the nanoparticles.

The lutein:C47 encapsulation was examined at different ratios and the mixture was prepared using a nanoemulsion method [2]. In all series, the concentration of lutein in water was 10 μ M and the concentration of C47 varied from 10 μ M to 100 μ M. The size of the particles, measured with DLS, ranged from 150 nm up to 500 nm and even larger aggregates, which settled at the bottom of the vial. The UV-Vis spectra of the lutein:C47 complexes showed both scattering of all wavelength from the particles, but also the characteristic peaks of solubilised lutein. This indicates that the lutein was encapsulated in an ACD-hydrophobic environment by maintaining the spectroscopic properties of lutein dissolved in organic solvent. Furthermore the complexes appeared stable over several days with only a slight decrease in absorption. The lower rim of the ACDs is still available to graft molecules relevant for e.g. targeting, prompt to recognise receptor protein on the surface of the nanoparticle.

[1] B. Perly et al. *Fine, Specialty & Performance Chemicals*, (2005).

[2] F. Quaglia, A. Mazzaglia et al. *Biomaterials* (2009), 30, 374.

[3] D. Gallois-Montbrun et al. *J Incl Phenom Macrocycl Chem* (2007) 57:131–135