

STUDY OF IMPACT OF LIFESTYLE MODIFICATION ON DIABETES AND PRE-DIABETES IN AN URBAN POPULATION

***¹Yogesh Gode, ²Swapnil Patond, ³Vishwajit Wankhade, ⁴Sarang Ghodki, ⁵Deepali Jadhav, ⁶Manish Ramdas Dhawade, ⁷Yashwant Wankhade**

¹Associate Professor Dept. of General Surgery Dr. Rajendra Gode Medical College, Amravati.

²Professor Dept. of Forensic Medicine Dr. Rajendra Gode Medical College, Amravati

³Associate Professor Dept. of Forensic Medicine Dr. Rajendra Gode Medical College, Amravati

⁴Assistant Professor Dept. of Pharmacology Dr. Rajendra Gode Medical College, Amravati

⁵Lecturer Dept. of Pharmacology Dr. Rajendra Gode Institute of Pharmacy, Amravati

⁶Assistant Professor Dept. of Mechanical Dr. Rajendra Gode Institute of Technology and Research, Amravati

⁷Associate Professor Dept. of Preventive and Social Medicine Dr. Rajendra Gode Ayurvedic College and Hospital, Amravati

ABSTRACT: Diabetes mellitus has emerged as a prevalent non-communicable disease worldwide. It ranks among the top causes of mortality in high-income countries and has reached epidemic proportions in newly developing nations.¹ Continuous medical care and monitoring is required in Diabetes mellitus along with multi-factorial risk reduction strategies along associated with glycemic control. Due to population aging, growth, urbanization, a lack of physical exercise, and a high incidence of obesity, there are more persons with DM. Lifestyle factors include eating patterns, exercise routines, alcohol consumption, and smoking. An improvement in these parameters would lead to better adherence to hypoglycemic medications. An epidemiological study on diabetes and pre-diabetes in an urban area with reference to lifestyle modification. An interventional community-based study was conducted in an urban setup. Personal interviews using a semi-structured, pretested questionnaire were the main technique of data gathering. Detailed information has been taken on the demographic and socio-economic characteristics at both the individual and household levels. All responders provided their willingly given consent. Every second household was visited by systematic random sampling procedure to select 400 respondents. Finally, analysis was carried out on 346 respondents. All respondents were questioned using structured a pretested, questionnaire after taking informed consent. Significant difference was observed with positive family history of diabetes, obesity, physical activities in male, high calorie intake, stress, chewable tobacco and blood pressure level with diabetes and pre-diabetes. Compared to responders with normal level of blood glucose, pre diabetics and diabetics individual showed more positive family histories. Compared to responders

* Corresponding Author : nikhilramy@outlook.com

with normal blood glucose levels, pre-diabetics and diabetics were more likely to be overweight. With regard to personal habits of the respondents, smoking was associated with more than 1-fold increase risk of obtaining serum glucose level >110 mg/dl as compared to non-smokers. The current research was an attempt to examine the effect of intervention with reference to life style modification. From the study, it is evident that overall awareness about diabetes has been found low and no awareness has been found about pre-diabetes. The present study also demonstrated that education is the fundamental tool to make the population aware of their health issues. Awareness about pre-diabetes and diabetes, which can make them beneficial for community and nation and so, they can play an innovative role for prevention of diabetes.

KEYWORDS: Diabetes mellitus, Non-communicable Diseases, life style modification, pre-diabetes.

1. INTRODUCTION:

Diabetes mellitus is the most prevalent non-communicable diseases worldwide which includes majority of developed nations, rank which is fourth or fifth biggest cause of death, and also significant evidence that it is an epidemic in many developing nations. Diabetes mellitus needs attention as well as measures to reduce the risk including glycemic management. Prevention of acute condition and reducing the risk of complication along with awareness of amongst patient.”[1]

The etiopathogenesis, symptomatology, consequences, and therapy of diabetes are all well explained in the Charaka Samhita. It is clear that the illness was prevalent even though there are no records of its prevalence in ancient India. This "Maharoga" has three therapy options: exercise, nutrition, and medication.[2,3]

Since there were no established diagnostic standards for DM in the first half of the 20th century, it was challenging to determine its prevalence. The majority of earlier studies used glycosuria as the diagnostic standard and was based on hospital records.

In their publication that was published in 1991, created a fantastic combination of investigations.[4]

A population-based survey conducted in 1966 revealed a prevalence of 2.3% in Chandigarh. A survey conducted in Puducherry that year that used blood glucose for diagnosis revealed a prevalence of 0.7%.[5]

In 1971, Cuttack recorded a prevalence of 1.2%.[6] In 1972 the prevalence of DM in urban cities like Hyderabad was 2.5% [7] and Metro Cities like Delhi had a prevalence of 2.3% [8]

There are so many non-communicable diseases prevalent in India like obesity and diabetes. Hypertension is increasing at an alarming rate among adolescents. Tobacco use, cigarette smoking, and chewable tobacco mostly start during early life i.e. adolescence which impacts health and can have disastrous consequences in later life. To put it in its appropriate context, it should be noted that the figures for the diabetes mellitus prevalence of in India come from a few dispersed studies carried out across the nation. Several multi-center studies have been done, including the ICMR studies from 1979, 1991 [9], and 2004, the National Urban Diabetes Survey (NUDS), 2001 [10], and the Prevalence of Diabetes in India Study (PODIS), 2004. [11]

It is possible that the quick changes in lifestyle that have taken place over the past 50 years are to blame for the substantial increase in the incidence of type 2 diabetes mellitus and associated illnesses including obesity, high blood pressure, and metabolic

syndrome. The Transition of epidemiology consists prevention of various communicable disease proper nutrition, good hygiene, along with improvement in quality healthcare access, has increased longevity, it has also sped up the emergence of new-age diseases like obesity, diabetes mellitus, and heart disease.[12]

By making changes to food choices, adopting stress management techniques, reducing alcohol and cigarette use, and increasing physical activity levels, it is possible to significantly prevent the onset of diabetes mellitus. Encouragingly, there have been recent advancements in the development of behavioral strategies aimed at modifying these lifestyle habits. To effectively promote a healthy lifestyle, public-private partnerships involving the government, partner organizations, health service providers, the community, and individuals with diabetes mellitus are emerging as a highly effective approach. Such collaborative efforts can contribute to the prevention of diabetes and its associated complications. There is an urgent need for efficient methods to lower the prevalence of diabetes mellitus and support for addressing the underlying problems.

There is a lack of information on diabetes awareness in emerging nations like India. Such information is crucial for developing public health plans for any future diabetes management initiatives. Few studies really address diabetes awareness among those with the condition, and there are almost no population-based statistics, according to a literature review on diabetes knowledge and awareness in developing nations.

2.MATERIAL AND METHODS:

Study Design and Area: A community-based intervention study. It was carried out in an urban area. The universe of the study comprises adults (22-60 years) residing in the ward, an urban area.

Exclusion Criterion

- Pregnant women
- Known diabetic patients

Sample size: As per 2020 census data, the total population of the area was 12,211 from 1245 households. So, the average household size was almost 5people. Also, the census data shows that about 50 percent of the population was between the ages of 22 and 60 years. Thus, on average every household has approximately3 adults between the ages of 22 and 60 years. If more than one eligible adult was present in the household, only one was selected by random sampling (lottery method). In the present study, households were chosen through systematic random sampling.

The study was carried out in three phases:

Pre-intervention phase:

Every second household was visited by systematic random sampling procedure to select 777 respondents. Due to non-response, inaccuracy, and/or unreliability of information, 71 cases were not included in the analysis. Further 6 known diabetic cases were also not included in the analysis. Finally, an analysis has been executed on 700 respondents. The next day morning between 6 am-8 am, the blood test was carried out for the purpose of estimating fasting blood glucose (FBG) levels. Respondents, who had forgotten to remain fasting on that day, were again explained about the importance of blood tests, and the test was conducted on the next day. Results were informed to all respondents.

Intervention:

During the pre-intervention phase (September 2020 to February 2021) interpersonal approach was used to educate all respondents. To maximize the impact of lifestyle changes on fasting blood glucose levels, a leaflet was also created. In the booklet, all information about the disease was simplified and given in a structured format in the

Hindi language. Following interviews and measurements of fasting blood glucose, blood pressure, weight, and height, each respondent received individualized instruction on pre-diabetes and diabetes mellitus. Importance was given to explaining risk factors, diet modification, physical activity, etc.

Post Intervention phase

After one and a half years from the pre-intervention phase, all respondents were interviewed again along with all measurements i.e. fasting blood sugar level, weight, height, and blood pressure were taken again.

Study Duration:

This investigation was carried out over the course of two years (July 2020 to June 2022). The initial period (July 2020 to December 2021) of the study was devoted to an extensive literature search followed by the preparation of the research design and appropriate tools for the investigations. The field data collection was carried out for a period of one year (September 2020 to October 2021).

- Pre-intervention phase of data collection was carried out for a period of five months (September 2020-January 2021).
- After 1.5 years from the pre-intervention phase, a four-month post-intervention phase of data collecting was also conducted (March-2020-June 2020). The entry of data was completed in one month (August 2020- September 2020). After that, it took more than six months for analyzing the data (October 2020 to march 2021)

Strengths of the research

The findings of the study may be useful in increasing awareness of prediabetes and diabetes. It will also help in suggesting measures to reduce risk factors in the community for the further progression of the disease.

Limitations

Because the prevalence of diabetes mellitus was calculated using fasting glucose values rather than OGTT, it is understated. Self-reported- physical activity. Single location study.

Statistical Method: Collected information was checked thoroughly to avoid errors and maintain consistency. Necessary editing was carried out before entering the information into the computer program. Thereafter, all the individual data form was entered into SPSS for appropriate data analysis. In the SPSS trial version 16.0, the data were re-coded into different categorical variables to make it easier for the further analysis process. Data were entered in the coded form was created for its rapid computerization.

RESULT: -

Table 1: Likelihood of respondents with risk factors vs diabetes mellitus

Risk factors	Likelihood of the respondents with risk factors			
	Unadjusted		Adjusted	
	B (Standard Error.)	OR (95% Confidence Interval)	B (Standard Error)	OR + (95% Confidence Interval)
Body Mass Index				
Normal weight		REF		REF
Under-weight	-0.5 (0.2)	0.2 (0.1-0.5)	-0.50 (0.3)	0.3 (0.2-1.2)
Over-weight and obese	0.3 (0.2)	1.2 (0.9-2.9)	0.1 (0.3)	1.0 (0.6-2.2)
Diet (Caloric consumption in Kcal)				
Below 2000		REF		REF
2001-2500	0.3 (0.3)	1.4 (0.9-3.3)	0.2 (0.3)	1.2 (0.8-3.3)
2501-3000	0.5 (0.3)	1.9 (1.2-5.0)	0.7 (0.4)	2.1 (1.1-5.9)
Above 3000	1.2 (0.3)	3.9 (2.7-9.1)	1.1 (0.3)	4.7 (2.8-12.2)

Physical activity				
Mild		REF		REF
Moderate and vigorous activity	-0.01 (0.2)	0.6 (0.5-1.5)	-0.3 (0.2)	0.4 (0.3-1.0)
Stress				
No Stressful events in their life		REF		REF
Stressful events in their life	2.2 (0.5)	20.5 (8.2-74.1)	28.4 (0.6)	26.0 (8.1-119.2)
Family History of diabetes mellitus				
No family history				
Positive family history	1.0 (0.2)	2.0 (2.4-6.7)	1.0 (0.3)	1.7 (1.5-5.0)
Smoking habit				
Non-smoker		REF		REF
Smoker	0.1 (0.5)	1.1 (0.4-3.6)	0.05 (0.5)	1.0 (0.3-3.4)
Chewable Tobacco				
No		REF		REF
Yes	0.2 (0.1)	1.3 (1.1-2.4)	0.03 (0.2)	1.0 (0.6-1.7)
Alcohol consumption				
No		REF		REF
Yes	-0.1 (0.5)	0.5 (0.2-2.3)	-0.2 (0.6)	0.3 (0.1-2.1)

The outcomes of logistic regression analysis for estimating the probability of having a fasting blood glucose level >110 mg/dl are shown in Table 1. In the analysis, fasting blood glucose level was taken dependent variable, while BMI, physical activity, stress, personal habits, and family history as predictors of having a fasting blood glucose level >110 mg/dl. One of the significant contributing variables to the chance of having a fasting blood glucose level >110 mg/dl was shown to be BMI. No discernible difference between mild and moderate-vigorous physical activity was found in the current research to account for fasting blood sugar value >110 mg/dl. Respondents who were taking tobacco were almost two times more likely to have fasting blood sugar value >110 mg/dl and the respondents who were consuming alcohol were less likely to have the risk of fasting blood glucose level >110 mg/dl as compared to non-users.

Table 2: Changes in mean fasting blood glucose level of respondents from the pre to post-intervention phase.

Fasting blood glucose level		Pre-intervention Mean \pm SD	Post-intervention Mean \pm SD
The Normal blood glucose level	446	82.5 \pm 16.4	82.4 \pm 13.2
Pre-diabetics	50	99.2 \pm 3.2	101.1 \pm 12.3
Diabetics	30	155.4 \pm 50.2	135.5 \pm 42.3

The analysis shows in table2 the mean fasting blood sugar value was significantly decreased from the pre- to post-intervention phase, where the mean fasting blood glucose in pre-diabetics and in diabetics in the post-intervention phase was decreased respectively.

Table 3: Changes in mean blood pressure of respondents from the pre- to post-intervention phase

		Blood pressure level	
Fasting blood glucose level		Pre-intervention Mean \pm SD	Post-intervention Mean \pm SD
The normal blood glucose level	446	110.3 \pm 14.1	106.7 \pm 11.2
Pre-diabetics	50	122.4 \pm 19.1	109.7 \pm 9.2
Diabetics	30	126.2 \pm 20.3	110.0 \pm 15.3

Table 3 shows that mean blood pressure was significantly decreased from the pre- to post-intervention phase. In the pre-intervention phase, the mean blood pressure of respondents belonged to normal blood glucose levels. While in the post-intervention phase, it decreased respectively.

3.DISCUSSION

A metabolic condition called diabetes mellitus affects the sugar-metabolizing property of the body and causes blood sugar to increase. It could be fatal if uncontrolled. However, treatment options for the disease are limited. Worldwide, the number of persons with diabetes mellitus has grown. In research released in September 2012, the World Health Organization predicted that there may be more than 300 million individuals worldwide who have diabetes mellitus, with that expected to rise by another 60–70% by the year 2030 [13].

The spurt in diabetes prevalence may be due to lifestyle changes and food habits, although the role of genetic components may not be ruled out. Lack of proper exercise and excessive sedentary life are thought to be major players in the onset of diabetes. Proper monitoring of diabetes mellitus is a serious issue in India. Due to this, the vast majority of the Indian population is unaware of diabetes mellitus and its underlying problems. Therefore, the fight against diabetes mellitus will have to be multidirectional [14,15].

The total prevalence of diabetes mellitus was 6.0% in the study by [16], with known diabetes mellitus being 4.0% of the study population and newly diagnosed diabetes mellitus being 2.0% of respondents. Another study was done by [17], where the prevalence of diabetes mellitus overall was 6.1%, with known diabetes prevalence being 5.0% and newly diagnosed diabetes mellitus prevalence being 1.1%.

World Health Organization (WHO) report shows About 32 million individuals in 2000 had diabetes mellitus. Currently, 387 million individuals worldwide have diabetes mellitus, and by 2035, that number is expected to reach 592 million. [18]

According to [19], diabetes mellitus ranks among the top five leading causes of death globally. Furthermore, individuals with diabetes are at a higher risk of developing various complications, as highlighted. These complications include heart disease, stroke, high blood pressure, vision impairment leading to blindness, kidney disease, disorders of the nervous system, amputations, dental problems, and complications during pregnancy.) [20].

Tobacco use increases hemoglobin A1C levels, an indicator of chronically-elevated blood glucose. [21]. Cigarette smoking and oral tobacco mostly start during early life i.e. adolescence which impacts health and can have disastrous consequences in later life. Significantly weak relationship was observed between systolic blood pressure and respondents' fasting blood glucose level. This is similar to the study conducted by [22] where diabetes mellitus and systolic blood pressure were substantially linked.

Chronic pancreatitis frequently results in diabetes mellitus and is mostly brought on by binge drinking. Chronic pancreatitis patients are one in three to acquire diabetes mellitus (NHS). An assessment of 15 prior research examining the link between alcohol consumption and diabetes mellitus discovered that individuals classified as moderate drinkers had a 33% reduced risk of developing type 2 diabetes compared to those who either abstained from alcohol or consumed it heavily. This finding suggests that moderate alcohol consumption is associated with a lower likelihood of developing type 2 diabetes in comparison to either heavy drinking or complete abstinence.. Heavy drinkers and teetotalers both have a significant risk of acquiring diabetes mellitus [23].

A study was done by [24] where the mean systolic blood pressure in the pre-intervention phase was 130.0+13.4 and in the post-intervention phase, it reached 126.9+15.8. This may be due to the respondents taking medicine to control high blood pressure and taking less salt in their diet and adopting some home remedies also. A study conducted by [25] and [26] Consuming more than 2-4 drinks per week was found to be associated with a lower prevalence of type 2 diabetes mellitus over the course of 12 years compared to individuals who did not drink. These associations remained significant even after taking into account factors such as BMI and other variables that are known to contribute to the risk of developing diabetes.

4.CONCLUSION:

The aim of this study was to investigate the impact of implementing lifestyle modifications as an intervention and it is evident that overall awareness about diabetes has been found low and no awareness has been found about pre-diabetes. The present study also demonstrated that education is the fundamental tool to make the population aware of their health issues. Awareness about pre-diabetes and diabetes can make them beneficial for the community and nation so, they can take a creative approach to the prevention of diabetes.

5.References: -

1. Diagnosis and Classification of Diabetes Mellitus, American Diabetes Association (ADA), *Diabetes Care*, 2014;37(1): 81-S90.
2. Parivallal T. Diabetes in Ancinet India. In: Type 2 Diabetes in South Asians: Epidemiology, Risk Factors and prevention. Mohan V, Gundu HR Rao (Eds), Under the Aegis of SASAT. Jaypee Brothers Medical Publishers. 2007:97-103.

3. Shah VK. Diabetes mellitus in Indian medicine. Jaikrishnadas Ayurveda series. Chaukhamba Oriental, Varanasi. 2001;2:83.
4. Ramaiya KL, Kodali VR, Alberti KG. Epidemiology of diabetes in Asians of the Indian subcontinent. *Diabetes Metab Rev* 1990;6:125-46.
5. Data SP, Verma NPS, Gopalakrishnan R, Ghosh BN: Survey of diabetes mellitus in Pondicherry. *Diabetes in Tropics*. Patel JC and Talwarkar NG, Eds Diabetic Association of India, Bombay, 1966;33.
6. Tripathy Bibhuti B. and Samal Kshitish C., Overview and Consensus Statement on Diabetes in Tropical Areas, *Diabetes/Metabolism Research and Reviews*, 1977; 13:1-3.
7. Vigg BL, Hrishikesh P, Sahay BK. Survey of diabetes mellitus in a rural area. *Madhumeh* 1972;12:171
8. Ahuja MMS. Epidemiological studies on diabetes mellitus in India. In: Ahuja MMS, editor. *Epidemiology of diabetes in developing countries*. New Delhi: Interprint; p. 1979;29-38.
9. Ahuja MMS. Recent contributions to the epidemiology of diabetes mellitus in India. *Int J Diab Developing Countries* 1991;11:5-9.
10. Ramachandran A, Snehalatha C, Kapur A, Vijay V, Mohan V, Das AK, Rao PV, Yajnik CS, Prasanna Kumar KM, Nair JD; Diabetes Epidemiology Study Group in India (DESI), High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey. *Diabetologia*. 2001; 44(9):1094-101.
11. Sadikot SM, Nigam A, Das S, Bajaj S, Zargar AH, Prasannakumar KM, et al. The burden of diabetes and impaired glucose tolerance in India using the WHO criteria: prevalence of diabetes in India study (PODIS). *Diabetes Res Clin Pract* 2004; 66: 301-7.
12. Mohan V, Deepa M, Farooq S, Datta M, Deepa R. Prevalence, awareness & control of Hypertension in Chennai - The Chennai Urban Rural Epidemiology Study (CURES – 52), *JAPI* 2007;55.
13. Holliday EG. Hints of unique genetic effects for type 2 diabetes in India. *Diabetes*, 2013; 62: 1369- 1370.
14. Mohan V, Radhika G, Sathya RM, Tamil SR, Ganesan A, et al. Dietary carbohydrates, glycaemic load, food groups and newly detected type 2 diabetes among urban Asian Indian population in Chennai, India (Chennai Urban Rural Epidemiology Study 59). *Br J Nutr*, 2009; 102: 1498-1506.
15. Sukla P, Shrivastava SR, Shrivastava PS. A longitudinal study to assess the impact of exercise on clinical, biochemical, and anthropometric parameters among the type 2 diabetes patients of South India. *Avicenna J Med*, 2015;5: 16-20.
16. Javid Ahmad, Ahmad Masoodi Muneer, Mohd Ashraf, Rauf Rashid , Rafiq Ahmad, Ashfaq Ahmad and Sheikh Dawood, Prevalence of Diabetes Mellitus and its Associated Risk Factors in Age Group of 20 Years and Above in Kashmir, India, *Al Ame en J Med Sci (An US National Library of Medicine Enlisted Journal)*, 2011; 4 (1): 38 -44
17. Hurol Aksu, Kayihan Pala, Huseyin Aksu, Prevalence and associated risk factors of type 2 diabetes mellitus in Nilufer District, Bursa, Turkey, *Int J Diabetes & Metabolism*, 2006; 14: 98-102.
18. International Diabetes Federation (IDF) *Diabetes Atlas*, 2013.
19. Caliskan D, Ozdemir O, Ocaktan E and Idil A. Evaluation of awareness of diabetes mellitus and associated factors in four health center areas. *Patient Educ Couns*, , 2006;62: 142 – 147.

20. Mun SK, Wong KH, Lo SB, Li Y, Bayarsaikhan S. Artificial Intelligence for the Future Radiology Diagnostic Service. *Front Mol Biosci.* 2021;7:614258.
21. Clair C et al. "Relationships of Cotinine and Self-Reported Cigarette Smoking with Hemoglobin A1c in the U.S." *Diabetes Care* 2011; (34), 2250-2255.
22. Valliyot Balakrishnan, Sreedharan Jayadevan, Muttappallymyalil Jayakumary , Valliyot Sudha Balakrishnan. Risk factors of type 2 diabetes mellitus in the rural population of North Kerala, India: A Case Control Study, *Diabetologia Croatica* 2013; 42-1.
23. Reyes M, Meier R, Pereira S, Silva CA, Dahlweid FM, von Tengg-Kobligk H, Summers RM, Wiest R. On the Interpretability of Artificial Intelligence in Radiology: Challenges and Opportunities. *Radiol Artif Intell.* 2020;2(3):e190043.
24. Obaidur Rahman and Rafiqul Islam Md., Association between Fasting of Ramadan and Risk Factors of Diabetes:A Study from Rajshahi City in Bangladesh, Accepted: Published: October 25, 2011, *Advance Journal of Food Science and Technology* 2011;3(5): 360-365.
25. Ak MF. A Comparative Analysis of Breast Cancer Detection and Diagnosis Using Data Visualization and Machine Learning Applications. *Healthcare (Basel).* 2020 Apr 26;8(2):111. doi: 10.3390/healthcare8020111.
26. Ajani UA, Hennekens CH, Spelsberg A, Manson JE. Alcohol consumption and risk of type 2 diabetes mellitus among US male physicians. *Archives of Internal Medicine,* 2000;160: 1025–30.