



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Unsatisfactory response to acute medications does not affect the medication overuse headache development in pediatric chronic migraine

Frattale, Ilaria; Ferilli, Michela Ada Noris; Ursitti, Fabiana; Sforza, Giorgia; Monte, Gabriele; Proietti Checchi, Martina; Tarantino, Samuela; Mazzone, Luigi; Valeriani, Massimiliano; Papetti, Laura

Published in:

The Journal of Headache and Pain

DOI (link to publication from Publisher):

[10.1186/s10194-024-01766-7](https://doi.org/10.1186/s10194-024-01766-7)

Creative Commons License

CC BY 4.0

Publication date:

2024

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Frattale, I., Ferilli, M. A. N., Ursitti, F., Sforza, G., Monte, G., Proietti Checchi, M., Tarantino, S., Mazzone, L., Valeriani, M., & Papetti, L. (2024). Unsatisfactory response to acute medications does not affect the medication overuse headache development in pediatric chronic migraine. *The Journal of Headache and Pain*, 25(1), Article 61. <https://doi.org/10.1186/s10194-024-01766-7>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

RESEARCH

Open Access



Unsatisfactory response to acute medications does not affect the medication overuse headache development in pediatric chronic migraine

Ilaria Frattale¹, Michela Ada Noris Ferilli², Fabiana Ursitti², Giorgia Sforza², Gabriele Monte², Martina Proietti Checchi², Samuela Tarantino², Luigi Mazzone¹, Massimiliano Valeriani^{2,3,4*} and Laura Papetti²

Abstract

Background Chronic migraine (CM) negatively impacts the quality of life of 2 to 4% of pediatric patients. In adults, CM is frequently linked to medication overuse headache (MOH), but there is a much lower prevalence of MOH in children. A suboptimal response to acute therapies may lead to their reduced use, thus preventing MOH development in children and adolescents. The frequency of patients with CM who do not respond to acute therapies was examined in the present study. We investigated whether the prevalence of MOH was different between responders and non-responders. We also examined whether patients receiving prophylactic therapy had an improved response to acute therapy. Finally, we investigated if there was a difference in the frequency of psychiatric comorbidities between responders and non-responders.

Methods We retrospectively analysed clinical data of all chronic pediatric migraineurs under the age of 18 referred to the Headache Centre at Bambino Gesù Children Hospital in June 2021 and February 2023. ICHD3 criteria were used to diagnose CM and MOH. We collected demographic data, including the age at onset of migraine and the age of the CM course. At baseline and after 3 months of preventive treatment, we evaluated the response to acute medications. Neuropsychiatric comorbidities were referred by the children's parents during the first attendance evaluation.

Results Seventy patients with CM were assessed during the chosen period. Paracetamol was tried by 41 patients (58.5%), NSAIDs by 56 patients (80.0%), and triptans by 1 patient (1.4%). Fifty-one participants (73%) were non-responder to the abortive treatment. The presence of MOH was detected in 27.1% of the whole populations. Regarding our primary aim, MOH was diagnosed in 29% of non-responder patients and 22% of responders ($p > 0.05$). All patients received preventative treatment. After 3 months of preventive pharmacological therapy, 65.4% of patients who did not respond to acute medications achieved a response, while 34.6% of patients who were non-responder remain non-responder ($p < 0.05$). Prophylactic therapy was also effective in 69% of patients who responded to acute

*Correspondence:
Massimiliano Valeriani
massimiliano.valeriani@opbg.net

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

medication ($p < 0.05$). Psychiatric comorbidities were detected in 68.6% of patients, with no difference between responders and non-responders (72.2% vs. 67.3%; $p = 0.05$).

Conclusions Despite the high prevalence of unresponsiveness to acute therapies in pediatric CM, it does not act as a protective factor for MOH. Moreover, responsiveness to acute drugs is improved by pharmacological preventive treatment and it is not affected by concomitant psychiatric comorbidities.

Keywords Childhood, Adolescents, Chronic migraine, Medication overuse headache, Acute treatment

Introduction

Migraine is the primary cause of pain in childhood and its frequency increases with age, overtaking 20% in adolescence [1]. Throughout life, the frequency, severity, and disability of this condition can fluctuate. The transition between episodic and chronic forms is possible [2–5], and the origin of this modification is not fully understood [6].

Around 2 to 4% of children with migraines experience chronic migraine (CM) [7–9]. According to the ICHD3 criteria, CM is defined by the presence of at least 15 headache days for a minimum of 3 months with at least 8 migraine days each month [10]. Chronic headaches can negatively impact one's quality of life and the ability to complete routine school and sports activities, as well as maintain social relationships [11].

Central sensitization changes can be caused by the increased use of painkillers, which can decrease the effectiveness of acute therapies [12–15]. This phenomenon may be responsible of frequent acute medication intakes, MOH development, and migraine chronification [6, 16–18].

The definition of response to acute treatment, which encompasses NSAIDs, triptans, and combinations, is pain relief within 2 h and a state of well-being that lasts for at least 24 h [19]. Oral formulation is the most used by migraineurs with a 2-hour pain free rate from 19 to 40% [20, 21], while the subcutaneous administration (e.g. triptans) leads to pain free between 60 and 65% of cases [22]. Factors like late intake, inadequate dosage, and formulations with inadequate absorption can lead to a failed response to treatment [23]. During the chronic course of migraine, acute therapy response is observed to be lower. The concomitant conditions of obesity and neuropsychiatric comorbidities can cause resistance to acute medications [24].

According to pediatric studies, MOH is less prevalent in children and adolescents than in adult patients [25–29]. Although the reason is not known, factors related to brain development and the progression of migraine could explain the different prevalence of MOH in pediatric migraines and adulthood [28]. It can be inferred that pediatric patients with CM may not always respond to pharmacological treatments for acute therapy, which could prevent them from taking analgesic drugs

frequently. If this were true, we would expect a lower prevalence of MOH in patients who do not respond than in those who respond to acute therapy.

The primary aims of our study were: (1) to calculate the frequency of pediatric patients with CM who exhibited resistance to the acute therapies, and (2) to investigate whether MOH prevalence was different between responders and non-responders to the analgesic drugs. As secondary aims, we examined: (1) the impact of prophylactic treatment on the response to acute treatment and (2) the different frequency of psychiatric comorbidities between responders and non-responders to acute therapy.

Methods

Patients

We retrospectively analyzed clinical data of all patients affected by CM who attended the Headache Centre at Bambino Gesù Children's Hospital between June 2021 and February 2023. Data was obtained from patients between 6 and 17 during the initial visit and a second follow-up visit after three months. We used ICHD3 criteria [10] to diagnose CM and MOH.

We analyzed demographic characteristics, including sex, age at onset of migraine, age at CM onset, frequency of the attacks, MOH diagnosis, and response to acute medications at baseline and after 3 months of preventive treatment. Neuropsychiatric comorbidities were referred by the children's parents during the first attendance evaluation.

Paracetamol, nonsteroidal anti-inflammatory drugs (NSAIDs), and triptans were considered as acute medication in their own category. A patient was defined as not responding to acute therapy when he did not respond to paracetamol, an NSAID, and a triptan. Also, for the definition of non-responder, we verified the correct use of acute therapy in terms of adequate dosage and adherence. We assumed that due to local legislative reasons and the age of patients, the prescription of triptans was not common. For this reason, we also considered those who had no response to paracetamol and two NSAIDs as not responders.

We only included subjects with CM for whom we have data on the use of acute therapies (dosage, class of

drug) at baseline and after three months of prophylactic treatment.

Statistical analysis

Statistical analysis was conducted using SPSS version 22.0 and consisted of three steps.

The initial step involved a descriptive analysis that reported the differences between the responders and non-responders to analgesic drugs. In particular, for the categorical variables (sex, response or not to prophylactic therapy, and the presence or absence of psychiatric comorbidities) we considered the χ^2 test, while for the ordinary variables (age) we used the analysis of variance (ANOVA).

The second step was a bivariate analysis to study how each variable, considered individually, correlated with either response or resistance to acute therapy. Different statistical tests were used depending on the nature of the covariates and the response. For ordinal categorical variables, we used the Mann-Whitney U-test, the Kruskal-Wallis test and the Spearman's Rho. For the numeric variables, we transformed the response using the log scale ratio and considered the t-test, ANOVA, and Spearman's Rho. The individual tests in the bivariate analysis did not consider the contemporary effect of other covariates.

The third step was a multivariate analysis where all the variables that in the bivariate analysis showed a p -value ≤ 0.2 in determining response or resistance to acute therapy were included. A generalized linear model (GLM) with cumulative link and proportional odds

assumption was utilized for multivariate analysis. A p -value being ≤ 0.05 was deemed significant.

The parents of the participants provided written informed consent. The study was approved by the Ethics Committee of Bambino Gesù Children Hospital.

Results

In the considered period of time considered, 1680 children and adolescents were visited at our Headache Center. According to ICHD 3 criteria, 81 patients (4.8%) were diagnosed with CM. Eleven patients were excluded from further analysis since we did not have enough data on the use of acute therapy (untried or assumed at inadequate dosages). The analysed population included 70 patients (58 females and 12 males) with an average age at the time of the first visit of 14.2 ± 2.4 years. Demographic features of the sample are summarized in Table 1.

The average age of migraine onset was 11.2 ± 3.2 years, while the average age of CM onset was 13.6 ± 2.3 years. Migraine lasted an average of 2.4 years from its onset before becoming chronic. Before having CM, 87% of patients had a high-frequency episodic migraine (EM) (more than 5 attacks per month). In 28.5% of cases, a prophylactic treatment had been tried and resulted ineffective.

Before arriving at our center, 41 out of 70 patients had tried paracetamol (58.6%), 56 NSAIDs (80%), and only 1 triptans (1.4%).

Fifty-one participants (73%) were non-responder to the acute treatment, while 19 (27%) were responsive. Response to acute treatment was independent of the drug (Fig. 1).

Response or resistance to the acute therapy did not depend on the age at migraine onset (mean age: 11.2 vs. 11.4 years; $p > 0.05$) and at CM onset (mean age: 13.8 vs. 13.5 years; $p > 0.05$), and the duration of the disease (32.3 vs. 27.5 months; $p > 0.05$).

Among the total subjects ($n=70$), in 19 (27.1%) of cases, there was a concomitant presence of MOH. In these patients, an average of 23.5 doses of acute medications per month were assumed. MOH prevalence was similar among non-responders and responders (29% vs. 22%; $p > 0.05$).

All patients were given prophylactic pharmacological therapy, and the response was confirmed at a follow-up visit three months after the start. Twelve patients did not assume the prescribed therapy or suspended it too early (within a month). A decrease in headache days per month $\geq 50\%$ was observed in 58% (34/58) of patients who received prophylaxis. After prophylaxis, the prevalence of patients responding to the analgesic drugs increased to 71.4% (41/58), while non-responder patients decreased to 28.6% (17/58). More in detail, 65.4% of subjects who failed to respond to acute therapy before initiating

Table 1 Demographic Characteristics of $n=70$ patients in the analysed period

Patients with follow up:	70
Female: male	58:12 (females 83% - males 17%)
Mean age at the visit	14.2 ± 2.4
Mean age at migraine onset	11.2 ± 3.2
Mean age at chronification onset	13.6 ± 2.3
High- frequency migraine before chronification (%)	$n=61$ (87%)
Preventative treatment before chronification (%)	$n=20$ (28.5%)
Acute Medication tried before (%)	Paracetamol $n=41$ (58.6%) NSAIDs $n=56$ (80%) Triptan $n=1$ (1.4%)
Responder vs. Non Responder Baseline (%)	$n=19/70$ vs. $n=51/70$ (27% vs. 73%)
Responder vs. Non Responder After preventive (%)	$n=41/58$ vs. $n=17/58$ (71.4% vs. 28.6%)
MOH	$n=19/70$ (27.1%)
MOH in Responder vs. Non Responder %	22% vs. 27%
Neuropsychiatric comorbidities	$n=48/70$ (69%)
Neuropsychiatric comorbidities in Responder vs. Non Responder %	72.2% vs. 67.3%




Fig. 1 Acute medications response rate at baseline and after 3-months-preventative treatment

prophylaxis achieved a response after 3 months of preventive pharmacological therapy. The remaining non-responder patients (34.6%) kept not responding to acute therapy ($p < 0.05$). Among the patients who responded to the acute therapy, 69% simultaneously responded to the prophylactic therapy, while 31% did not experience a significant change in headache days per month ($p < 0.01$). The bivariate analysis identified only prophylactic therapy as a factor associated with response to acute therapy (R 0.9; C.I. 0.7–0.95; $p < 0.05$).

Psychiatric comorbidities were referred in 48 of patients (69%). Anxiety and depressive disorders were the most prevalent psychiatric conditions. The frequency of psychiatric comorbidities did not differ significantly between responders and non-responders to analgesic drugs (72.2% vs. 67.3%; $p = 0.05$).

Discussion

In both pediatric and adult migraine patients, resistance to acute therapy is a significant issue. In this context, our study in a pediatric cohort of CM patients produced the following significant findings:

1. Most CM patients found acute medications to be ineffective.
2. Both responders and non-responders had the same MOH prevalence, indicating that MOH development is not affected by failure to abort medication.
3. After receiving preventive treatment for three months, most patients, particularly those with good outcomes after prophylaxis, returned to respond to acute medication.
4. The response to acute medication was not influenced by psychiatric comorbidities.

Poor response to acute therapy in children and adolescents with CM

Acute medications were found to be ineffective for the majority of CM patients. The results confirm our hypothesis that CM patients have a decreased response to acute therapy. Whether poor response to acute therapy is a risk factor for the development of CM or merely a consequence of migraine severity still remains a debated question [6]. In patients with episodic migraine, suboptimal response to acute treatment was associated with a risk of developing CM within one year, independently of other migraine features, such as disability and frequency of the attacks [23]. Acute treatment failure can result in more frequent and longer attacks, as well as greater disability, which can lead to the onset of CM [6, 12–15]. In this view, more effective acute treatments should decrease the chances of developing CM. Multiple treatments are often necessary to overcome resistance to acute drugs in CM patients [23]. Our findings, showing that non-responders are resistant to at least two categories of drugs, including paracetamol, NSAIDs, and triptans, agree with previous results. The main reason for the resistance to analgesic drugs in CM is the severity of the disease, as evidenced by the weak response to multiple acute therapies [6]. Migraine progression leads to central sensitization, which in turn could reduce the response to acute therapies [6]. Poor response to acute therapy and the overuse of acute medications both contribute to migraine progression and the onset of MOH [16–18, 23, 30, 31].

Might poor response to acute therapy reduce the risk of MOH?

A primary aim of this study was to investigate whether the poor response to analgesic drugs could contribute

to the lower prevalence of MOH in children and adolescents, as compared to adults [25]. Indeed, patients with suboptimal response to acute therapy may be discouraged from using it frequently. Compared to the adult population with chronic headache in which MOH has a prevalence of 64% [32], in pediatric age MOH is far rarer, ranging from 20 to 50% of chronic patients [27–29]. Since the possibility of MOH development raises with increasing age, the parental control over drug intake could partly explain the difference between children and adulthood [33]. MOH pathophysiology is very complex and not fully understood. Beyond the amount of analgesic drug intake, other factors, such as medication effects [34], genetics factors [35] and headache-specific pain pathways [36], are involved.

In our sample, MOH was diagnosed in 25% of patients with a similar prevalence in both responders and non-responders. This result did not confirm our initial hypothesis of a key role played by the poor response of analgesic drugs in explaining the rather low prevalence of MOH in children compared to adulthood. Therefore, we can conclude that in pediatric age the unresponsiveness to the acute therapy does not have a protective role for MOH development.

One may wonder why non-responders keep assuming drugs which, however, do not help them in reducing pain. Though not completely known, this behavior suggests the possibility of an addictive psychopathological profiles of either the patient or her/his parents who administer the drug [37, 38].

Are prophylactic therapies useful in pediatric migraine?

Our study revealed that more than 70% of patients who received prophylactic therapy experienced a significant improvement in their response to acute medication. The number of headache days per month decreased by more than 50% for 69% of them, compared to the three months before treatment.

In children and adolescents, the use of pharmacological prophylactic treatment is not as widely accepted as for adults. In 2014, topiramate was approved by the Food and Drug Administration for the prophylactic treatment of migraine in children over 12 years based on robust clinical results [39, 40]. Unfortunately, there are only limited data for other drugs which are not licenced for pediatric age [33, 41–45]. Monoclonal antibodies and gepants are still subject to clinical trials and cannot be prescribed. Second, the usefulness of the prophylactic pharmacological treatments in children and adolescents has been challenged by the CHAMP study, which failed in showing a superiority of either topiramate or amitriptyline over placebo [41]. Powers et al. recommended that psychological treatments should be the preferred treatment over

medication because of the possible side effects and high placebo efficacy rate.

Although our study was not designed to investigate the efficacy of pharmacological prophylactic treatments and we did not consider a control group with placebo, more than half of our patients undergone prophylaxis showed a significant reduction in headache days per month. However, what is most noteworthy within the present results is that more than two third of our patients improved their response to analgesic drugs after prophylaxis. Though needing confirmation in appropriately designed trial, our present findings suggest that the response to acute therapy should be considered in future clinical studies on migraine prophylaxis in children and adolescents [41].

Our study was not designed to investigate the efficacy of pharmacological prophylactic treatments in pediatric CM. Despite this and the absence of a control group with placebo, we found that more than half of our patients undergone prophylaxis showed a significant reduction in headache days per month. Our findings indicate that over two-thirds of our patients have improved their ability to respond to acute therapies after prophylaxis. Though needing confirmation in an appropriately designed trial, we suggest that the response to acute therapy should be considered in future clinical studies on migraine prophylaxis in children and adolescents.

The effect of psychiatric comorbidities on the response to acute therapy

Anxiety and depression are highly prevalent in children and adolescents with migraines [46, 47]. The presence of these disorders can predict a poor response to acute and preventive therapy and a greater disability [48, 49]. On the other hand, headache chronification may lead to reduced quality of life and disability, which can be represented as factors for the onset of mood disorders [50]. From this perspective, it is possible that neuropsychiatric disturbances could play a role in determining the response to acute therapy.

In our study, almost 70% of patients had psychiatric comorbidities, which caused the same poor response to analgesic drugs as those without psychiatric symptoms. In conclusion, our data suggest that in pediatric patients with CM, the response to analgesic drug is not affected by neuropsychiatric comorbidities.

Limitations

The small number of patients and retrospective nature of our study result in some limitations. The retrospective design of the study increases the risk of recall bias and incomplete data. Since most patients came to our attention already showing a CM, we do not have detailed information about the previous migraine time course. Furthermore, we lack knowledge about the response

to acute therapy for most patients prior to migraine chronification. It's impossible to determine with certainty whether the poor response to acute medications is caused by chronification mechanisms or if it's not dependent on them. Finally, in our patients, neuropsychiatric comorbidities were diagnosed based on what was referred by the patients and their parents. Validated tools for diagnosing psychiatric disturbances are necessary to determine their impact on the response to acute medications.

Further longitudinal studies would provide more reliable evidence on the relationship between acute therapy response and MOH development.

Conclusion

In conclusion, our results suggest that the poor response to analgesic drugs observed in our patients with CM does not explain the rather low prevalence of MOH. Furthermore, we discovered that using a preventative treatment may enhance the therapeutic response to acute medication, thus reducing the disability related to the migraine attack. Lastly, although psychiatric comorbidities should be considered in the whole assessment of CM patients, they do not affect the response to acute therapy.

Author contributions

Conceptualization, L.P. and M.V.; Methodology, M.A.N.F.; Validation, L.P. and M.V.; Writing—Original Draft Preparation, I.F.; Writing—Review & Editing, L.P. and M.V.; Figure and table: F.U.; Data setting: G.M.; M.P.C. and S.T.; Visualization L.P.; M.V., L.M.; M.A.N.F.; G.M.; G.S.; F.U.; I.F.; S.T.; M.P.C.; Supervision, L.P.

Funding

This work was supported by the Italian Ministry of Health with Current Research funds and FINALIZED RESEARCH Grant 2019 (code: GR-2019-12369766).

Data availability

The data that support the findings of this study are not openly available due to reasons of sensitivity and are available from the corresponding author upon reasonable request. Data are located in controlled access data storage at Bambino Gesù Children's Hospital, Rome.

Declarations

Clinical trial number

not available.

Competing interests

The authors declare no competing interests.

Author details

¹Child Neurology and Psychiatry Unit, Department of Wellbeing of Mental and Neurological, Dental and Sensory Organ Health, Policlinico Tor Vergata Foundation Hospital, Rome, Italy

²Developmental Neurology, Bambino Gesù Children's Hospital, IRCCS, Piazza di Sant'Onofrio 4, 00165 Rome, Italy

³System Medicine Department, Tor Vergata University of Rome, Rome, Italy

⁴Center for Sensory-Motor Interaction, Aalborg University, Aalborg, Denmark

Published online: 23 April 2024

References

1. Abu-Arafeh I, Razak S, Sivaraman B, Graham C (2010) Prevalence of headache and migraine in children and adolescents: a systematic review of population-based studies. *Develop Med Child Neuro* Dicembre 52(12):1088–1097
2. Serrano D, Lipton RB, Scher AI, Reed ML, Stewart WF, Adams AM et al (2017) Fluctuations in episodic and chronic migraine status over the course of 1 year: implications for diagnosis, treatment and clinical trial design. *J Headache Pain* Dicembre 18(1):101
3. Bigal ME, Lipton RB (2008) Clinical course in migraine: conceptualizing migraine transformation. *Neuro* 9 Settembre 71(11):848–855
4. Bigal ME, Lipton RB (2008) Concepts and mechanisms of Migraine Chronification. *Headache* Gennaio 48(1):7–15
5. Bigal ME, Lipton RB (2006) When migraine progresses: transformed or chronic migraine. *Expert Rev Neurother* Marzo 6(3):297–306
6. Lipton RB, Buse DC, Nahas SJ, Tietjen GE, Martin VT, Löff E et al (2023) Risk factors for migraine disease progression: a narrative review for a patient-centered approach. *J Neurol* Dicembre 270(12):5692–5710
7. Zwart JA, Dyb G, Holmen T, Stovner L, Sand T (2004) The prevalence of migraine and tension-type headaches among adolescents in Norway. The Nord-Trøndelag Health Study (Head-Hunt-Youth), a large Population-based epidemiological study. *Cephalalgia* Maggio 24(5):373–379
8. Kernick D, Campbell J (2009) Measuring the impact of Headache in children: a critical review of the literature. *Cephalalgia* Gennaio 29(1):3–16
9. Arruda MA, Guidetti V, Galli F, Albuquerque RCP, Bigal ME (2010) Frequent headaches in the preadolescent pediatric population: a population-based study. *Neuro* 16 Marzo 74(11):903–908
10. Headache Classification Committee of the International Headache Society (IHS) (2018);38(1):1–211
11. Lipton RB, Manack A, Ricci JA, Chee E, Turkel CC, Winner P (2011) Prevalence and burden of chronic migraine in adolescents: results of the Chronic Daily Headache in adolescents Study (C-dAS). *Headache* Maggio 51(5):693–706
12. De Felice M, Ossipov MH, Porreca F (2011) Persistent medication-induced neural adaptations, descending facilitation, and medication overuse headache. *Curr Opin Neurol* Giugno 24(3):193–196
13. De Felice M, Ossipov MH, Wang R, Lai J, Chichorro J, Meng I et al (2010) Triptan-induced latent sensitization: a possible basis for medication overuse headache. *Annals Neurol* Marzo 67(3):325–337
14. Felice MD, Porreca F (2009) Opiate-Induced Persistent Nociceptive Trigeminal neural adaptations: potential relevance to Opiate-Induced Medication Overuse Headache. *Cephalalgia* Dicembre 29(12):1277–1284
15. Bigal ME, Lipton RB (2008) Excessive acute migraine medication use and migraine progression. *Neuro* 25 Novembre 71(22):1821–1828
16. Katsarava Z, Schneeweiss S, Kurth T, Kroener U, Fritsche G, Eikermann A et al (2004) Incidence and predictors for chronicity of headache in patients with episodic migraine. *Neuro* 9 Marzo 62(5):788–790
17. Louter MA, Bosker JE, Van Oosterhout WJP, Van Zwet EW, Zitman FG, Ferrari MD et al (2013) Cutaneous allodynia as a predictor of migraine chronification. *Brain* Novembre 136(11):3489–3496
18. Wang SJ, Fuh JL, Lu SR, Juang KD (2007) Outcomes and predictors of chronic daily headache in adolescents: a 2-year longitudinal study. *Neuro* 20 Febbraio 68(8):591–596
19. Sacco S, Lampl C, Amin FM, Braschinsky M, Deligianni C, Uludüz D et al (2022) European Headache Federation (EHF) consensus on the definition of effective treatment of a migraine attack and of triptan failure. *J Headache Pain* Dicembre 23(1):133
20. Ferrari M, Goadsby P, Roon K, Lipton R (2002) Triptans (serotonin, 5-HT_{1B/1D} agonists) in migraine: detailed results and methods of a Meta-analysis of 53 trials. *Cephalalgia* Ottobre 22(8):633–658
21. Ferrari MD, Roon KI, Lipton RB, Goadsby PJ (2001) Oral triptans (serotonin 5-HT_{1B/1D} agonists) in acute migraine treatment: a meta-analysis of 53 trials. *Lancet* Novembre 358(9294):1668–1675
22. Cady RK, Wendt JK, Kirchner JR (1991) Treatment of acute migraine with subcutaneous sumatriptan. *JAMA*; (265):2831–2835
23. Lipton RB, Munjal S, Buse DC, Fanning KM, Bennett A, Reed ML (2016) Predicting inadequate response to Acute Migraine Medication: results from the American migraine prevalence and Prevention (AMPP) Study. *Headache* Novembre 56(10):1635–1648

24. Saracco MG, Allais G, Tullo V, Zava D, Pezzola D, Reggiardo G et al (2014) Efficacy of frovatriptan and other triptans in the treatment of acute migraine of normal weight and obese subjects: a review of randomized studies. *Neurol Sci Magg* 35(5):115–119
25. Papetti L, Salfa I, Battan B, Moavero R, Termine C, Bartoli B et al (2019) Features of primary chronic headache in children and adolescents and Validity of ICHD 3 Criteria. *Front Neurol* 15 Febbraio 10:92
26. Moavero R, Stornelli M, Papetti L, Ursitti F, Ferilli MAN, Balestri M et al (2020) Medication overuse withdrawal in children and adolescents does not always improve headache: a cross-sectional study. *Front Neurol* 19 Agosto 11:823
27. Stevens J, Hayes J, Pakalnis A (2014) A randomized trial of telephone-based motivational interviewing for adolescent chronic headache with medication overuse. *Cephalalgia Magg* 34(6):446–454
28. Langdon R, DiSabella MT (2017) Pediatric Headache: an overview. *Curr Probl Pediatr Adolesc Health Care* Marzo 47(3):44–65
29. Heyer GL, Idris SA (2014) Does Analgesic Overuse Contribute to Chronic Post-traumatic Headaches in Adolescent Concussion Patients? *Pediatric Neurology*. maggio;50(5):464–8
30. Bahra A, Walsh M, Menon S, Goadsby PJ (2003) Does Chronic Daily Headache Arise De Novo in Association With Regular Use of Analgesics? *Headache*. marzo;43(3):179–90
31. Wilkinson SM, Becker WJ, Heine JA (2001) Opiate Use to Control Bowel Motility May Induce Chronic Daily Headache in Patients With Migraine. *Headache*. 30 marzo;41(3):303–9
32. Consortium COMOESTAS, Find NL, Terlizzi R, Munksgaard SB, Bendtsen L, Tassorelli C et al (2016) Medication overuse headache in Europe and Latin America: general demographic and clinical characteristics, referral pathways and national distribution of painkillers in a descriptive, multinational, multi-center study. *J Headache Pain* Dicembre 17(1):20
33. Papetti L, Ursitti F, Moavero R, Ferilli MAN, Sforza G, Tarantino S et al (2019) Prophylactic Treatment of Pediatric Migraine: Is There Anything New in the Last Decade? *Front Neurol*. 16 luglio;10:771
34. Kristoffersen ES, Lundqvist C (2014) Medication-overuse headache: epidemiology, diagnosis and treatment. *Therapeutic Adv Drug Saf* Aprile 5(2):87–99
35. Cevoli S, Sancisi E, Grimaldi D, Pierangeli G, Zanigni S, Nicodemo M et al (2009) Family History for Chronic Headache and Drug Overuse as a risk factor for Headache Chronification. *Headache* Marzo 49(3):412–418
36. Cupini LM, De Murtas M, Costa C, Mancini M, Eusebi P, Sarchielli P et al (2009) Obsessive-compulsive disorder and migraine with medication-overuse headache. *Headache* Luglio 49(7):1005–1013
37. Odegard MN, Ceasar RC, Hijaz D, Obinelo A, Rosales A, Bhanvadia S et al Adolescent and parent perceptions of postoperative opioid use: a qualitative, thematic analysis. *Journal of Pediatric Surgery*. dicembre 2023;50(22):34682300756X.
38. Sarchielli P, Corbelli I, Messina P, Cupini LM, Bernardi G, Bono G et al (2016) Psychopathological comorbidities in medication-overuse headache: a multi-centre clinical study. *Euro J Neurol* Gennaio 23(1):85–91
39. Winner P, Pearlman EM, Linder SL, Jordan DM, Fisher AC, Hulihan J et al (2005) Topiramate for Migraine Prevention in children: a Randomized, Double-Blind, Placebo-Controlled Trial. *Headache* Novembre 45(10):1304–1312
40. FDA approves Topamax for migraine prevention in adolescents (2014) *J Pain Palliat Care Pharmacother* Giugno 28(2):191
41. Powers SW, Coffey CS, Chamberlin LA, Ecklund DJ, Klingner EA, Yankey JW et al (2017) Trial of Amitriptyline, Topiramate, and Placebo for Pediatric Migraine. *N Engl J Med*. 12 gennaio;376(2):115–24
42. Ludvigsson J, PROPRANOLOL, USED IN PROPHYLAXIS OF MIGRAINE IN CHILDREN (2009) *Acta Neurol Scand* 29 Gennaio 50(1):109–115
43. Sorge F, Simone RD, Marano E, Nolano M, Orefice G, Carrieri P (1988) Flunarizine in Prophylaxis of Childhood Migraine: a Double-Blind, Placebo-Controlled, crossover study. *Cephalalgia* Marzo 8(1):1–6
44. Ashrafi MR, Salehi S, Malamiri RA, Heidari M, Hosseini SA, Samiei M et al (2014) Efficacy and safety of Cinnarizine in the Prophylaxis of Migraine in children: a double-blind placebo-controlled Randomized Trial. *Pediatr Neurol* Ottobre 51(4):503–508
45. Battistella PA, Ruffilli R, Moro R, Fabiani M, Bertoli S, Antolini A et al (1990) A placebo-controlled crossover trial of Nimodipine in Pediatric Migraine. *Headache* Aprile 30(5):264–268
46. Gesztelyi G (2004) [Primary headache and depression]. *Orv Hetil* 28 Novembre 145(48):2419–2424
47. Tarantino S, Papetti L, Di Stefano A, Messina V, Ursitti F, Ferilli MAN et al (2020) Anxiety, Depression, and Body Weight in Children and adolescents with Migraine. *Front Psychol* 28 Ottobre 11:530911
48. Guidetti V, Galli F, Fabrizi P, Giannantoni A, Napoli L, Bruni O et al (1998) Headache and psychiatric comorbidity: clinical aspects and outcome in an 8-year follow-up study. *Cephalalgia* Settembre 18(7):455–462
49. Lantéri-Minet M, Radat F, Chautard MH, Lucas C (2005) Anxiety and depression associated with migraine: influence on migraine subjects' disability and quality of life, and acute migraine management. *Pain* Dicembre 118(3):319–326
50. Lewandowski AS, Palermo TM, Peterson CC (2006) Age-Dependent relationships among Pain, depressive symptoms, and functional disability in Youth with recurrent headaches. *Headache* Aprile 46(4):656–662

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.