

The Effect of Methylmercury Exposure on Astrocyte of Cerebellar Cortex of Wgite Rats (*Rattus novergicus*)

by Paulus Sugianto

Submission date: 29-Aug-2020 09:17PM (UTC+0800)

Submission ID: 1375952287

File name: rocyte_of_Cerebellar_Cortex_of_Wgite_Rats_Rattus_novergicus.pdf (202.34K)

Word count: 2836

Character count: 16069

THE EFFECT OF METHYLMERCURY EXPOSURE ON ASTROCYTE OF CEREBELLAR CORTEX OF WHITE RATS (*Rattus norvegicus*)

Paulus Sugianto^{1,2}, Sabrina Melisa Pardede³, Ngakan Made Rai Widjaja⁴, Widjiati⁵

¹Department Neurology Faculty of Medicine, Universitas Airlangga, ²Medical Staff of Dr. Soetomo Hospital, ³Faculty of Veterinary Medicine, ⁴Department of Basic Veterinary Medicine, ⁵Department of Anatomy Veterinary, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia

11 ABSTRACT

The aim of this research was to investigate the effect of different dose methylmercury (II) chloride on astrocyte in cerebellar cortex of white rat (*Rattus norvegicus*) exposed. This study used randomized control design using 15 adult female Wistar rats weight 180-200 g of body weight. Before treatment the white rats was adapted in a week, then randomly divided into 3 groups each consist of 5 rats. P0 as control were given 0.5 ml aquades, P1 and P2 were given 0.1 and 0.2 mg/kg/day respectively. All groups were given treatment per oral in 30 days with sonde. The data was analyzed by ANOVA, Duncan's multiple range (Duncan's Multiple Range Test). White rats exposed by methylmercury (II) chloride, had a significant differences in the percentage of necrotic astrocyte ($p < 0.05$). Methylmercury chloride exposure increases the number of necrotic astrocytes on white rat.

Keywords: Methylmercury; astrocyte; necrosis; cerebellar cortex

ABSTRAK

Penelitian ini bertujuan untuk mengetahui efek paparan berbagai dosis dari metilmerkuri klorida terhadap astrosit dari korteks serebellum tikus putih (*Rattus norvegicus*). Penelitian ini merupakan penelitian acak terkontrol yang menggunakan 15 ekor tikus Wistar betina dengan berat badan 180-200 gram. Tikus diberikan adaptasi selama seminggu sebelum perlakuan, kemudian dibagi secara acak menjadi 3 group dan masing-masing group terdiri dari 5 tikus P0 sebagai control diberikan perlakuan dengan aquades sebanyak 5 ml. P1 dan P2 diberikan dengan dosis 0,1 mg/kg BB/hari. Semua group diberikan perlakuan dengan sonde selama 30 hari. Data dianalisa dengan ANOVA, Duncan's multiple range. Tikus putih yang terexpose dengan metilmerkuri klorida mempunyai perbedaan yang signifikan dalam persentase astrosit yang mengalami nekrosis ($p < 0.05$). Paparan metilmerkuri klorida meningkatkan jumlah sel astrosit tikus putih yang mengalami nekrosis.

Kata kunci: Metil merkuri; astrosit; nekrosis; korteks serebellum

Correspondence: Paulus Sugianto, Department Neurology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. Phone : +628989359888. E-mail : paulus.sugianto@gmail.com

pISSN:2355-8393 • eISSN: 2599-056x • doi: <http://dx.doi.org/10.20473/fmi.v55i2.14343>

• Fol Med Indones. 2019;55:122-126 • Received 30 Jan 2019 • Accepted 23 May 2019

• Open access under CC-BY-NC-SA license • Available at <https://e-journal.unair.ac.id/FMI/>

INTRODUCTION

Mercury is toxic pollutant, prevalent widespread and persistent in the environment. Mercury is characterized as a highly malleable liquid at normal temperature and pressure (Bernhoft 2012). Its use in many products and its emission from combustion processes have resulted in well-documented instances of population poisonings, high-level exposures of occupational groups, and worldwide chronic, low-level environmental exposures (NRC 2000).

Methylmercury is structurally the simplest form of the organic mercurial; it bioaccumulates in certain species

of fish, some of which are important human and wildlife foods. There are three primary forms: elemental, inorganic and organic, each with its own distinctive character (Peterson & Talcott 2006). There are several organic mercury compounds; however, by far the most common in the environment and in the food chain is methylmercury. Methylmercury is organic mercury which is always a serious concern in toxicology. Among organic forms, the most toxic is methylmercury (Alexander et al 2008). Most human exposure to mercury is caused by outgassing of mercury from dental amalgam, ingestion of contaminated fish, or occupational exposure (Richardson 1996). Exposure to mercury – even small amounts – may cause serious

health problems, and is a threat to the development of the child in utero and early in life.

The original environmental concerns about mercury related to the outbreaks of neurological disease in Japan in the village of Minamata, where between 70-150 tons of mercury were discharged into the coastal fishing water of Minamata bay. In fresh and salt water, bacteria convert mercury into toxic methylmercury, which primarily attacks the central nervous system. As levels of mercury ingested from fish become a concern, the Food and Drug Association (FDA) set a maximum level of one part mercury per million part seafood (1ppm), which is the same as 1µg mercury in every gram of seafood (Timberlake 2009).

Fish based diets therefore represent a potential source of mercury exposure for dogs and cats. There are many commercial and prescription veterinary diets that contain fish. Fish oil supplement are also frequently used in therapeutic regimens for a variety of conditions for dogs and cats (Peterson & Talcott 2006). Methyl mercury is easily absorbed through the gut and deposits in many tissues, but does not cross the blood-brain barrier as efficiently as elemental mercury; however, on entering the brain it is progressively demethylated to elemental mercury (Berlin et al 2007).

Brain tissue is a heterogeneous system comprising of two distinct compartments known as neurons and glia. The glial cells play a significant role by supplying neurons with a number of metabolites and precursors (Kaur 2008). They also have a role in maintaining tight junction of capillaries that form the blood-brain barrier (Ross & Pawlina 2010). In mammalian CNS, astrocytes are known as a preferential site of methylmercury accumulation and a main target of toxicity. Methylmercury preferentially accumulates in astrocytes and inhibits uptake systems for glutamate and cysteine transport, both of which will compromise glutathione (GSH) synthesis and redox status in astrocytes (Hurtado et al 2008). Reduced GSH only neutralizes the ROS overflow. Presumably, this occurs downstream from the ROS activity responsible for the critical step in necrotic cell death (Fiers et al 1999).

Methylmercury depolarizes the presynaptic membrane which increases the Na²⁺ and decreases K⁺ ion concentration. This causes disruption of Ca²⁺ homeostasis leading to increased intracellular Ca²⁺ concentration (Kaur 2008).

The cerebellum is responsible for involuntary control of balance, posture and coordination of movement. It attempt to maintain equilibrium and balance on receiving sensory information from portions of the inner

ear, as well as visual and proprioceptive input. The bulk of cerebellum is composed of white matter, with a thin layer of gray matter on the surface (Christenson 2009). The gray matter of the cerebellum (cerebellar cortex) forms the periphery of this portion of the brain. Neuron within this tissue is involved in directing activities associated with vestibulation (balance), skeletal muscle and joints as well as spontaneously from cerebellar nuclei.

Methylmercury generally accumulates in the central nervous system and is found most widely in the cortex and cerebellum. The most sensitive part of brain to methylmercury are cerebral cortex (especially visual cortex) and granular membrane of cerebellum (Ganiswara et al 1995). Brain of patient with Minamata disease in Japan demonstrated unintegrated cerebellar cortex neuron because of consumption of raw fish exposed to mercury (Widowati et al 2008). Methylmercury exposure symptoms are mostly neurological, such as impaired vision, ataxia, paresthesia, neurasthenia, hearing loss, dysarthria, mental retardation, tremor, motor impairment, paralysis and death (Ganiswara et al 1995, Bernhoft, 2012). Some patients also experience delayed reaction time, poor fine motor control, and deficits in mental concentration, vocabulary, task switching, and the One Hole test, as well as mood lability (Echeverria et al 1995)

Based on the background described above the researcher the researcher conducted a study about the percentage of necrotic astrocyte in the cerebellar cortex of white rats (*Rattus norvegicus*) exposed to different dose of methylmercury chloride.

MATERIAL AND METHODS

Setting

This study was conducted at 2 sites, i.e. Departement of Embryology, Faculty of Veterinary and Department of Anatomy and Hystology, Faculty of Medicine, Universitas Airlangga, Surabaya.

Fifteen adult female Wistar rats (*Rattus norvegicus*) weighed 180-200g purchased at a breeder in Batu, Malang. Fifteen adult female Wistar rats were placed into 3 plastic cages of 30 x 35 x 35 cm in size. Each had five animals, and were adapted in the same relative condition for a week, and fed with chicken pellet and given drink of tap water. One week later the subjects were weighed and randomly divided into three groups, consist of: P0: the rats were given 0.5 ml aquades orally in 30 days, P1: methylmercury (II) chloride

0.1mg/kg/day, P2: ²⁹ methylmercury (II) chloride 0.2mg/kg/day

Methylmercury (II) chloride was administered orally via Nasogastric tube once daily at 12.00 for 30 days to see the damage from low-chronic dose in white rat brain (adapted from (Huang et al 2011)).

After 30 days, 15 adult female Wistar rats were euthanized with inhalation method by putting them inside a jar that was filled with chloroform sprayed cotton. Brain samples were collected after performing dissection on each subject brain, then inserted into plastic pot contained formalin 10%. Histopathological slides were made using HE stain at Department, school of medicine, University of Airlangga. The presence of necrotic (pyknosis) astrocyte in the cerebellar cortex was assessed and calculated on 5 different field of view, with magnification 400 times. Then the percentage of necrotic (pyknosis) astrocytes against necrotic cells was measured. This research used percentage parameter to reduce errors that could occur in the calculation using this formula as follows:

$$\left[\frac{\text{The amount of necrotic astrocytes}}{\text{The amount of normal + necrotic astrocytes}} \right] \times 100 \%$$

Experimental design and data analysis

This was a completely randomized designed study with 3 groups of white rats, with 3 types of treatment. Each treatment repeated 5 times. Sample size was $t(n-1) = 15$, with t as the number of treatments and n as the number of repetitions (Kusriningrum 2010).

RESULTS AND DISCUSSION

Data obtained from research about the percentage of necrotic astrocyte in the cerebellar cortex of white rats

(*Rattus norvegicus*) exposed to different dose of methylmercury (II) chloride can be seen in Table 1.

Fig. 1 shows that treatment P2 has the highest necrotic astrocytes mean, which is 59.43%. The lowest necrotic astrocyte mean is P0 0.65%. Therefore, among all three treatments there are significant differences ($p < 0.05$).

Histopathological view of cerebellar cortex from white rats (*Rattus norvegicus*) exposed by different dose of methylmercury (II) chloride showed a change, that was the increase of necrotic astrocytes. Statistical analysis showed effect of methylmercury (II) chloride on necrotic astrocytes ($p < 0.05$). There was a significant difference between treatment and control. The higher dose of methylmercury (II) chloride, the greater effect could be seen on histopathological view of astrocyte in cerebellar cortex of white rats. White rats that received treatment P2 had the highest necrotic astrocytes, which was 59.43 ± 10.30 . The rats in P2 group also showed several clinical sign, like incoordination movement and difficulty in walking. As mentioned by Waldichuk (1974) that cited by Ulum (Ulum 2005), the toxic effects of mercury on the body of organisms, depend on the amount of that substance in the body, the entry point of substances and period of mercury exposure with the sensitive organ in the body. After long-term subclinical methylmercury exposure, may be a proximate toxic form of mercury responsible for the changes within the astrocyte and microglial.

The changes in the structure of the nucleus can be in the form of pyknosis, karyorrhexis and karyolysis. Karyolysis occurs when the chromatin diffuses out through the nuclear membrane so the nucleus is only seen as an empty space. Karyorrhexis is characterized by a fragmented nucleus. Pyknosis is a state where chromatin condenses, so in HE staining, the nucleus will appear darker (Herawati & Handari 2004). In this research, author calculated the percentage of necrotic astrocytes which had pyknosis.

Table 1. The mean of necrotic astrocyte in cerebellar cortex of white rats exposed by methylmercury (II) chloride in Arc sin vy (%)

Replications	Necrotic Astrocyte		
	P0	P1	P2
1	0	26.92	44.58
2	0	20.87	53.71
3	3.25	22.57	62.51
4	0	20.85	70.23
5	0	24.53	66.17
Mean	0.65	23.14	59.43
SD	1.45	2.59	10.30
(X±SD)	²⁰ 0.65±1.45	23.14 ^b ±2.59	59.43 ^a ±10.30

Annotation: Different superscripts in the every columns show significant differences ($p < 0.05$)




Fig. 1. Necrotic astrocyte mean in cerebellar cortex of white rats exposed to methylmercury (II) chloride (%).

The necrotic astrocyte in the cerebellar cortex of white rats occurs because of methylmercury induce ROS and increases intracellular Ca^{2+} . Calcium and reactive oxygen species (ROS) are the main players during the propagation and execution phases of necrotic cell death, directly or indirectly provoking damage to proteins, lipids and DNA, which culminates in disruption of organelle and cell integrity (Festjens et al 2006). ROS can react with biomolecules and cause oxidative damage and even necrosis (Maurino & Flugge 2008).

Methylmercury causes a depletion of neuronal glutathione that increases the intracellular concentration of methylmercury in neurons and therefore influences the generation of ROS (Kaur et al 2006). Excessive production of ROS leads to oxidative stress, damage of intracellular molecules and organelles, and ultimately necrosis (Zong & Thompson 2006).

In P0 groups necrotic astrocyte occurred in small amounts, even though the white rats in the groups not exposed to methylmercury (II) chloride. Necrosis will occur in the development and maintenance of organismal homeostasis, and also in cellular stress, resulting from nutrient starvation or reduced oxygen supply (Zong & Thompson 2006). According to Festjens et al. (Festjens et al 2006) necrotic cell death also occur in immune defense mechanism and physico-chemical stress such as freeze-thawing or severe hyperthermia.

CONCLUSION

Methylmercury (II) chloride, can be concluded that methylmercury (II) chloride exposure can increase the number of necrotic astrocyte in cerebellar cortex of

white rats. The percentage of necrotic astrocyte among all treatment showed significant differences, the highest percentage of necrotic astrocytes was obtained in group exposed to methylmercury (II) chloride 0.2mg/kg/day, and the lowest in control group.

28

REFERENCES

- Alexander J, Audunsson GA, Benford DA, Cockburn JP, Cravedi E, et al (2008). Mercury as undesirable substance in animal feed. *Efsa* 654, 1-76
- Berlin M, Zalups RK, Fowler BA (2007). Mercury in handbook on the toxicology of metals. New York, Elsevier
- Bernhoft RA (2012). Mercury toxicity and treatment: a review of the literature. *J environ public health* 2012, 460508
- Christenson DE (2009). Veterinary medical terminology. Canada, WB Saunder Company
- Echeverria D, Heyer NJ, Martin MD, Naleway CA, Woods JS, Bittner AC jr (1995). Behavioral effects of low-level exposure to elemental hg among dentists. *Neurotoxicol Teratol* 17, 161-8
- Festjens N, Vanden Berghe T, Vandenabeele P (2006). Necrosis, a well-orchestrated form of cell demise: signalling cascades, important mediators and concomitant immune response. *Biochim Biophys Acta* 1757, 1371-87
- Fiers W, Beyaert R, Declercq W, Vandenabeele P (1999). More than one way to die: apoptosis, necrosis and reactive oxygen damage. *Oncogene* 18, 7719-30
- Ganiswara SG, Setiabudy R, Suyatna FD, Purwastyastuti, Nafrialdi (1995). Farmakologi dan terapi. 4th Ed. Jakarta, Fakultas Kedokteran Universitas Indonesia

- Herawati E, Handari S (2004). Pengaruh infus jamur ling zhi (*Ganoderma lucidum* (*leys ex fr.*) Karst) terhadap mencit (*Mus musculus*, L.) yang diberi timbal: kajian struktur mikroanatomi Ren. Enviro 4, 55
- Huang CF, Liu SH, Hsu CJ, Lin-Shiau SY (2011). Neurotoxicological effects of low-dose methylmercury and mercuric chloride in developing offspring mice. Toxicol Lett 201, 196-204
- Hurtado O, Pradillo JM, Fernandez-Lopez D, Morales JR, Sobrino T, Castillo J, Alborch E, Moro MA, Lizasoain I (2008). Delayed post-ischemic administration of cdp-choline increases eaat2 association to lipid rafts and affords neuroprotection in experimental stroke. Neurobiol dis 29, 123-31
- Kaur P (2008). Cellular and molecular mechanisms behind methylmercury-induced neurotoxicity. Doctor, Norwegian University of Science and Technology
- Kaur P, Aschner M, Syversen T (2006). Glutathione modulation influences methyl mercury induced neurotoxicity in primary cell cultures of neurons and astrocytes. Neurotoxicology 27, 492-500
- Kusriningrum RS (2010). Perancangan percobaan. Surabaya, Airlangga University Press
- Maurino VG, Flugge UI (2008). Experimental systems to assess the effects of reactive oxygen species in plant tissues. Plant Signal Behav 3, 923-8
- NRC (2000). Toxicological effects of methylmercury. Washington DC, National Academy Press
- Peterson ME, Talcott PA (2006). Small animal toxicology. St. Louis, Missouri, United States, Elsevier Inc
- Richardson M (1996). The safety of dental amalgam. Canada, Minister of Health
- Ross MH, Pawlina W (2010). Hystology a text and atlas. Philadelphia, Lippincot Williams & Willkins
- Timberlake KC (2009). Chemistry: an introduction to general, organic and biological chemistry. Los Angeles, Prentice Hall
- Ulum R (2005). Pengaruh pemberian merkuriklorida (hgcl2) terhadap gambaran histopathologi otak mencit (*Mus musculus*). Surabaya, Universitas Airlangga
- Widowati W, Sastiono A, Raymond J (2008). Efek toksik logam: Pencegahan dan penanggulangan. Yogyakarta, CV Andi Offset
- Zong WX, Thompson CB (2006). Necrotic death as a cell fate. Genes Dev 20, 1-15

The Effect of Methylmercury Exposure on Astrocyte of Cerebellar Cortex of Wgite Rats (Rattus novergicus)

ORIGINALITY REPORT

21%

SIMILARITY INDEX

14%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	Michael Gilbertson, David O Carpenter. "An ecosystem approach to the health effects of mercury in the Great Lakes basin ecosystem", Environmental Research, 2004 Publication	2%
2	www.nap.edu Internet Source	1%
3	www.toxicswatch.org Internet Source	1%
4	genesdev.cshlp.org Internet Source	1%
5	journal.unair.ac.id Internet Source	1%
6	www.explainmedicine.com Internet Source	1%
7	"Necrotic Cell Death", Springer Science and Business Media LLC, 2014 Publication	1%

8	www.nature.com Internet Source	1%
9	Yin, Z.. "Methylmercury induces oxidative injury, alterations in permeability and glutamine transport in cultured astrocytes", Brain Research, 20070202 Publication	1%
10	semspub.epa.gov Internet Source	1%
11	repository.unair.ac.id Internet Source	1%
12	web-archive-collection.blogspot.com Internet Source	1%
13	doaj.org Internet Source	1%
14	Ling Wang, Haiyan Jiang, Zhaobao Yin, Michael Aschner, Jiyang Cai. "Methylmercury Toxicity and Nrf2-dependent Detoxification in Astrocytes", Toxicological Sciences, 2009 Publication	1%
15	eprints.umsida.ac.id Internet Source	1%
16	Guo Lin Wu, Jing Cui, Ling Tao, Hong Yang. "Fluroxypyr triggers oxidative damage by producing superoxide and hydrogen peroxide in	1%

rice (*Oryza sativa*)", Ecotoxicology, 2009

Publication

17

f1000research.com

Internet Source

1%

18

J.M.C. Sousa, L.A. Matos, D.F.A. Alcântara, H.F. Ribeiro et al. "Cellular responses induced in vitro by pestheic acid, a fungal metabolite, in a gastric adenocarcinoma cell line (PG100)", *Genetics and Molecular Research*, 2013

Publication

<1%

19

repositorio.ufsm.br

Internet Source

<1%

20

Bhawna Kushawaha, Rajkumar Singh Yadav, Dilip Kumar Swain, Pradeep K Rai, Satish Kumar Garg. "Mercury-Induced Inhibition of Tyrosine Phosphorylation of Sperm Proteins and Altered Functional Dynamics of Buck Spermatozoa: an In Vitro Study", *Biological Trace Element Research*, 2020

Publication

<1%

21

www.omicsonline.org

Internet Source

<1%

22

"Abstracts 2007", *Diabetologia*, 2007

Publication

<1%

23

www.exeley.com

Internet Source

<1%

24	www.scribd.com Internet Source	<1%
25	uin-suka.ac.id Internet Source	<1%
26	www.scielo.br Internet Source	<1%
27	Francesca Gorini, Filippo Muratori, Maria Aurora Morales. "The Role of Heavy Metal Pollution in Neurobehavioral Disorders: a Focus on Autism", Review Journal of Autism and Developmental Disorders, 2014 Publication	<1%
28	Allan W. Carson, P. Michael Rutherford, Philip J. Burton. "Desulphurized tailings serve as a useful soil supplement for mine reclamation", Canadian Journal of Soil Science, 2014 Publication	<1%
29	Pedro Espitia-Pérez, Suelen Marin Albino, Helen Tais da Rosa, Alexandre Kleber Silveira et al. "Effects of methylmercury and retinol palmitate co-administration in rats during pregnancy and breastfeeding: Metabolic and redox parameters in dams and their offspring", Ecotoxicology and Environmental Safety, 2018 Publication	<1%
30	"Toxicological Profile for Mercury", ATSDR s	

Toxicological Profiles Web Version, 2002.

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The Effect of Methylmercury Exposure on Astrocyte of Cerebellar Cortex of Wgite Rats (Rattus novergicus)

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
