

Urban-rural disparity of preventive healthcare utilisation among children under the universal health insurance coverage in Taiwan: a national birth cohort analysis

by C. L. Lu

Submission date: 18-Aug-2020 11:36AM (UTC+0800)

Submission ID: 1370864044

File name: ion_among_children_under_the_UH_Insurance_Coverage_in_Taiwan.pdf (317.65K)

Word count: 7682

Character count: 39924



Original Research

1
Urban–rural disparity of preventive healthcare utilisation among children under the universal health insurance coverage in Taiwan: a national birth cohort analysis



C.L. Lu^a, Y.H. Hsu^b, W.L. Su^b, N.A. Damayanti^c, C.W. Chen^d, Y.J. Lin^e, K.S. Tsai^{f,i}, C.Y. Li^{b,c,g,h,*}

^a Graduate Institute of Food Safety, College of Agriculture and Natural Resources, National Chung Hsing University, Taichung, Taiwan

^b Department of Public Health, College of Medicine, National Cheng Kung University, Tainan, Taiwan

^c Department of Health Policy and Administration, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

^d School of Nursing, National Yang-Ming University, Taipei, Taiwan

^e Department of Pediatrics, College of Medicine and Hospital, National Cheng Kung University, Tainan, Taiwan

^f Department of Pediatrics, Tainan Sinlau Hospital, Tainan, Taiwan

^g Department of Public Health, College of Public Health, China Medical University, Taichung, Taiwan

^h Department of Healthcare Administration, College of Medical and Health Science, Asia University, Taichung, Taiwan

16
ARTICLE INFO

Article history:

Received 25 July 2019

Received in revised form

18 January 2020

Accepted 14 February 2020

Available online 1 April 2020

Keywords:

Low birth weight

Child

Rural health services

Healthcare disparities

Cohort studies

ABSTRACT

Objective: In the context of universal health insurance coverage, this study aimed to determine whether urban–rural inequality still exists in preventive health care (PHC) amongst children in Taiwan.

Study design: Prospective cohort study.

Methods: A total of 184,117 mothers and their children born in 2009 were identified as the study cohort. The number of children born in urban, satellite and rural areas was 40,176, 57,565 and 86,805, respectively. All children were followed for 7 years, before which a total of seven times PHC were provided by Taiwan's National Health Insurance (NHI) programme. Ordinal logistic regression models were used to associate urbanisation level with the frequency of PHC utilisation. Stratified analyses were further performed in accordance with the children's birth weight and the mothers' birthplace.

Results: Children from satellite areas had higher utilisation for the first four scheduled PHC visits. Children living in urban areas received more PHC for the fifth and sixth scheduled visits. Compared with those from rural areas, children in satellite areas exhibited a small but significant increase in odds in PHC utilisation, with a covariate-adjusted odds ratio (aOR) of 1.04 and 95% confidence interval (CI) of 1.02–1.06. By contrast, no significant difference was observed between rural and urban areas (aOR = 1.01). Further stratified analyses suggest more evident urban–rural difference in PHC utilisation amongst children with low birth weight and foreign-born mothers.

Conclusions: Given a universal health insurance coverage and embedded mechanisms in increasing the availability of healthcare resources in Taiwan, a slight urban–rural difference is observed in PHC utilisation amongst children. Hence, sociodemographic inequality in utilisation of PHC still exists. This issue should be addressed through policy intervention.

© 2020 The Royal Society for Public Health. Published by Elsevier Ltd. All rights reserved.

15
Introduction

Preventive health care (PHC) plays a crucial role in children's health through the early detection of health problems. The children's PHC programme of Taiwan, also known as well-child visits, provides scheduled primary immunisations and health checkups for physical, developmental and behavioural problems; it also offers counselling to caregivers.¹ Regular health checkups for children

* Corresponding author. Department and Graduate Institute of Public Health, College of Medicine, National Cheng Kung University, #1, University Rd., Tainan, 70101, Taiwan. Tel.: +886 6 2353535x5862.

E-mail address: cyli99@mail.ncku.edu.tw (C.Y. Li).

¹ K–S Tsai and C–Y Li contributed equally to this article.

have been effective in reducing neonatal and infant mortality, decreasing the likelihood of hospitalisation and improving children's health.^{2–4} Despite all these benefits, the utilisation rate of health checkups varies worldwide. In the USA, for example, 58.9% of children took recommended well-child visits in 2007–2008.⁵ An Australian survey conducted in 2012–2013 showed that 70.3% of parents with children aged under 6 months and 48.6% of parents with children aged 3–5 years reported that they regularly visit child and family health nurses.⁶ In Taiwan, the utilisation rate of PHC for children under 7 years was approximately 70% in 2002 and increased to 78.3% in 2015.³

Disparity in healthcare utilisation in urban and rural areas amongst children has been reported in previous studies. The determinants of such disparity can result from inadequate healthcare resources, low socio-economic status of families, lack of insurance coverage, poor health literacy of parents and negative attitude towards health checkup in rural areas.^{7,8} Previous studies have shown an evident urban–rural variation in morbidity and mortality amongst children from developing countries;^{9,10} meanwhile, disparity still exists in health problems between urban and rural areas amongst children from developed countries. These health problems include hospitalisation due to physical abuse, emergency department reliance, drug-related poisoning and underdiagnosis of obesity, which are more prevalent in children living in rural areas than in urban areas.^{11–14} Although well-child visit can play varying roles in different countries, reducing the gap in PHC utilisation between urban and rural areas is expected to effectively improve the disparity in children's health problems.

Health insurance coverage is a pivotal factor that determines whether a baby will use PHC.¹ The low insurance coverage in rural areas may result from the poor socio-economic status of people from such areas.¹⁵ The government of Taiwan launched its universal health insurance coverage in 1994, and over 99% of Taiwan residents have been insured since 2009.¹⁶ The present study aimed to determine if urban–rural disparity in PHC utilisation still exists in Taiwan after nearly two decades of universal health insurance coverage.

29 Methods

Data source

The data analysed in this study included Taiwan's national birth notification (BN), national health insurance database (NHID) and parents–child pair dataset (PCPD). BN includes various pieces of information about mothers, including personal identification number (PID), birthday, birthplace, delivery date and residential city district/township at the time of delivery. BN also includes information regarding gestational week and birth weight for each live birth. NHID comprises various subdatasets, including beneficiary registry (BR) and inpatient/outpatient medical claims. BR includes the demographic characteristics, insurance premium and PID of each individual covered by Taiwan's National Health Insurance (NHI) programme. NHID covers claim data for nearly all (>99%) Taiwanese residents.¹⁶ To ensure the accuracy of claim data, the National Health Insurance Administration (NHIA) performs quarterly expert reviews on a random sample for every 50 to 100 ambulatory and inpatient claims (NHIA, 2018), and the information available from NHIA is considered complete and accurate.¹⁷

To obtain the link between newborns and their parents, we used PCPD to identify parental PIDs for each newborn. By linking PCPD and BN, we determined the gestational weeks and birth weight of each newborn. In addition, paternal age at the time of delivery can be obtained through the link between PCPD and BR. The various datasets and the study variables derived from these datasets are illustrated in Appendix A. We also used the national death registry (DR) to identify

newborns that died under the age of 7 years and were unable to complete the seven PHC visits provided by the NHI programme.

The current study protocol was approved by the Institutional Review Board of National Cheng Kung University Hospital (No. B-ER-107-014). Access to the aforementioned databases was approved by the Health and Welfare Data Science Centre (HWDSC) of the Ministry of Health and Welfare. To guard the data, the data management and statistical analyses involved in this study were conducted on-site at HWDSC.

Study design

This work is a nationwide cohort study of all neonates born in 2009, who were followed up until the age of 7 years. We initially identified 186,044 newborns from BR in 2009. After excluding the newborns whose mothers' PIDs were not found in BN ($n = 37$) and whose mothers' residential city district/township was missing ($n = 509$), 185,498 children were left in the study cohort. We further excluded children who died during follow-up (i.e. prior to the age of 7 years) ($n = 954$). A total of 184,544 children were finally included in the current analysis.

Level of urbanisation

On the basis of the information regarding the residential city district/township of a mother at the time of delivery, we classified maternal residential area into different urbanisation levels: rural, satellite and urban areas. The classification scheme for determining urbanisation level was proposed by Liu et al. who categorised all 316 city districts and townships in Taiwan into seven clusters in accordance with population density, proportion of residents with college or higher education levels, percentage of elderly (>65 years old) people, proportion of the agricultural workforce and number of physicians per 10⁵ people.¹⁸ In this study, we regarded the most and second most urbanised clusters as 'urban' and 'satellite' areas, respectively, and the remaining ones as 'rural' areas.¹⁹

PHC services

In addition to the treatment-oriented medical care provided by the NHI programme, the Health Promotion Administration (HPA) of the Ministry of Health and Welfare also provides every child with seven PHC services before his or her 7th birthday. PHC services include primary regular physical examinations, dental checkup, assessment of physical and mental development and screening for various disorders, such as hearing loss, inherited metabolic disorders and ophthalmological problems.³ Regular PHC is expected to increase the probability of early detection of certain disorders, and thus, provide timely referrals for clinical treatment. These services are directly provided by paediatricians under contract with the NHI programme.

Every child may receive PHC service when he or she reaches the ages of 0–2 months (1st), 2–4 months (2nd), 4–10 months (3rd), 10 months to 1.5 years (4th), 1.5–2 years (5th), 2–3 years (6th) and 3–7 years (7th). The utilisation of PHC service during the aforementioned periods can be identified from the medical order codes in NHID, which were IC11/IC71, IC12/IC72, IC13/IC73, IC15/IC75, IC16/IC76, IC17/IC77 and IC19/IC79, respectively.

Covariates

A number of parental and children's characteristics were considered, including paternal and maternal ages at the time of delivery, birth weight and mother's birthplace. Birth weight was classified as normal birth weight (NBW, ≥ 2500 g), low birth weight (LBW, 1500–2499 g) and very low birth weight (VLBW, <1500 g).

Paternal and maternal ages were categorised as <18, 18 to <25, 25 to <35 and ≥ 35 years. Maternal birthplace was grouped under 'otherwise' if a mother was not a native-born Taiwanese.

Statistical analysis

Sociodemographic characteristics were compared amongst children with maternal residential city/district/township in rural, satellite and urban areas. We used a Chi-squared test to compare the differences of discrete variables and Student's *t*-test to compare continuous variables. The prevalence of PHC utilisation was calculated by dividing the total number of utilisation by the study's children with the total number of children. A Chi-squared test was also performed to compare the prevalence of PHC utilisation at each scheduled visit amongst children living in rural, satellite or urban areas.

We used an ordinal logistic regression model to estimate the odds of higher PHC utilisation in association with urbanisation level and various covariates. Crude odds ratio (OR), adjusted OR and their corresponding 95% confidence intervals (CIs) were estimated from the regression coefficients derived from the ordinal logistic regression. We tested the assumption of proportional odds for the ordinal logistic regression model and determined that the assumption holds.²⁰ All data management and analyses were performed using statistical software SAS for Windows (version 9.4; SAS Institute, Cary, NC). A *P*-value of 0.05 or less was considered statistically significant.

Results

Table 1 shows and compares the characteristics of the study's children in rural, satellite and urban areas. A higher number of boys were reported in the three study groups. The prevalence of LBW children was higher in rural areas (8.09%) than in satellite (7.62%) and urban (7.64%) areas. Most mothers had given birth at the age of

25–34 years (72%–74%). Mothers under 18 years old were slightly more common in rural areas (0.65%) than in satellite (0.35%) and urban (0.25%) areas. Similarly, fathers in rural areas were slightly younger. Approximately 6.03% of the study's children had foreign-born mothers, with higher prevalence amongst children from rural areas (7.07%) than from satellite and (5.39%) urban (5.64%) areas.

Fig. 1 compares PHC utilisation for each scheduled visit amongst the three study groups. Within-group and between-group variations were observed in utilisation rate. In all three study groups, the 1st scheduled visit was associated with the lowest utilisation rate, and the 3rd and 4th scheduled visits tended to have higher rates. In addition, whilst the difference in utilisation rate was small between satellite and urban areas, children from rural areas tended to have lower utilisation rates for most of the scheduled visits. The detailed information about utilisation and comparison amongst the study groups are presented in Appendix B.

Table 2 provides the crude and covariate-adjusted ORs of the high utilisation of PHC services in association with urbanisation level and various covariates. Compared with children living in rural areas, those from satellite areas had significantly higher odds of utilising PHC service (adjusted OR: 1.04, 95% CI: 1.02–1.06). Such significantly elevated OR, however, was not observed amongst children from urban areas. Apart from lower urbanisation, a number of covariates were also associated with significantly lower utilisation, including lower birth weight (adjusted OR: 0.85, 95% CI: 0.82–0.88 for LBW; adjusted OR: 0.20, 95% CI: 0.18–0.22 for VLBW), younger parents [adjusted OR: 0.72, 95% CI: 0.61–0.88 for mothers <18 years; adjusted OR: 0.51 (95% CI 0.33–0.78) and 0.85 (95% CI 0.81–0.89) for fathers <18 years and 18–24 years, respectively] and foreign-born mothers (adjusted OR: 0.64, 95% CI: 0.62–0.67).

We assessed the potential effect modification by each covariate and noted significant ($P < 0.05$) interactions of urbanisation level with birth weight and maternal birthplace. We then performed stratified analyses in accordance with the two effect modifiers. The

Table 1
Comparison of characteristics among study subjects with different residential levels of urbanisation.

Characteristics	Level of urbanisation						P-value
	Rural		Satellite		Urban		
	n	%	n	%	n	%	
Gender							0.13
Male	45,350	52.24	29,787	51.74	20,824	51.83	
Female	41,454	47.76	27,777	48.25	19,352	48.17	
Birthweight							<0.01
Normal	79,782	91.91	53,181	92.39	37,106	92.36	
Low	6434	7.41	4005	6.96	2801	6.97	
Very low	588	0.68	378	0.66	269	0.67	
Mean \pm SD	3066.6	450.59	3079.5	447.71	3086.5	452.39	<0.01
Maternal age (yrs)							<0.01
<18	564	0.65	201	0.35	100	0.25	
18–24	13,880	15.99	5031	8.74	3037	7.56	
25–34	62,499	72.00	42,396	73.65	29,526	73.49	
≥ 35	9861	11.36	9936	17.26	7513	18.70	
Mean \pm SD	29.41	4.63	30.89	4.45	31.22	4.39	<0.01
Paternal age (yrs)							<0.01
<18	52	0.06	6	0.01	3	<0.01	
28–24	5356	6.17	1704	2.96	940	2.34	
25–34	56,795	65.47	34,832	60.52	23,765	59.15	
≥ 35	24,601	28.36	21,022	36.52	15,468	38.50	
Mean \pm SD	32.48	5.35	33.72	5.05	34.03	5.05	<0.01
Mother's birthplace							<0.01
Taiwan	80,670	92.93	54,461	94.61	37,912	94.36	
Otherwise	6134	7.07	3103	5.39	2264	5.64	
Total	86,804	47.04	57,564	31.19	40,176	21.77	

^a Normal birthweight: ≥ 2500 g, low birthweight: 1500–2499g, very low birthweight: <1500g.

^b Based on χ^2 or Student's *t* test.

^c Inconsistency between total number and summation of specific numbers for some variables was due to missing data.




Fig. 1. Graphic illustration of urbanisation level specific prevalence of preventive healthcare visit for each scheduled visit at age 0–6 years.

urban–rural difference in OR of high PHC utilisation was only significant amongst LBW infants and those with foreign-born mothers. Amongst LBW infants, satellite (adjusted OR: 1.09, 95% CI: 1.01–1.18) and urban (adjusted OR: 1.17, 95% CI: 1.08–1.27) areas were associated with a significantly higher adjusted OR of PHC utilisation, with a significant trend. A similar trend was observed amongst VLBW children (corresponding adjusted OR = 1.09 and 1.22, $P = 0.13$) and those with foreign-born mothers (adjusted OR = 1.09 and 1.19, $P < 0.01$), as shown in Table 3.

Discussion

Only a slight urban–rural difference in PHC utilisation was observed amongst young children in the context of universal health insurance coverage in Taiwan. However, urban–rural difference was more evident in certain segments of children’s population, including children with lower birth weight and foreign-born mothers. In addition, some sociodemographic characteristics were associated with lower PHC utilisation, including lower birth weight, younger or older parents and foreign-born mothers.

The small disparity across different urbanisation levels noted in our findings slightly differs from observations made in other countries that frequently do not have universal health coverage. A US study reported a covariate-adjusted difference of 12.2% in well-child visit adherence ratio in favour of children from metropolitan statistical areas (MSAs) compared with those living in non-MSA areas in 2007–2008.⁵ Although this US study determined that the utilisation of well-child visits increased through years 1996–98 to 2007–08, the increase was smaller in uninsured subgroups than in insured ones. In the uninsured children population, the increase in adherence ratio was from 4.8% to 32.0%; by contrast, the increase was from 9.7% to 56.6% during the same period amongst publicly insured children and from 14.6% to 63.6% in the privately insured

Table 2
Odds ratio of higher frequency of preventive healthcare utilisation in association with level of urbanisation and other sociodemographic characteristics.

	Total number of preventive healthcare utilisation								Crude estimate			Adjusted estimate			
	0	1	2	3	4	5	6	7	OR ^a	95% CI ^a	P	OR ^a	95% CI ^a	P	
Level of urbanisation															
Rural	1554	2403	3799	6233	10,506	17,284	24,094	20,931	Ref.			Ref.			
Satellite	785	1322	2162	3752	6873	11,988	17,082	13,600	1.06	1.04–1.08	<0.01	1.04	1.02–1.06	<0.01	
Urban	549	995	1556	2654	4825	8498	12,025	9074	1.02	1.00–1.04	0.05	1.01	1.00–1.04	0.21	
Gender															
Male	1514	2394	3837	6632	11,728	19,749	27,618	22,489	Ref.			Ref.			
Female	1374	2326	3680	6007	10,476	18,021	25,583	21,116	1.02	1.00–1.03	0.06	1.02	1.00–1.04	0.02	
Birthweight^b															
Normal	2451	4156	6711	11,365	20,369	35,050	49,417	40,550	Ref.			Ref.			
Low	307	466	638	1092	1646	2529	3584	2978	0.84	0.82–0.87	<0.01	0.85	0.82–0.88	<0.01	
Very low	130	98	168	182	189	191	200	77	0.20	0.18–0.22	<0.01	0.20	0.18–0.22	<0.01	
Maternal age (yrs)															
<18	39	59	95	110	125	159	177	102	0.46	0.40–0.51	<0.01	0.72	0.61–0.85	<0.01	
18–24	472	784	1123	1787	2876	4406	5703	4797	Ref.			Ref.			
25–34	1769	3046	4982	8692	15,741	27,484	39,670	33,036	1.30	1.27–1.33	<0.01	1.17	1.13–1.20	<0.01	
≥35	608	831	1317	2050	3462	5721	7651	5670	1.03	1.00–1.06	0.06	1.01	0.97–1.05	0.58	
Paternal age (yrs)															
<18	3	8	5	5	8	12	12	8	0.35	0.23–0.53	<0.01	0.51	0.33–0.78	<0.01	
18–24	190	271	436	669	1025	1568	2102	1739	0.76	0.73–0.80	<0.01	0.85	0.81–0.89	<0.01	
25–34	1392	2530	4234	7480	13,571	23,623	33,915	28,647	Ref.			Ref.			
≥35	1303	1911	2842	4485	7600	12,567	17,172	13,211	0.82	0.80–0.83	<0.01	0.88	0.87–0.90	<0.01	
Mother’s birthplace															
Taiwan	2412	4053	6746	11,589	20,718	35,656	50,488	41,381	Ref.			Ref.			
Otherwise	476	667	771	1050	1486	2114	2713	2224	0.60	0.58–0.62	<0.01	0.64	0.62–0.67	<0.01	
Total	2888	4720	7517	12,639	22,204	37,770	53,201	43,605							

^a OR: odds ratio; CI: confidence interval.

^b Normal birthweight: ≥2500g, low birthweight: 1500–2499g, very low birthweight: <1500g.

Table 3

Odds ratio of higher frequency of preventive healthcare utilisation in association with levels of urbanisation stratified by children's birthweight and birthplaces of mothers.

	Total number of preventive healthcare utilisation								aOR ^a	95% CI ^b	P-value for interaction	P-value for trend in OR
	0	1	2	3	4	5	6	7				
Birthweight^b												
<i>P</i> < 0.01												
Normal												
Rural	1335	2118	3364	5567	9591	15,963	22,336	19,508	Ref.			
Satellite	663	1171	1938	3398	6318	11,116	15,902	12,675	1.04	1.02–1.06		
Urban	453	867	1409	2400	4460	7971	11,179	8367	1.00	0.98–1.02		<i>P</i> = 0.87
Low												
Rural	158	239	349	569	823	1233	1671	1392	Ref.			
Satellite	82	118	174	305	497	806	1121	902	1.09	1.01–1.18		
Urban	67	109	115	218	326	490	792	684	1.17	1.08–1.27		<i>P</i> < 0.01
Very low												
Rural	61	46	86	97	92	88	87	31	Ref.			
Satellite	40	33	50	49	58	66	59	23	1.09	0.86–1.39		
Urban	29	19	32	36	39	37	54	23	1.22	0.93–1.60		<i>P</i> = 0.13
Mother's birthplace												
<i>P</i> < 0.01												
Taiwan												
Rural	1284	2042	3377	5658	9690	16,159	22,689	19,771	Ref.			
Satellite	665	1137	1954	3480	6474	11,420	16,332	12,999	1.04	1.02–1.06		
Urban	463	874	1415	2451	4554	8077	11,467	8611	1.00	0.98–1.02		<i>P</i> = 0.63
Otherwise												
Rural	270	361	422	575	816	1125	1405	1160	Ref.			
Satellite	120	185	208	272	399	568	750	601	1.09	1.01–1.18		
Urban	86	121	141	203	271	421	558	463	1.19	1.09–1.30		<i>P</i> < 0.01
Total	2888	4720	7517	12,639	22,204	37,770	53,201	43,605				

^a aOR: adjusted odds ratio; CI: confidence interval.^b Normal birthweight: ≥2500g, low birthweight: 1500–2499g, very low birthweight: <1500g.

children population.⁵ Another US study observed a significant increase in well-child visit adherence by 20%, which was accompanied by a 28% reduction in the uninsured rate after the execution of the Patient Protection and Affordable Care Act.²¹ Reasons that may account for the dissimilarity in findings between our study and those of others can be multifaceted, and insurance coverage may be one of the crucial issues. A lack of insurance has been associated with increased morbidity and mortality due to financial barriers to health care.^{22,23} The implementation of the NHI programme in Taiwan effectively removed such financial barriers and increase the affordability of using healthcare services.¹⁹

Taiwan launched its NHI programme, which covers nearly all Taiwanese residents, in 1995. Such universal health insurance coverage considerably benefits several disadvantaged populations, such as children, the elderly, non-working adults and low-income individuals, who may thereafter afford to receive health care at a reasonable cost.²⁴ Under the policy of Taiwan's NHI programme, seven PHC visits are available to all children whose parents are only required to pay a small amount of registration fee (approximately USD 5) for each visit. PHC services are provided by general practitioners or paediatrics specialists who can obtain subsidies from the Health Promotion Administration via NHI's reimbursement.^{2,25} A universal health insurance coverage may help parents with low socio-economic status eliminate financial barriers to seeking PHC services.

Apart from the affordability issue, improved accessibility through the NHI policy may be another reason that accounts for the slight urban–rural disparity in PHC utilisation by children. One of the health disparity-eliminating strategies of the NHI programme is to increase medical service supply in rural and remote areas.²⁶ The NHI programme implements various integrated healthcare programmes in rural Taiwan with high reimbursement, increasing physicians' incentive to practice in rural areas.²⁶ In addition, the local public health stations in each of the 316 city districts/townships around Taiwan also provides well-baby clinic services to

parents, all of which has effectively improved healthcare resources in rural Taiwan.³ Huang et al. analysed 5-year data since the initiation of Taiwan's NHI programme and reported that the rural–urban disparity in the rates of ruptured appendix apparently narrowed down; meanwhile, the rural–urban disparity in medical utilisation exhibited greater improvement amongst children than amongst adults.²⁶ To further explore the slight urban–rural difference in PHC services observed in our study, we analysed the data of two most socially deprived areas, i.e. aboriginal townships and offshore islands in Taiwan.^{27,28} Table 4 shows that 5.49% and 2.03% of preschool children from aboriginal townships and off-shore islands, respectively, did not make any visit, which were higher than the figure (1.38%) for the remaining children. Moreover, only 15.19% of children from aboriginal townships completed seven visits. Meanwhile, 91.72% had at least one visit. These values were considerably lower than the corresponding figures for children from off-shore islands and other areas. The preceding analysis suggested that some subgroups of children from rural Taiwan may still receive noticeably inadequate PHC services. The considerably low PHC utilisation observed in aboriginal townships may suggest factors other than accessibility that limit PHC utilisation in aboriginal townships. This issue warrants further investigations

Table 4

Preventive healthcare visit in aboriginal townships and off-shore islands as compared to the other areas.

Preventive healthcare visit	Geographic areas					
	Aboriginal townships		Off-shore islands		Others	
	N = 5795		N = 8378		N = 170,371	
	n	%	n	%	n	%
Complete 7 visits	880	15.19	2053	24.50	40,900	24.01
At least one visit	5315	91.72	8152	97.30	167,584	98.36
No visit	318	5.49	170	2.03	2350	1.38

because aboriginal townships and off-shore islands are considered areas with the most limited healthcare accessibility in Taiwan.

In addition, we noted that the frequencies of PHC for the 1st (1 month after birth) and 7th (3–5 years old) scheduled visits were relatively low in all the study groups. This observation could have originated from different scenarios. Low utilisation in LBW babies may be partly explained by the low overall utilisation of the 1st scheduled visit. Some LBW babies were struggling with physical prematurity and frequently receiving medical care during the 1st month after birth. Parents of LBW infants could have paid more attention to medical care than regular PHC for their babies. In addition, Taiwan's NHI subsidises the mental and physical assessments of VBLW babies in accordance with their corrected ages. However, we might have underestimated PHC utilisation for VLBW babies because the corrected age of VBLW babies was unavailable in the National Health Insurance Research Database (NHIRD).

Moreover, mothers and babies staying in postpartum care centres for 1 month-long postpartum recuperation is prevalent in Taiwan; some centres do not have contract paediatricians for PHC services. Therefore, we speculate that some babies may miss PHC for the 1st month after birth. The low utilisation of PHC for the seventh scheduled visit might have another reason. For the study birth cohort born in 2009, all PHC visits were matched to the schedules of various vaccinations, except for the 7th scheduled visit. Parents may be less motivated to utilise PHC if it is not aligned with a scheduled vaccination.²⁹ However, vaccinations for young children have been rescheduled in Taiwan since 2012. At present, children aged 3 years or older are scheduled to receive several vaccines before they attend elementary school at 7 years old. Thus, the utilisation rate of the 7th visit is expected to improve considerably. Policy input changes dynamically; hence, continuously monitoring PHC utilisation and regularly examining the entire system should be strongly considered by health policy makers.

Although only a slight rural–urban disparity in PHC utilisation is observed in Taiwan, considerable improvement can still be achieved, particularly in certain segments of children's population, such as LBW babies, babies with foreign-born mothers and babies from aboriginal townships. Cultural convictions may be a barrier to parents bringing their children to receive PHC.³⁰ A parent coach model has been reported to improve the receipt of well-child care in the US.³¹ Integrating culture into service provisions may be an effective means to strengthen parents' belief in receiving PHC.³⁰ For example, trained aboriginal nurses or those familiar with diverse ethnic cultures may help parents perceive threats from babies' disorders and the benefits of early detection. Elevating clinical professional's self-efficacy also eliminates the barrier of PHC.³⁰ Delivering a multifaceted intervention for practical skills to clinical practitioners increased the frequency and quality of Healthy Kids Check in Australia.³² We suggest that public health authorities should develop tailored information for target parents and provide useful materials about practical skills to clinical practitioners in paediatrics and obstetrics, enhancing the belief in and self-efficacy to utilise PHC of parents and clinicians.

A previous US study¹ reported that children with special needs had higher compliance rate (86.6%) to well-child visits recommended by the American Academy of Paediatrics compared with children without special needs (56.4%). Another research on pre-term baby's preventive care utilisation showed an increased OR of good compliance if children's birth weight were 1500–2500 g (1.51), 1000–1500 g (1.42) or under 1000 g (1.67).³³ In contrast with the aforementioned findings, our study presented the opposite

result, in which children with lower birth weight received less PHC services in Taiwan. One of the potential reasons may be related to the greater need of children with low birth weight to frequently seek curative treatments because of their generally poor health conditions. Physicians and parents may have overlooked PHC and paid more attention to treating various health disorders suffered by LBW children. In addition, a lack of norms and valid indicators for making evaluation of LBW children's development may impede physicians and parents from the optimal utilisation of PHC services.⁶ Given the higher prevalence of potential developmental problems among LBW children throughout their childhood years, the utilisation of scheduled PHC services should be emphasised in LBW infants, particularly amongst those with VLBW.³⁴

Mother's birthplace was another factor that was significantly associated with PHC utilisation. Children of foreign-born mothers received less PHC utilisation. The number of children of foreign-born mothers has been increasing in Taiwan. In 2017, 10% of children in elementary schools had foreign-born mothers.³⁵ Most foreign-born mothers were from Asian countries, particularly from China (64%) and Vietnam (19%).³⁶ The majority of foreign-born mothers married Taiwanese men with low socio-economic status and were reported to less likely receive proper prenatal care and health care.³⁷ As such, children of foreign-born mothers were likely to come from socio-economically disadvantaged families and their parents could have inadequate health literacy with regard to the importance of attending scheduled PHC services.

Apart from their roles as predictors of PHC utilisation, birth weight and maternal birthplace were found to modify the magnitude of urban–rural disparity in PHC services. For NBW children, minimal variation in PHC utilisation was observed across urbanisation levels. However, for LBW and VLBW children from urban areas, a significant increase of 17% and 22%, respectively, was noted in the OR of higher PHC utilisation. In addition, for children of foreign-born mothers in urban areas, a 9% and 19% significantly higher OR of higher PHC utilisation was observed amongst children from satellite and urban areas, respectively; but such urban–rural difference was less evident in children with Taiwan-born mothers. Certain non-financial factors may be responsible for effect modification by low birth weight and maternal birthplace. LBW children or those born to foreign mothers are believed to be associated with low socio-economic families. Rural areas are typically associated with inadequate public transportation supply, affecting more people with low socio-economic status.³⁸ In addition, rural healthcare workers usually face barriers to information access and distribution due to lack of funds, inadequate hardware and infrastructure problems.³⁹ Such limitation in disseminating health information to local people also likely affects low socio-economic families. People with high socio-economic status will find it easier to have a wide range of means of locating and accessing health information resources even if they are living in rural areas. Tailored strategies in improving the PHC visit of low socio-economic families by enhancing home visiting programmes may help reduce urban–rural disparity in PHC utilisation amongst children from low socio-economic families.⁴⁰

Our study exhibits several strengths. First, this study was the first to use a prospective design to investigate young children's utilisation of PHC. The information was based on the latest national cohort, which may provide up-to-date information of PHC utilisation by young children in Taiwan. Second, most previous studies have attributed the urban–rural difference in health care to economic barriers. The current study was based on Taiwan's NHI,

largely eliminating affordability reasons. We found a slight association of urbanisation level with children's PHC utilisation in a setting with universal health coverage. This finding suggests the importance of health insurance policy on the utilisation of PHC services. Thirdly, we identified certain specific subgroups that are more vulnerable to urban–rural disparity in PHC utilisation. The results of the stratified analyses highlighted the need for additional attention on these subgroups whilst designing PHC programmes.

Despite the aforementioned strengths, several limitations should be noted in this study due to the use of secondary administrative datasets. First, we considered the city district/township of mothers at the time of delivery to determine the urbanisation level of residence of children aged 0–7 years. This method can be subject to misclassification if the children are not living with their mothers or if they have moved to other areas after birth. However, we do not have sufficient information to clarify this issue. Second, other factors identified as associated with PHC utilisation, such as household income, language used by parents and the health literacy of caregivers, were not considered and analysed in this study.^{41,42} Incomplete consideration of these factors might have entailed confounding, at least to a certain extent, in our study.

Conclusion

Urbanisation level posed only a slight influence on the PHC utilisation of children under Taiwan's universal health coverage setting. However, the urban–rural disparity in PHC utilisation was more apparent in certain segments of children's population, including LBW children and children with foreign-born mothers. Policy makers should provide additional attention to improving PHC utilisation amongst LBW children and children of younger mothers, younger/older fathers and foreign-born mothers.

Author statements

Ethical approval

This research has been approved by National Cheng Kung University Governance Framework of Human Research Ethics [project number: 107–116].

Funding

This work was supported by Health Promotion Administration from Ministry of Health and Welfare in Taiwan [project number C1070310, 2018].

Competing interests

None declared.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.puhe.2020.02.011>.

References

- Selden TM. Compliance with well-child visit recommendations: evidence from the medical expenditure panel survey, 2000–2002. *Pediatrics* 2006;**118**: e1766–78. <https://doi.org/10.1542/peds.2006-0286>.
- Chen CS, Liu TC, Lin HC, Tian WH. The effect of Taiwan's National Health Insurance on infants' preventive care use and inpatient care use. *Health Pol* 2007;**80**: 432–43. <https://doi.org/10.1016/j.healthpol.2006.04.003>.
- Health Promotion Administration, Ministry of Health and Welfare, Taiwan. *Annual report of health promotion administration 2016 Part 2 Healthy birth and*

growth. 2016. https://health99.hpa.gov.tw/educZone/edu_detail.aspx?CatId=21937. [Accessed 22 July 2019].

- Tom JO, Tseng CW, Davis J, Solomon C, Zhou C, Mangione-Smith R. Missed well-child care visits, low continuity of care, and risk of ambulatory care-sensitive hospitalizations in young children. *Arch Pediatr Adolesc Med* 2010;**164**: 1052–8. <https://doi.org/10.1001/archpediatrics.2010.201>.
- Abdus S, Selden TM. Adherence with recommended well-child visits has grown, but large gaps persist among various socioeconomic groups. *Health Aff* 2013;**32**:508–15. <https://doi.org/10.1377/hlthaff.2012.0691>.
- Rossiter C, Fowler C, Hesson A, Kruske S, Homer CSE, Schmied V. Australian parents' use of universal child and family health services: a consumer survey. *Health Soc Care Community* 2019;**27**:472–82. <https://doi.org/10.1111/hsc.12667>.
- Farmer J, Iversen L, Campbell NC, Guest C, Chesson R, Deans G, et al. Rural/urban differences in accounts of patients' initial decisions to consult primary care. *Health Place* 2006;**12**:210–21. <https://doi.org/10.1016/j.healthplace.2004.11.007>.
- Ijarotimi IT, Fatiregun AA, Adebisi OA, Ilesanmi OS, Ajumobi O. Urban–rural differences in immunisation status and associated demographic factors among children 12–59 months in a south-western state, Nigeria. *PLoS One* 2018;**13**. <https://doi.org/10.1371/journal.pone.0206086>. e0206086.
- McClure K, Jun JK, Johnson CR, Fischer PR, Lu L, Vy S, et al. Disparities in rural-urban achievement of millennium development goals in Cambodia: implications for current and future child health. *Paediatr Int Child Health* 2018;**38**: 235–43. <https://doi.org/10.1080/20469047.2018.1515811>.
- Beatriz ED, Molnar BE, Griffith JL, Salhi C. Urban-rural disparity and urban population growth: a multilevel analysis of under-5 mortality in 30 sub-Saharan African countries. *Health Place* 2018;**52**:196–204. <https://doi.org/10.1016/j.healthplace.2018.06.006>.
- Puls HT, Bettenhausen JL, Markham JL, Walker JM, Drake B, Kyler KE, et al. Urban-rural residence and child physical abuse hospitalizations: a national incidence study. *J Pediatr* 2018;**205**:230–5. <https://doi.org/10.1016/j.jpeds.2018.09.071>.
- SanGiovanni CMJ, Morella K, Basco Jr W. Underdiagnosis and lower rates of office visits for overweight/obese pediatric patients in rural compared with urban areas. *South Med J* 2017;**110**:480–5. <https://doi.org/10.14423/SMJ.0000000000000672>.
- Uva JL, Wagner VL, Gesten FC. Emergency department reliance among rural children in Medicaid in New York state. *J Rural Health* 2011;**28**:152–61. <https://doi.org/10.1111/j.1748-0361.2011.00377.x>.
- Xiang Y, Zhao W, Xiang H, Smith GA. ED visits for drug-related poisoning in the United States, 2007. *Am J Emerg Med* 2012;**30**:293–301. <https://doi.org/10.1016/j.ajem.2010.11.031>.
- Coburn AF, McBride TD, Ziller EC. Patterns of health insurance coverage among rural and urban children. *Med Care Res Rev* 2002;**59**:272–92. <https://doi.org/10.1177/1077558702059003003>.
- National Health Insurance Administration Taiwan. *Statistics and surveys: the national health insurance statistics in taiwan*. 2009. https://www.nhi.gov.tw/english/Content_List.aspx?n=616897F8DF2C3614&topn=616897F8DF2C3614; 2009. [Accessed 22 July 2019].
- Chen HF, Li CY. Effect-modifications by age and sex on the risks of coronary artery disease and revascularization procedures in relation to diabetes. *Diabetes Res Clin Pract* 2007;**75**:88–95. <https://doi.org/10.1016/j.diabres.2006.05.020>.
- Liu CY, Hung YT, Chuang YL, Chen YJ, Weng WS, Liu JS. Incorporating development stratification of Taiwan townships into sampling design of large scale health interview survey. *J Health Manag* 2006;**4**:1–22. <https://www.scienceopen.com/document?vid=d9313108-0e01-4100-a9a1-60ee152d8266>.
- Ko MC, Lin WH, Martini S, Chang YH, Chiu CT, Li CY. A cohort study of age and sex specific risk of pyogenic liver abscess incidence in patients with type 2 diabetes mellitus. *Medicine* 2019;**98**. <https://doi.org/10.7326/0003-4819-148-4-200802190-00004>. e14849.
- Kim JH. Assessing practical significance of the proportional odds assumption. *Stat Probab Lett* 2003;**65**:233–9. <https://doi.org/10.1016/j.spl.2003.07.017>.
- Ortega AN, McKenna RM, Chen J, Alcalá HE, Langellier BA, Roby DH. Insurance coverage and well-child visits improved for youth under the Affordable Care Act, but latino youth still lag behind. *Acad Pediatr* 2018;**18**:35–42. <https://doi.org/10.1016/j.acap.2017.07.006>.
- Baker DW, Sudano JJ, Albert JM, Borawski EA, Dor A. Lack of health insurance and decline in overall health in late middle age. *N Engl J Med* 2001;**345**: 1106–12. <https://doi.org/10.1056/NEJMs002887>.
- Lau JS, Adams SH, Park MJ, Boscardin WJ, Irwin Jr CE. Improvement in preventive care of young adults after the affordable care act: the affordable care act is helping. *JAMA Pediatr* 2014;**168**:1101–6. <https://doi.org/10.1001/jamapediatrics.2014.1691>.
- Cheng TM. Taiwan's new national health insurance program: genesis and experience so far. *Health Aff* 2003;**22**:61–76. <https://doi.org/10.1377/hlthaff.22.3.61>.
- Chen I, Yang WS, Lee SD, Chang HC, Yeh CL. Utilization of well-baby care visits provided by Taiwan's National Health Insurance program. *Soc Sci Med* 2004;**59**: 1647–59. <https://doi.org/10.1016/j.socscimed.2004.02.008>.
- Huang N, Yip W, Chang HJ, Chou YJ. Trends in rural and urban differentials in incidence rates for ruptured appendicitis under the National Health Insurance

- in Taiwan. *Publ Health* 2006;**120**:1055–63. <https://doi.org/10.1016/j.puhe.2006.06.011>.
27. Leu HI, Chang WT, Lin MH, Chen TJ, Hwang SJ, Chou LF, et al. Urban-Rural Disparity in geographical and temporal availability of pediatric clinics: a nationwide survey in Taiwan. *Pediatr Neonatol* 2017;**58**:344–9.
 28. Chang YC, Lin JD, Tung HJ, Chiang PH, Hsu SW. Outpatient physical therapy utilization for children and adolescents with intellectual disabilities in Taiwan: a population-based nationwide study. *Res Dev Disabil* 2014;**35**:498–505.
 29. Garg P, Eastwood J, Liaw ST, Jalaludin B, Grace R. A case study of well child care visits at general practices in a region of disadvantage in Sydney. *PLoS One* 2018;**13**: e0205235.
 30. Kataoka-Yahiro MR, Munet-Vilaro F. Barriers to preventive health care for young children. *J Am Acad Nurse Pract* 2002;**14**:66–72.
 31. Coker TR, Chacon S, Elliott MN, Bruno Y, Chavis T, Biely C, et al. A parent coach model for well-child care among low-income children: a randomized controlled trial. *Pediatrics* 2016;**137**: e20153013.
 32. Alexander KE, Brijnath B, Mazza D. The challenges of trying to increase preventive healthcare for children in general practice: results of a feasibility study. *BMC Fam Pract* 2015;**16**:94.
 33. D'Agostino JA, Passarella M, Saynisch P, Martin AE, Macheras M, Lorch SA. Preterm infant attendance at health supervision visits. *Pediatrics* 2015;**136**: e794–802. <https://doi.org/10.1542/peds.2015-0745>.
 34. Stolt S, Lind A, Matomaki J, Haataja I, Lapinleimu H, Lehtonen L. Do the early development of gestures and receptive and expressive language predict language skills at 5;0 in prematurely born very-low-birth-weight children? *J Commun Disord* 2016;**61**:16–28. <https://doi.org/10.1016/j.jcomdis.2016.03.002>.
 35. Ministry of Education Taiwan. *Statistics of new immigrant children entering elementary school*. 2017 [Chinese], http://stats.moe.gov.tw/files/analysis/son_of_foreign_105.pdf; 2017. [Accessed 22 July 2019].
 36. Liao CC, Chan WY. Concerns over the life and health of alien brides from southeastern Asia and mainland China. *Tzu Chi Nursing Journal* 2005;**4**:12–6. <http://www.airitiilibrary.com/Publication/alDetailedMesh?docid=16831624-200503-4-1-12-16-a>.
 37. Huang CC, Li CY, Yang CH. Cultural implications of differing rates of medically indicated and elective cesarean deliveries for foreign-born versus native-born Taiwanese mothers. *Matern Child Health J* 2012;**16**:1008–14. <https://doi.org/10.1007/s10995-011-0824-7>.
 38. Ogunbodede EO, Kida IA, Madjapa HS, Amedari M, Ehizele A, Mutave R, et al. Oral health inequalities between rural and urban populations of the African and Middle East region. *Adv Dent Res* 2015;**27**:18–25. <https://doi.org/10.1177/0022034515575538>.
 39. Samuel S, Bayissa G, Asaminewu S, Alaro T. Electronic information sources access and use for healthcare services in governmental and non-governmental hospitals of western Oromia, Ethiopia: a cross sectional study. *Ethiop J Health Sci* 2016;**26**:341–50. <https://doi.org/10.4314/ejhs.v26i4.6>.
 40. Doi L, Jepson R, Hardie S. Realist evaluation of an enhanced health visiting programme. *PLoS One* 2017;**12**. <https://doi.org/10.1371/journal.pone.0180569>.
 41. Cohen AL, Christakis DA. Primary language of parent is associated with disparities in pediatric preventive care. *J Pediatr* 2006;**148**:254–8. <https://doi.org/10.1016/j.jpeds.2005.10.046>.
 42. Lee JY, Divaris K, DeWalt DA, Baker AD, Gizlice Z, Rozier RG, et al. Caregivers' Health literacy and gaps in children's medicaid enrollment: findings from the Carolina Oral Health Literacy Study. *PLoS One* 2014;**9**. <https://doi.org/10.1371/journal.pone.0110178>. e110178.

Urban-rural disparity of preventive healthcare utilisation among children under the universal health insurance coverage in Taiwan: a national birth cohort analysis

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

read.qxmd.com

Internet Source

2%

2

www.mdpi.com

Internet Source

2%

3

Wen-Ling Su, Chin-Li Lu, Santi Martini, Yuu-Hueih Hsu, Chung-Yi Li. "A population-based study on the prevalence of gestational diabetes mellitus in association with temperature in Taiwan", *Science of The Total Environment*, 2020

Publication

1%

4

bmcpublichealth.biomedcentral.com

Internet Source

1%

5

link.springer.com

Internet Source

1%

6

Hua-Fen Chen, Ching-An Ho, Chung-Yi Li. "Risk of heart failure in type 2 diabetes population:

1%

Comparison with non-diabetes subjects with and without coronary heart diseases", Diabetes, Obesity and Metabolism, 2018

Publication

7 Chin-Li Lu, Ya-Hui Chang, Yu Sun, Chung-Yi Li. "A population-based study of epilepsy incidence in association with type 2 diabetes and severe hypoglycaemia", Diabetes Research and Clinical Practice, 2018 1%

Publication

8 Chi-Chu Liu, Chin-Li Lu, Hari Basuki Notobroto, Chiang-Chin Tsai, Pei-Hung Wen, Chung-Yi Li. "Individual and neighborhood socioeconomic status in the prediction of liver transplantation among patients with liver disease", Medicine, 2019 1%

Publication

9 www.pubfacts.com <1%

Internet Source

10 boris.unibe.ch <1%

Internet Source

11 journals.plos.org <1%

Internet Source

12 www.med.unic.ac.cy <1%

Internet Source

13 global-qhr.org

Internet Source

<1%

14

research-repository.st-andrews.ac.uk

Internet Source

<1%

15

worldwidescience.org

Internet Source

<1%

16

escholarship.org

Internet Source

<1%

17

mafiadoc.com

Internet Source

<1%

18

Victoria H. Buelow. "Timely Immunization Series Completion among Children of Immigrants", *Journal of Immigrant and Minority Health*, 12/06/2007

Publication

<1%

19

pubmed.ncbi.nlm.nih.gov

Internet Source

<1%

20

hub.hku.hk

Internet Source

<1%

21

Lu, Chin-Li, Hsiu-Nien Shen, Hua-Fen Chen, and Chung-Yi Li. "Epidemiology of childhood Type 1 diabetes in Taiwan, 2003 to 2008", *Diabetic Medicine*, 2014.

Publication

<1%

www.degruyter.com

22

Internet Source

<1%

23

www.science.gov

Internet Source

<1%

24

hdl.handle.net

Internet Source

<1%

25

Chun-Ying Wu, Ming-Shiang Wu, Ken N. Kuo, Chang-Bi Wang, Yi-Ju Chen, Jaw-Town Lin. " Effective Reduction of Gastric Cancer Risk With Regular Use of Nonsteroidal Anti-Inflammatory Drugs in –Infected Patients ", Journal of Clinical Oncology, 2010

Publication

<1%

26

Chiang, Po-Huang, Yu-Chia Chang, Jin-Ding Lin, Ho-Jui Tung, Lan-Ping Lin, and Shang-Wei Hsu. "Healthcare utilization and expenditure analysis between individuals with intellectual disabilities and the general population in Taiwan: A population-based nationwide child and adolescent study", Research in Developmental Disabilities, 2013.

Publication

<1%

27

vermontforsinglepayer.org

Internet Source

<1%

28

www.dovepress.com

Internet Source

<1%

29	alzres.biomedcentral.com Internet Source	<1%
30	Joanne E. Croker, John L. Campbell. "Satisfaction with access to healthcare: qualitative study of rural patients and practitioners", Primary Health Care Research & Development, 2009 Publication	<1%
31	Julie L. Hudson, Salam Abdus. "Coverage And Care Consequences For Families In Which Children Have Mixed Eligibility For Public Insurance", Health Affairs, 2015 Publication	<1%
32	www.ncbi.nlm.nih.gov Internet Source	<1%
33	www.repository.cam.ac.uk Internet Source	<1%
34	lipidworld.biomedcentral.com Internet Source	<1%
35	www.cambridge.org Internet Source	<1%
36	www.jabfm.org Internet Source	<1%
37	Ning Lu. "Rural-Urban Differences in Health Insurance Coverage and Patterns Among	<1%

Working-Age Adults in Kentucky", The Journal of Rural Health, 03/2010

Publication

38

www.researchgate.net

Internet Source

<1%

39

Chang, J. S., C.-R. Tsai, Y.-W. Tsai, and J. L. Wiemels. "Medically diagnosed infections and risk of childhood leukaemia: a population-based case-control study", International Journal of Epidemiology, 2012.

Publication

<1%

40

Azeem Tariq Malik, Usman Younis Panni, Bassam A. Masri, Shahryar Noordin. "The impact of surgeon volume and hospital volume on postoperative mortality and morbidity after hip fractures: A systematic review", International Journal of Surgery, 2018

Publication

<1%

41

Jessica Y. Lee, Kimon Divaris, Darren A. DeWalt, A. Diane Baker, Ziya Gizlice, R. Gary Rozier, William F. Vann. "Caregivers' Health Literacy and Gaps in Children's Medicaid Enrollment: Findings from the Carolina Oral Health Literacy Study", PLoS ONE, 2014

Publication

<1%

42

Chris Rossiter, Cathrine Fowler, Amiee Hesson, Sue Kruske, Caroline S. E. Homer, Virginia

<1%

Schmied. "Australian parents' use of universal child and family health services: A consumer survey", Health & Social Care in the Community, 2018

Publication

43

Genevieve M. Kenney. "The Effects of Medicaid and CHIP Policy Changes on Receipt of Preventive Care among Children : Medicaid/CHIP Policy Changes and Receipt of Preventative Care", Health Services Research, 02/2011

Publication

44

I-Lin Hsu, Chia-Ming Chang, Deng-Chi Yang, Ya-Hui Chang, Chia-Chun Li, Susan C. Hu, Chung-Yi Li. "Socioeconomic Inequality in One-Year Mortality of Elderly People with Hip Fracture in Taiwan", International Journal of Environmental Research and Public Health, 2018

Publication

45

May Nawal Lutfiyya, Martin S. Lipsky, Jennifer Wisdom-Behounek, Melissa Inpanbutr-Martinkus. "Is Rural Residency a Risk Factor for Overweight and Obesity for U.S. Children?*", Obesity, 2007

Publication

46

Chen, H.S.. "Increased Risk of Urinary Tract

<1%

<1%

<1%

<1%

Calculi Among Patients With Diabetes Mellitus- A Population-based Cohort Study", Urology, 201201

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Urban-rural disparity of preventive healthcare utilisation among children under the universal health insurance coverage in Taiwan: a national birth cohort analysis

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
