

Barriers to diagnosis and management of CNS infections in Indonesia

by Darma Imran

Submission date: 09-Apr-2021 11:15AM (UTC+0800)

Submission ID: 1554230141

File name: s_to_diagnosis_and_management_of_CNS_infections_in_Indonesia.pdf (135.05K)

Word count: 2288

Character count: 13008

Barriers to diagnosis and management of CNS infections in Indonesia¹⁰

Darma Imran, MD, Sekar Satiti, MD, PhD, Paulus Sugianto, MD, PhD, Riwanti Estiasari, MD, PhD, Kartika Maharani, MD, David Pangeran, MD, Putri Widya Andini, MD, MSc, Badrul Munir, MD, Arthur H.P. Mawuntu, MD, Ni Made Susilawathi, MD, PhD, Kiking Ritarwan, MD, PhD, O.S. Hartanto, MD, PhD, Meity Frida, MD, Ahmad Rizal Ganiem, MD, PhD, Dede Gunawan, MD, Sofiati Dian, MD, MSc, A.A.R. Sudewi, MD, PhD, and Reinout van Crevel, MD, PhD

Correspondence

Dr. Imran
darma_imran@yahoo.com

Neurology[®] 2019;92:104-106. doi:10.1212/WNL.0000000000006732

Little is known about the management and outcome of CNS infections in Indonesia (population: approximately 261 million), and the burden of CNS infections is increasing as result of a rapidly growing HIV epidemic. Indeed, in a recent study in Jakarta,¹ 54% of patients with a CNS infection were HIV-infected, vs 25% in a similar study conducted in Bandung, the second largest city on Java.² Establishing the etiology of CNS infections is important for clinical management and outcome, but anecdotal evidence suggests that there are major gaps in clinical management, including a failure to recognize CNS infections, perform appropriate diagnostics, or initiate necessary drug treatment. In this study, we address some of these barriers by interviewing a group of Indonesian neurologists.

On behalf of the Indonesian Neurologic Association, neurologists who attend the National Neurology Conference (July 27–30, 2017, Yogyakarta) were given questionnaires to complete with respect to the frequency and type of CNS infections and possible barriers in diagnosis and treatment. The questionnaire was developed and piloted in Cipto Mangunkusumo Hospital, Jakarta, and adjusted with feedback from the neuroinfection working group.

The anonymous paper questionnaire (available as supplementary material, links.lww.com/WNL/A775) was completed by 288 neurologists (median age 41 years, 56% female) from tertiary (19%), regional (48%), or district (30%) hospitals in Java/Bali (62%), Sumatra (22%), Sulawesi (7%), Kalimantan (6%), and the eastern islands including Papua (3.2%). Most (67%) had worked fewer than 10 years since their neurology training. Almost all neurologists (93%) had seen presumptive CNS infections in the last 3 months; however, only 15% had performed lumbar puncture (LP) during this same period. More recently qualified neurologists and those working in regional hospitals reported a higher number of CNS infections and LPs.¹ Neurologists' considerations not to perform LP included "unfavorable condition of the patient," logistical issues like lack of LP kits, and concerns regarding possible blame from the family if complications would occur (table). Fear of complications, such as pain, paralysis, and death, was also reported as a reason for patient reluctance to undergo LP. Most neurologists mentioned shock (90%), raised intracranial pressure (89%), brain herniation (87%), and papilledema (86%) as contraindications for LP, but 35% also mentioned HIV infection as contraindication for LP. When asked what blood examinations they would like to order in approaching suspected CNS infection cases, only 15 (5%) spontaneously mentioned HIV testing.

With regard to specific diagnosis and treatment, serious barriers were identified. The majority of neurologists reported working in a hospital where CT scans (81%) and routine CSF examination (73%) are available. However, only a small number mentioned having access to specific microbiological testing for tuberculous meningitis, cryptococcal meningitis, and viral

From the Faculty of Medicine (D.I., R.E., K.M., D.P., P.W.A.), Universitas Indonesia, Jakarta; Faculty of Medicine (S.S.), Universitas Gadjah Mada, Yogyakarta; Faculty of Medicine (P.S.), Universitas Airlangga, Surabaya; Faculty of Medicine (B.M.), Universitas Brawijaya, Malang; Faculty of Medicine (A.H.P.M.), Universitas Sam Ratulangi, Manado; Faculty of Medicine (N.M.S., A.A.R.S.), Universitas Udayana, Denpasar; Faculty of Medicine (K.R.), Universitas Sumatera Utara, Medan; Faculty of Medicine (O.S.H.), Universitas Negeri Sebelas Maret, Solo; Faculty of Medicine (M.F.), Universitas Andalas, Padang; Faculty of Medicine (A.R.G., D.G., S.D.), Universitas Padjadjaran, Bandung, Indonesia; Department of Internal Medicine and Radboud Center for Infectious Diseases (R.v.C.), Radboud University Medical Center, Nijmegen, the Netherlands; and Centre for Tropical Medicine and Global Health (R.v.C.), Nuffield Department of Medicine, University of Oxford, UK.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Table Experience of Indonesian neurologists (n = 288) regarding diagnosis and management of CNS infection

	Percent
Patients with suspected CNS infections in the last 3 months	
0	7
1-5	60
6-10	20
>10	13
LP performed in the last 3 months	
None	40
1-3	41
>3	19
Reasons for not performing LP	
Patient reluctance for LP	
Fear of complication	78
Financial constraints	15
Doctors' concerns regarding LP	
Unfavorable condition of the patient	64
No supporting facilities (LP kit)	29
LP infrequently performed	7
Fear of being blamed by the family in case of complications	25
Perceived contraindications for LP	
Increased intracranial pressure	89
HIV infection	35
Intracranial mass	82
Availability of CT scan/MRI	81
Availability of CSF laboratory tests	
Routine CSF analysis (protein, leukocytes, glucose)	
Tuberculosis diagnostics	71
Microscopy	56
GeneXpert MTB/RIF	26
Solid culture	34
Microscopy (India ink) for <i>Cryptococcus</i>	39
PCR for HSV, VZV, or CMV	20
Availability of specific medication for CNS infections	
Tuberculous meningitis (FDC tuberculosis drugs)	90
Cerebral toxoplasmosis	
Pyrimethamine	61
Clindamycin	80

Table Experience of Indonesian neurologists (n = 288) regarding diagnosis and management of CNS infection (continued)

	Percent
Cryptococcal meningitis	
Fluconazole	67
Amphotericin B	43
Viral encephalitis	
Acyclovir	91
Valacyclovir	43

Abbreviations: CMV = cytomegalovirus; FDC = fixed-dose combination; HSV = herpes simplex virus; LP = lumbar puncture; VZV = varicella-zoster virus. Paper-based questionnaires were completed during the annual Indonesian Neurology meeting (July 27-30, 2017, Yogyakarta); 98%-100% of questions included in the table were complete.

encephalitis (table). Similarly, many reported a lack of drug treatment for cerebral toxoplasmosis or cryptococcal meningitis (table). In daily practice, the situation might be even more difficult, as supply of certain tests (e.g., Xpert MTB/RIF cartridges) or medication (e.g., amphotericin B) is often interrupted, or costs are not covered by patients' insurance.

Based on our findings, several conclusions can be drawn. First, it seems that neurologists in Indonesia frequently encounter patients with suspected CNS infections, but often fail to do LP. A previous report from Uganda described how educational material helped decrease patient refusal to undergo LP.³ Our data suggest that it may be at least as important to improve doctors' knowledge and skills. Experienced groups with an interest in CNS infections in Bandung and Jakarta who have reported high rates of large-volume LPs, with <5% refusal rate, might take the lead in educating other neurologists in Indonesia.^{1,2} Second, many neurologists mentioned having no access to appropriate diagnostics and specific microbiological testing, something that could be defined as a health systems constraint. WHO has defined 6 building blocks of a health system (service delivery, health workforce, information, medical products and technologies, financing, and leadership), which were used recently for constraints analysis for neurologic disease in Timor-Leste, a low-income country neighboring Indonesia.⁴ One could apply a similar approach specifically for CNS infections. Our third conclusion is that HIV testing does not seem standard for patients with suspected CNS infection, even though a patient's HIV status is relevant in terms of the etiology and management of CNS infections, and that many neurologists consider HIV infection as a contraindication for doing LP. This is worrying, as in a recent cohort study in Jakarta we found that 54% of adult patients with CNS infections were HIV-infected, with half of this group diagnosed with HIV infection previously, but only a minority receiving antiretroviral therapy, and with a median CD4 cell count of 30 μ L both among newly and previously

13 diagnosed patients.¹ These findings and those of the present study underscore the urgent need to address the many barriers to providing good care for patients with suspected or proven CNS infections in Indonesia, as in many low- or middle-income countries. We have reported our findings to the Indonesian Ministry of Health and National Neurology Association. As a next step, based on our findings, we intend to perform a more detailed qualitative and quantitative assessment to prioritize and address amendable gaps in care for patients with CNS infections, as was previously done for management of child contacts of tuberculosis patients in Indonesia.⁹

Author contributions

9 Darma Imran: study concept and design, acquisition of data, analysis and interpretation, critical revision of the manuscript for important intellectual content, study supervision. Sekar Satiti: study concept and design, acquisition of data. Paulus Sugianto: study concept and design, acquisition of data. Riwanti Estiasari: study concept and design, critical revision of the manuscript for important intellectual content, study supervision. Kartika Maharani: analysis and interpretation, critical revision of the manuscript for important intellectual content. David Pangeran: acquisition of data, analysis and interpretation, critical revision of the manuscript for important intellectual content. Putri Widya Andini: analysis and interpretation, critical revision of the manuscript for important intellectual content. Badrul Munir: study concept and design, acquisition of data. Arthur H.P. Mawuntu: study concept and design, acquisition of data. Ni Made Susilawati: study concept and design, acquisition of data. Kiking Ritarwan: study concept and design, acquisition of data. O.S. Hartanto: study concept and design, acquisition of data. Meity Frida: study concept and design, acquisition of data. Ahmad

11 Rizal Ganiem: study concept and design, acquisition of data, critical revision of the manuscript for important intellectual content, study supervision. Dede Gunawan: study concept and design. Sofiaty Dian: study concept and design, acquisition of data. A.A.R. Sudewi: study concept and design, acquisition of data, study supervision. Reinout van Crevel: study concept and design, analysis and interpretation, critical revision of the manuscript for important intellectual content, study supervision.

Study funding

The study was supported by the Indonesian Neurological Association.

Disclosure

13 D. Imran received a grant from the Indonesian Neurological Association. S. Satiti, P. Sugianto, R. Estiasari, K. Maharani, D., P. Widya Andini, B. Munir, A. Mawuntu, N. Susilawati, K. Ritarwan, O. Hartanto, M. Frida, A. Rizal Ganiem, D. Gunawan, S. Dian, A. Sudewi, and R. van Crevel report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

1. Imran D, Estiasari R, Maharani K, et al. Presentation, etiology and outcome of brain infections in an Indonesian hospital: a cohort study. *Neurol Clin Pract Epub* 2018 Sep 13.
2. Ganiem AR, Parwati I, Wisaksana R, et al. The effect of HIV infection on adult meningitis in Indonesia: a prospective cohort study. *AIDS* 2009;23:2309–2316.
3. Thakur KT, Mateyo K, Hachambwa L, et al. Lumbar puncture refusal in sub-Saharan Africa: a call for further understanding and intervention. *Neurology* 2015;84:1988–1990.
4. Mateen FJ, Martins N. A health systems constraints analysis for neurologic diseases: the example of Timor-Leste. *Neurology* 2014;82:1274–1276.
5. Rutherford ME, Ruslami R, Anselmo M, et al. Management of children exposed to *Mycobacterium tuberculosis*: a public health evaluation in West Java, Indonesia. *Bull World Health Organ* 2013;91:932–941A.

2 Disputes & Debates: Rapid online correspondence

The editors encourage comments on recent articles through Disputes & Debates:

Access an article at Neurology.org/N and click on “COMMENT” beneath the article header. Responses will be posted within 3 business days.

Before submitting a comment to Disputes & Debates, remember the following:

- Disputes & Debates is restricted to comments about studies published in *Neurology* within the last eight weeks
- Read previously posted comments; redundant comments will not be posted
- Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
- You can include a maximum of five authors (including yourself)

Neurology[®]

5 Barriers to diagnosis and management of CNS infections in Indonesia

Darma Imran, Sekar Satiti, Paulus Sugianto, et al.

Neurology 2019;92:104-106

DOI 10.1212/WNL.0000000000006732

This information is current as of January 7, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/92/2/104.full
References	6 This article cites 4 articles, 2 of which you can access for free at: http://n.neurology.org/content/92/2/104.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/92/2/104.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All global neurology http://n.neurology.org/cgi/collection/all_global_neurology Encephalitis http://n.neurology.org/cgi/collection/encephalitis HIV http://n.neurology.org/cgi/collection/hiv Meningitis http://n.neurology.org/cgi/collection/meningitis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology [®] is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.



Barriers to diagnosis and management of CNS infections in Indonesia

ORIGINALITY REPORT

20%

SIMILARITY INDEX

17%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Luuk Wieske, Ulrich Beuers, Matthijs C. Brouwer. "Clinical Reasoning: Accumulating endocrinopathies in a car salesman", *Neurology*, 2018
Publication 2%
- 2 Monica E. Lemmon, Charlene Gamaldo, Rachel Marie E. Salas, Ankita Saxena, Tiana E. Cruz, Renee D. Boss, Roy E. Strowd. "Education Research: Difficult conversations in neurology", *Neurology*, 2018
Publication 2%
- 3 Kate M. Daniello, Daniel J. Weber. "Education Research: The current state of neurophysiology education in selected neurology residency programs", *Neurology*, 2018
Publication 2%
- 4 bmcneurol.biomedcentral.com
Internet Source 2%
- 5 www.neurology.org
Internet Source 2%

2%

6

cebp.aacrjournals.org

Internet Source

1%

7

Bhaskar Roy, Nizar Chahin. "Clinical Reasoning: A 14-year-old boy with fatigue and episodic worsening of weakness", *Neurology*, 2017

Publication

1%

8

nn.neurology.org

Internet Source

1%

9

Rehana Z. Hussain, William A. Miller-Little, Richard Doelger, Gary R. Cutter, Nicolas Loof, Petra D. Cravens, Olaf Stüve. "Defining standard enzymatic dissociation methods for individual brains and spinal cords in EAE", *Neurology - Neuroimmunology Neuroinflammation*, 2018

Publication

1%

10

cp.neurology.org

Internet Source

1%

11

www.nature.com

Internet Source

1%

12

www.tandfonline.com

Internet Source

1%

13

F. Cresswell, C. Lange, R. van Crevel.

"Improving the diagnosis of tuberculous meningitis: good, but not good enough", *Clinical Microbiology and Infection*, 2020

Publication

1%

14

blogs.neurology.org

Internet Source

1%

15

www.ncbi.nlm.nih.gov

Internet Source

1%

16

Hitoshi Shinotoh, Hitoshi Shimada, Yasumasa Kokubo, Kenji Tagai et al. "Tau imaging detects distinctive distribution of tau pathology in ALS/PDC on the Kii Peninsula", *Neurology*, 2018

Publication

<1%

17

C. Chabala, G. Chongwe, E. Jumbe-Marsden, S. W. Somwe. "Missed opportunities for screening child contacts of smear-positive tuberculosis in Zambia, a high-prevalence setting", *The International Journal of Tuberculosis and Lung Disease*, 2017

Publication

<1%

18

Meng-sha Yao, Sheng Chen, Jun Liu. "Mystery Case: Cowden syndrome presenting with paraneoplastic encephalitis", *Neurology*, 2019

Publication

<1%

19

epub.ub.uni-muenchen.de

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Barriers to diagnosis and management of CNS infections in Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
