

Comparison between baby led weaning and traditional spoon-feeding on iron status and growth in breastfed infants

by Meta Herdiana Hanindita

Submission date: 16-Apr-2021 02:35PM (UTC+0800)

Submission ID: 1560743063

File name: spoon-feeding_on_iron_status_and_growth_in_breastfed_infants.pdf (177.23K)

Word count: 2197

Character count: 11077



COMPARISON BETWEEN BABY LED WEANING AND TRADITIONAL SPOON-FEEDING ON IRON STATUS AND GROWTH IN BREASTFED INFANTS

Meta Herdiana Hanindita^{1*}, Nur Aisyah Widjaja¹, Roedi Irawan¹, Boerhan Hidayat¹

¹Department of Child Health, Dr. Soetomo Hospital Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

*hanindita.meta@gmail.com

<https://doi.org/10.34302/crpjfst/2019.11.5.14>

Article history:

Received:

9 March 2019

Accepted:

20 September 2019

Keywords:

Complementary feeding;

Baby-led weaning;

Faltering growth;

Iron deficiency anemia

ABSTRACT

Background: The current guideline recommends infants are initially offered smoothly pureed foods, known as the traditional method of spoon feeding. Currently, an alternative method known as 'baby-led weaning' (BLW) has been really popular. With BLW, infants are allowed to self-feed family foods in their whole form instead special-prepared foods. Infants following BLW may be at increased risk of faltering growth.

Objective: To compare traditional and BLW methods in the risk of iron deficiency anemia (IDA) and growth faltering of breastfed infants.

Methods: A cross-sectional study was undertaken in 9 – 15 months breastfed infants admitted to Bedah Hospital Surabaya from August – October 2017. Exclusion criteria were infants suffering from chronic conditions. Biochemical assessment of Haemoglobin (Hb), Serum Transferrin (ST) and Serum Ferritin (SF) was measured. Diagnosis of IDA and growth chart interpretation was made based on WHO criteria.

Results: Out of all, 12/30 boys and 15/30 following BLW. Mean age was 12.6±2.14 months old. Mean Hb level, ST level and SF in BLW group was 10.9±0.55 g/dl, 11.6±7.13%, 19.1±18.40ug/dl prospectively, while in the traditional group were 12.5±0.75 g/dl, 24.6±7.92%, 57.6±18.78 ug/dl prospectively. IDA was higher in BLW group than in the traditional group. (13/15 vs 3/15, OR 26.000 95%CI 3.686-183.418, p <0.001). Underweight was higher in BLW. (13/15 vs 3/15, OR 26.000 95%CI 3.686-183.418, p <0.001). Stunted were higher in BLW (2/15 vs none, OR 1.667 95%CI 1.103-2.519, p=0.017).

Conclusion: In breastfed infants, those who were following BLW are in higher risk of IDA, underweight and stunted than traditional spoonfeeding.

1. Introduction

The current guideline of the World Health Organization recommends that infants are initially offered smoothly pureed foods, and gradually increased food consistency and variety as the infant gets older, adapting to the infant's requirements and abilities. The guideline also recommends practicing responsive

feeding, specifically feed infants directly and assist older children when they feed themselves. This is known as the traditional method of spoon-feeding (WHO, 2001). Over the last 10 years, an alternative method known as 'baby-led weaning' (BLW) has been really popular in Indonesia.

Baby-led weaning is a method for introducing complementary foods in which

the infant feeds themselves hand-held foods or finger foods instead of being spoon-fed by an adult. Infants following BLW share family food and are offered breast milk on demand until they wean. (Cameron *et al.*, 2012; Brown *et al.*, 2017)

Baby-led weaning was introduced with the publication of the book of *Baby-Led Weaning* by Gill Rapley and Tracey Murkett. It has been suggested that BLW may encourage greater acceptance of foods with a variety of textures and flavors and that this may result in higher intakes of “healthier” foods such as vegetables and unprocessed foods as the child grows, so it could lower the risk of obesity. In BLW, infants share family food and meal times. The food is offered as “graspable” pieces that they can pick up and they learn to feed themselves. The infant is in control of exactly what and how much they eat, They never hurried or forced to eat food. Combined method of BLW and traditional spoon-feeding is not defined as BLW (Rapley, 2008).

Weaning period is a peak period for faltering in a child’s growth. (Abeshu *et al.*, 2016) Infants following BLW may be at increased risk of growth faltering, based on the assumption that not all infants will have the motor skills to feed themselves the amount of food they require, and that many of the first foods offered will be low in energy and iron.

2. Materials and methods

2.1. Materials

All breastfed infants between 9 months old to 15 months old were prospectively enrolled. Exclusion criteria included chronic conditions such as cerebral palsy, congenital deformities and chromosomal disorders. Infants with urinary tract infection, hypothyroid, and tuberculosis were also excluded.

2.2. Methods

This study was approved by the institutional review board of the Bedah Hospital Surabaya. The present study was carried out in the Outpatient Clinic Bedah

Hospital Surabaya, during the period from August 2017 until October 2017. Based on the anamnesis, we classified all the subjects into 2 groups based on their feeding methods, the BLW group and the traditional group. Assessment of Haemoglobin (Hb), Serum Transferrin (ST) and Serum Ferritin (SF) was measured, as well body weight and body length. Diagnosis of IDA and growth anthropometry was interpreted based on WHO criteria. SPSS for Mac was used for analyses. To compare continuous variables, we used a two-sample *t* test or Mann-Whitney rank sum test, and to compare categorical variables we used the Chi-square or Fisher’s exact test, as indicated.

3. Results and discussions

There were 30 infants included in the study. Of these, 12 infants were boys. All the infants started the complementary feeding at 6 months old. There was 15 infants who were following BLW, and there was 15 infants who were following traditional spoon-feeding method. Mean age was 12.6±2.14 months old. There was no significant difference between two groups with respect to age or sex (see Table 1). There was significant difference between two groups on Hb level, Serum Transferrin and Serum Ferritin ($P < 0.001$). Iron Deficiency Anemia was higher in BLW group than in the traditional group. (13/15 vs 3/15, OR 26.00 95%CI 3.686-183.418, $p < 0.001$) (see Table 2).

3.1. Comparison on Iron Status

Iron Deficiency Anemia was higher in BLW group than in the traditional group. (13/15 vs 3/15, OR 26.00 95%CI 3.686-183.418, $p < 0.001$).

3.2. Comparison on Growth Status

Underweight was higher in BLW group than in the traditional group. (13/15 vs 3/15, OR 26.00 95%CI 3.686-183.418, $p < 0.001$). Severe underweight was higher in BLW group than in the traditional group. (2/15 vs 0/15). Stunted were higher in BLW group (2/15 vs none, OR 1.67 95%CI 1.103-2.519, $p = 0.017$) than in the traditional group, meanwhile wasted

was higher in BLW group then in the traditional group (11/15 vs 2/15).

To date, there are limited researches about BLW. Our study showed there was no difference noted with both groups with respect to gender and age. This was in line with some previous study (Morison et al., 2016; Cameron et al., 2013).

A significant difference was noted in iron status between the two groups ($P < 0.001$). There was no consideration about iron status before study. Until now, there were no studies examining iron status in infants following BLW. But, there was a study revealed that infants following BLW appeared to have significantly lower intakes of iron, zinc and vitamin B12 (Morison et al., 2016). This was consistent with another study revealed that infants following BLW ate more fruits and vegetables which were low-iron foods while the infants following traditional spoon-feeding eat more baby rice cereal (Cameron et al., 2013).

Food presentation significantly influences food preferences (Blossfeld et al., 2007), so it is possible that differences in the presentation of foods between the two groups impacted on preferences. Meat may be easier to eat when pureed and spoon-fed (Morison et al., 2016). This can be explained by the fact that infants' early chewing is a primitive up-down munching pattern (Morris et al., 2000; Rudolph et al., 2002). For small pieces of soft foods, this pattern may be adequate. But for hard foods such as meat or strips, it is inadequate (Morris et al., 2000). BLW may not be suitable for age and infant's feeding skill development.

Our study noted a significant difference in growth status between both groups. Infants following BLW were at increased risk to have underweight, stunted and wasted. Consistent with this, some previous studies also showed underweight was higher in infants following BLW compared to those who were following traditional spoon-feeding (Townsend et al., 2012; Brown et al., 2015). According to one study in the United Kingdom which compared body weight for age in infants following BLW and infants following traditional spoon-feeding. For the BLW group, 86.5% were of normal weight, 8.1% overweight and 5.4% underweight. In comparison, 78.3% of those in the traditional spoon-feeding group were normal weight, 19.2% overweight and 2.5% underweight (Brown et al., 2015).

Mothers following BLW method estimated that their babies ate more milk feeds and less solid food compare to those following a traditional spoon-feeding. This may provide inadequate nutrient intake such as energy and iron requirements for infants from 6 months of age onwards (Brown et al., 2015).

Since in BLW approach, the infants are in full control to eat, there are some risks of inadequate intake. When the infants do not have interest in eating, when foods of inappropriate consistency are offered, and infants with poor self-feeding skills will not be able to have enough foods to meet their needs for rapid growth. This risk of inadequate energy and nutrient intake could lead to failure to thrive.

Table 1. Characteristic of samples

Variable	BLW (n=15)	Traditional (n=15)	P
Age (mo) 12.6±2.14	13±1.78	12.2±2.30	0.396
Sex			0.143
Boys	12	18	
Girl	18	12	

Iron Status			
Hb (g/dl)	10.9±0.55	12.5±0.75	<0.001*
ST (%)	11.6±7.13	24.6±7.92	<0.001*
SF (ng/ml)	19.1±18.40	57.6±18.78	<0.001*

Table 2. Logistic Regression Analysis

	Odds Ratio	P	95% CI
Iron Deficiency Anemia	26.00	<0.001	3.686-183.418
Underweight	26.00	<0.001	3.686-183.418
Stunted	1.67	0.017	1.103 -2.519

* Chi-square test

Table 3. Comparison of growth parameter between two groups

Growth Parameter	BLW	Traditional
Weight for age		
Normal	2	12
Underweight	11	3
Severe underweight	2	0
Length for age		
Normal	9	15
Stunted	6	0
Weight for length		
Normal	4	13
Wasted	11	2

4. Conclusions

Those who were following BLW are in higher risk of IDA, underweight and stunted than those who were fed traditionally. There are still some controversies of BLW that need further research.

5. References

- Abeshu MA., Lelisa A., Geleta B. (2016) Complementary feeding: Review of recommendations, feeding practices, and adequacy of homemade complementary food preparations in developing countries. Lessons from Ethiopia. *Frontier Nutrition*, 17, 3,41.
- Blossfeld I., Collins A., Kiely M., Delahunty C. (2007) Texture preferences of 12-month-old infants and the role of early experiences. *Food Quality and Preference*, 18(2), 396-404.
- Brown A., Jones SW., Rowan H. (2017) Baby-led weaning: the evidence to date. *Current Nutrition Reports*, 6(2), 148-56.
- Brown A., Lee MD. (2015) Early influences on child-satiety-responsiveness: the role of weaning style, *Pediatric Obesity*, 10(1), 57-66.
- Cameron S., Taylor R., Heath AL. Parent-led or baby-led? Associations between complementary feeding practices and health-related behaviours in a survey of New Zealand families. *BMJ Open*, 3, 1- 9.
- Cameron SL., Heath AL., Taylor RW. (2012) How feasible is baby-led weaning as an approach to infant feeding? A review of the evidence. *Nutrients*, 4, 1575-609.
- Morison BJ., Taylor RW., Haszard JJ., Schramm CJ., Erickson LW., Fangupo LJ., Fleming EA., Luciano A., Heath AM. (2016) How different are baby-led weaning and conventional complementary feeding? A cross-sectional study of infants aged 6-8 months. *BMJ Open*, 6(5), 1-11.
- Morris E., Klein D. (2000) Pre-feeding skills, 2nd ed. USA: *Therapy skills builders*.
- Rapley G., Murkett T. (2008) *Baby-led weaning: Helping your baby to love good food.* (1st ed). London: Vermillion.
- Rudolph C., Link T. (2002) Feeding disorders in infants and children. *Pediatric Clinics of North America*, 49, 97-112.
- Townsend E., Pitchford NJ. (2012) Baby knows

best? The impact of weaning style on food preferences and body mass index in early childhood in a case- controlled sample. *BMJ Open*, 2(1), 1-6.

World Health Organization. (2001) *Guiding Principles for Complementary Feeding of the Breastfed Child*. Geneva: WHO Press.

Comparison between baby led weaning and traditional spoon-feeding on iron status and growth in breastfed infants

ORIGINALITY REPORT

18%

SIMILARITY INDEX

8%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Erkan Dogan, Gonca Yilmaz, Nilgun Caylan, Mahmut Turgut, Gulbin Gokcay, Melahat Melek Oguz. "Baby-led complementary feeding: Randomized controlled study", *Pediatrics International*, 2018
Publication 2%
- 2 J. A. Y. Cichero. "Introducing solid foods using baby-led weaning vs. spoon-feeding: A focus on oral development, nutrient intake and quality of research to bring balance to the debate", *Nutrition Bulletin*, 2016
Publication 2%
- 3 Ellen Townsend, Nicola J Pitchford. "Baby knows best? The impact of weaning style on food preferences and body mass index in early childhood in a case-controlled sample", *BMJ Open*, 2012
Publication 2%
- 4 Amy Brown, Sara Wyn Jones, Hannah Rowan. "Baby-Led Weaning: The Evidence to Date", 2%

Current Nutrition Reports, 2017

Publication

5	"Handbook of Food Fortification and Health", Springer Science and Business Media LLC, 2013 Publication	1 %
6	Simon Langley-Evans. "Nutrition, health and disease", Wiley, 2015 Publication	1 %
7	bmjopen.bmj.com Internet Source	1 %
8	www.sjkdt.org Internet Source	1 %
9	journals.lww.com Internet Source	1 %
10	trialsjournal.biomedcentral.com Internet Source	1 %
11	www.mummyinprovence.com Internet Source	1 %
12	mafiadoc.com Internet Source	1 %
13	pediatricianuae.com Internet Source	1 %
14	E. Ross. "Eating Development in Young Children: Understanding the Complex	<1 %

Interplay of Developmental Domains", Elsevier BV, 2017

Publication

15

Chantal L. Verhage, Marleen Gillebaart,
Shelley M.C. van der VEEK, Carolus M.J.L.
Vereijken. "The relation between family meals
and health of infants and toddlers: A review",
Appetite, 2018

Publication

<1 %

16

Lisa Daniels, Rachael W Taylor, Sheila M
Williams, Rosalind S Gibson et al. "Impact of a
modified version of baby-led weaning on iron
intake and status: a randomised controlled
trial", BMJ Open, 2018

Publication

<1 %

17

etheses.dur.ac.uk

Internet Source

<1 %

18

kclpure.kcl.ac.uk

Internet Source

<1 %

19

livrepository.liverpool.ac.uk

Internet Source

<1 %

20

pdfs.semanticscholar.org

Internet Source

<1 %

21

www.frontiersin.org

Internet Source

<1 %

22

www.ideals.illinois.edu

Internet Source

<1 %

23

Sonya L Cameron, Rachael W Taylor, Anne-Louise M Heath. "Parent-led or baby-led? Associations between complementary feeding practices and health-related behaviours in a survey of New Zealand families", BMJ Open, 2013

Publication

<1 %

24

Brittany J Morison, Rachael W Taylor, Jillian J Haszard, Claire J Schramm et al. "How different are baby-led weaning and conventional complementary feeding? A cross-sectional study of infants aged 6–8 months", BMJ Open, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Comparison between baby led weaning and traditional spoon-feeding on iron status and growth in breastfed infants

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
