

Cardiovascular & Cardiometabolic Journal

e-journal.unair.ac.id/CCJ



Editor-in-Chief

Yudi Her Oktaviono, MD, PhD

Deputy Editors

Dyana Sarvasti, MD, PhD
J. Nugroho E.P, MD, PhD

Original Research

- The Correlation between Serum TG/HDL-c ratio and Arterial Stiffness Using the Cardio-ankle Vascular Index in Overweight or Obese Patients

Case Report

- Long Term Survival of Ventricular Septal Rupture (VSR) Closure Concomitant with CABG in Post-Acute Myocardial Infarction Patient



Vol. 1 No. 2 (2020): Cardiovascular and Cardiometabolic Journal

Current Issue



Vol. 1 No. 2 (2020): Cardiovascular and Cardiometabolic Journal

Published: 2020-09-19

Original Research

The Correlation between Serum TG/HDL-c Ratio and Arterial Stiffness Using The Cardio-ankle Vascular Index in Overweight or Obese Patients

Andrianus Oktovianto ⁽¹⁾, Ni Putu Anggun Laksmi ⁽²⁾, Johannes Nugroho Eko Putranto ⁽³⁾, Raden Mohammad Yogiarto ⁽⁴⁾
(1) Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia ,

(3) Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, RSUD Dr. Soetomo, Surabaya, Indonesia ,

(4) Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya

31-37

Abstract : 491

PDF : 43



DOI : 10.20473/ccj.v1i2.2020.31-37

Moderate Cardiovascular Risk Factor among Indonesian: Do Carotid Intima-media Thickness (CIMT) Predict Further?

Rina Mawarti ⁽¹⁾, Denny Suwanto ⁽²⁾, Johannes Nugroho Eko Putranto ⁽³⁾, Djoko Soemantri ⁽⁴⁾
(1) Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia ,

(2) Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia ,


(3) Department of Cardiology and Vascular Medicine, Dr. Soetomo General Hospital, Surabaya, Indonesia ,

(4) Department of Cardiology and Vascular Medicine, Dr. Soetomo General Hospital, Surabaya, Indonesia

38-44

Abstract : 547

PDF : 40




DOI : 10.20473/ccj.v1i2.2020.38-44

Pulmonary Hypertension in Patient with Acyanotic Congenital Heart Defects in Abdul Wahab Sjahranie General Hospital, Samarinda from 2015-2016



 Abstract : 877

 PDF : 81



 DOI : 10.20473/ccj.v1i2.2020.45-49

Case Report

Long Term Survival of Ventricular Septal Rupture (VSR) Closure Concomitant with CABG in Post-Acute Myocardial Infarction Patient


 Irien E. Hernawati ⁽¹⁾

(1) Airlangga University

 50-56

 Abstract : 635

 PDF : 90



 DOI : 10.20473/ccj.v1i2.2020.50-56

Difficult Atrial Fibrillation Rate-Control and Digitalis Toxicity in Mitral-Valve Prolapse Patient with Hyperthyroidism

 Suryo Ardi Hutomo ⁽¹⁾, Agus Subagjo ⁽²⁾

(1) Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga ,


(2) Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga

 57-68


 Abstract : 949

 PDF : 132



 DOI : 10.20473/ccj.v1i2.2020.57-68

Acute Coronary Syndrome in Well Trained Young Adult: A Rare Cases

 Aprila Aqmarina Indah ⁽¹⁾, Praditya Virza Ramadhan ⁽²⁾, Vireza Pratama ⁽³⁾, Haikal Haikal ⁽⁴⁾, Irnizarifka Irnizarifka ⁽⁵⁾

(1) National Brain Centre, Jakarta Departmenet of Cardiology and Vascular Medicine, Gatot Soebroto Amry Hospital ,

(2) Tengku Rafi'an Hospital, Riau, Indonesia ,

(3) Department Cardiology and Vascular Medicine, Gatot Soebroto Army Hospital, Indonesia ,

(4) Department Cardiology and Vascular Medicine, Gatot Soebroto Army Hospital, Indonesia ,


(5) Cardiologist, Department of Cardiology and Vascular Medicine, Sebelas Maret University, Solo, Indonesia

 69-76

 Abstract : 1404

 PDF : 78



 DOI : 10.20473/ccj.v1i2.2020.69-76

Review Article

Lymphatic Disease in Cardiology: The Forgettable One

 Sidhi Laksono Purwowiyoto ⁽¹⁾, Novi Anggriyani ⁽²⁾, Steven Philip Surya ⁽³⁾

(1) Faculty of medicine, Universitas Muhammadiyah Prof. DR. Hamka, Tangerang ,

(2) Department Cardiology and Vascular Medicine, RSUP dr. Karyadi, Universitas Diponegoro, Semarang, Central Java ,





Login

Username *

Password *

[Forgot your password?](#)

Keep me logged in

[Login](#)

[Register](#)

Instruction For Author

[Author's Guidelines](#)

[Online Submission](#)

[Article Templates](#)

[Submission Guidelines](#)

Journal Policies

[Focus and Scope](#)

[Author Fees](#)

[Publication Ethics](#)

[Publication Frequency](#)

[Peer Review Process](#)

[Open Access Policy](#)

[Archiving](#)

[Plagiarism Check](#)

[Deposit Policy](#)

[ORCID ID Policy](#)

[Copyright Notice](#)

[License Term](#)

[Journal History](#)



People

Editorial Team

Peer Reviewer

Meet Our Editorial Team



Prof. Yudi Her Oktaviono, MD, PhD
Editor in Chief
Universitas Airlangga, Indonesia
Scopus Yudi Her Oktaviono



Dyana Sarvasti, MD, PhD
Deputy Editor
Universitas Katolik Widya Mandala, Indonesia
Scopus Dyana Sarvasti



Prof. Pieter A. Doevendans, MD
International Editors
Department of Cardiology, UMC Utrecht, Netherlands
Scopus Pieter A. Doevendans

[Read More](#)

Social Share



Indexed In



Visitors

Visitors





[View CCJ Stats](#)


Current Issue

ATOM 1.0

RSS 2.0

RSS 1.0

Keywords



In Collaboration with



Address

Department of Cardiology and Vascular Medicine

Faculty of Medicine, Universitas Airlangga

Jalan Mayjend Prof. Dr. Moestopo 6-8 Surabaya (60286) - East Java - INDONESIA

Contact Info:

Phone: +62-31-5501011

+628883208113

Email: ccj@journal.unair.ac.id

PRESENTED BY
DEPARTMENT OF CARDIOLOGY AND VASCULAR MEDICINE
AIRLANGGA UNIVERSITY, SOETOMO GENERAL HOSPITAL

IN COLLABORATION WITH
PERKI (INDONESIAN HEART ASSOCIATION)
SURABAYA



Cardiovascular and Cardiometabolic Journal is a scientific peer-reviewed article which freely available to be accessed, downloaded and used for research purposes. Manuscripts submitted to CCJ are published under the terms of the Creative Commons License.





Editorial Team



Prof. Yudi Her Oktaviono, MD, PhD

Editor in Chief

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

0000-0002-2350-2789 hRnBUVIAAAAJ **Scopus** 56431152100 6061731



Dyana Sarvasti, MD, PhD

Deputy Editor

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Katolik Widya Mandala, Indonesia

0000-0003-3276-7970 GI904UAAAAJ **Scopus** Dyana Sarvasti 6708528



J. Nugroho, MD, PhD

Deputy Editor

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

0000-0002-3501-1872 SKgGMXcAAAAJ **Scopus** 34971787500 -



Prof. Pieter A. Doevendans, MD

International Editors

Department of Cardiology, UMC Utrecht, Netherlands

0000-0002-6257-7169 - **Scopus** 7004918581 -



Maarten J. M. Cramer, MD, Ph.D

International Editors

Department of Cardiology, UMC Utrecht, Netherlands

0000-0002-6041-6759 04rXDTEAAAAJ **Scopus** 7102305152 -



Anwar Santoso, MD, PhD

International Editors

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Indonesia, Indonesia

0000-0001-8247-4151 KYPnrJoAAAAJ **Scopus** 36905206100 -



Prof. Budi Susetio Pikir, MD, PhD

Editorial Board

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

0000-0003-0705-9462 zZA8uNUAAAAJ **Scopus** 6507011164 60520



Prof. Rachmad Romdoni, MD, PhD

Editorial Board

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

0000-0002-5982-8983



Scopus[®]

6504356190



Prof. R. Muhammad Yogiarto, MD

Editorial Board

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

0000-0003-4372-6875



c9MGvgkAAAAJ

Scopus[®]

57190580716



6049463



Achmad Lefi, MD, PhD

Editorial Board

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Indonesia

-



YIP5d0YAAAAJ

Scopus[®]

57215562536



Trisulo Wasyanto, MD

Editorial Board

Sebelas Maret University, Indonesia

0000-0001-9900-0497



F9zeuEsAAAAJ

Scopus[®]

57191879710



6728138

Login

Username *

Password *

[Forgot your password?](#)

Keep me logged in

Login

[Register](#)





Original Research

The Correlation between Serum TG/HDL-c ratio and Arterial Stiffness Using the Cardio-ankle Vascular Index in Overweight or Obese PatientsA. Oktovianto¹, N. P. A. Laksmi¹, R. M. Yogiarto^{1,2*}, and J. N. E. Putranto^{1,2}¹Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.²Department of Cardiology and Vascular Medicine, Soetomo General Hospital, Surabaya, Indonesia.

ARTICLE INFO

Article history:

Submitted August 2020

Reviewed September 2020

Accepted September 2020

Available online September 2020

Corresponding author:myogiarto@yahoo.com**Keywords:*

Arterial stiffness

TG/HDL serum ratio

CardioAnkle Vascular Index

ABSTRACT

Background. The effect of increasing TG/HDL-c serum ratio, as a result of insulin resistance, is considered to have an important role in the occurrence of arterial stiffness. Increased arterial stiffness is an important marker in the development of cardiovascular disease and is a predictor of heart attack and stroke. **Aims.** This research was conducted to determine the relationship between the arterial stiffness assessed through non-invasive cardio-ankle vascular index (CAVI) examination and the TG/HDL-c ratio in overweight and obese individuals. Cardio-Ankle Vascular Index (CAVI) is a non-invasive clinical measurement to evaluate arterial stiffness manner and independent of blood pressure. **Methods.** This study is considered as correlational study using purposive sampling as the technique. Thirty two subjects participated in this research and each of them was subjected a CAVI examination to assess arterial stiffness. Their blood samples were collected for lipid profile measurement. The data were analyzed using Pearson Correlation Test. **Results.** The correlation between TG/HDL-c serum level and arterial stiffness using CardioAnkle Vascular Index (CAVI) in overweight or obese patients ($p < 0,01$) proved to be positive and significant.

Introduction

Overweight or obesity could change the structure and function of the blood vessels that lead to an increase in the blood vessels stiffness (arterial stiffness). Arterial stiffness occurs in the early phase and contributes to the process of cardiovascular premature dysfunction [1,2]. The mechanism of arterial stiffness in overweight or obese populations involving some complex process

has not been known completely. Furthermore, the last study is still controversial and advanced research is needed to clarify the factors that initiate and hasten the progression of this phenomenon. Therefore, a deeper understanding of the mechanism of arterial stiffness in overweight/obesity has a great clinical significance [1]. In this research, we will discuss the relationship

between the arterial stiffness assessed through non-invasive cardio-ankle vascular index (CAVI) examination and the TG/HDL-c ratio in overweight and obese individuals, to give understanding about the management [1,2].

Methods

This research used analytic correlational design. The samples of this research were overweight or obese individuals gathered during October 2016 until December 2016, who met the research criteria. Inclusion criteria in this research were overweight or obese man or woman, aged 20-60 years, and willing to participate by signing the informed consent. Exclusion criteria in this research were patient with a history of cardiovascular disease, coronary atherosclerotic disease, stroke, peripheral arteries disease/PAD), history of diabetes mellitus, history of chronic kidney disease, or history of smoking. CAVI examination used an automatic machine called VaSera VS-1500 (Fukuda Denshi, Tokyo, Japan). Thirty-two subjects participating in this research underwent CAVI examination to assess arterial stiffness and their blood sample were collected for lipid profile measurement. Relationship between serum TG/HDL-c ratio and the arterial stiffness was analyzed using the Pearson correlation test if the data were distributed normally, or the Spearman rank correlation test as an alternative.

Results

The study was done in Cardiology Clinic Dr. Soetomo Hospital, Surabaya, during September 2016- December 2017. We enrolled 32 overweight or obese subjects who fulfilled the research criteria. The subjects' ages were ranging from 20 to 60 years old, with the mean age of 42.69 ± 8.42 years old. Based on sex, subjects were mostly female (75%). The average of subject's body mass index (BMI) was 30.83 ± 5.07 , with the lowest was 25.8 kg/m² and the highest was 47.8 kg/m². Based on the BMI category, 18 subjects were overweight (56,3%), meanwhile 14 subjects were obese (43,7%). There were 7 subjects with obese grade I (21,9%), 6 subjects with obese grade II (18,8%), and 1 subject with extreme obese (3,0%). There were 30 subjects (93,7%) with abnormal waist circumference. The subject with increasing TG (>150 mg/dL) were 16 subjects (50,0%), meanwhile subject with low HDL-c (<35 mg/dL) was 16 subjects (50,0%). The lowest ratio of TG/ serum HDL-c was 1.39, and the highest was 7.47, with an average of 3.83 ± 1.39 . There were 2 subjects (6,3%) with normal TG/HDLc ratio (≤ 2.0), 17 subjects (53,1%) with high TG/HDL-c ratio (2.0–4.0), and 13 subjects (40,6%) with high TG/HDL-c ratio (>4). We also collected the blood pressure, fasting blood glucose, and 2 hours postprandial blood glucose data. The baseline characteristics were shown in Table 1.

Table 1. Baseline characteristic of the subject (N=32).

Variable	n (%) or Mean \pm SD
Age(years old)	42.69 \pm 8.42
20 – 29	4 (12.5)
30 – 39	5 (15.6)
40 – 49	15 (46.9)
50 – 59	8 (25.0)

Sex		
	Male	8 (25.0)
	Female	24 (75.0)
Height (cm)		158.84 ± 7.48
Weight (kg)		78.37 ± 17.35
Body Mass Index (kg/m ²)		30.83 ± 5.07
	Overweight (25 – 29,9)	18 (56.3)
	Obese grade I (30 – 34,9)	7 (21.9)
	Obese grade II (35 – 39,9)	6 (18.8)
	Extremely Obese (≥ 40)	1 (3.1)
Waist Circumference (cm)		101.36 ± 14.81
Systolic Blood Pressure (mmHg)		122.50 ± 10.98
Diastolic Blood Pressure (mmHg)		80.06 ± 8.01
Right-Ankle Brachial Index		1.04 ± 0.12
Left-Ankle Brachial Index		1.05 ± 0.13

The study found that the lowest TG/ HDL-c ratio was 1.39, meanwhile the highest was 7.47 with the average number of $3,83 \pm 1,39$. TG/serum HDL-c ratio is classified into three categories: normal if the TG/ serum HDL-c ratio is ≤ 2 , high TG/ serum HDL-c if it is in the range $2,0 - \leq 4$, and very high if it is >4 . About 6,3% of subjects have normal TG/ HDL-c ratio and the rest have high TG/serum HDL-c ratio. CAVI is classified based on the index score: <8 is categorized as normal; $>8- <9$ as borderline; and >9 as abnormal/atherosclerotic suspected. In this study, the minimum index score for Right-CAVI (R-CAVI) was 3.40 and the maximum was 11.10, with the mean score of $6,69 \pm 2,24$. Meanwhile, the minimum index score for Left-CAVI (L-CAVI) was 3.30 and the maximum was 11.60, with the mean score of 6.45 ± 2.26 . Mean-CAVI (M-CAVI) was obtained by calculating the mean index score of R-CAVI and L-CAVI for each subject.

Distribution of normality tests was done for the R-CAVI, L-CAVI, M-CAVI, and TG/HDL-c ratio by using Kolmogorov-Smirnov test. The result of normality test showed p value of ≥ 0.05 . The

Pearson correlation test found that there were positive and significant correlations between the TG/HDL-c ratio and the CAVI scores, as shown in Table 2. The TG/HDL-c ratio had a positive and significant correlation with the R-CAVI score ($r=0.455$, $p=0.004$). The TG/HDL-c ratio had a positive and significant correlation with the L-CAVI score ($r=0.528$, $p=0.001$). The TG/HDL-c ratio had a positive and significant correlation with the M-CAVI score ($r=0.497$, $p=0.002$).

Table 2. Correlation results using Spearman analysis

Variable	p	r
TG/HDL-c serum ratio and R-CAVI	0,004	0,455
TG/HDL-c serum ratio and L-CAVI	0,001	0,528
TG/HDL-c serum ratio and M-CAVI	0,002	0,497

^aTG/HDL-c : Triglyceride/High density lipoprotein-c
^bR: right

Discussion

Obesity was already a health epidemic in the last two decades. Some studies said that the increase of BMI is correlated with increasing risk of morbidity and mortality of the cardiovascular diseases. Although the correlation is clearly showed, most of the studies did not give the clear pathophysiology about BMI increase contributing to the increase of cardiovascular diseases incidence. However, some of the literature said that the correlation of overweight or obesity with cardiovascular diseases occurs through metabolic profile changes or cardiac and blood vessel's structural and functional changes. The metabolic profile changes include glucose intolerant, insulin resistant, increasing of inflammation marker, and the lipid metabolism changes. Meanwhile, the structure changes, also the cardiac and blood vessel function changes, occur through the increasing of the arterial stiffness and left ventricle hypertrophy. Wang et al. and Di Bonito et al. studies found a strong correlation between arterial stiffness with the proatherogenic lipid ratio in teenagers or adults [3,4]. Our study was intended to evaluate the correlation between TG/serum HDL-c ratio with arterial stiffness assessed with CardioAnkle Vascular Index (CAVI) in overweight or obese patient.

a. Basic characteristics of the subject

The subjects in this study were overweight or obese patient who visited the cardiology clinic during October 2016 – December 2016. We enrolled 32 subjects with overweight or obese who were selected by consecutive sampling. Most of the respondent is female (75%), with the average age in the fourth decade, and mean BMI of 30.83 ± 5.07 kg/m². Almost all of the subject had abnormal waist circumference (93,7%). The increased CAVI value above normal was found in 9 subjects, and all of them were older than 40 years old. In the subject with abnormal CAVI (>8), only one subject (11,1%)

had a normal TG/serum HDL-c ratio, while the others had TG/serum HDL-c ratio above normal.

The previous study had shown that there was an effect from some of the risk factors which correlated to the arterial stiffness progressivity, such as age, hypertension, smoking, and diabetes. However, other factors such as dyslipidemia and obesity also play a role in the arterial stiffness. The study by Holewijn et al., showed that non-HDL-c usage is a predictor in the identification of arterial stiffness compared to the LDL-c. The study by Di Bonito et al., said that TG/serum HDL-c ratio is the independent risk factor of the arterial stiffness in teenagers and adults population [3]. In this study, from all of the subject with normal BMI, there were 16 subjects (50,0%) with increasing TG, while 16 subjects (50,0%) had low HDL levels.

b. TG/serum HDL-c ratio

The study by Gaziano et al., was the first that reported 16 times increase of the risk to a myocardial infarction in the highest TG/serum HDL-c ratio quartile group compared to the lowest quartile [5]. Another study reported that high TG/serum HDL-c ratio is correlated independently with coronary artery disease, which is assessed by angiography (defined as stenosis >50%) and cardiovascular mortality. The latest study by Yuji, et al., said that high TG/HDL-c ratio is the risk to atherosclerosis and significantly increases the arterial stiffness in Japanese male population with diabetes, which was assessed by carotid artery intima-media thickness (CIMT) examination and CAVI.

Based on the data from Wan et al., TG level is an independent variable which inversely proportional to predict LDL particle size. Meanwhile, HDL-c level is an independent variable which directly proportional to the LDL particle size. Because of that, TG/ HDL-c ratio can be more

accurate to predict small sd-LDL. Low TG/ HDL-c ratio presents big size LDL particle or non-atherogenic LDL particle. However, high TG/ HDL-c ratio present small size pro-atherogenic LDL particle, which is very correlated to the atherosclerosis initiation and progression. Dobiasova et al., found that the increasing of TG/ HDL-c log parameter as the atherogenic index showed the high concentration of plasma triglyceride-rich lipoprotein and small size LDL-c particle [4,6]. The study by Shimizu et al., involving 2.431 subjects in Japan, showed that BMI has a positive correlation to the TG/serum HDL-c ratio, also the correlation between BMI and diabetes is influenced by the role of TG/serum HDL-c ratio [7].

In this study, we obtained the minimum the TG/serum HDL-c ratio of 1.39 and the maximum of 7.47, with the mean ratio of 3.83 ± 1.39 . From this result, we did not find any significant correlation between TG/serum HDL-c ratio with LDL-c level. This can be caused by quantitatively measured LDL-c level without considering the quality of the LDL-c particle size, such as mentioned in the study by Wan et al. This study also found that there was no correlation between TG/serum HDL-c ratio with BMI. This phenomenon can be explained by Grundy et al. and Festa et al. studies, where they found that insulin resistant prevalence in the overweight or obese population was about 30-37% [8]. However, the interesting finding here was that we obtained a significant correlation between BMI and LDL-c level ($p < 0,05$; $r = 0,395$). This finding supports the study by Lauvicius et al., which showed that the proatherogenic profile lipid metabolism changes and the increase of the cardiovascular incidence are in line with the increasing of BMI [9].

c. Cardio-ankle vascular index (CAVI) number

Arterial stiffness is one of the cardiovascular morbidity and mortality predictor. A study by Luo et al., showed that there was a significant positive correlation between arterial stiffness with the risk of coronary heart disease incident [10]. The progressive structural changes, such as elastin fragmentation and degeneration, increasing of collagen, artery wall thickness, and endothelium damage cause the increase of arterial stiffness incidence.

In this study, CAVI examination was done in all subjects using VaSera VS-1000 machine. CAVI was used to assess the arterial stiffness and to evaluate the risk of cardiovascular disease incidence. From 32 subjects, there were nine subjects (28,2%) with abnormal CAVI. The increased CAVI score showed the progressivity of the arterial stiffness. However, from this data, we could not find the significant correlation between CAVI and the blood pressure, neither systolic ($p = 0.686$) nor diastolic ($p = 0.408$), also the correlation between CAVI with ABI ($p = 0.504$).

These results were in line with the study by Kotani et al., which proved that the benefit of arterial stiffness measurement with CAVI depended on the stiffness- β parameter and was not influenced by blood pressure [11].

d. Correlation between TG/serum HDL-c ratio with arterial stiffness which is assessed by CAVI

This study showed that there was a positive and significant correlation between TG/serum HDL-c ratio level and the arterial stiffness assessed with CAVI in overweight or obese patient. The higher the TG/serum HDL-c ratio means the higher the arterial stiffness is.

Some potential mechanism can explain the correlation between TG/serum HDL-c ratio and arterial stiffness. Insulin resistant is believed to have a significant role in cardiovascular disease pathogenesis and has been correlated independently with brachial-ankle Pulse Wave Velocity, even in a healthy person. Insulin resistant causes the increasing of synthesis and secretion of TRL, whilst decreasing the HDL cholesterol. McLaughlin et al., said that TG/serum HDL-c ratio could be identified if there is insulin resistant in overweight person with normal glucose tolerant and it serves as an insulin resistant marker that has the same sensitivity and specificity with the plasma insulin concentration in fasting condition [12]. In a cross-sectional study from 6.546 Korean adults who had done the routine health examination, it was found that TG/serum HDL-c ratio had a significant correlation with the insulin resistant in person without metabolic syndrome abnormality. Because the TG/HDL-c ratio examination is a simple clinical examination used as an indicator of insulin resistance, this examination can predict arterial stiffness related to insulin resistance.

Next, the correlation between TG/ serum HDL-c ratio and arterial stiffness may be related to the increased of small dense-LDL (sd-LDL) particle, which is considered as more atherogenic than normal size LDL particle because the sd-LDL particle is easier to enter the blood vessel wall, as well as more vulnerable to oxidative modification, which in turn will be increasing its atherogenicity [12]. These findings showed that high TG/serum HDL-c ratio could be used as the clinical indicator of insulin resistance and the increasing of sd-LDL particle can be used to evaluate the early damage phase of vascular. Further study is needed to explain the right mechanism responsible for that correlation.

This study was in line with the study in 2012 by Urbina et al. that showed TG/serum HDL-c ratio was positively correlated to the arterial stiffness in healthy teenagers and adults population. In the study by Laucevicius et al., it was showed that subject group with high CAVI had worse metabolic profile compared to the subject group with normal CAVI, and after further follow up (average $3.8 \pm 1,7$ years) CAVI was correlated significantly to the cardiovascular and myocardial infarction incidence [9]. With the data above and based on the literature, CAVI measurement can be used to evaluate the subclinical cardiovascular disease that can predict the cardiovascular incidence in the future. The correlation between obesity and CAVI is strengthened by the study that showed that weight loss in the obesity subject can improve the arterial stiffness assessed with CAVI [13]. Therefore, healthy lifestyle changes to achieve the ideal weight can decrease the cardiovascular risk. Some of the limitations in this study were its sampling technique and small scope population. Samples were taken by purposive sampling because of the time limit, number of patients, and the budget. This study conducted in a small scope of population and non-multicentre.

Conclusion

There is a positive and meaningful correlation between TG/serum HDL-c ratio level and arterial stiffness which is assessed by Cardio-Ankle Vascular Index (CAVI) in overweight or obese patients ($p < 0,01$; $r = 0,455 - 0,528$). Further study about the correlation between TG/ serum HDL-c ratio and arterial stiffness is needed for an overweight or obese patient with the bigger numbers of sample to get the result that can be generalized.

Acknowledgement

There is no conflict of interest.

References

1. Brillante DG, O'Sullivan AJ, and Howes LG. Arterial Stiffness in Insulin Resistance: The Role of Nitric Oxide and Angiotensin II Receptors. *Vasc Health Risk Manag*; 2009. 5: 73–8.
2. Villacorta L, and Chang DL. The Role of Perivascular Adipose tissue in Vasoconstriction, Arterial Stiffness, and Aneurysm. *Horm Mol Biol Clin Investig*; 2015. 21: 137.
3. DiBonito P, et al. Usefulness of The High Triglyceride-to-HDL Cholesterol Ratio to Identify Cardiometabolic Risk Factors and Preclinical Signs of Organ Damage in Outpatient Children. *Diabetes Care*; 2012. 35: 158–62.
4. Wang F, Ye P, Luo L, Xiao W, Qi L, Bian S, Wu H, Sheng L, Xiao T, and Xu R. Association of Serum Lipids with Arterial Stiffness in a Population-based Study in Beijing. *Eur J Clin Invest*; 2011. 41: 929–36.
5. Gaziano JM, Hennekens CH, O'Donnell CJ, Breslow JL, and Buring JE. Fasting Triglycerides, High-density Lipoprotein, and Risk of Myocardial Infarction. *Circulation*; 1997. 96: 2520–5.
6. Diraison F, Moulin P, and Beylot M. Contribution of Hepatic de Novo Lipogenesis and Re-esterification of Plasma Non Esterified Fatty Acids to Plasma Triglyceride Synthesis during Non-alcoholic Fatty Liver Disease. *Diabetes Metab*; 2003. 29: 478–85
7. Schwarz J-M, Linfoot P, Dare D, and Aghajanian K. Hepatic de Novo Lipogenesis in Normoinsulinemic and Hyperinsulinemic Subjects Consuming High-fat, Low-carbohydrate and Low-fat, High-carbohydrate Isoenergetic Diets. *Am J Clin Nutr*; 2003. 77: 43–50.
8. Grundy SM. Small LDL, Atherogenic Dyslipidemia, and the Metabolic Syndrome Circulation; 1997. 95: 1–4.
9. Laucevičius A, Ryliškyte L, Balsyte J, Badariene J, Purnaite R, Navickas R, and Solovjova S. Association of Cardio-Ankle Vascular index with Cardiovascular Risk Factors and Cardiovascular Events in Metabolic Syndrome Patients. *Med*; 2015. 51: 152–8.
10. Luo K-Q, Feng X-W, Xu B-C, and Long H-B. Association Between Arterial Stiffness and Risk of Coronary Artery Disease. *Pakistan J Med Sci*; 2014. 30: 1314–8.
11. Kotani K. Cardio-Ankle Vascular Index and Its Potential Clinical Implications for Sleep Apnea Pulse (Basel, Switzerland). 2014. 1: 139–42.
12. McLaughlin T, Abbasi F, Cheal K, Chu J, Lamendola C, and Reaven G. Use of Metabolic markers to identify overweight individuals who are insulin resistant. *Ann Intern Med*; 2003. 139: 802–9.
13. Nagayama D, et al. Effects of Body Weight Reduction on Cardio-Ankle Vascular Index (CAVI). *Obes Res Clin Pract*; 2013. 7: e139–45