

In vitro Antioxidant and Anticholinesterase Activities of Extracts from the Leaves of Cassia moschata Kunth

by Suciati Suciati

Submission date: 28-Jun-2022 10:37AM (UTC+0800)

Submission ID: 1863962526

File name: 59_RJPT_15_4_2022_Journal_Suciati.pdf (225.19K)

Word count: 4833

Character count: 26495

RESEARCH ARTICLE

***In vitro* Antioxidant and Anticholinesterase Activities of Extracts from the Leaves of *Cassia moschata* Kunth**

Suciati Suciati^{1,2*}, Wachidatur Rizqiyah², Dwiki Nur Inayah², Retno Widayati¹, Wiwied Ekasari¹, Nunggruthai Suphrom³

¹Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya 60115, East Java, Indonesia.

²Center for Natural Product Medicine Research and Development, Institute of Tropical Diseases, Universitas Airlangga, Surabaya, 60115, East Java, Indonesia.

³Department of Chemistry, Faculty of Science, Naresuan University, Phitsanulok, 65000, Thailand.

*Corresponding Author E-mail: suciati@ff.unair.ac.id

ABSTRACT:

Alzheimer's disease (AD) is a neurodegenerative disorder, which is the most common cause of dementia. This disease commonly occurs in elderly people. The increase in life expectancy means that the number of people suffering from AD is expected to rise each year if there is no effective treatment found. The relation of cholinesterase and oxidative stress to Alzheimer's disease has been reported. In our previous study, we have investigated the potency of the ethanolic extract of *Cassia moschata* leaves as an anticholinesterase. The current study aimed to investigate the antioxidant and anticholinesterase properties of the ethanolic and aqueous extracts of *C. moschata* as well as to determine the total phenolic content (TPC). Two different methods were used to evaluate the antioxidant activity by 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assays. The anticholinesterase assay was carried out against acetylcholinesterase (AChE) according to the modified Ellman's method. The TPC was determined by a colorimetric method using Folin-Ciocalteu's phenol reagent, and employing gallic acid as a reference. The ethanolic and aqueous extracts of *C. moschata* demonstrated antioxidant activity in both DPPH and ABTS assays. There were statistically significant differences in the IC₅₀ values of the ethanolic and aqueous extracts in both DPPH and ABTS assays. The aqueous extract exhibited a lower IC₅₀ value compared to the ethanolic extract. The IC₅₀ value for the aqueous extract was 36.46 µg/mL in the DPPH assay, and 10.61 µg/mL in the ABTS method compared to IC₅₀ 38.74 µg/mL and 17.17 µg/mL for the ethanolic extract, respectively. Meanwhile, the ethanolic extract showed higher potency as anticholinesterase with the IC₅₀ value of 44.43 µg/mL compared to the aqueous extract with an IC₅₀ value of 114.60 µg/mL. The TPC measurement revealed that the aqueous extract has a higher amount of phenolic than the ethanolic extract. These data suggest that the aqueous extract from the leaves of *C. moschata* has a higher ability to scavenge free radicals compared to the ethanolic extract, which also contains a higher amount of phenolic compounds. However, the high content of phenolic compounds in the aqueous extract did not correspond to the anticholinesterase activity. The presence of non-phenolic compounds may also contribute to the anticholinesterase activity in the ethanolic extract.

KEYWORDS: Alzheimer's disease, *Cassia moschata*, Medicinal plant, Anticholinesterase, Antioxidant.

INTRODUCTION :

Alzheimer's disease (AD) is a progressive neurodegenerative disorder that has been known as the common cause of dementia¹. The low level of neurotransmitters especially acetylcholine is one of the common features in this disease which is caused by the loss of cholinergic nerves as well as the presence of senile

plaques and neurofibrillary tangles²⁻⁶. Therefore one of the strategies in the treatment of this disease is the use of anticholinesterases such as tacrine, donepezil, rivastigmine, and galantamine. The relation of oxidative stress and neurological diseases, such as Alzheimer's disease has been well documented. Numerous experimental and clinical studies have shown that oxidative stress causes the loss of neurons and the progression of the disease to dementia⁷⁻¹⁰. The presence of a toxic peptide, β -amyloid in the brain of patients with Alzheimer's disease is also caused by oxidative stress^{10,11}. Antioxidants are components that can scavenge free radicals so that can prevent and repair the damage caused by oxidative stress^{10,12}.

Medicinal plants have shown a great contribution to the treatment of many diseases. Herbal medicines have been reported to show significant effects in the treatment of Alzheimer's disease, such as the well-known *Ginkgo biloba* and lately Chinese medicinal plant *Huperzia serrata*¹³⁻¹⁷. Plant from the genus *Cassia*, the family Caesalpiniaceae consists of more than 500 species worldwide. These flowering plants are distributed in tropical and subtropical regions¹⁸⁻²⁰. The genus *Cassia* is not only known as an ornamental plant but also known for its use in folk medicine, such as for skin diseases, gastrointestinal problems, cough, and cardiac disorders^{18,20,21}. Metabolites with promising bioactivities have also been reported from several *Cassia*, such as terpenoid, flavonoid, alkaloid, xanthone, and anthraquinone²¹⁻²⁴. In our continuing study on the search of medicinal plants and marine sources for neurodegenerative diseases, we have found that several *Cassia* species, including the ethanolic extract of *Cassia moschata* have potency as cholinesterase inhibitor²⁵. Several *Cassia* species have also been reported as antioxidants, such as the ethanolic and aqueous extracts of *C. angustifolia* seed²⁶. The antioxidant activity of four species of *Cassia*, namely *C. auriculata*, *C. siamea*, *C. uniflora*, and *C. italica* have also been reported²⁷. Other *Cassia* species that have shown potency as antioxidants including *C. tora*, *C. occidentalis*, *C. javanica*, *C. glauca*, and *C. grandis*²⁸⁻³². To the best of our knowledge, there is no report on the antioxidant potency of the bronze shower plant, *C. Moschata*, as well as its phytochemical contents. Therefore, the current study focus on the evaluation of the antioxidant and anticholinesterase activities of the ethanolic and aqueous extracts of *Cassia moschata* leaves as well as the determination of the phenolic content.

MATERIALS AND METHODS:

Materials:

2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), gallic acid, quercetin, and potassium persulfate were

purchased from Sigma. Folin-Ciocalteu's phenol reagent, aluminum chloride, sodium carbonate, sodium nitrite were purchased from Merck. All chemicals and solvents used were of analytical grade. The leaves of *Cassia moschata* were obtained from Purwodadi Botanic Garden, East Java, Indonesia on March 9th, 2019. The voucher specimen (PWD 04) was deposited at the Institute of Tropical Diseases Universitas Airlangga. The plant was identified by Purwodadi Botanic Garden, Indonesian Institute of Sciences with identification letter number: 0371/IPH.06/HM/III/2019.

Preparation of extracts:

The leaves of *Cassia moschata* were air-dried at room temperature for seven days, then powdered in a grinder. Ten grams of the powdered leaves were each extracted with 80 mL ethanol and water by using ultrasonic extraction for 3 x 10 mins, then the filtrate was separated by filtration. The residue was re-extracted with each solvent using the same procedure three times. All collected filtrates were evaporated in a rotary evaporator to obtain the ethanolic (1.40 g) and aqueous (0.77 g) extracts.

DPPH radical scavenging assay:

The assay was carried out in triplicates based on modified protocols^{33,34}. The DPPH solution (0.25 mM) was prepared by dissolving DPPH powder in methanol. The extracts were dissolved in methanol at a concentration of 100 μ g/mL, this was then further diluted to series of concentrations 2.5 – 60 μ g/mL. Gallic acid was used as a standard. The samples (100 μ L) were mixed with 0.25 mM DPPH reagent (100 μ L) in 96 well plates. DPPH reagent (100 μ L) was mixed with methanol (100 μ L) to serve as a control, while methanol (200 μ L) was used as blank. The reaction mixtures were incubated in the dark at room temperature for 30 mins. The solutions were shaken for 30 s, followed by measurement of absorbance at 517 nm in a microplate reader (Bio-Tek Instrument, USA). The DPPH scavenging effect was calculated by the following formula.

$$\text{DPPH Radical Scavenging activity (\%)} = \frac{(\text{abs control} - \text{abs sample})}{\text{abs control}} \times 100$$

where *abs control* is the absorbance of DPPH radical + methanol and *abs sample* is absorbance DPPH radical + extract/standard.

ABTS radical scavenging assay:

The bioassay was performed based on the previous method with some modifications³⁴. Briefly, to produce ABTS radical, 5 mL ABTS (7 mM) was mixed with 88 μ L potassium persulfate (140 nM), and was kept in the dark at room temperature for 16 h. Samples at the concentration range 2.5 – 60 μ g/mL were prepared in

methanol. The samples (100 μL) were then mixed with 100 μL of ABTS in a 96-well microplate and allowed for incubation for 6 mins in the dark at room temperature. The absorbance was measured at 734 nm by using a microplate reader (set to shake for 30 s before reading). Gallic acid was used as standard. Experiments were done in triplicate. The ABTS radical scavenging activity was calculated by using the following equation.

$$\text{ABTS Radical Scavenging activity (\%)} = \frac{(\text{abs control} - \text{abs sample})}{\text{abs control}} \times 100$$

abs control is the absorbance of ABTS radical + methanol and *abs sample* is absorbance ABTS radical + extract/standard.

Anticholinesterase Assay:

The assay was carried out according to the modified Ellman's method³⁵⁻³⁷. The extracts were dissolved in methanol at a concentration of 10mg/mL and were then diluted with 50mM Tris buffer to obtain series of concentrations containing not more than 10% of methanol. The final test concentrations in the wells were: 500, 400, 200, 100, 50, 40, 20, 10, 5, 2 and 1 $\mu\text{g/mL}$. Sample solutions were then added to a 96-well microplate, followed by the addition of 1.5 mM ATCI or 1.5 mM BTCl (25 μL), 3 mM DTNB (125 μL), and Tris buffer (50 μL). The enzyme *EeAChE* (25 μL of 0.22 U/mL) was then added. The solutions were shaken for 30 s in a microplate reader (Bio-Tek Instrument, USA) before measurement. The presence of yellow color due to the formation of 5-thio-2-nitrobenzoate was monitored at 405 nm every 5 s for 2 mins. Every experiment was carried out in triplicates. Galantamine was used as a positive control, and 10% methanol was used as a negative control. The inhibitory activity was calculated by using the equation below:

$$\% \text{Inhibition} = \frac{(\text{Mean velocity of control} - \text{Mean velocity of sample})}{\text{Mean velocity of control}} \times 100$$

29

Determination of total phenolic content (TPC):

The TPC in the samples was determined according to the method Herald et al (2012) Zhang et al (2006) with slight modification^{33,38}. In brief, twenty-five microlitres of gallic acid standard (25–500 $\mu\text{g/mL}$) or samples (1000 $\mu\text{g/mL}$) were added to 96-well microplate, followed by water (75 μL) and Folin & Ciocalteu's phenol reagent (25 μL). The mixture was incubated for 6 mins at room temperature. After incubation, 100 μL Na_2CO_3 solution (75 g/L) was added to each well, followed by incubation for 90 mins in the dark at room temperature. The mixtures were shaken for 30 s before reading the absorbance at 765 nm in a microplate reader.

Statistical analysis:

The 50% inhibitory concentration (IC_{50}) was determined using GraphPad Prism 6.0 software by plotting log concentrations as axis and % inhibition as ordinate. Results were expressed as mean \pm standard deviation (SD) of three parallel measurements. The IC_{50} values from DPPH and ABTS assays were analyzed using independent-sample T-tests. The correlation between the antioxidant capacity and the total phenolic content was determined using Pearson's correlation test using IBM SPSS statistics 21 software. Difference and correlation were regarded as statistically significant when $p < 0.05$.

RESULT:

Antioxidant Activity:

The dose-response inhibition of extracts in DPPH and ABTS assays can be seen in Figures 1 and 2. The data showed that in the DPPH assay the radical quenching abilities of both extracts were increased in a dose-dependent manner (Figure 1). At a concentration of 5 $\mu\text{g/mL}$, the ethanolic and aqueous extracts scavenged 4.30% and 16.52% of DPPH radical, respectively. These values increased to 66.15% and 66.65%, respectively at a concentration of 50 $\mu\text{g/mL}$. The same trend was observed in the ABTS assay, the inhibition of ABTS radical for both ethanolic and aqueous extracts increased more than four and two folds, respectively at concentration 50 $\mu\text{g/mL}$ compared to at concentration 5 $\mu\text{g/mL}$ (Figure 2).




Figure 1. Percentage scavenging of DPPH at different concentrations of *C. moschata* extracts




Figure 2. Percentage scavenging of ABTS at different concentrations of *C. moschata* extracts

The IC₅₀ values were calculated and presented in table 1 which shows that the aqueous extract gave lower IC₅₀ values at 36.46 and 10.61 µg/mL in DPPH and ABTS assays, respectively compared to the aqueous extract.

Table 1. Antioxidant and anticholinesterase of *C. moschata* extracts

Samples	IC ₅₀ (µg/mL) ^a		
	DPPH	ABTS	AChE
Ethanolic extract	38.74 ± 0.26	17.17 ± 0.41	44.43 ± 3.58
Aqueous extract	36.46 ± 0.09	10.61 ± 0.50	114.60 ± 2.88
Gallic acid	2.76 ± 0.02	0.97 ± 0.03	n.d
Galantamine	nd	nd	0.63 ± 0.05

^aData presented as mean ± SD of three experiments, each done in triplicate. nd: not determine

Anticholinesterase Activity:

The evaluation of the anticholinesterase activity of extracts was carried out against the AChE enzyme. The results as can be seen in Table 1 and Figure 3 showed that the ethanolic extract gave higher potency as anticholinesterase with an IC₅₀ value of 44.43 µg/mL compared to the aqueous extract with an IC₅₀ value of 114.60 µg/mL.




Figure 3. Dose-dependent response of *C. moschata* ethanolic and aqueous extracts against AChE

Total Phenolic Content (TPC):

The total phenolic contents in the ethanolic and aqueous extracts were calculated based on a standard curve of gallic acid ($y = 0.005x + 0.029$, $R^2 = 0.999$). The results as can be seen in Table 2 showed that the aqueous extract contains higher phenolic contents (260.0 mg GAE/g extract) compared to that of the ethanolic extract (238.2 mg GAE/g extract).

Table 2. Extract yield and total phenolic content (TPC) of *C. moschata* extracts

Samples	Extract Yield (%)	TPC (mg GAE/g extract) ^a
Ethanolic extract	7.70	238.2 ± 0.06
Aqueous extract	13.97	260.0 ± 0.25

^aData presented as mean ± SD of three experiments, each done in triplicates.

DISCUSSION:

Cassia species are higher plants native to Southeast Asia and sub-Saharan Africa. These plants are commonly used as food as well as various traditional medicines. Several *Cassia* species have been reported to possess antioxidant properties in many experimental studies²⁶⁻³². However, there are *Cassia* species, such as *Cassia moschata*, that have limited reports both for biological activities as well as chemical composition.

In this study, the radical-scavenging activities of the ethanolic and aqueous extracts of *C. moschata* leaves were determined using the DPPH and ABTS assays. In the DPPH assay, the antioxidant provides a hydrogen atom that will react with DPPH radical (diphenylpicrylhydrazyl) to form a non-radical diphenylpicrylhydrazin, which has a yellow color. The degree of discoloration indicates the radical-scavenging potential of the sample^{39,40}. The principle of ABTS assay is similar to that of DPPH assay, in which the antioxidant acts as a hydrogen donor to form a non-radical ABTS. The reduction of a dark-bluish color of ABTS radical can be monitored by spectrophotometer^{41,42}.

The data presented in Figures 1 and 2 suggested that the radical quenching abilities of both extracts are increased in a dose-dependent manner in both ethanolic and aqueous extracts. The potency of samples to scavenge DPPH and ABTS radicals are expressed as IC₅₀ value, which is the concentration of sample required to scavenge 50% DPPH or ABTS free radicals. The value was calculated from the inhibition curve. Based on the IC₅₀ values (Table 1), it can be seen that the aqueous extract has higher potency as an antioxidant compared to the ethanolic extract. Analysis by using an unpaired t-test suggested that there were statistically significant differences observed between the IC₅₀ values for the ethanolic and aqueous extracts in both DPPH and ABTS assays with *p* values of 0.0001 (DPPH) and <0.0001 (ABTS), respectively.

The anticholinesterase activity of the ethanolic extract of *C. moschata* has been reported in our previous study. In this study, the evaluation of anticholinesterase activity was carried out in both ethanolic and aqueous extracts in order to compare the potency. The results as can be seen in Table 1 and Figure 3 showed that the ethanolic extract higher potency as anticholinesterase with an IC₅₀ value of 44.43 µg/mL compared to the aqueous extract with an IC₅₀ value of 114.60 µg/mL.

The higher contents of phenolic in the aqueous extract compare to the ethanolic extract suggested that the phenolic presents in the leaves of *C. moschata* possibly are polar compounds, such as in the glycoside form. In addition, the results also showed that there is a significant correlation between the IC₅₀ values in both DPPH and

ABTS assays and the total phenolic contents in the extracts (Table 3). The higher phenolic content is responsible for the higher antioxidant activity in the aqueous extract. These results are in agreement with the previous studies, which prove that there is a strong correlation between the total phenolic content and the antioxidant activity of some medicinal plants²⁸. However, the higher phenolic content in the aqueous extract does not correspond to the anticholinesterase activity. Therefore, this suggests that the presence of other compounds, such as alkaloids that are commonly present in the plant from the genus *Cassia*²¹⁻²⁴, may also be responsible for the anticholinesterase activity in the ethanolic extract.

Table 3. Correlations between the IC₅₀ values of antioxidant assays and phenolic content

Assay	Correlation (r) TPC
IC ₅₀ of DPPH radical scavenging ability	-0.991*
IC ₅₀ of ABTS radical scavenging ability	-0.994*

r: correlation coefficient. *indicates significance at $p < 0.01$

Phenolics are aromatic secondary plant metabolites that are known for their extensive profitable biological activities^{43,44}. These compounds are always considered as the major contributor to antioxidants in plants due to their ability to scavenge free radicals, active oxygen species as well as metal chelators^{43,45}. Studies revealed that the antioxidant capacity of phenolic compounds depends on the number and arrangement of the hydroxyl groups in the phenolic compound. Studies on the phenolic content of *Cassia* spp. have been documented^{26-28,30,46}, however, there is no report for *C. moschata*. Mehta et al (2017) reported the total phenolic contents of various parts of *C. javanica* and *C. siamea*, which stated that *C. javanica* leaves contain higher phenolic compared to other tested extracts³⁰. The antioxidant potency, as well as phenolic contents of seven *Cassia* species, have also been reported. *C. glauca* was reported to show the strongest antioxidant capacity compared to the other six *Cassia* species, which was also related to its high content of phenolic compounds²⁸. The study on the chemistry of *C. moschata* is very limited, however tracing on the chemistry of the plant from the same genus revealed that anthraquinone, terpenoid, xanthone, flavonoid as well as alkaloid presence in *Cassia* spp.²¹⁻²⁴. Considering higher anticholinesterase potency in the ethanolic extract, the non-phenolic compounds may contribute to this activity, which needs further study.

CONCLUSION:

The aqueous extract of *Cassia moschata* leaves gave higher free radical scavenging activity compared to the ethanolic extract due to higher phenolic content, however, the ethanolic extract gave better anticholinesterase activity. The presence of other compounds, such as

alkaloids in the ethanolic extracts may be responsible for its anticholinesterase activity. Further study is needed to investigate the chemistry of *C. moschata* leaves.

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this investigation.

ACKNOWLEDGMENTS:

Authors acknowledge the Faculty of Pharmacy, Universitas Airlangga for research grant PUF 2020.

REFERENCES:

- Bird TD. Genetic Aspects of Alzheimer Disease. *Genetics in Medicine*. 2008; 10(4): 231-239. doi:10.1097/gim.0b013e31816b64dc.
- Patel CK, Panigrahi B, Badmanaban R, Patel CN. Biochemical Origins of Alzheimer's Disease with Treatment Techniques. *Research Journal of Pharmacology and Pharmacodynamics*. 2010; 2(1): 33-38.
- Kumar DR, Shankar MS, Reddy PP, Kumar BRS, Sumalatha N. A Review on Alzheimer's Disease. *Research Journal of Pharmacology and Pharmacodynamics*. 2014; 6(1): 59-63.
- Aanandhi MV, Niventhi A, Rujaswini T, Hemalatha CN, Praveen DA. Comprehensive Review on the Role of Tau Proteins in Alzheimer's Pathology. *Research Journal of Pharmacology and Technology*. 2018; 11(2): 788-790.
- Choudhury S, Vellapandian C. Alzheimer's Disease Pathophysiology and its Implications. *Research Journal of Pharmacology and Technology*. 2019; 12(4): 2045-2048.
- Dhinakaran S, Tamilanban T, Chitra V. Targets for Alzheimer's Disease. *Research Journal of Pharmacology and Technology*. 2019; 12(6): 3073-3077.
- Christen Y. Oxidative Stress and Alzheimer Disease. *The American Journal of Clinical Nutrition*. 2000; 71(2): 621S-629S. doi:10.1093/ajcn/71.2.621s.
- Halliwell B. Role of Free Radicals in Neurodegenerative Diseases: Therapeutic Implications for Antioxidant Treatment. *Drugs Aging*. 2001; 18(9): 685-716. doi:10.2165/00002512-200118090-00004.
- Singh RP, Sharad S, Kapur S. Free Radicals and Oxidative Stress in Neurodegenerative Diseases: Relevance of Dietary Antioxidants. *Journal Indian Academy of Clinical Medicine*. 2004; 5(3): 218-225.
- Pham-Huy LA, He H, Pham-Huy C. Free Radicals, Antioxidants in Disease and Health. *International Journal of Biomedical Science*. 2008; 4(2): 89-96.
- Butterfield DA. Amyloid Beta-peptide (1-42)-Induced Oxidative Stress and Neurotoxicity: Implications for Neurodegeneration in Alzheimer's Disease Brain. A Review. *Free Radical Research*. 2002; 36(12): 1307-1313. doi:10.1080/1071576021000049890.
- Krishnaiah D, Sarbatly R, Nithyanandam R. A Review of the Antioxidant Potential of Medicinal Plant Species. *Food Bioprocess Processing*. 2011; 89(3): 217-233. doi:10.1016/j.fbp.2010.04.008
- Vakhariya RR, Talokar SS, Salunkhe VR, Magdum CS. Cognitive Disorders and its Herbal Remedies. *Research Journal of Pharmacognosy and Phytochemistry*. 2017; 9(1): 42-46.
- Chitra V, Narayanan J. In vitro Screening for Anti-Cholinesterase and Antioxidant Activity of Extract of *Garcinia hanburyi*. *Research Journal of Pharmacology and Technology*. 2018; 11(7): 2918-2921.
- Velraj M, Lavaniya N. Alzheimer Disease and a Potential Role of Herbs- A Review. *Research Journal of Pharmacology and Technology*. 2018; 11(6): 2695-2700.
- Jadhav RP, Kengar MD, Narule OV, Koli VW, Kumbhar SB. A Review on Alzheimer's Disease (AD) and its Herbal Treatment of Alzheimer's Disease. *Asian Journal of Research in Pharmaceutical Sciences*. 2019; 9(2):112-122.
- Patil SV, Patil VK, Patil PA. Review on Herbal medicines of Alzheimer's Disease. *Asian Journal of Research in Pharmaceutical Sciences*. 2020; 10(3):171-177.
- Ganapaty S, Thomas PS, Ramana KV, Vidyadhar KN, Chakradhar V. A Review of Phytochemical Studies of *Cassia* Species. *Journal of Natural Remedies*. 2002; 2(2): 102-120.

19. Dave H, Ledwani L. A Review on Anthraquinones Isolated from *Cassia* Species and Their Applications. *Indian Journal of Natural Products and Resources*. 2012; 3(3): 291-319.
20. Sundaramoorthy S, Gunasekaran S, Arunachalam S, Sathivelu M. A Phytopharmacological Review on *Cassia* Species. *Journal of Pharmaceutical Sciences and Research*. 2016; 8(5): 260-264.
21. Jothy SL, Torey A, Darah I, Choong YS, Saravanan D, Chen Y, Latha LY, Deivanai S, Sasidharan S. *Cassia spectabilis* (DC) Irwin et Barn: A Promising Traditional Herb in Health Improvement. *Molecules*. 2012; 17(9): 10292-10305. doi: 10.3390/molecules170910292.
22. Torey A, Sasidharan S, Yeng C, Latha LY. Standardization of *Cassia spectabilis* with Respect to Authenticity, Assay and Chemical Constituents Analysis. *Molecules*. 2010; 15(5): 3411-3420. doi: 10.3390/molecules15053411.
23. Deshpande HA, Bhalsing SR. Recent Advances in the Phytochemistry of Some Medicinally Important *Cassia* Species: A Review. *International Journal of Pharma Medicine and Biological Sciences*. 2013; 2(3): 60-78.
24. Ekasari W, Wahyuni TS, Arwaty H, Putri NT. Determination of Effective Dose of Antimalarial from *Cassia spectabilis* Leaf Ethanol Extract in *Plasmodium berghei*-Infected Mice. *African Journal of Infectious Diseases*. 2018; 12(1 Suppl): 110-115. doi: 10.2101/Ajid.12v1S.16
25. Suciati, Laili ER, Poerwanto D, Hapsari AP, Gifanda LZ, Rabgay K, Ekasari W, Ingkaninan K. Evaluation of Cholinesterase Inhibitory Activity of Six Indonesian *Cassia* Species. *Journal of Research Pharmacy*. 2020; 24(4): 472-478. doi:10.35333/jrp.2020.195
26. Ahmed SI, Hayat MQ, Tahir M, Mansoor Q, Ismail M, Keck K, Bates RB. Pharmacologically Active Flavonoids from the Anticancer, Antioxidant and Antimicrobial Extracts of *Cassia angustifolia* Vahl. *BMC Complementary and Alternative Medicine*. 2016; 16(1): 460. doi: 10.1186/s12906-016-1443-z.
27. Kolar FR, Gogi CL, Khudavand MM, Choudhari MS, Patil SB. Phytochemical and Antioxidant Properties of Some *Cassia* Species. *Natural Product Research*. 2017; 32(11): 1324-1328. doi: 10.1080/14786419.2017.1342085.
28. El-Hashash MM, Abdel-Gawad MM, El-Sayed MM, Sabry WA, Abdel-Hameed el-SS, Abdel-Lateef Eel-S. Antioxidant Properties of Methanolic Extracts of the Leaves of Seven Egyptian *Cassia* Species. *Acta Pharmaceutica*. 2010; 60(3): 361-367. doi: 10.2478/v10007-010-0030-y.
29. Arya V, Yadav JP. Comparative Assessment of Relative Antioxidant Activity of Sequential Leaf Extracts of *Cassia occidentalis* L and *Cassia tora* L. *Pharmacologyonline*. 2011; 1: 529-543.
30. Mehta JP, Parmar PH, Vadia SH, Patel MK, Tripathi CB. In-Vitro Antioxidant and In-Vivo Anti-Inflammatory Activities of Aerial Parts of *Cassia* Species. *Arabian Journal of Chemistry*. 2017; 10(supp. 2): S1654-S1662. doi:10.1016/j.arabjc.2013.06.010
31. Kumar RS, Narasingappa RB, Joshi CG, Girish TK, Prasada Rao UJ, Danagoudar A. Evaluation of *Cassia tora* Linn. against Oxidative Stress-Induced DNA and Cell Membrane Damage. *Journal of Pharmacy and BioAllied Sciences*. 2017; 9(1): 33-43. doi: 10.4103/0975-7406.206215
32. Singh VV, Jain J, Mishra AK. Determination of Antipyretic and Antioxidant Activity of *Cassia occidentalis* Linn Methanolic Seed Extract. *Pharmacognosy Journal*. 2017; 9(6): 913-916. doi:10.5530/pj.2017.6.143.
33. Herald TJ, Gadgil P, Tilley M. High-Throughput Micro Plate Assays for Screening Flavonoid Content and DPPH-Scavenging Activity in Sorghum Bran and Flour. *Journal of the Science of Food and Agriculture*. 2012; 92(11): 2326-2331. doi:10.1002/jsfa.5633.
34. Lee KJ, Oh YC, Cho WK, Ma JY. Antioxidant and Anti-Inflammatory Activity Determination of One Hundred Kinds of Pure Chemical Compounds Using Offline and Online Screening HPLC Assay. *Evidence-Based Complementary and Alternative Medicine*. 2015; 2015: 165457. doi:10.1155/2015/165457.
35. Ellman GL, Courtney KD, Andres V Jr, Featherstone RM. A New and Rapid Colorimetric Determination of Acetylcholinesterase Activity. *Biochemical Pharmacology*. 1961; 7(2): 88-95. doi:10.1016/0006-2952(61)90145-9
36. Ingkaninan K, Temkithawon P, Chuenchom K, Yuyaem T, Thongnoi W. Screening for Acetylcholinesterase Inhibitory Activity in Plants Used in Thai Traditional Rejuvenating and Neurotonic Remedies. *Journal of Ethnopharmacology*. 2003; 89(2-3): 261-264. doi:10.1016/j.jep.2003.08.008.
37. Suciati, Rabgay K, Fachrunniza Y, Saesong T, Hadi TA, Wahyuni TS, Widyawaruyanti A, Ingkaninan K. Enzyme Inhibitory Activities of Marine Sponges against Cholinesterase and 5 α -Reductase. *Malaysian Applied Biology Journal*. 2019; 48(3): 77-83.
38. Zhang Q, Zhang J, Shen J, Silva A, Dennis DA, Barrow CJ. A Simple 96-Well Microplate Method for Estimation of Total Polyphenol Content in Seaweeds. *Journal of Applied Phycology*. 2006; 18: 445-450. doi:10.1007/s10811-006-9048-4
39. Blois MS. Antioxidant Determinations by The Use of A Stable Free Radical. *Nature*. 1958; 181: 1199-1200. doi:10.1038/1811199a0
40. Molyneux P. The Use of The Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity. *Songklanakarin Journal of Science and Technology*. 2004; 26(2): 211-219.
41. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant Activity Applying an Improved ABTS Radical Cation Decolorization Assay. *Free Radical Biology and Medicine*. 1999; 26(9-10): 1231-1237. doi:10.1016/S0891-5849(98)00315-3
42. Aadesariya MK, Ram VR, Dave PN. Evaluation of Antioxidant Activities by Use of Various Extracts from *Abutilon pinnosum* and *Grewia tenax* Leaves in the Kachchh Region. *MOJ Food Processing and Technology* 2017; 5(1): 216-230. doi:10.15406/mojfpt.2017.05.00116
43. Shahidi F, Ambigaipalan P. Phenolics And Polyphenolics in Foods, Beverages and Spices: Antioxidant Activity and Health Effects – A Review. *Journal of Functional Foods*. 2015. 18(part B): 820-897. doi:10.1016/j.jff.2015.06.018
44. Wang TY, Li Q, Bi KS. Bioactive Flavonoids in Medicinal Plants: Structure, Activity and Biological Fate. *Asian Journal of Pharmaceutical Sciences*. 2018; 13(1):12-23. doi:10.1016/j.ajps.2017.08.004.
45. Dzoyem JP, Eloff JN. Anti-inflammatory, Anticholinesterase and Antioxidant Activity of Leaf Extracts of Twelve Plants Used Traditionally to Alleviate Pain and Inflammation in South Africa. *Journal of Ethnopharmacology*. 2015; 160:194-201. doi:10.1016/j.jep.2014.11.034
46. Irshad Md, Zafaryab Md., Singh M, Rizvi MM. Comparative Analysis of the Antioxidant Activity of *Cassia fistula* Extracts. *International Journal of Medicinal Chemistry*. 2012; 2012: 157125. doi: 10.1155/2012/157125.

In vitro Antioxidant and Anticholinesterase Activities of Extracts from the Leaves of Cassia moschata Kunth

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

8 %
INTERNET SOURCES

8 %
PUBLICATIONS

0 %
STUDENT PAPERS

PRIMARY SOURCES

1 www.mdpi.com 1 %
Internet Source

2 digiresearch.vut.ac.za 1 %
Internet Source

3 Monika Stojanova, Milena Pantić, Mitko Karadelev, Biljana Čuleva, Miomir Nikšić. "Antioxidant potential of extracts of three mushroom species collected from the Republic of North Macedonia", Journal of Food Processing and Preservation, 2020 1 %
Publication

4 psasir.upm.edu.my 1 %
Internet Source

5 Qing Zhang, Junzeng Zhang, Jingkai Shen, Angelica Silva, Dorothy A. Dennis, Colin J. Barrow. "A Simple 96-Well Microplate Method for Estimation of Total Polyphenol Content in Seaweeds", Journal of Applied Phycology, 2006 1 %
Publication

6	bmccomplementmedtherapies.biomedcentral.com Internet Source	1 %
7	estudogeral.sib.uc.pt Internet Source	<1 %
8	R. K. Toor, C.E. Lister, G.P. Savage. "Antioxidant activities of New Zealand-grown tomatoes", International Journal of Food Sciences and Nutrition, 2009 Publication	<1 %
9	Retno Widyowati, Neny Purwitasari, Rice Disi Oktarina, Wiwied Ekasari, Saarah Khairunnisa, Hsin-I. Chang. "Exploration of several plants from Baung Forest on bone formation cell models", Journal of Basic and Clinical Physiology and Pharmacology, 2021 Publication	<1 %
10	ipnacs2017.weebly.com Internet Source	<1 %
11	amb-express.springeropen.com Internet Source	<1 %
12	www.jbsdonline.com Internet Source	<1 %
13	Maher El-hashash, Mahfouz Abdel-Gawad, Mortada El-Sayed, Waffa Sabry, El-Sayed Abdel-Hameed, Ezzat Abdel-Lateef. "Antioxidant properties of methanolic extracts	<1 %

of the leaves of seven Egyptian Cassia species", Acta Pharmaceutica, 2010

Publication

14

S. Badami, K.P. Channabasavaraj. " .
Antioxidant Activity of Thirteen Medicinal
Plants of India's Western Ghats ",
Pharmaceutical Biology, 2008

Publication

15

Letricia Barbosa-Pereira, Alessandro
Guglielmetti, Giuseppe Zeppa. "Pulsed Electric
Field Assisted Extraction of Bioactive
Compounds from Cocoa Bean Shell and
Coffee Silverskin", Food and Bioprocess
Technology, 2018

Publication

16

www.acarindex.com

Internet Source

<1 %

17

www.ijcpr.org

Internet Source

<1 %

18

www.science.gov

Internet Source

<1 %

19

Nantaka Khorana. "Evaluation of a new lead
for acetylcholinesterase inhibition", Medicinal
Chemistry Research, 03/2009

Publication

<1 %

20

dspace.nwu.ac.za

Internet Source

<1 %

21	ijpsr.com Internet Source	<1 %
22	innovareacademics.in Internet Source	<1 %
23	mafiadoc.com Internet Source	<1 %
24	repository.up.ac.za Internet Source	<1 %
25	www.degruyter.com Internet Source	<1 %
26	PO-CHUEN CHAN, QINGSU XIA, PETER P. FU. "Leave Extract: Biological, Medicinal, and Toxicological Effects ", Journal of Environmental Science and Health, Part C, 2007 Publication	<1 %
27	"Plant and Human Health, Volume 3", Springer Science and Business Media LLC, 2019 Publication	<1 %
28	Aysegul Peksel, Inci Arisan, Refiye Yanardag. "Radical scavenging and anti-acetylcholinesterase activities of aqueous extract of wild pistachio (<i>Pistacia atlantica</i> Desf.) leaves", Food Science and Biotechnology, 2013 Publication	<1 %

29

Mun Fei Yam, Rusliza Basir, Mohd. Zaini Asmawi, Rosidah, Mariam Ahmad, Gabriel Akyirem Akowuah. " Antioxidant and Hepatoprotective Activities of . Ethanol Extract ", Pharmaceutical Biology, 2008

Publication

<1 %

30

İlhami GÜLÇİN, Arif Daştan. " Synthesis of dimeric phenol derivatives and determination of antioxidant and radical scavenging activities ", Journal of Enzyme Inhibition and Medicinal Chemistry, 2008

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

In vitro Antioxidant and Anticholinesterase Activities of Extracts from the Leaves of *Cassia moschata* Kunth

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
