

Age, neutrophil lymphocyte ratio, and radiographic assessment of the quantity of lung enema (RALE) score to predict in-hospital mortality in COVID-19 patients: a retrospective study

by Anggraini Dwi Sensusiaty

Submission date: 14-Sep-2021 12:02AM (UTC+0800)

Submission ID: 1647512393

File name: Age,_Neutrophyl.pdf (1.22M)

Word count: 6308

Character count: 32613



RESEARCH ARTICLE

REVISED Age, neutrophil lymphocyte ratio, and radiographic assessment of the quantity of lung edema (RALE) score to predict in-hospital mortality in COVID-19 patients: a retrospective study [version 2; peer review: 2 approved]

Anggraini Dwi Sensusiaty^{1,3}, Muhammad Amin^{2,4}, Nasronudin Nasronudin^{3,5}, Alfian Nur Rosyid^{3,4}, Nanda Aulya Ramadhan^{2,3}, Rofida Lathifah⁶, Eva Puspitasari², Ria Indah Wahyuningtyas², Erika Soebakti^{1,3}

¹Department of Radiology, Universitas Airlangga Hospital, Surabaya, East Java, 60115, Indonesia

²Department of Research, Planning and Development, Universitas Airlangga Hospital, Surabaya, East Java, 60115, Indonesia

³Medical Faculty, Universitas Airlangga, Surabaya, East Java, 60115, Indonesia

⁴Department of Pulmonology, Universitas Airlangga Hospital, Surabaya, East Java, 60115, Indonesia

⁵Department of Internal Medicine, Universitas Airlangga Hospital, Surabaya, East Java, 60115, Indonesia

⁶Department of School of Health Science, City, University of London, London, EC1V 0HB, UK

V2 First published: 30 Oct 2020, 9:1286
<https://doi.org/10.12688/f1000research.26723.1>
 Latest published: 06 Jan 2021, 9:1286
<https://doi.org/10.12688/f1000research.26723.2>

Abstract

Background: Available data suggest that case fatality rate of COVID-19 patients in Surabaya is higher than global cases. Thus, it is important to identify risk factors to prevent the mortality. This study aimed to assess the factors associated with hospital mortality of COVID-19 patients, and develop a prediction score based on these findings.

Methods: We analyzed 111 patients, who were diagnosed with COVID-19 based on reverse-transcriptase polymerase chain reaction. The following patient characteristics were obtained from records: age, gender, type of symptoms, onset of symptoms, neutrophil lymphocyte ratio (NLR), absolute lymphocyte count, chest x-ray abnormalities, lung involvement, type of lesion, radiographic assessment of the quantity of lung edema (RALE) score, and mortality. Data were analyzed using SPSS 25.0.

Results Multivariate analysis showed that age >50 years ($p=0.043$), NLR score >5.8 ($p=0.016$) and RALE score >2 ($p=0.002$) can predict the mortality of COVID-19 patients in the hospital. ROC curve analysis of the score ability to predict mortality showed an area under the curve of 0.794. The cut-off point is 4.5, with a sensitivity of 96.7% and specificity of 49.4% to predict the mortality of COVID-19 patient in the hospital.

Conclusions Age, NLR score and RALE score were associated with

Open Peer Review

Reviewer Status ✓ ✓

	Invited Reviewers	
	1	2
version 2 (revision) 06 Jan 2021		✓ report
version 1 30 Oct 2020	✓ report	? report

1. **Hamzaini Abdul Hamid**, Universiti Kebangsaan Malaysia Medical Centre (UKMMC), Kuala Lumpur, Malaysia


2. **Wiwien Heru Wiyono**, Universitas Indonesia, Jakarta, Indonesia

Any reports and responses or comments on the article can be found at the end of the article.

mortality of COVID-19 patients in the hospital and might be used as a predictor for mortality of COVID-19 patients in health care centre where radiologists are available. The prediction score may be useful for frontline physicians to effectively manage patients with a higher score to prevent mortality.


Keywords

COVID-19 mortality, NLR, RALE Score



25

This article is included in the [Disease Outbreaks gateway](#).



This article is included in the [Coronavirus collection](#).

Corresponding author: Anggraini Dwi Sensusiati (anggrainids.unair@gmail.com)

Author roles: **Sensusiati AD:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Review & Editing; **Amin M:** Project Administration, Resources, Supervision; **Nasronudin N:** Project Administration, Resources, Supervision; **Rosyid AN:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Resources; **Ramadhan NA:** Conceptualization, Formal Analysis, Investigation, Methodology, Resources, Validation, Writing – Review & Editing; **Lathifah R:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation; **Puspitasari E:** Data Curation, Investigation, Project Administration, Resources, Writing – Original Draft Preparation; **Wahyuningtyas RI:** Data Curation, Investigation, Project Administration, Resources, Writing – Original Draft Preparation; **Soebakti E:** Data Curation, Investigation, Methodology, Resources

Competing interests: No competing interests were disclosed.

Grant information: This article is supported by Airlangga University.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Sensusiati AD *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Sensusiati AD, Amin M, Nasronudin N *et al.* [Age, neutrophil lymphocyte ratio, and radiographic assessment of the quantity of lung edema \(RALE\) score to predict in-hospital mortality in COVID-19 patients: a retrospective study \[version 2; peer review: 2 approved\]](#) F1000Research 2021, 9:1286 <https://doi.org/10.12688/f1000research.26723.2>

First published: 30 Oct 2020, 9:1286 <https://doi.org/10.12688/f1000research.26723.1>

REVISED Amendments from Version 1

Abstract: we have added the sentence "might be used as a predictor for mortality of COVID-19 patients in health care centre where radiologists are available" on the conclusions section.

Introduction: we have added the sentences "Several factors may associate with the higher CFR such as low testing, low contact tracing and poor COVID-19 register (Sorci *et al.*, 2020). However, finding characteristics and risk factors to predict mortality is of utmost importance in order to prevent the mortality. It is also important to develop a prediction score to assess patients during early stages, when patients receive treatment at tertiary hospitals" on the second paragraph and "to assess the demographical characteristic, laboratory and radiological findings associated with COVID-19 mortality in hospital patients" on the last paragraph.

Discussion: we have added the sentences "Another study in Indonesia found a similar result where the average age of COVID-19 patients who died is 58.2 years old (Rozaliyani *et al.*, 2020)" and "This study also showed that the mortality risk is three times higher in people age ≥ 50 years. This finding is similar to a study in the United States, which found that the mortality risk of COVID-19 was 8.1 times higher among those ≥ 55 years compared with individuals ages < 54 years (Yanez *et al.*, 2020)." On the second paragraph. We also added term "RALE score" on the sixth paragraph.

Conclusions: we have added the sentence "It may be used as a predictor for mortality of COVID-19 patients in the health care centre where radiologists are available".

Any further responses from the reviewers can be found at the end of the article

Introduction

Cases of pneumonia of unknown etiology were first reported in the city of Wuhan, China, at the end of December 2019. After identification, the etiology of these cases was a new type of coronavirus (Wang *et al.*, 2020). The World Health Organization officially called this coronavirus disease 2019 (COVID-19) (WHO, 2020). COVID-19 is caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) virus (Gorbalenya *et al.*, 2020).

Since December 2019, more than 20 million people have been diagnosed with COVID-19, and more than 700,000 have died (Worldometers, 2020). In Indonesia, 128,776 people have been diagnosed with COVID-19, and 83,710 have died since March 2020 (Gugus Tugas Percepatan Penanganan COVID-19, 2020). The global case fatality rate (CFR) of COVID-19 in August was 3.7%. In Indonesia, the CFR is slightly higher (4.5%). Meanwhile, the CFR in Surabaya (the second biggest city in Indonesia) is almost twice that of Indonesia for the same period (8.9%). Several factors may associate with the higher CFR such as low testing, low contact tracing and poor COVID-19 register (Sorci *et al.*, 2020). However, finding characteristics and risk factors to predict mortality is of utmost importance in order to prevent the mortality. It is also important to develop a prediction score to assess patients during early stages, when patients receive treatment at tertiary hospitals.

Several studies have found risk factors related to COVID-19 mortality (Galloway *et al.*, 2020; Gupta *et al.*, 2020; Jalili *et al.*, 2020; Lippi & Plebani, 2020; Zhao *et al.*, 2020).

However, currently, there is no study that analyzes risk factors for COVID-19 mortality in hospitals for the Indonesian population. Existing prediction scores also based on complex laboratory findings, which are less feasible to be used in low resource health centers. Therefore, the purpose of this study was to assess the demographical characteristic, laboratory and radiological findings associated with COVID-19 mortality in hospital patients and develop a prediction score based on these findings.

Methods**Study design and setting**

This was a retrospective study that was conducted in the Emergency Department of Airlangga University Teaching Hospital. We included all the patients who fulfilled the inclusion and exclusion criteria, who were admitted for COVID-19 from March 13, 2020 to May 15, 2020. A total of 111 patients met the criteria and included in this study.

We collected data regarding clinical symptoms and the date of onset of symptoms from emergency department medical records.

Patients

The inclusion criteria for this retrospective study were: (1) patients with a chief complaint of one COVID-19-related symptom, including: fever, dry cough, tiredness, aches and pains, nasal congestion, headache, conjunctivitis, sore throat, diarrhea, loss of taste or smell, a rash on skin, or discoloration of fingers or toes (WHO, 2020); (2) confirmed SARS-CoV-2 infection by reverse-transcriptase polymerase chain reaction (RT-PCR) using nasopharyngeal and oropharyngeal specimens; (3) patients who underwent chest x-ray (CXR) on the day of admission to the hospital.

The following patients were excluded: (1) asymptomatic patients; (2) those with negative results for SARS-CoV-2 infection by RT-PCR.

All patients underwent CXR and laboratory examination on the day of admission to the emergency department.

Subgrouping of patients

Patients were divided into two groups based on the outcome of the patient: group I (discharged with negative results for SARS-CoV-2 infection by RT-PCR test); and group II (died).

Data collection

The following patient characteristics were obtained from the medical records: age, gender, type of symptoms, onset of symptoms, neutrophil lymphocyte ratio (NLR), absolute lymphocyte count (ALC), CRX abnormalities, lung involvement, type of lesion, radiographic assessment of the quantity of lung edema (RALE) score, and mortality.

Image acquisition and evaluation. Radiology data was collected from the radiology department. All the patients underwent an anteroposterior projection chest radiography at full inspiration where possible. The results were reviewed by two

radiologists (A.D.S, a radiologist with 25 years of experience and E.S, a radiologist with four years of experience) based on consensus.

CXRs were evaluated for the presence of pulmonary alterations, type of pulmonary alterations, and their distribution. CXR alterations that were found specifically in COVID-19 patients were defined according to the Fleischner Society's nomenclature, available in the Glossary of Terms for Thoracic Imaging (Hansell *et al.*, 2008), as follows:

- Reticular alteration: a reticular pattern is a collection of innumerable small linear opacities that, by summation, produce an appearance resembling a net
- Consolidation: as a homogeneous increase in pulmonary parenchymal attenuation that obscures the margins of vessels and airway walls
- Ground-glass opacity (GGO): an area of hazy increased lung opacity, usually extensive, within which margins of pulmonary vessels may be indistinct

The distribution of pulmonary alterations was classified as lung involvement unilateral (right/left) or bilateral. Other features, such as pleural effusion, were also recorded.

Radiograph scoring. A severity score was calculated to quantify the extent of the infection by adapting and simplifying the RALE score proposed by Warren *et al.* (2018). A score of 0-4 was assigned to each lung depending on the extent of involvement by consolidation or GGO (0 = no involvement; 1 = <25%; 2 = 25-50%; 3 = 50-75%; 4 = >75% involvement). The scores for each lung were summed to produce the final RALE score.

Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics Version 25.0. Continuous variables were expressed as mean \pm standard deviation values. The frequency of symptoms, laboratory findings and CXR findings was shown as the number of incidence and percentage per cluster of groups. The correlation between the patient characteristic, symptoms, laboratory findings and CXR findings with the outcome was analyzed by logistic regression. We also conducted the Hosmer-Lemeshow test to evaluate the goodness of fit of the scoring model and conducted receiver operating characteristic (ROC) analysis to evaluate the sensitivity and specificity of the model.

Ethical considerations

This study received ethical approval from the Ethical Committee of Airlangga University Teaching Hospital (approval number 147/KEP/2020). Consent from the participants was waived by the committee.

Results

Patient characteristics

A total of 111 COVID-19 patients were evaluated. In total, 72.9% (n=81) of patients were discharged from the hospital, while 27.1% (30) patients died during their hospitalization. There were 45 men (48.6%) and 47 women (51.4%). Average

patient age was 51 \pm 14.2 years old. The mean age for the patients who died was higher (55 \pm 12.8 years) compared to the patients who were discharged (48 \pm 14.8 years). Fever (32.6%) and shortness of breath (26%) were the most frequent symptoms for all patients. However, among the patients who died, shortness of breath was the most frequent symptom (40.7%). The mean NLLR score is higher on the patient who died (10.1 \pm 10.5) compared to patients who were discharged (5.3 \pm 5.4). In contrast, the mean ALC score was lower in patients who died (1130 \pm 252) compared to patients who were discharged (1349 \pm 702). Most of the patients showed GGO on their CXR (67.6%). Mean RALE score was higher in patients who died (5.3 \pm 2.5) compared to patients who were discharged (2.7 \pm 2.7) (Table 1).

Table 1. Patient characteristics.

Variables	All patients (n=111)		Discharged (n=81)		Died (n=30)	
	n	%	n	%	n	%
Mean age \pmSD	50 \pm 14.6		48 \pm 14.8		55 \pm 12.8	
Gender						
Men	54	48.6	39	48.1	15	50.0
Women	57	51.4	42	51.9	15	50.0
Clinical symptoms						
Fever	36	32.4	30	37.0	6	20.0
Cough	22	19.8	20	24.7	2	6.7
Fatigue	11	9.9	8	9.9	3	10.0
Shortness of breath	29	26.1	16	19.8	13	43.3
Headache	3	2.7	0	0.0	3	10.0
Abdominal pain	3	2.7	2	2.5	1	3.3
Chest pain	2	1.8	1	1.2	1	3.3
Vomiting	1	0.9	1	1.2	0	0.0
Hemoptysis	1	0.9	1	1.2	0	0.0
Nausea	2	1.8	2	2.5	0	0.0
Unconscious	1	0.9	0	0.0	1	3.3
Laboratory findings						
Mean NLR \pm SD	6.6 \pm 7.2		5.3 \pm 5.4		10.1 \pm 10.5	
Mean ALC \pm SD	1290 \pm 664		1349 \pm 702		1130 \pm 252	
Type of lesion						
GGO	75	67.6	53	65.4	22	73.3
Reticular alteration	10	9.0	6	7.4	4	13.3
Fibrotic	1	0.9	0	0.0	1	3.3
Pleural effusion	6	5.4	4	4.9	2	6.7
Mean RALE score \pmSD	3.4 \pm 2.8		2.7 \pm 2.7		5.3 \pm 2.5	

Risk factors of mortality

From **Table 2**, no significant relationship was observed between the outcome of the patients based on gender, type of symptoms, onset of symptoms, ALC score, x-ray abnormalities, and type of lesion. This suggested that those variables have no predictive value for mortality outcome for COVID-19

Table 2. Risk factors of COVID-19 hospital mortality.

Variables	Discharged (n=81)		Died (n=30)		p-Value
	n	%	n	%	
Age					0.009
≤50 years	47	58.0	9	30.0	
>50 years	34	42.0	21	70.0	
Gender					0.862
Male	39	48.1	15	50.0	
Female	42	51.9	15	50.0	
Type of symptoms					0.548
Respiratory	43	53.1	14	46.7	
Non-respiratory	36	44.4	16	53.3	
Onset of the symptoms					0.473
≤ 3 days	37	45.7	16	53.3	
>3 days	44	54.3	14	46.7	
Laboratory findings					
NLR Score					0.002
≤5.8	60	74.1	12	40.0	
>5.8	21	25.9	18	60.0	
ALC Score					0.093
≤1000	24	29.6	14	46.7	
>1000	57	70.4	16	53.3	
CXR abnormalities					0.055
Negative chest radiography	22	27.2	3	10.0	
Positive chest radiography	59	72.8	27	90.0	
Lung involvement					0.007
Unilateral	24	29.6	4	13.3	
Bilateral	35	43.2	23	76.7	
Type of lesion					0.430
GGO	53	65.4	22	73.3	
Non-GGO	28	34.6	8	26.7	
RALE Score					0.001
≤2	45	55.6	4	13.3	
>2	36	44.4	26	86.7	

patients. There is a significant relationship between age, NLR score, lung involvement and RALE score towards the outcome of the patients (p -value <0.05). All of these variables were analyzed using logistic regression, which revealed there are three variables with $p<0.05$: age, NLR score and RALE score (**Table 3**).

Based on the odds ratio (OR) value of each variable, the mortality risk for patients who were of an older age is 2.787 times higher than patients who were of a younger age. A higher NLR score has a 3.246 times higher mortality risk than patients who have lower NLR score. Mortality risk for patients who have a higher RALE score is 6.826 times higher than patients who have lower RALE score. It should be noted that three patients died while showing no abnormalities on their CXR.

Prediction of mortality

We form a scoring model based on the OR of each variable, which can be seen in **Table 4**. If the variable were present, we gave the value of “1”, and if absent we gave the value of “0”. We also performed the Hosmer-Lemeshow test and concluded that the model is fit ($p=0.802$). Based on our scoring model, each score from 111 patients was calculated. The percentage of mortality was then calculated for each score. **Table 5** shows that high mortality (>60%) was seen for a total score of 13, while low mortality (<10%) was seen for a total score ≤ 3 .

The total score was then analyzed with ROC analysis to predict the sensitivity and specificity for the probability of mortality of COVID-19 patients in the hospital. **Figure 1** shows that based on the ROC curve, the area under the curve (AUC) of the score is 0.794. The score has a cut off point of 4.5, with a sensitivity of 96.7% and specificity of 49.4%, to predict the mortality of COVID-19 patients in the hospital.

Table 3. Logistic regression.

	p-Value	OR	95% CI
Age	0.043	2.787	1.032-7.531
NLR Score	0.016	3.246	1.241-8.487
Lung involvement	0.965	1.024	0.356-2.946
RALE Score	0.002	6.826	2.076-22.444

Table 4. Scoring model to predict the mortality of the patient.

Age >50 years	3
NLR Score >5.8	3
RALE Score >2	7

Table 5. Probability of mortality based on the total score.

Total Score	Probability of mortality
0	0%
3	6.7%
6	37.5%
7	26.7%
10	38.7%
13	62.5%




Figure 1. ROC curve analysis for the scoring model to predict the mortality in COVID-19 patients.

Discussion

This is the first study to analyze the radiologic and laboratory findings of COVID-19 patients in Indonesia. The main result of our study was that age, NLR score and RALE score have a strong relationship for mortality risk of COVID-19 patients in the hospital.

This study found that the average age of COVID-19 patients who died is 55 ± 12.8 years. Another study in Indonesia found a similar result where the average age of COVID-19 patients who died is 58.2 years old (Rozaliyani *et al.*, 2020). This finding is younger than a study in Iraq, who found that the

average age of COVID-19 patients who died in hospital is 67.49 ± 15.28 years (Jalili *et al.*, 2020). The younger average age of the patients who died might be due to the co-morbidities that also develop in the relatively younger age of the Indonesian population, especially for type 2 diabetes (Arifin *et al.*, 2019). This study also showed that the mortality risk is three times higher in people age ≥ 50 years. This finding is similar to a study in the United States, which found that the mortality risk of COVID-19 was 8.1 times higher among those ≥ 55 years old compared with individuals ages < 54 years (Yanez *et al.*, 2020).

Based on our findings, there is no correlation between gender and patient mortality. The result is different from studies in the United States and Iran, which conclude that men has a higher risk of death than women (Gupta *et al.*, 2020; Jalili *et al.*, 2020). This could be due to the characteristic of the co-morbidities of the Indonesian population, which affect both men and women at an almost similar number (Kemenkes, 2019). We also found that there is no correlation between type and onset of symptoms to the outcome of the patients.

There is a correlation between NLR score and the patient outcome. Previously, it was concluded that higher NLR score related to the severe illness of the patients and could be an early identification for patient assessment in the early stage for intensive care unit (Liu *et al.*, 2020; Yang *et al.*, 2020). Our study showed that the higher NLR score (> 5.8) would increase the risk of mortality by three times. In contrast, we found no correlation between ALC score and patient mortality in the hospital. However, various other research shows that low ALC score is related to disease severity for patients admitted to the hospital for COVID-19 (Wagner *et al.*, 2020) and a higher risk of mortality (Lippi & Plebani, 2020).

This study showed that there is no correlation for CXR abnormalities, lung involvement and type of lesion to COVID-19 mortality in the hospital. Interestingly, we also found that three patients (10%) who died showed normal CXR. This could be due to the lower sensitivity of CXR to detect lung lesions compare to CT scan. However, it also shows that COVID-19 could become severe rapidly even though patients have normal CXR on the day of the admission.

Interestingly, we found that RALE score is correlated with patient mortality, and a higher score (> 2) will increase mortality risk by seven times. This finding is consistent with another study in the United Kingdom with a larger sample size, which showed that higher radiological severity score (> 3) increased the incidence of critical admission or death (Galloway *et al.*, 2020). RALE score in CXR could become a key predictor of mortality because of the simplicity of modality and is broadly available across health care provider.

The prediction score developed in this study demonstrated good accuracy (AUC of 79.4) to predict the discharged probability of COVID-19 patients in the hospital with only three

simple parameters. Another score showed good accuracy (AUC of .83) with more complicated parameters, such as health failure, procalcitonin, lactate dehydrogenase, chronic obstructive pulmonary disease, pulse oxygen saturation, heart rate, and age (Zhao *et al.*, 2020). This suggests that our score can be used in a broader setting such as locations with low resource of health care service.

Limitations

This study only analyzes patient characteristics, and laboratory and radiology findings related to COVID-19 mortality in the hospital. We do not evaluate the co-morbidities of the patient and the hospital setting, such as the location of the treatment where the patient died (emergency room, intensive care, low care), or the resource (facilities, human resource, intervention).

References

12

- Arifin B, van Asselt ADI, Setiawan D, *et al.*: **Diabetes distress in Indonesian patients with type 2 diabetes: a comparison between primary and tertiary care.** *BMC Health Serv Res.* 2019; **19**(1): 773.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Galloway JB, Norton S, Barker RD, *et al.*: **A clinical risk score to identify patients with COVID-19 at high risk of critical care admission or death: An observational cohort study.** *J Infect.* 2020; **81**(2): 282–288.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gorbalenya AE, Baker SC, Baric RS, *et al.*: **The species *Severe acute respiratory syndrome-related coronavirus*: classifying 2019-nCoV and naming it SARS-CoV-2.** *Nat Microbiol.* 2020; **5**(4): 536–544.
[Publisher Full Text](#)
- Gugus Tugas Percepatan Penanganan COVID-19: **Peta Sebara.** 2020.
[Reference Source](#)
- Gupta S, Hayek SS, Wang W, *et al.*: **Factors Associated With Death in Critically Ill Patients With Coronavirus Disease 2019 in the US.** *JAMA Intern Med.* 2020; e203596.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hansell DM, Bankier AA, MacMahon H, *et al.*: **Fleischner Society: Glossary of Terms for Thoracic Imaging.** *Radiology.* 2008; **246**(3): 697–722.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jalili M, Payandemehr P, Saghaei A: **Characteristics and Mortality of Hospitalized Patients With COVID-19 in Iran: A National Retrospective Cohort Study.** *Ann Intern Med.* 2020; M20–2911.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kemenkes RI: **Hasil Utama Riskesdas 2018.** 2019.
[Reference Source](#)
- Lippi G, Plebani M: **Laboratory abnormalities in patients with COVID-19 infection.** *Clin Chem Lab Med.* 2020; **58**(7): 1131–1134.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liu J, Liu Y, Xiang P, *et al.*: **Neutrophil-to-lymphocyte ratio predicts critical illness patients with 2019 coronavirus disease in the early stage.** *J Transl Med.* 2020; **18**(1): 206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rozaliyani A, Savitri AI, Setianingrum F, *et al.*: **Factors Associated with Death**

Conclusions

Age, NLR score and RALE score were associated with mortality of COVID-19 patients in our hospital setting in an Indonesian population. It may be used as a predictor for mortality of COVID-19 patients in the health care centre where radiologists are available. The prediction score may be useful for physicians to determine the mortality risk of a patient with COVID-19.

Data availability

Underlying data

Figshare: datasheet f1000research 2909.xlsx, <https://doi.org/10.6084/m9.figshare.13017899.v1> (Sensusiati, 2020).

Data are available under the terms of the [Creative Commons Attribution 4.0 International license \(CC-BY 4.0\)](#).

in COVID-19 Patients in Jakarta, Indonesia: An Epidemiological Study. *Acta Med Indones.* 2020; **52**(3): 246–254.
[PubMed Abstract](#)

Sensusiati AD: **datasheet f1000research 2909.xlsx.** figshare. Dataset. 2020.
<http://www.doi.org/10.6084/m9.figshare.13017899.v1>

Sorci G, Falvre B, Morand S: **Explaining among-country variation in COVID-19 case fatality rate.** *Sci Rep.* 2020; **10**(1): 18909.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wagner J, DuPont A, Larson S, *et al.*: **Absolute lymphocyte count is a prognostic marker in Covid-19: A retrospective cohort review.** *Int J Lab Hematol.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wang C, Horby PW, Hayden FG, *et al.*: **A novel coronavirus outbreak of global health concern.** *Lancet.* 2020; **395**(10223): 470–473.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Warren MA, Zhao Z, Koyama T, *et al.*: **Severity scoring of lung oedema on the chest radiograph is associated with clinical outcomes in ARDS.** *Thorax.* 2018; **73**(9): 840–846.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Worldometers: **COVID-19 Coronavirus Pandemic.** 2020.
[Reference Source](#)

World Health Organization: **Statement on the Second Meeting of the International Health Regulations, Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV).** 2020.
[Reference Source](#)

Yanez ND, Weiss NS, Romand JA, *et al.*: **COVID-19 mortality risk for older men and women.** *BMC Public Health.* 2020; **20**(1): 1742.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yang AP, Liu JP, Tao WQ, *et al.*: **The diagnostic and predictive role of NLR d-NLR and PLR in COVID-19 patients.** *Int Immunopharmacol.* 2020; **84**: 106504.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhao Z, Chen A, Hou W, *et al.*: **Prediction model and risk scores of ICU admission and mortality in COVID-19.** *PLoS One.* 2020; **15**(7): e0236618.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 25 January 2021

<https://doi.org/10.5256/f1000research.33024.r76665>

© 2021 Wiyono W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Wiwien Heru Wiyono

Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

I have read the revised version and I agree to all revisions so I approved this article.

8

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Pulmonary medicine

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 20 November 2020

1

<https://doi.org/10.5256/f1000research.29506.r74043>

© 2020 Wiyono W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



22

Wiwien Heru Wiyono

Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

There are some statements in this paper that are not so accurate.

Introduction:

My comment:

To reduce the mortality rate in some hospitals in Surabaya is not by finding characteristics and risk factors to predict mortality. As I know at that time tracing in Surabaya still low in subjects, so the CFR in this city is still higher than Jakarta and the mean CFR in Indonesia.

"the purpose of this study was to assess the factors associated with COVID-19 mortality in hospital patients and develop a prediction score based on these findings."

My comment:

To assess the factors associated with mortality even to develop prediction score should use all factors. But this study actually only uses some parameters not include commorbidities.

Discussion:

The statement:

1. "this finding is similar to a study in the US, which found that an older age (> 60 years) is associated with a higher risk of death is not so accurate, because:"

My comments:

1. The difference between mean ages of >50 and >60 is so significant.
2. Criteria for elder people in Indonesia is >60 years old.
3. The younger average of the patients in this study is not representative of Indonesia Covid patients.

The statement :

1. "CXR could become a key predictor of mortality because....."

My comment:

1. This statement is over confident and is in contradiction with the statement: "This study showed that there is no correlation for CXR abnormalities....."

Conclusion:

The statement:

1. "It may be used as a predictor for mortality of COVID 19 patients in low health care resource setting."

My comment :

1. It is not so true. The RALE score should be made by radiologists who have many experiences, so it only applicable in a referral hospital setting.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Pulmonary medicine

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 21 Dec 2020

Anggraini Dwi Sensusiati, Airlangga University Teaching Hospital, Surabaya, Indonesia

The authors thank the Honorable Reviewer 2 for the excellent review of the paper and in-depth suggestions made to improve the quality of the paper. Followings are the responses to the suggestions of the Honorable Reviewer. The valuable suggestions and corrections are incorporated cautiously, and the paper is made clearer while revising the manuscript.

Introduction:

Comment:

To reduce the mortality rate in some hospitals in Surabaya is not by finding characteristics and risk factors to predict mortality. As I know at that time tracing in Surabaya still low in subjects, so the CFR in this city is still higher than Jakarta and the mean CFR in Indonesia. *We acknowledged the possible cause of high fatality rate and added the sentence "Several factors may associate with the higher CFR such as low testing, low contact tracing and poor COVID-19 register (Sorci *et al.*, 2020). However, finding characteristics and risk factors to predict mortality is of utmost importance in order to prevent the mortality."

"the purpose of this study was to assess the factors associated with COVID-19 mortality in hospital patients and develop a prediction score based on these findings."

Comment:

To assess the factors associated with mortality even to develop prediction score should use all factors. But this study actually only uses some parameters not include commorbidities. *We acknowledged the study limitation which did not evaluate the commorbidities in the limitations section. We focused the research on the demographical characteristic, laboratory and radiological findings associated with COVID-19 mortality in hospital patients and develop a prediction score based on these findings.

Discussion:

The statement:

"this finding is similar to a study in the US, which found that an older age (> 60 years) is associated with a higher risk of death is not so accurate, because:"

Comments:

1. The difference between mean ages of >50 and >60 is so significant.
2. Criteria for elder people in Indonesia is >60 years old.
3. The younger average of the patients in this study is not representative of Indonesia Covid patients.

*We found similar finding in another research in Indonesia where the average age of COVID-19 patients who died is 58.2 years old (Rozaliyani *et al.*, 2020). We also found that this finding is similar to a study in the United States, which found that the mortality risk of COVID-19 was 8.1 times higher among those ≥ 55 years compared with individuals ages <54 years (Yanez *et al.*, 2020).

The statement :

"CXR could become a key predictor of mortality because....."

Comment:

1. This statement is over confident and is in contradiction with the statement: "This study showed that there is no correlation for CXR abnormalities....."

*We meant that the RALE score from CXR might become a key predictor

Conclusion:

The statement:

"It may be used as a predictor for mortality of COVID 19 patients in low health care resource setting."

Comment:

1. It is not so true. The RALE score should be made by radiologists who have many experiences, so it only applicable in a referral hospital setting.

*We changed the "low health care resource setting" into "health care centre where radiologists are available."

We thank Prof. Wiwien for raising these points which we feel have improved the paper considerably.

Sensusiati AD *et al.*

31

Competing Interests: No competing interests were disclosed.

Reviewer Report 20 November 2020

<https://doi.org/10.5256/f1000research.29506.r74045>

10

© 2020 Hamid H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Hamzaini Abdul Hamid**

Department of Radiology, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre (UKMMC), Kuala Lumpur, Malaysia

Congratulation and I think the paper has been presented very well. The statistical analysis and tools were done appropriately.

This paper definitely will contribute to the management of COVID 19 patients i.e. in predicting which patient may deteriorate and need to be monitor closely.

Although in this paper there is no correlation for CXR abnormalities, lung involvement and type of lesion to COVID-19 mortality. However, I think all the above CXR changes would also contribute collectively to the RALE score.

Overall, well done and congratulation to the author. Just a minor error in spelling.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: General Radiology, Paediatric Radiology

1

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research

Age, neutrophil lymphocyte ratio, and radiographic assessment of the quantity of lung enema (RALE) score to predict in-hospital mortality in COVID-19 patients: a retrospective study

ORIGINALITY REPORT

19%

SIMILARITY INDEX

10%

INTERNET SOURCES

19%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Sunil Shrestha, Subish Palaian. "Respiratory concerns of gabapentin and pregabalin: What does it mean to the pharmacovigilance systems in developing countries?", F1000Research, 2021
Publication 2%
- 2** Salima Meherali, Mehnaz Rehmani, Arzoo Rafiq, Neelam Punjani, Helen Vallianatos, Claudia Romeu.
"Empowering young immigrant girls in Canada through the 'Girls' Voices' Curriculum: A qualitative study", F1000Research, 2021
Publication 1%
- 3** Sergio Giuseppe Vancheri, Giovanni Savietto, Francesco Ballati, Alessia Maggi et al.
"Radiographic findings in 240 patients with COVID-19 pneumonia: time-dependence after the onset of symptoms", European Radiology, 2020 1%

4

pubs.rsna.org

Internet Source

1 %

5

G.V. Ishara Geelaka Bandara Jayarathna, G.K. Chaminda Jayalath, Ramya Pathiraja. "Case Report: severe hemoperitoneum due to spontaneous rupture of uterine vessels during the second trimester of pregnancy", F1000Research, 2021

Publication

1 %

6

Giovanni Savietto, Sergio Vancheri, Francesco Ballati, Alessia Maggi et al. "Radiographic findings in 240 patients with COVID-19 pneumonia", Research Square, 2020

Publication

1 %

7

global.oup.com

Internet Source

1 %

8

Charles Murphy, Danit Arad. "Case Report: Hepatopulmonary syndrome as the first clinical manifestation of cirrhosis in a patient with underlying chronic lung disease", F1000Research, 2019

Publication

1 %

9

Huan Han, Linlin Xie, Rui Liu, Jie Yang, Fang Liu, Kailang Wu, Lang Chen, Wei Hou, Yong Feng, Chengliang Zhu. "Analysis of heart injury laboratory parameters in 273 COVID-19

1 %

patients in one hospital in Wuhan, China",
Journal of Medical Virology, 2020

Publication

10

Million Abraha, Shiden Solomon. "Case Report: A rare cause of vaginal bleeding at Keren Zonal Referral Hospital, Eritrea", F1000Research, 2019

Publication

11

Alka Singh, Hemant Saxena, Jitendra Nath Rai. "Adaptive Spline Based PLL for Synchronization and Power Quality Improvement in Distribution System", IET Generation, Transmission & Distribution, 2020

Publication

12

Abraham Kebede Wolde. "Diabetic Distress Among Diabetic Patients in the Amhara Regional State, Ethiopia", International Quarterly of Community Health Education, 2021

Publication

13

N. David Yanez, Noel S. Weiss, Jacques-André Romand, Miriam M. Treggiari. "COVID-19 mortality risk for older men and women", BMC Public Health, 2020

Publication

14

Jie Xu, Wenwei Xiao, Xuan Liang, Li Shi, Peihua Zhang, Ying Wang, Yadong Wang, Haiyan Yang. "A meta-analysis on the risk factors

1 %

1 %

1 %

<1 %

<1 %

adjusted association between cardiovascular disease and COVID-19 severity", BMC Public Health, 2021

Publication

15

www.garda.com

Internet Source

<1 %

16

journals.plos.org

Internet Source

<1 %

17

"Abstracts", *Respirology*, 11/2011

Publication

<1 %

18

"Poster session 2", EP Europace, 2019

Publication

<1 %

19

Mirjam Groger, Hannah S. Fischer, Luzia Veletzky, Albert Lalremruata, Michael Ramharter. "A systematic review of the clinical presentation, treatment and relapse characteristics of human *Plasmodium ovale* malaria", *Malaria Journal*, 2017

Publication

<1 %

20

François M. Beloncle, Bertrand Pavlovsky, Christophe Desprez, Nicolas Fage et al. "Recruitability and effect of PEEP in SARS-Cov-2-associated acute respiratory distress syndrome", *Annals of Intensive Care*, 2020

Publication

<1 %

21

Brandon D Pickett, Sheena Talma, Jessica R Glass, Daniel Ence, Paul D Cowley, Perry G

<1 %

Ridge, John SK Kauwe. " Genome Assembly of the Roundjaw Bonefish (), a Vulnerable Circumtropical Sportfish ", Cold Spring Harbor Laboratory, 2021

Publication

22

"2-C2: Pulmonary Circulation", Respirology, 2013

Publication

<1 %

23

"Poster Presentation - Asthma", Respirology, 2014

Publication

<1 %

24

bmccomplementalalternmed.biomedcentral.com

Internet Source

<1 %

25

Stephen Senn. "Misunderstanding publication bias: editors are not blameless after all ", F1000Research, 2012

Publication

<1 %

26

ideas.repec.org

Internet Source

<1 %

27

ir.ymlib.yonsei.ac.kr

Internet Source

<1 %

28

m.f1000research.com

Internet Source

<1 %

29

www.degruyter.com

Internet Source

<1 %

30

Alok Atreya, Ritesh G. Menezes, Ashal Timalisina, Geeta Bashyal, Lokaratna Gyawali, Sushila Gyawali. "Case Report: Penetrating Thoracic Trauma by A Gunshot Involving the Heart", F1000Research, 2021

Publication

<1 %

31

Freedman, Jane E, Bahadir Ercan, Kristine M Morin, Ching-Ti Liu, Lulufer Tamer, Lokman Ayaz, Mehmet Kanadasi, Dilek Cicek, Ali Ihsan Seyhan, Rabia Eker Akilli, Celalettin Camci, Beyhan Cengiz, Serdar Oztuzcu, and Kahraman Tanriverdi. "The distribution of circulating microRNA and their relation to coronary disease", F1000 Research, 2012.

Publication

<1 %

32

Hamed Hosseini, Anahita Sadeghi, Payam Tabarsi, Azin Etemadimanesh et al. "Another step toward final call on Remdesivir efficacy as a treatment for hospitalized COVID-19 patients: a multicenter open-label trial", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

33

Natalie Marte, John Kileci, M. Faisal Siddiqi, Martha Catalina Morales Alvarez, Risako Shirane, Young Im Lee. "CORRELATION OF RADIOGRAPHIC ASSESSMENT OF THE QUANTITY OF LUNG EDEMA (RALE) SCORES AND ARDS SEVERITY", Chest, 2019

<1 %

34

onlinelibrary.wiley.com

Internet Source

<1 %

35

"Coronavirus Disease 2019 (COVID-19)",
Springer Science and Business Media LLC,
2020

Publication

<1 %

36

Madone Mandina, Jean-Robert Makulo, Roger
Wumba, Ben Bepouka et al. "Epidemiological,
clinical Characteristics and mortality of
patients Infected with SARS-CoV-2 Admitted of
Kinshasa University Hospital, Democratic
Republic of the Congo from March 24, 2020 to
January 30, 2021: Two waves, two faces?",
Cold Spring Harbor Laboratory, 2021

Publication

<1 %

37

Surabhi Madan, Manish Rana, Rohan Gajjar,
Nitesh Shah et al. "Evaluating the Efficacy of
Tocilizumab in Moderate to Severe COVID-19
with Progressive Illness despite Steroids:
Identifying the Optimal Timing of its
Administration in C3G study", Cold Spring
Harbor Laboratory, 2020

Publication

<1 %

38

en.wikipedia.org

Internet Source

<1 %

39

www.cambridge.org

Internet Source

<1 %

40

"Clinical Synopsis of COVID-19", Springer Science and Business Media LLC, 2020

Publication

<1 %

41

&NA;, . "European Society of Thoracic Imaging : Abstracts From the 2012 20th Anniversary Congress, London, UK June 22â€"24, 2012", Journal of Thoracic Imaging, 2012.

Publication

<1 %

42

Deng Pan, Dandan Cheng, Yiwei Cao, Chuan Hu, Fenglin Zou, Wencheng Yu, Tao Xu. "A Predicting Nomogram for Mortality in Patients With COVID-19", Frontiers in Public Health, 2020

Publication

<1 %

43

Jason Neidleman, Xiaoyu Luo, Ashley F. George, Matthew McGregor et al. "Distinctive features of SARS-CoV-2-specific T cells predict recovery from severe COVID-19", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

44

Jie Xu, Wenwei Xiao, Xuan Liang, Li Shi, Peihua Zhang, Ying Wang, Yadong Wang, Haiyan Yang. "A meta-analysis on the risk factors adjusted association between cardiovascular

<1 %

disease and COVID-19 severity", Research Square Platform LLC, 2021

Publication

45

Uluhan Sili, Pinar Ay, Ahmet Topuzoglu, Huseyin Bilgin et al. "Factors associated with progression to critical illness in 28 days among COVID-19 patients: results from a tertiary care hospital in Istanbul, Turkey", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

46

Veronika Seebacher, Stefanie Aust, David D'Andrea, Christoph Grimm et al. "Development of a tool for prediction of ovarian cancer in patients with adnexal masses: Value of plasma fibrinogen", PLOS ONE, 2017

Publication

<1 %

47

Yoshiaki Kinoshita, Hiroshi Ishii, Hisako Kushima, Kentaro Watanabe, Masaki Fujita. "High-dose steroid therapy for acute respiratory distress syndrome lacking common risk factors: predictors of outcome", Acute Medicine & Surgery, 2018

Publication

<1 %

48

www.researchsquare.com

Internet Source

<1 %

49

Chi Zhang, Ling Qin, Kang Li, Qi Wang et al. "A Novel Scoring System for Prediction of

<1 %

Disease Severity in COVID-19", *Frontiers in Cellular and Infection Microbiology*, 2020

Publication

50

Louisa Fadjri Kusuma Wardhani, Ivana Purnama Dewi, Denny Suwanto, Meity Ardiana. "Case Report: Heparin-induced thrombocytopenia during COVID-19 outbreak: the importance of scoring system in differentiating with sepsis-induced coagulopathy", *F1000Research*, 2021

Publication

<1 %

51

Muhamad Erza Aminanto, Bahrul Ilmi Nasution, Andi Sulasikin, Yudhistira Nugraha, Juan Kanggrawan, Alex L. Suherman. "Contributing Clinical Attributes to COVID-19 Mortality in Jakarta: Machine Learning Study", 2021 9th International Conference on Information and Communication Technology (ICoICT), 2021

Publication

<1 %

52

Tom L. Schmidt, Nicholas H. Barton, Gordana Rašić, Andrew P. Turley, Brian L. Montgomery, Inaki Iturbe-Ormaetxe, Peter E. Cook, Peter A. Ryan, Scott A. Ritchie, Ary A. Hoffmann, Scott L. O'Neill, Michael Turelli. "Local introduction and heterogeneous spatial spread of dengue-suppressing *Wolbachia* through an urban

<1 %

population of Aedes aegypti", PLOS Biology, 2017

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Age, neutrophil lymphocyte ratio, and radiographic assessment of the quantity of lung enema (RALE) score to predict in-hospital mortality in COVID-19 patients: a retrospective study

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13
