

PAIN REDUCTION AFTER WET AND DRY CUPPING THERAPIES ROLES OF INTEGRIN AND OPIOID RECEPTOR IN ANIMAL MODELS

by Imam Subadi

Submission date: 08-Sep-2022 12:15PM (UTC+0800)

Submission ID: 1894893201

File name: APIES_ROLES_OF_INTEGRIN_AND_OPIOID_RECEPTOR_IN_ANIMAL_MODELS.pdf (168.57K)

Word count: 2457

Character count: 13835

PAIN REDUCTION AFTER WET AND DRY CUPPING THERAPIES: ROLES OF $\alpha 2\beta 1$ INTEGRIN AND μ -OPIOID RECEPTOR IN ANIMAL MODELS

¹Imam Subadi, ²BoyaNugraha, ³AbdulhadiSuwandi, *¹Nur Sulastr¹*, ⁴Imam Susilo

ABSTRACT---Background: Wet cupping therapy (WCT) and dry cupping therapy (DCT) are complementary therapies in pain management. Stretching keratinocyte by cupping seems to stimulate the expression of $\alpha 2\beta 1$ -integrin. Its benefit in pain reduction could be mediated by the expression of μ -opioid receptor (MOR).

Objective: This study aims to determine pain thresholds, $\alpha 2\beta 1$ -integrin and MOR after WCT and DCT.

Method: Thirty-two adult male Wistar rats were divided into negative control, positive control, DCT, and WCT groups. In DCT group and WCT group, cupping was conducted at left and right paralumbar region 48h after CFA injection. Hot plate test was conducted 24h after therapy to assess pain threshold and immunohistochemistry were conducted from the skin for $\alpha 2\beta 1$ -integrin and the spinal cord for MOR.

Result: WCT showed the highest expression of $\alpha 2\beta 1$ -integrin in the keratinocytes and MOR in the lamina II spinal cord than the other groups. On immunohistochemical examination, the expression of $\alpha 2\beta 1$ integrin showed positive cells (keratinocytes) with brown stain on dry cupping and wet cupping. Significant differences were shown between negative control group showed significant differences with positive control group, dry cupping therapy group, and wet cupping therapy group.

Conclusion: WCT is more effective than DCT in reducing pain. The benefit of both therapies in reducing pain in rats could be mediated by the expression of $\alpha 2\beta 1$ -integrin and MOR.

Keywords---Dry cupping, pain reduction, wet cupping,

I. INTRODUCTION

Cupping therapy is one of the complementary and alternative therapies that is widely used to reduce pain (Kim *et al.*, 2011). Previous study showed that cupping therapy can be used for acute and chronic pain (Cao *et al.*, 2004). In Indonesia there are two methods of cupping therapy, the dry cupping therapy and wet cupping therapy. Dry cupping therapy cups the skin without bleeding. In wet cupping therapy, the skin is injured and cupped, resulting in blood loss (Arslan, Gökğöz and Dane, 2016). The difference in the effectiveness of each cupping therapy method is not

¹Department of Physical Medicine and Rehabilitation, Faculty of Medicine/Universitas Airlangga Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

²Department of Rehabilitation Medicine, Hannover Medical School, Hannover 30625, Germany

³Institute of Medical Microbiology and Hospital Epidemiology, Hannover 30625 Medical School, Hannover, Germany

⁴Department of Pathology Anatomy, Faculty of Medicine, Universitas Airlangga, Surabaya 60131, Indonesia

*Corresponding Author: Nur Sulastr¹, MD

Department of Physical Medicine and Rehabilitation, Hospital of Airlangga University, Campus C Mulyorejo, Surabaya 60115, Indonesia

Phone: +62315916290

E-mail: nuri.rehab@gmail.com

widely known. In both cupping methods negative pressure is applied which causes the skin to stretch. Stretched cells cause stimulation of mechanoreceptors of degenerin/epithelial sodium channel (DEG/ENaC), acid-sensing ion channels ASICs, transient receptor potential vanilloid 4 TRPV4 and integrins (Delmas, Hao and Rodat-Despoix, 2011).

Integrins are transmembrane receptors that connect the extracellular matrix with the cytoskeleton (Hynes, 2004). Integrins function as mechanoreceptors which transmit forces between the extracellular matrix and the cytoskeleton to maintain tissue structure integrity (Baker and Zaman, 2010). Cell strain causes activation of signal pathways which include mitogen-activated protein (MAP) kinase, Rho guanosine triphosphate (GTP)ase, increased cytoplasmic calcium, and reactive oxygen. These events cause changes in gene expression (Schwartz, 2010). Puncture on the skin on wet cupping is a stressor that can stimulate the expression of endorphin (Subadi *et al.*, 2017). The hypothesis of this study was that wet cupping is more effective than dry cupping because there is an addition of puncture which causes endorphin expression. This study aims to determine pain thresholds, $\alpha 2\beta 1$ -integrin and MOR after WCT and DCT.

II. METHODS

This study used post-test control group design. Thirty-two Wistar, 3 months, 250±50 grams, male rats were randomized into 4 groups: negative control (n=8); positive control (n=8); dry cupping (n=8); wet cupping (n=8). In the negative control group, the rats had no exposure at all. In the positive control group, the rats were given complete Freund's adjuvant (CFA) (Sigma-Aldrich, Saint Louis, Missouri, USA). In the dry cupping group, the rats received CFA and dry cupping. In the wet cupping therapy group, the rats received CFA and wet cupping therapy.

Adult Wistar rats were put into plastic cage (4-5 rats per cage). They were acclimatized for 7 days with 12:12 h (dark/light; the light was turned off every 19:30) with temperature maintained at 24°C. CFA (100 µL) was injected in the center of the right plantar footpad in the three groups of the rats (Fehrenbacher, Vasko and Duarate, 2012). In the groups of dry cupping and wet cupping, 48 hours after the injection, the rats were treated with dry cupping or wet cupping therapy (Subadi *et al.*, 2017). In the dry cupping group, the skin of the right and left paralumbal areas was cupped (2 cm in diameter) with a negative pressure of 200 mmHg for 5 minutes (Subadi *et al.*, 2017). In the wet cupping group, the rats were pricked with lancet 28G before being cupped. Twenty-four hours after wet and dry cupping, all rats were examined for pain threshold using a hot plate (Hot/Cold Plate Cat 35100, Ugo Basile, Varese, Italy). The temperature of the hot plate was 51°C [15]. After examination of the pain threshold, the rats were terminated by cervical dislocation and the paralumbal skin was examined with immunohistochemistry.

Pain was examined using hot plate in bright conditions. The pain threshold was calculated from the moment the rats were placed on the hot plate surface until the nociceptive response occurred. Response showing discomfort (Nocifensive) behaviors include forepaw withdrawal or licking, hind paw withdrawal or licking, stamping, leaning posture and jumping (Deuis, Dvorakova and Vetter, 2017), with a maximum time limit of 30 seconds. One measurement used a stopwatch for each rats to obtain the average value. All the nocifensive behavior responses were conducted by a trained and experienced person in order to have homogenous results.

Expression of $\alpha 2\beta 1$ integrin in keratinocytes and μ -opioid receptors in nerve cells in the spinal cord lamina II was immunohistochemically examined using anti-integrin monoclonal antibody $\alpha 2\beta 1$ (Cat. No MAB2141Z, Chemicon, Merck Millipore co., Darmstadt, Germany) and μ -opioid anti-receptor monoclonal antibody (ab64746, Abcam,

Cambridge, USA). Cells that were positive for $\alpha 2\beta 1$ integrins and μ -opioid receptors were calculated using a microchemical light (Olympus CX21, New York, USA).

III. RESULT

Pain threshold was determined 24 hours after dry cupping therapy and wet cupping therapy, while in positive control group it was performed 72 hours after CFA injection. Figure 1 demonstrates the results of pain threshold experiment. Significant differences were observed between negative control and dry cupping therapy; and negative control and wet cupping therapy. Similar significant differences were also observed between positive control and dry cupping therapy; and positive control and wet cupping therapy. Interestingly, wet cupping therapy and dry cupping therapy showed significant difference. In this study there were two samples in wet cupping group which had to be stopped because it exceeded 30 seconds.

On immunohistochemical examination, the expression of $\alpha 2\beta 1$ integrin showed positive cells (keratinocytes) with brown stain on dry cupping and wet cupping (Figure 2). Negative reactions of $\alpha 2\beta 1$ integrin expression did not show brown stain. Number of keratinocytes expressing $\alpha 2\beta 1$ integrin is demonstrated on Figure 3. Significant differences were shown between negative control group and dry cupping therapy group, negative control group and wet cupping therapy group, and positive control group and wet cupping therapy.

On immunohistochemical examination, the expression of μ -opioid receptors showed positive cells (neurons) with brown stain on dry cupping and wet cupping (Figure 4). Negative reactions of μ -opioid receptors did not show brown stain. Number of nerve cells in the spinal cord lamina II expressing μ -opioid receptors is demonstrated on Figure 4. Significant differences were shown between negative control group showed significant differences with positive control group, dry cupping therapy group, and wet cupping therapy group. Similar trend was also observed in positive control group. Interestingly, dry cupping therapy group showed significant difference with wet cupping therapy group.

IV. DISCUSSION

In this study, dry cupping therapy and wet cupping therapy increased the expression of $\alpha 2\beta 1$ integrins in keratinocytes and μ -opioid receptors in spinal cord nerve cells. It seems that the negative pressure applied in dry cupping therapy and wet cupping therapy triggers the expression of $\alpha 2\beta 1$ integrins. In this study, $\alpha 2\beta 1$ integrin expression in wet cupping therapy was higher than that in dry cupping therapy. The difference between wet and dry cupping therapy methods is the puncture. The puncture causes damage to the keratinocyte membrane. Cell damage causes inflammatory reactions and secretion of inflammatory mediators such as bradykinin, prostaglandin, leukotriene, serotonin, histamine, and substance P (Amaya *et al.*, 2013). The integrins which contained subunit $\beta 1$ play a role in nociceptor sensitization by inflammatory mediators (Dina *et al.*, 2004, 2005). Additionally, they also modulate neuronal excitability and injury after mechanical stimulation to the surrounding extra cellular matrix (Wang *et al.*, 2006; Hemphill *et al.*, 2011). Mechanical stimulation is resulted by both dry and wet cupping therapy.

This study shows that dry cupping therapy and wet cupping therapy can reduce pain. These findings were in accordance with those of other researchers that dry cupping therapy and wet cupping therapy reduced pain (Dalton and BJ, 2017). Pain threshold test in this study was limited to 30 seconds. This study was conducted to prevent tissue

damage to the rat's paws. There were 25% (2 out of 8) samples in wet cupping therapy group exceeded 30 seconds. It could be caused by effectiveness of wet cupping therapy in increasing pain thresholds.

The expression of μ -opioid receptors in wet cupping therapy was higher than that in dry cupping therapy. μ -opioid receptor activation inhibits pain transmission in the spinal cord (Al-Hasani and Bruchas, 2011). In a previous study, β -endorphin expression was found after wet cupping therapy (Subadi *et al.*, 2017). β -endorphin causes analgesia through bonding with μ -opioid receptors. In central nervous system, β -endorphin binds with μ -opioid receptor, causing a cascade of interactions that inhibits the expression of substance P and GABA that contribute to pain transmission (Sprouse-Blum *et al.*, 2010).

In this study, the expression of opioid receptor in wet cupping therapy was higher than that in dry cupping therapy. It was allegedly because puncture was applied in wet cupping therapy, which caused cell injury. Cells affected by lesion will express inflammatory cytokines that can trigger endorphins through the POMC in the hypothalamus (Slominski *et al.*, 2000).

V. CONCLUSION

This study provided evidence-based data that wet cupping therapy is more effective in reducing pain as compared to dry cupping therapy. Pain reduction mechanism in both therapies was mediated by the expression of $\alpha 2\beta 1$ integrin and μ -opioid receptors which are higher in wet cupping.

REFERENCE

- [1] Al-Hasani, R. and Bruchas, M. (2011) 'Molecular mechanisms of opioid receptor-dependent signaling and behavior', *Anesthesiology*, 115(6), pp. 1363–81.
- [2] Amaya, F. *et al.* (2013) 'Tissue Injury and Related Mediators of Pain Exacerbation', *Curr Neuropharmacol*, 11(6), pp. 592–597.
- [3] Arslan, M., Gökgöz, N. and Dane, S. (2016) 'The effect of traditional wet cupping on shoulder pain and neck pain: A pilot study', *Complement Ther Clin Pract*, 23, pp. 30–3.
- [4] Baker, E. and Zaman, M. (2010) 'The biomechanical integrin', *J Biomech*, 43(1), pp. 38–44.
- [5] Cao, H. *et al.* (2004) 'Cupping therapy for acute and chronic pain management: a systematic review of randomized clinical trials', *J Tradit Chinese Med Sci*, 1(1), pp. 49–61.
- [6] Dalton, E. and BJ, V. (2017) 'Cupping Therapy: An Alternative Method of Treating Pain', *Public Health - Open Journal*, 2(2), pp. 59–63.
- [7] Delmas, P., Hao, J. and Rodat-Despoix, L. (2011) 'Molecular mechanisms of mechanotransduction in mammalian sensory neurons', *Nat Rev Neurosci*, 12(3), pp. 139–53.
- [8] Deus, J., Dvorakova, L. and Vetter, I. (2017) 'Methods Used to Evaluate Pain Behaviors in Rodents', *Front Mol Neurosci*, 10, p. 284.
- [9] Dina, O. *et al.* (2004) 'Integrin signaling in inflammatory and neuropathic pain in the rat', *Eur J Neurosci*, 19(3), pp. 634–42.
- [10] Dina, O. *et al.* (2005) 'Primary afferent second messenger cascades interact with specific integrin subunits in producing inflammatory hyperalgesia', *Pain*, 115(1–2), pp. 191–203.
- [11] Fehrenbacher, J., Vasko, M. and Duarate, D. (2012) 'Models of inflammation: Carrageenan- or complete Freund's Adjuvant (CFA)-induced edema and hypersensitivity in the rat', *Curr Protoc Pharmacol*.
- [12] Hemphill, M. *et al.* (2011) 'A Possible Role for Integrin Signaling in Diffuse Axonal Injury', *PLOS ONE*, 6(7), p. e22899.
- [13] Hynes, R. (2004) 'The emergence of integrins: a personal and historical perspective', *Matrix Biol*, 23(6), pp. 333–40.
- [14] Kim, J. *et al.* (2011) 'Cupping for treating pain: a systematic review', *Evid Based Complement Alternat Med*, 2011, p. 467014.
- [15] Schwartz, M. (2010) 'Integrins and extracellular matrix in mechanotransduction', *Cold Spring Harb Perspect Biol*, 2(12), p. a005066.

- [16] Slominski, A. *et al.* (2000) 'Corticotropin releasing hormone and proopiomelanocortin involvement in the cutaneous response to stress', *Physiol Rev*, 80(3), pp. 979–1020.
- [17] Sprouse-Blum, A. *et al.* (2010) 'Understanding Endorphins and Their Importance in Pain Management', *Hawaii Med J*, 69(3), pp. 70–71.
- [18] Subadi, I. *et al.* (2017) 'Pain Relief with Wet Cupping Therapy in Rats is Mediated by Heat Shock Protein 70 and β -Endorphin', *Iran J Med Sci*, 42(4), pp. 384–391.
- [19] Wang, X. *et al.* (2006) 'Role of TGF beta-mediated inflammation in cutaneous wound healing', *J Investig Dermatol Symp Proc*, 11(1), pp. 112–7.




Figure 1: Results of Pain Threshold Experiment




Figure 2: Expression of $\alpha 2\beta 1$ Integrin with Keratinocytes




Figure 3: Number of Keratinocytes Expressing $\alpha 2\beta 1$ Integrin




Figure 4: Result of Immunohistochemical Examination The Expression of μ -opioid Receptors with Neurons

PAIN REDUCTION AFTER WET AND DRY CUPPING THERAPIES ROLES OF INTEGRIN AND OPIOID RECEPTOR IN ANIMAL MODELS

ORIGINALITY REPORT

17%

SIMILARITY INDEX

10%

INTERNET SOURCES

14%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Udayana University Student Paper	2%
2	www.researchgate.net Internet Source	2%
3	journal.stikeskendal.ac.id Internet Source	1%
4	link.springer.com Internet Source	1%
5	www.ncbi.nlm.nih.gov Internet Source	1%
6	Mehmet Cavlak, Alper Özkök, Serhat Sarı, Ahmet Dursun, Taner Akar, Mustafa Karapirli, Birol Demirel. "Iatrogenic artefacts attributable to traditional cupping therapy in a shotgun fatality", Journal of Forensic and Legal Medicine, 2015 Publication	1%

7

Xiu-Ling Wang, Ai-Ying Zhang, Wan-Shan Zhou. "Cupping therapy for treating ankylosing spondylitis: Protocol for a systematic review", *European Journal of Integrative Medicine*, 2018

Publication

1 %

8

Sweetey Malik, Pooja Anand, Pooja Bhati, M. Ejaz Hussain. "Effects of dry cupping therapy on pain, dynamic balance and functional performance in young female with recreational runners chronic plantar fasciitis", *Sports Orthopaedics and Traumatology*, 2022

Publication

1 %

9

Xian-jun Xiao, Lei-xiao Zhang, Yun-zhou Shi, Jun-peng Yao et al. "Cupping therapy for patients with chronic urticaria: A systematic review and meta-analysis", *Journal of Integrative Medicine*, 2020

Publication

1 %

10

www.neurona.web.id

Internet Source

1 %

11

Ilana Cruz-Silva, Viviane Abreu Nunes, Andrezza Justino Gozzo, Priscila Praxedes-Garcia et al. " Protease Inhibitors Extracted from Lam. Affect Kinin Release during Lung Inflammation ", *Pulmonary Medicine*, 2016

Publication

1 %

12	res.mdpi.com Internet Source	1 %
13	"Heat Shock Proteins in Human Diseases", Springer Science and Business Media LLC, 2021 Publication	<1 %
14	ijriar.com Internet Source	<1 %
15	www.nature.com Internet Source	<1 %
16	Deldar Morad Abdulah, Hawar Abdulrazaq Mohammedsadiq, Ary Habeeb Mohammed. "Effectiveness of wet cupping therapy on relieving pain in patients with chronic migraine: an observational study", Journal of Complementary and Integrative Medicine, 2021 Publication	<1 %
17	Eva Mercado, Nancy Paniagua, Eva M Sánchez-Robles, Rocío Girón, Diego Alvarez de la Rosa, Teresa Giraldez, Carlos Goicoechea. "SGK1.1 isoform is involved in nociceptive modulation, offering a protective effect against noxious cold stimulus in a sexually dimorphic manner", Pharmacology Biochemistry and Behavior, 2022 Publication	<1 %

18

Hugo Jário de Almeida Silva, Bruno T Saragiotto, Rodrigo Scattone Silva, Caio Alano de Almeida Lins, Marcelo Cardoso de Souza. "Dry cupping in the treatment of individuals with non-specific chronic low back pain: a protocol for a placebo-controlled, randomised, double-blind study", *BMJ Open*, 2019

Publication

<1 %

19

Hümeyra ÇELİK, Ayhan ÇETİNKAYA, Bihter Gökçe BOZAT, Selma ERDOĞAN DÜZCÜ, Murat ALIŞIK, Özgür Mehmet YİS. "The Effect of Wet Cupping Therapy on Anxiety and Thiol/Disulfide Homeostasis in Premature Ovarian Failure: An Experimental Study", *Journal of Traditional Medical Complementary Therapies*, 2022

Publication

<1 %

20

Khawla Al-luhaidan, M. S. Prarthana. "Hijama Practices and the Perceived Health Benefits among the People of Al-Qassim Region, Saudi Arabia", *Journal of Complementary and Alternative Medical Research*, 2020

Publication

<1 %

21

Stephan Kellenberger, Laurent Schild. "Epithelial Sodium Channel/Degenerin Family of Ion Channels: A Variety of Functions for a

<1 %

Shared Structure", Physiological Reviews, 2002

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

PAIN REDUCTION AFTER WET AND DRY CUPPING THERAPIES ROLES OF INTEGRIN AND OPIOID RECEPTOR IN ANIMAL MODELS

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
