



Volume 119 June 2023

ISSN 0889-1575

JOURNAL OF FOOD COMPOSITION AND ANALYSIS



Journal of Food Composition and Analysis

EDITOR-IN-CHIEF

J.S. Elmore
*University of Reading, Reading,
United Kingdom*

ASSISTANT EDITOR

C. Girardot

CONSULTING AND FOUNDING EDITOR

K. Stewart
University of Texas System, Austin, Texas, United States

EDITORS

F. J. Barba
*University of Valencia
Burjassot, Spain*

D. de Beer
*ARC Infruitec-Nietvoorbij,
Stellenbosch, South Africa*

L. Bell
*University of Reading, Reading,
United Kingdom*

R. Chisté
*Federal University of Para,
Belém, Brazil*

F.Y. Chye
*University Malaysia Sabah,
Kota Kinabalu, Malaysia*

A. Ipeaiyeda
*University of Ibadan, Ibadan,
Nigeria*

P. Onianwa
University of Ibadan, Ibadan, Nigeria

D. Restuccia
*University of Calabria,
Rende, Italy*

EDITORIAL BOARD

M. Campo
Zaragoza, Spain

E. Cantos-Villar
Jerez de la Frontera, Spain

C. Cardoso
Lisboa, Portugal

I. Castanheira
Lisboa, Portugal

C. Champagne
Louisiana, United States

S. Chung
Hong Kong, China

D. Cozzolino
Queensland, Australia

M. Dueñas Patón
Salamanca, Spain

G. Duodu
Pretoria, South Africa

T. Gallina Toschi
Bologna, Italy

V. Gökmen
Ankara, Turkey

J.L. Guil-Guerrero
Almeria, Spain

D. Haytowitz
Maryland, United States

D. Hornero-Méndez
Seville, Spain

L. Hu
Zhengzhou, China

P. Hulshof
Wageningen, Netherlands

L. Husáková
Pardubice, Czech Republic

P. van Jaarsveld
Tygerberg, South Africa

J. Jakobsen
Søborg, Denmark

J. Lee
Idaho, United States

L. Mariutti
São Paulo, Brazil

A.J. Meléndez-Martínez
Sevilla, Spain

B. Min
Maryland, United States

M. Netzel
Queensland, Australia

P. Pehrsson
Maryland, United States

J. Porubska

E. Rodrigues
Porto Alegre, Brazil

R.M. Schweiggert
Stuttgart, Germany

W. Sun
Guangzhou, China

D. Thavarajah
South Carolina, United States

T. Zimmerman
*Maryland, United States of
America*

[Submit your article](#)

[Menu](#)



[Search in this journal](#)

Volume 115

January 2023

[← Previous vol/issue](#)

[Next vol/issue >](#)

Receive an update when the latest issues in this journal are published

[Set up journal alerts](#)

[Full text access](#)

[Editorial Board](#)

[Article 105058](#)

[↓ Download PDF](#)

[Review Articles](#)

[FEEDBACK](#) 

Submit your article



[Download PDF](#) [Article preview](#)

[Review article](#) [Full text access](#)

The concentration of potentially toxic elements (PTEs) in human milk: A systematic review, meta-analysis, and health risk assessment

Aliasghar Neshat, Ali Oghazyan, Fatemeh Kariminejad, Trias Mahmudiono, ... Amin Mousavi Khaneghah
Article 104933

[Download PDF](#) [Article preview](#)

[Review article](#) [Full text access](#)

Sample preparation and spectrometric methods for elemental analysis of milk and dairy products – A review

Samara Soares, Liz Mary B. Moraes, Fábio R.P. Rocha, Alex Virgilio
Article 104942

[Download PDF](#) [Article preview](#)

[Review article](#) [Full text access](#)

The greater yam (*Dioscorea alata* L.): A review of its phytochemical content and potential for processed products and biofortification

Vincent Lebot, Floriane Lawac, Laurent Legendre
Article 104987

[Download PDF](#) [Article preview](#)

[Review article](#) [Full text access](#)

Submit your article



Original Research Articles

Research article Full text access

Non-carcinogenic risk assessment of cadmium and lead in raw milk from dairy production farms in Iran, using Monte Carlo simulation approach

Neematollah Jaafarzadeh, Kamran Tari, Mohammad Reza Samarghandi, Masoud Panahi Fard, ... Maryam Mehrbakhsh

Article 104864

[Download PDF](#) Article preview

Research article Full text access

Simultaneous determination of 20 phenolic compounds in propolis by HPLC-UV and HPLC-MS/MS

Yan Zhang, Cuiling Cao, Zhiwei Yang, Guangqun Jia, ... Adan Li

Article 104877

[Download PDF](#) Article preview

Research article Full text access

Selenate fertilization of sorghum via foliar application and its effect on nutrient content and antioxidant metabolism

Patriciani Estela Cipriano, Rodrigo Fonseca da Silva, Fabio Aurélio Dias Martins, Alexandre Boari de Lima, ... Luiz Roberto Guimarães Guilherme

Article 104865

[Download PDF](#) Article preview

Research article Full text access

Submit your article



Research article Full text access

Is *Tritordeum* (~~×~~*Tritordeum martinii* A. Pujadas, nothosp. nov.) grain a potentially useful source of essential minerals in the human diet?

Elżbieta Suchowilska, Wolfgang Kandler, Marian Wiwart, Rudolf Krska

Article 104874

[Download PDF](#) Article preview 

Research article Full text access

Inexpensive and simple tool for quality control of nutraceutical and tonic products with capillary electrophoresis and contactless conductivity detection: Some developments in Vietnam

Quang Huy Nguyen, Thi Huyen My Dang, Thi Phuong Quynh Le, Thi Huyen Trang Luu, ... Thanh Duc Mai

Article 104882

[Download PDF](#) Article preview 

Research article Full text access

A dilute-and-shoot based SWATH-MS approach for rapid analysis of 23 synthetic dyes in spices

Guangyun He, Xue Hou, Mei Han, Shiting Qiu, ... Mei Liang

Article 104878

[Download PDF](#) Article preview 

Research article Full text access

Submit your article



Research article Full text access

Assessment of interspecies (*Capsicum annuum* X *Capsicum frutescens*) recombinant inbred lines (RIL) for fruit nutritional traits

Hasan Pinar, Mahmut Kaplan, Kevser Karaman, Beyza Ciftci

Article 104848

[Download PDF](#) Article preview

Research article Full text access

Potential for increasing folate contents of traditional African fermented sorghum gruel (*Motoho*) using presumptive probiotic lactic acid bacteria

Olanrewaju E. Fayemi, Gabriel B. Akanni, Samuel S. Sobowale, Andre Oelofse, Elna M. Buys

Article 104850

[Download PDF](#) Article preview

Research article Full text access

Metal-organic framework with dual-loading of nickel/nitrogen-doped carbon dots and magnetic nanoparticles for fluorescence detection of fenitrothion in food samples

Elnaz Delnavaz, Mohammad Amjadi, Mir Ali Farajzadeh

Article 104873

[Download PDF](#) Article preview

Research article Full text access

The most consumed beef cuts in Brazil: Prices versus metabolic profile

Submit your article



Elemental composition of rice and lentils from various countries: A probabilistic risk assessment of multiple life stages

Mom TatahMentan, Syprose Nyachoti, Onoja F. Okwori, Tewodros R. Godebo

Article 104852

[Download PDF](#) Article preview

Research article Full text access

Elemental profile of food aids and mineral provision for pregnant and lactating refugee women

Desta Woldetsadik, Hilette Hailu, Solomie Gebrezgabher, Mary Njenga, ... Eulogio J. Llorent-Martínez

Article 104881

[Download PDF](#) Article preview

Research article Full text access

Effect of fermentation stages on the nutritional and mineral bioavailability of cacao beans (*Theobroma cacao* L.)

Cristopher G. Millena, Aubrey Rosebud R. Balonzo, Jem R. Rentoy, Sylvester S. Ruivivar, Samuel C. Bobiles

Article 104886

[Download PDF](#) Article preview

Research article Full text access

Polysaccharides from pinhão seeds of *Araucaria angustifolia*: Extraction, isolation and structural characterization

Shayla Fernanda Barbieri, Lucas Trevisan França de Lima, Jéssica Caroline Scholze Baum, Keylla Lençone Mischiatti, ... Joana Léa Meira Silveira

Submit your article



Supalak Kongsri, Chunyapuk Kukusamude

Article 104883

[Download PDF](#) Article preview 

Research article Full text access

Characteristics, water quality index and human health risk from nitrate and fluoride in Kakhk city and its rural areas, Iran

Mehdi Qasemi, Mostafa Darvishian, Hojjat Nadimi, Mehdi Gholamzadeh, ... Ahmad Zarei

Article 104870

[Download PDF](#) Article preview 

Research article Full text access

Analysis of residual monepantel sulfone in milk using icELISA based on a monoclonal antibody developed from an easy-synthesized and low-cost hapten

Yanfang Zhang, Pimiao Zheng, Zile Wang, Zhuoren Ling, ... Haiyang Jiang

Article 104897

[Download PDF](#) Article preview 

Research article *Open access*

Lipidomics of common octopus' (*Octopus vulgaris*) arm muscle using untargeted high-resolution liquid chromatography-mass spectrometry

Luís Gaspar, Fernando Ricardo, Tânia Melo, Pedro Domingues, ... Felisa Rey

Article 104871

[Download PDF](#) Article preview 

Submit your article



[Download PDF](#) [Article preview](#)

Research article Full text access

The content of biologically active compounds in potato tubers of Ismena (yellow flesh) and Provita (purple flesh) varieties – A comparison

A. Kita, E. Rytel, J. Miedzianka, W.A. Turski, ... T. Lenartowicz

Article 104898

[Download PDF](#) [Article preview](#)

Research article Full text access

Characterization of key odor-active compounds in pure chicken powder and the effect of yeast extract, using instrumental and sensory techniques

Dong-Yu Shen, Meng-Ke Li, Mu Zhao, Jie Li, ... Ku Li

Article 104879

[Download PDF](#) [Article preview](#)

Research article Full text access

Levels of selected trace metals in enset (*Ensete ventricosum* (Welw.), *Cheesman*) (Unprocessed and processed) and soil of siltie zone, Southern Ethiopia.

Misgana Negera, Tesfahun Bufebo, Dereje Abebe

Article 104905

[Download PDF](#) [Article preview](#)

Research article Full text access

Submit your article



Research article Full text access

Pre-column derivatization with trimethylsilyl diazomethane coupled with ASE-SPE-GC-MS/MS method for the quantification and validation of penicillin G residues in poultry tissues and pork

Chujun Liu, Xing Xie, Yawen Guo, Bo Wang, ... Wenbin Bao

Article 104866

[Download PDF](#) Article preview 

Research article Full text access

Comprehensive characterization of phytochemicals in edible sea fennel (*Crithmum maritimum* L., Apiaceae) grown in central Italy

Diletta Piatti, Simone Angeloni, Filippo Maggi, Giovanni Caprioli, ... Gianni Sagratini

Article 104884

[Download PDF](#) Article preview 

Research article Full text access

Identification of insect sources of honey in China based on real-time fluorescent LAMP technology

Jie Gao, Xiue Jin, Bo Gong, Jingjing Li, ... Jun Wang

Article 104875

[Download PDF](#) Article preview 

Research article Full text access

Submit your article




Research article Full text access

Development of non-destructive NIRS models to predict oil and major fatty acid contents of Ethiopian sesame

Girmay Tsegay, Yibrah Ammare, Samuel Mesfin

Article 104908

[Download PDF](#) Article preview 

Research article Full text access

In vitro bioaccessibility of copper, iron, and zinc from common meat substitutes, influence of exogenously added garlic/onion and contribution to the diet

Fiorella laquinta, Nataly Rodríguez, Ignacio Machado

Article 104910


[Download PDF](#) Article preview 

Research article Full text access

Effect of infant's gender and lactation period on biochemical and energy breast milk composition of lactating mothers from Algeria

Naima Khelouf, Khadidja Haoud, Samira Meziani, Meriem Fizir, ... Nadjat Kadi

Article 104889

[Download PDF](#) Article preview 

Research article Full text access

Submit your article



Research article Full text access

Multi-residue method validation, processing factor and monitoring of thirteen targeted fungicide residues in the process of wine making

TP Ahammed Shabeer, Ramhari Somkuwar, Ajay Kumar Sharma, Umakant Deshmukh, Sandip Hingmire
Article 104912

[Download PDF](#) Article preview 

Research article Full text access

Risk assessment of oral exposure to phthalates from coffee samples marketed in Turkey

Gursel Isci, Elif Feyza Topdas, Elif Dagdemir, Huseyin Efe Genis
Article 104913

[Download PDF](#) Article preview 

Research article Full text access

Standardization of proanthocyanidin extract's mean degree of polymerization using eutectic solvents

Rodrigo T. Neto, Sónia A.O. Santos, Joana Oliveira, Armando J.D. Silvestre
Article 104887

[Download PDF](#) Article preview 

Research article Full text access

Vitamins, minerals and heavy metals profiling of seaweeds from Mauritius and Rodrigues for food security

Submit your article



halophytes – Saltbush, Seablite and Seapurslane

Sukirtha Srivarathan, Anh Dao Thi Phan, Hung Trieu Hong, Gabriele Netzel, ... Michael E. Netzel

Article 104876

[Download PDF](#) Article preview 

Research article Full text access

Delineating unique and discriminant chemical traits in Brazilian and Italian extra-virgin olive oils by quantitative 2D-fingerprinting and pattern recognition algorithms

Federico Stilo, Eugenio Alladio, Simone Squara, Carlo Bicchi, ... Humberto Ribeiro Bizzo

Article 104899

[Download PDF](#) Article preview 

Research article Full text access

SERS paper sensor based on three-dimensional ZnO@Ag nanoflowers assembling on polyester fiber membrane for rapid detection of florfenicol residues in chicken

Xiaodi Li, Hualan Zhou, Lehui Wang, Huiwen Wang, ... Jianguo Zhang

Article 104911

[Download PDF](#) Article preview 

Research article Full text access

Do vine cropping and breeding practices affect the biogenic amines' content of produced wines?

Giovanna Loredana La Torre, Archimede Rotondo, Andrea Salvo

Article 104901

[Download PDF](#) Article preview 

Submit your article



[Download PDF](#) [Article preview](#)

Research article Full text access

Triacylglycerol composition and thermodynamic profiles of fractions from dry fractionation of anhydrous milk fat

Xin Si, Huiquan Zhu, Panpan Zhu, Yunna Wang, ... Shuwen Zhang

Article 104916

[Download PDF](#) [Article preview](#)

Research article Full text access

Residue levels and risk assessment of pesticides in litchi and longan of China

Lixue Kuang, Zhiqiang Wang, Yang Cheng, Yinping Li, ... Guofeng Xu

Article 104921

[Download PDF](#) [Article preview](#)

Research article Full text access

Isolation of quercetin-3-O-sulfate and quantification of major compounds from *Psidium guajava* L. from Vietnam

Phuc-Dam Nguyen, Marie-France Hérent, Thi-Bach Le, Thi-Buu-Hue Bui, ... Joëlle Quetin-Leclercq

Article 104928

[Download PDF](#) [Article preview](#)

Research article Full text access

Identification and comparison of palmitoleic acid (C16:1 n-7)-derived lipids in marine fish by-products by UHPLC-Q-exactive orbitrap mass spectrometry

Submit your article



by HPLC/UV method

Kenan Dost, Mertcan Öksüz, Mustafa Cittan, Büşra Mutlu, Betül Tural

Article 104926

[Download PDF](#) Article preview

Research article Full text access

Comparison of measured and declared vitamin D concentrations in Australian fortified foods

Katya M. Clark, Anthony P. James, Hio Ian Ip, Eleanor Dunlop, ... Lucinda J. Black

Article 104922

[Download PDF](#) Article preview

Research article Full text access

Monitoring of pesticide and some metal residues in Mazafati date fruit cultivar and risk assessment to the health

Parisa Jafarian Asl, Razieh Niazmand, Bibi Marzieh Razavizadeh, Monireh Alsadat Shakeri, Moslem Jahani

Article 104917

[Download PDF](#) Article preview

Research article Full text access

Effect of bio-nano-selenium on yield, nutritional quality and selenium content of radish

Sisi Huang, Kan Yu, Qiushui Xiao, Banghui Song, ... Wei Zheng

Article 104927

[Download PDF](#) Article preview

Submit your article



[Download PDF](#) Article preview

Research article Full text access

Novel amide- and imide-co-polymers modified with sulfathiazole as efficient chelator for selective extraction, pre-concentration and determination of trace inorganic antimony (as Sb(III)) from edible vegetable oils by ultrasound assisted-cloud point extraction coupled to micro-volume UV-spectrophotometry

H.B. Zengin, R. Gürkan

Article 104931

[Download PDF](#) Article preview

Research article Full text access

Quantitative lipidomic analysis of egg yolk, yolk granule, and yolk plasma

Xiaoqi He, Jinqiu Wang, Yi Wang, Beibei Wang, ... Fang Geng

Article 104880

[Download PDF](#) Article preview

Research article Full text access

Development of a mutton powder certified reference material for the analysis of ractopamine

Jian Zhou, Min Wang, Liyuan Zhang, Mengrui Yang, ... Qifan Sun

Article 104935

[Download PDF](#) Article preview

Research article Full text access

Submit your article




Research article Full text access

Mineral oil contamination in basil pesto from the Italian market: Ingredient contribution and market survey

Ana Srbinovska, Lara Gasparotto, Chiara Conchione, Luca Menegoz Ursol, ... Sabrina Moret

Article 104914

[Download PDF](#) Article preview 

Research article Full text access

Transmissibility of anisakid allergenic peptides from animal feed to chicken meat: Proof of concept

Ganna Saelens, Sören Planckaert, Bart Devreese, Sarah Gabriël

Article 104939

[Download PDF](#) Article preview 

Research article Full text access

Groundwater quality assessment for drinking and irrigation purposes in the Hangu District, Pakistan

Imran Ud Din, Said Muhammad, Inayat ur Rehman

Article 104919

[Download PDF](#) Article preview 

Research article *Open access*

Three *Pleurotus* mushroom species cultivated in a mixed *Phragmites australis* substrate differ in nutrient utilization capacity

Submit your article



chemical characterization and quality evaluation

Jun Peng, Rahima Abdulla, Yi Li, Xiao-yan Liu, ... Haji Akber Aisa

Article 104930

[Download PDF](#) Article preview

Research article Full text access

Evaluation of different extraction procedures for the quantification of seven cannabinoids in cannabis-based edibles by the use of LC-MS

Marios C. Christodoulou, Atalanti Christou, Ioannis J. Stavrou, Constantina P. Kapnissi-Christodoulou

Article 104915

[Download PDF](#) Article preview

Research article Full text access

Fresh *Brassicaceae* sprouting broccoli (Bimi®) glucosinolates profile characterization and bioaccessibility through an *in vitro* dynamic digestion study

Juan Antonio Nieto, Pilar Hellín, Beatriz Pérez, Blanca Viadel, ... Agatha Agudelo

Article 104941

[Download PDF](#) Article preview

Research article Full text access

Qualitative and quantitative analysis of curcumin in dried ginger by the resonance Rayleigh scattering technique and absorption spectroscopy

Yan Liu, Hong Jiang

Article 104923

[Download PDF](#) Article preview

Submit your article



[Download PDF](#) [Article preview](#)

Research article [Full text access](#)

The effects of pollen source on the fatty acid profile of Macadamia kernels

Wei Hu, Melissa Fitzgerald, Bruce Topp, Mobashwer Alam, ... Tim J. O'Hare

Article 104943

[Download PDF](#) [Article preview](#)

Research article [Full text access](#)

Effects of pasteurization and high-pressure processing on the fatty acids, triacylglycerol profile, Dornic acidity, and macronutrients in mature human milk

Luciana Pelissari Manin, Adriela Albino Rydlewski, Jessica Santos Pizzo, Victor Hugo Maldonado da Cruz, ... Jesui Vergilio Visentainer

Article 104918

[Download PDF](#) [Article preview](#)

Research article [Full text access](#)

Aroma-active compounds related to Maillard reaction during roasting in Wuyi Rock tea

Ping Yang, Haili Wang, Qingqing Cao, Huanlu Song, ... Yanping Lin

Article 104954

[Download PDF](#) [Article preview](#)

Research article [Open access](#)

Submit your article



Research article Full text access

Non-acylated and acylated anthocynins in red wines of different ages: Color contribution and evaluation

Zhaoxiang Wang, Lin Zhang, Yuyu Li, Qianqian Liu, Chunlong Yuan
Article 104951

[Download PDF](#) Article preview 

Research article Full text access

A novel platform based on MnO₂ nanoparticles and carboxylated multi-walled carbon nanotubes composite for accurate and rapid determination of curcumin in commercial food products

Peihong Deng, Yanping Wei, Wei Li, Shuting Shi, ... Quanguo He
Article 104940

[Download PDF](#) Article preview 

Discussion Full text access

Commentary: “Quantitative” vs quantitative Headspace Solid-Phase Microextraction (HS-SPME) in food volatile and flavor compounds analysis

Henryk H. Jeleń, Martyna N. Wiczorek
Article 104955

[Download PDF](#) Article preview 

Research article *Open access*

Submit your article



Research article Full text access

An ultrahigh-performance liquid chromatography-fluorescence detection (UHPLC-FLD) method for simultaneous determination of albendazole and its three metabolites in poultry eggs

Fanxun Guan, Zhaoyuan He, Yayun Tang, Pengfei Gao, ... Kaizhou Xie

Article 104959

[Download PDF](#) Article preview 

Research article Full text access

Comprehensive characteristic of lipid fraction as a distinguishing factor of three lupin seed species

Aleksander Siger, Anna Grygier, Jaroslaw Czubinski

Article 104945


[Download PDF](#) Article preview 

Research article Full text access

Assessment of the concentrations of ochratoxin A, zearalenone, and deoxynivalenol during cracker production

Amin Mousavi Khaneghah, Parisa Mostashari, Carlos A.F. Oliveira, Fernanda M. Vanin, ... Anderson S. Sant'Ana

Article 104950

[Download PDF](#) Article preview 

Research article Full text access



Research article Full text access

Discrimination based on commercial/craft origin and on lager/ale fermentation of undiluted Spanish beer samples: front-face excitation-emission matrices and chemometrics

Mónica Palomino-Vasco, María Isabel Rodríguez-Cáceres, Nielene Mora-Díez

Article 104946

[Download PDF](#) Article preview 

Research article Full text access

The influence of high pressure processing and germination on anti-nutrients contents, in vitro amino acid release and mineral digestibility of soybeans

Xinyi Wu, Mei Tan, Yinfei Zhu, Hanying Duan, ... Chao Wang

Article 104953


[Download PDF](#) Article preview 

Research article Full text access

Determination and dietary risk assessment of 52 pesticide residues in vegetable and fruit samples by GC-MS/MS and UHPLC-QTOF/MS from Gujarat, India

Divya Kottadiyil, Tejal Mehta, Rupal Thasale, Sivaperumal, P

Article 104957

[Download PDF](#) Article preview 

Research article Full text access

Submit your article



Research article Full text access

Chemical composition and nutraceutical characterization of *Balanites roxburghii* seed oil

Guggalada Govardhana Yadav, Vallamkondu Manasa, Hosakatte Niranjana Murthy, Ajay W. Tumaney
Article 104952

[Download PDF](#) Article preview 

Research article Full text access

Determination of fatty acids in vegetable oils by GC-MS, using multiple-ion quantification (MIQ)

Moisés Guerrero-Esperanza, Katarzyna Wrobel, Kazimierz Wrobel, José Juan Ordaz-Ortiz
Article 104963

[Download PDF](#) Article preview 

Research article Full text access

Fatty acids and minor functional compounds of hemp (*Cannabis sativa* L.) seeds and other Cannabaceae species

José Ignacio Alonso-Esteban, María José González-Fernández, Dmitri Fabrikov, María de Cortes Sánchez-Mata, ... José Luis Guil-Guerrero
Article 104962

[Download PDF](#) Article preview 

Research article Full text access

Submit your article




Research article Full text access

Uptake of heavy metal in wheat from application of different phosphorus fertilizers

Sabiha-Javied, N. Siddque, S. Waheed, Q. uz Zaman, ... R. Nasir

Article 104958

[Download PDF](#) Article preview 

Research article Full text access

Water distribution, physicochemical and microstructural properties of scallop adductors as affected by different drying methods

Jing Liu, Ya Zhao, Qilong Shi, Xiaotian Wu, Zhongxiang Fang

Article 104966

[Download PDF](#) Article preview 

Research article Full text access

Combined effects of metals (Cr^{6+} , Hg^{2+} , Ni^{2+} , Zn^{2+}) and calcium on the serum biochemistry and food quality of the Nile fish (*Oreochromis niloticus*)

Sezen Özçelik, Mustafa Canli

Article 104968

[Download PDF](#) Article preview 

Research article *Open access*

Mercury in fish tissues from the Galapagos marine reserve: Toxic risk and health implications

Submit your article



Microwave assisted effective synthesis of CdS nanoparticles to determine the copper ions in artichoke leaves extract samples by flame atomic absorption spectrometry

Meltem Şaylan, Berfin Metin, Hilal Akbıyık, Fatma Turak, ... Sezgin Bakırdere

Article 104965

[Download PDF](#) Article preview

Research article Full text access

Comparison of the triglyceride composition of vegetable samples with ultra-high efficiency / low-pressure supercritical fluid chromatography – mass spectrometry

Quentin Gros, Marta Wolniaczyk, Johanna Duval, Caroline West, ... Eric Lesellier

Article 104960

[Download PDF](#) Article preview

Research article Full text access

Human health risk assessment of pesticides in lettuce and spinach grown in urban backyard garden soils

Islam Md Meftaul, Kadiyala Venkateswarlu, Aney Parven, Prasath Annamalai, Mallavarapu Megharaj

Article 104977

[Download PDF](#) Article preview

Research article Full text access

¹³C-NMR-based MixONat strategy coupled with 2D NMR for rapid dereplication and identification of new secondary metabolites from *Aloe vera*

Tianpeng Yin, Yi Yu, Qinghua Liu, Guoyuan Zhu, ... Zhihong Jiang

Article 104975

Submit your article



[Download PDF](#) [Article preview](#)

Research article [Full text access](#)

Carbon dioxide spectrophotometric determination in whole roasted coffee beans using a total analysis system after super-modified simplex optimization

Margarida Q.M. Silva, Inês M. Valente, Paulo J. Almeida, João Rodrigo Santos
Article 104978

[Download PDF](#) [Article preview](#)

[< Previous vol/issue](#)

[Next vol/issue >](#)

ISSN: 0889-1575

Copyright © 2023 Elsevier Inc. All rights reserved

Journal of Food Composition and Analysis

Supports *open access*


[Submit your article](#)





Contents lists available at ScienceDirect

Journal of Food Composition and Analysis

journal homepage: www.elsevier.com/locate/jfca

Review article

The concentration of potentially toxic elements (PTEs) in human milk: A systematic review, meta-analysis, and health risk assessment

Aliasghar Neshat^a, Ali Oghazyan^{b,*}, Fatemeh Kariminejad^c, Trias Mahmudiono^d, Yadolah Fakhri^e, Amir Mohammad Sheikh Asadi^f, Ali Atamaleki^{g,*}, Amin Mousavi Khaneghah^h^a Department of Public Health, Esfarayen Faculty of Medical Sciences, Esfarayen, Iran^b Department of Environmental Health, Faculty of Health, Sabzevar University of Medical Sciences, Sabzevar, Iran^c Student Research Committee, Department of Environmental Health Engineering, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran^d Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia^e Food Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran^f Gdańsk University of Technology, Faculty of Civil and Environmental Engineering, Department of Sanitary Engineering, Gdańsk, Poland^g Department of Environmental Health Engineering, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran^h Department of Fruit and Vegetable Product Technology, Prof. Waclaw Dąbrowski Institute of Agricultural and Food Biotechnology – State Research Institute, 36 Rakowiecka St., 02-532 Warsaw, Poland

ARTICLE INFO

Keywords:

Human milk
Health risk assessment
Carcinogenic risk
Potentially toxic elements (PTEs)
Food safety

ABSTRACT

Human milk has an important role in infants' psychological and immunological development. In addition to providing vital substances, some environmental contaminants, such as potentially toxic elements (PTEs), can be transmitted by human breast milk to infants. However, some studies monitored PTEs concentration in human breast milk; no meta-analysis was conducted to estimate the concentration of PTEs in human breast milk. Therefore, this review aimed to determine PTEs concentrations in human breast milk and consumption-related health effects worldwide via meta-analysis and health risk assessment. After searching among Scopus, Web of Science, and PubMed databases, 32 studies were included in this work. Based on the results, the rank order of PTEs was Fe (258.44 µg/kg) > Zn (205.16 µg/kg) > Cu (32.29 µg/kg) > Mn (4.30 µg/kg) > Cr (2.62 µg/kg) > Hg (0.44 µg/kg) > As (0.21 µg/kg) > Cd (0.16 µg/kg) > Pb (0.03 µg/kg). Moreover, Egypt, Pakistan, Brazil, Jordan, and Turkey for non-carcinogenic risk (n-CR) and Egypt, Jordan, Brazil, and Romania for carcinogenic risk (CR) have shown unsafe levels, respectively. Since the lactating mothers' diet can directly affect their milk's content, monitoring the feeding behavior (especially supplements taken during pregnancy) and the quality of foods is recommended.

1. Introduction

Chemical and microbial pollution of the environment (Yang et al., 2021; Yu et al., 2022), followed by food (Ke et al., 2022; Lei et al., 2022; Sun et al., 2022), can endanger human health in the long term. Human breast milk contains numerous macro and micronutrients such as carbohydrates, fats, proteins, antibodies, and hormones (Kılıç Altun et al., 2018) which play a fundamental role in infants' psychological and immunological development (Salmani et al., 2016). The outstanding composition of breast milk makes it a unique source of nourishment for infants; hence, mothers are highly recommended to exclusively breast-feed their children for at least six months (Ecsedi-Angyal et al., 2020). Breast, due to containing lipophilic tissue and lipids (such as

triacylglycerols, phospholipids, fatty acids, and sterols), is susceptible to the accumulation of compounds with lipophilic nature like potentially toxic elements (PTEs). In other words, humans are exposed to various types of environmental contaminants through several exposure routes such as inhalation, ingestion, and dermal absorption, resulting in the accumulation of PTEs in the body tissues and, consequently, different levels of health risks (Bounar et al., 2020; Chen et al., 2022; Gao et al., 2022; Heshmati et al., 2020; Khanjani et al., 2018; Salmani et al., 2018; Wongsasuluk et al., 2020).

Breast milk can transfer such substances from the mother to the infant, who is biologically more vulnerable to PTEs than adults (Salmani et al., 2016). High elemental absorption due to an immature intestinal system and undeveloped immune system and detoxification mechanisms

* Corresponding authors.

E-mail addresses: aoghazyan@yahoo.com (A. Oghazyan), Aliatamaleki69@yahoo.com (A. Atamaleki).<https://doi.org/10.1016/j.jfca.2022.104933>

Received 12 May 2022; Received in revised form 17 September 2022; Accepted 20 September 2022

Available online 23 September 2022

0889-1575/© 2022 Elsevier Inc. All rights reserved.

Table 1
Meta-analysis concentration of Pb ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Austria	2	0.200	0.181	0.219	33.23	28.62	17	0.038	40.60%
Saudi Arabia	1	4.018	2.474	5.562	0.19	6.78	1	0.009	85.30%
Pakistan	1	11.391	5.956	16.827	0.39	336.84	2	< 0.001	99.40%
Greece	1	0.041	0.000	0.083	4.47	40.97	1	< 0.001	97.60%
India	1	0.328	0.250	0.405	13.45	224.86	11	< 0.001	95.10%
Iran	4	5.233	4.774	5.691	6.61	907.29	47	< 0.001	94.80%
Turkey	4	0.307	0.274	0.340	15.42	2339.08	28	< 0.001	98.80%
Sweden	1	0.195	0.165	0.225	2.17	0.00	0	.	.%
Taiwan	1	1.231	0.550	1.913	5.06	293.91	3	< 0.001	99.00%
Egypt	1	39.011	5.162	72.860	0	25.88	2	< 0.001	92.30%
Iraq	1	1.509	0.936	2.082	1.32	9.86	1	0.002	89.90%
Poland	1	0.107	0.096	0.118	4.45	1.13	1	0.288	11.40%
Nigeria	2	4.719	3.910	5.527	2.41	110.24	4	< 0.001	96.40%
Cyprus	1	0.155	0.100	0.210	2.02	0.00	0	.	.%
Brazil	2	35.155	0.000	101.174	0.53	176.07	1	< 0.001	99.40%
Romania	1	0.026	0.000	0.077	2.05	0.00	0	.	.%
Hungary	1	0.253	0.230	0.277	6.11	0.34	2	0.844	0.00%
USA	1	1.123	0.406	1.840	0.11	0.00	0	.	.%
Jordan	1	10.062	8.069	12.055	0.02	0.00	0	.	.%
Overall	28	0.611	0.586	0.635	100	15,569.09	139	< 0.001	99.10%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO "normal condition levels" in human breast milk ($\mu\text{g}/\text{l}$): 2–5(1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

by the kidney and liver make infants more susceptible to the adverse effects of these toxicants (Ekeanyanwu et al., 2020; Fakhri et al., 2019). Therefore, although the benefits of human breast milk make it the best nutrition for newborns and infants, it can be a route of exposure for infants to toxic substances (Al-Saleh et al., 2003; Tahboub et al., 2021).

PTEs are of considerable interest due to their toxicity and widespread use caused by fossil fuels, industrial by-products (Fakhri et al., 2021; Hu et al., 2022; Luo et al., 2022; Nematollahi et al., 2021; Xiao et al., 2010), atmospheric deposits (Malakootian et al., 2021), agricultural lands, pesticides, herbicides, insecticides, fertilizers applications, waste incinerators, and sewage sludge (Rezaei et al., 2020). Moreover, several studies identified them in human breast milk, suggesting the exposure of infants to these toxicants and consequent health effects during lactation (Ekeanyanwu et al., 2020; Kunter et al., 2017; Olszowski et al., 2016; Samiee et al., 2019). Additionally, some parameters can affect the level of PTEs in human milk, including place of residence (higher in industrial and urban areas), maternal age (higher in older mothers), stage of lactation (higher in colostrum), smoking habits, maternal dietary intakes, and parity (Cherkani-Hassani et al., 2019). A significant correlation between PTEs concentration in breastfeeding mothers and sociodemographic characteristics such as mother's age, educational level, fish, fruit and vegetable consumption, family income, lipstick use, employment status, and smoking habits was noted (Vahidinia et al., 2019).

Based on previous studies, heavy metals such as Cd, Pb, As, and Cr result in the incidence of autism, rickets, and anemia, affect neurological development and infants growth, and decrease the intelligence quotient (IQ) score (Ekeanyanwu et al., 2020; Szukalska et al., 2021; Wongsasuluk et al., 2020; Zoghi et al., 2022). Accordingly, exposure to PTEs can affect the mother's health, breast milk quality, and infant health status (Wongsasuluk et al., 2020).

Assessment of PTEs concentration in human breast milk was investigated in different studies. The study of Pb and Cd levels in the breast milk of lactating mothers in Lebanon showed a range of 0.87 ± 1.18 and $18.18 \pm 13.31 \mu\text{g}/\text{kg}$ for Cd and Pb, respectively (Bassil et al., 2018). In another work in Slovak, results showed that the average Cd, Pb, and Hg concentrations in breast milk samples were 0.43, 4.7, and 0.94 mg/kg, respectively (Ursinyova and Masanova, 2005). Also, PTEs such as Se, Zn, Cu, and Mn have been detected in breast milk samples from Korea while the Mn concentration exceeds daily recommended requirements (Kim

et al., 2012). The toxicological analysis resulting from the determination of breast milk levels from Jordanian mothers indicated high intakes of As and Pb than their provisional tolerable weekly intake (PTWI) (Tahboub et al., 2021).

Thus, Identification and quantification of these contaminants in the mother milk and systematic investigation of the correlation between sociodemographic characteristics and PTEs can be used to assess the mother's exposure as well as respective health risks in infants (Bastos et al., 2018; Olowoyo et al., 2021; Samiee et al., 2019). Prolonged contact with PTEs can cause adverse effects on infant health (Ecse-di-Angyal et al., 2020; Samiee et al., 2019). Safety and quality assurance of breast milk is of great importance and has been the subject of many surveys. However, no meta-analysis study regarding PTEs in human breast milk and following health risks of infants was published. Accordingly, the present work is conducted to review the concentration of potentially toxic elements in human milk and estimate the consequent health risk in the infant.

2. Methods

2.1. Study identification and selection

According to the PRISMA protocol, a search was conducted on international databases, including Scopus, PubMed, and Web of Sciences, for 1 January 2000–10 August 2021. The search was conducted using the following terms: "metal OR heavy metal OR elements OR trace element OR potential toxic element or potential hazard element" AND "breast milk OR human milk OR women's milk".

2.2. Inclusion and exclusion criteria and extraction of data

The inclusion criteria were English language articles and descriptive and cross-sectional studies that measured PTEs concentration in human milk. The studies with unclear data, systematic reviews, narrative reviews, clinical trials, experimental studies, and dissertations were excluded during the selection process. The data were extracted, including the year of publication, country, sample size, and concentration statistics (Mean, standard deviation/ standard error). The concentrations based on fresh weight were included in the meta-analysis. To convert the weights, Eq. 1 (Table 1S) was used. Studies on mother milk




Fig. 1. Flow chart for searching process.




Fig. 2. statistics of investigated PTEs among countries.

with positive smoking were excluded. [Table 1](#).

2.3. Meta-analysis

The data was extracted using concentration statistics (mean and SE), and STATA 14.00 software was used for the meta-analysis ([Aranega and Oliveira, 2022](#); [De Souza et al., 2021](#)). The parameter of SE was calculated according to Eq. 2 ([Table 1S](#)). The heterogeneity of data is a crucial parameter for determining the used model. Depending on the I^2 index, a random effect model ($> 50\%$) or fixed-effect model ($< 50\%$) for meta-analysis are used. The fixed-effect model assumes the same effect size for all studies, while the random effect model assumes various effect sizes among studies using Cochrane's Q-test and I^2 statistical analysis ([Atamaleki et al., 2021](#); [Borenstein et al., 2011](#); [Higgins et al., 2019](#)).

2.4. Health risk assessment

In the current study, non-carcinogenic and carcinogenic risks due to PTEs in human milk were estimated in infants according to the equation in [Table 1s](#).

3. Results and discussion

3.1. Selection process and characteristics of studies

From those databases (Scopus, Web of Science, and PubMed) searched systematically between 2000 and 2021, 32 articles with 181 data reports were included. The study selection process is illustrated in [Fig. 1](#). The number of studies conducted on of PTEs in human milk are shown in [Fig. 2](#). Accordingly, the percentage of countries based on number of study was: Turkey (12.12%) $>$ Brazil (8.33%) $>$ Iran $>$ (7.58%) $>$ Poland (6.82%) \sim Nigeria (6.82%) $>$ Sweden (6.06%) \sim

Table 2
Meta-analysis concentration of Cd ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Saudi Arabia	1	0.216	0.092	0.34	5.19	34.11	1	< 0.001	97.10%
Pakistan	1	6.886	3.589	10.183	0.06	53.02	2	< 0.001	96.20%
Greece	1	0.022	0.016	0.028	5.92	8.16	1	0.004	87.70%
India	1	0.113	0.084	0.142	20.06	153.1	11	< 0.001	92.80%
Iran	3	0.299	0.258	0.34	12.37	18.94	7	0.008	63.00%
Turkey	1	3.484	2.102	4.867	0.01	2.40	2	0.301	16.60%
Sweden	1	0.011	0.01	0.013	2.96	0.00	0	.	%.%
Taiwan	1	0.083	0.061	0.104	22.11	135.76	7	< 0.001	94.80%
Egypt	1	1.638	0.785	2.492	0.06	3.93	2	0.14	49.10%
Iraq	1	0.998	0.349	1.647	0.52	11.98	1	0.001	91.70%
Poland	2	0.011	0.005	0.017	5.93	31.82	1	< 0.001	96.90%
Nigeria	2	2.953	0.835	5.070	3.13	805.85	4	< 0.001	99.50%
Cyprus	1	0.058	0.050	0.067	2.93	0.00	0	.	%.%
Brazil	2	4.833	0.000	14.213	3.21	5688.05	1	< 0.001	100.00%
Romania	1	0.052	0.016	0.088	4.27	1.18E+ 00	1	0.277	15.30%
Hungary	1	0.025	0.023	0.028	8.88	3.53	2	0.171	43.30%
USA	1	0.040	0.000	0.095	1.99	0.00	0	.	%.%
Jordan	1	0.820	0.616	1.025	0.38	0.00	0	.	%.%
Overall	23	0.161	0.147	0.174	100	11,182.75	60	< 0.001	99.50%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO “normal condition levels” in human breast milk ($\mu\text{g}/\text{l}$): < 1 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

Table 3
Meta-analysis concentration of Hg ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Austria	2	0.212	0.190	0.234	26.34	55.08	17	< 0.001	69.10%
Saudi Arabia	1	0.408	0.159	0.658	2.89	20.96	1	< 0.001	95.20%
Pakistan	2	1.003	0.945	1.061	27.01	5468.54	25	< 0.001	99.50%
India	1	0.244	0.183	0.305	11.24	202.37	11	< 0.001	94.60%
Iran	1	0.120	0.097	0.142	1.64	0.00	0	.	%.%
Turkey	1	0.237	0.198	0.276	28.61	473.72	19	< 0.001	96.00%
Brazil	1	0.669	0.000	1.893	2.28	96.84	1	< 0.001	99.00%
Overall	9	0.444	0.419	0.468	100	7488.04	80	< 0.001	98.90%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO “normal condition levels” in human breast milk ($\mu\text{g}/\text{l}$): 1.4–1.7 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

Table 4
Meta-analysis concentration of Cu ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Greece	1	50.490	47.930	53.040	10.97	0.12	1	0.729	0.00%
Turkey	2	60.566	51.259	69.874	12.19	7.87	3	0.049	61.90%
Sweden	1	61.230	58.763	63.697	6.11	0.00	0	.	%.%
Egypt	1	31.580	21.420	41.739	11.94	11.19	2	0.004	82.10%
Iraq	1	13.293	5.153	21.434	12.53	45.01	1	< 0.001	97.80%
Poland	1	54.790	44.459	65.121	9.39	4.46	1	0.035	77.60%
Nigeria	1	4.747	3.857	5.637	25.63	72.34	3	< 0.001	95.90%
Romania	1	76.267	61.101	91.433	4.17	1.76	1	0.184	43.30%
USA	1	1.299	1.122	1.475	6.42	0.00	0	.	%.%
Jordan	1	122.200	89.173	155.227	0.64	0.00	0	.	%.%
Overall	11	32.287	29.502	35.071	100	6321.88	21	< 0.001	99.70%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO “normal condition levels” in human breast milk ($\mu\text{g}/\text{l}$): 180–310 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

Pakistan (6.06%) ~ Jordan (6.06%) ~ Egypt (6.06%) > Romania (5.30%) > USA (4.55%) ~ Austria (4.55%) ~ Greece (4.55%) > India (3.79%) > Taiwan (2.27%) ~ Saudi Arabia (2.27%) ~ Iraq (2.27%) ~ Hungary (2.27%) ~ Cyprus (2.27%).

3.2. Meta-analysis finding

3.2.1. The concentration of PTEs based on the countries

According to meta-analysis results (Tables 1–9), the concentrations of each PTE in human milk varied widely from one country to another, which can be correlated with the difference in the sources of these pollutants in different countries (Guo et al., 2022). PTEs contamination

Table 5
Meta-analysis concentration of As ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Pakistan	2	0.261	0.220	0.302	44.95	4171.5	25	< 0.001	99.40%
India	1	0.181	0.139	0.224	15.08	311.97	11	< 0.001	96.50%
Iran	2	0.114	0.098	0.130	15.6	39.90	9	< 0.001	77.40%
Turkey	1	0.130	0.091	0.169	1.68	0.00	0	.	.%
Sweden	1	0.072	0.048	0.095	1.75	0.00	0	.	.%
Taiwan	1	0.079	0.009	0.148	6.15	21.24	3	< 0.001	85.90%
Egypt	1	22.818	0.794	44.842	0.02	19.40	2	< 0.001	89.70%
Poland	1	0.227	0.129	0.325	3.45	21.77	1	< 0.001	95.40%
Cyprus	1	0.095	0.074	0.116	1.76	0.00	0	.	.%
Brazil	2	4.016	0.000	12.018	1.8	41.29	1	< 0.001	97.60%
Romania	1	0.595	0.379	0.811	2.35	6.80	1	0.009	85.30%
Hungary	1	0.054	0.043	0.064	5.35	6.66	2	0.036	70.00%
Jordan	1	4.082	3.381	4.783	0.08	0.00	0	.	.%
Overall	16	0.207	0.186	0.227	100	6053.96	67	< 0.001	98.90%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO "normal condition levels" in human breast milk ($\mu\text{g}/\text{l}$): 0.2–0.6 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

Table 6
Meta-analysis concentration of Fe ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Pakistan	1	393.556	346.901	440.211	55.94	733.85	22	< 0.001	97.00%
Greece	1	65.411	54.469	76.353	7.45	4.31	1	0.038	76.80%
Turkey	2	154.698	122.357	187.04	9.13	5.99	3	0.112	49.90%
Sweden	1	44.070	39.662	48.478	3.77	0.00	0	.	.%
Egypt	1	3990.000	1096.800	6880.000	0.1	17.27	2	< 0.001	88.40%
Poland	1	708.000	86.710	1330.000	3.34	96.75	1	< 0.001	99.00%
Nigeria	1	6.128	4.434	7.823	15.15	27.61	3	< 0.001	89.10%
Brazil	1	87.562	77.950	97.173	3.69	0.00	0	.	.%
Jordan	1	421.200	348.425	493.975	1.43	0.00	0	.	.%
Overall	10	258.443	247.398	269.487	100	9387.37	40	< 0.001	99.60%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO "normal condition levels" in human breast milk ($\mu\text{g}/\text{l}$): 350–720 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

Table 7
Meta-analysis concentration of Cr ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Pakistan	1	2.874	2.623	3.125	69.12	325.65	22	< 0.001	93.20%
Turkey	1	17.393	9.552	25.234	1.11	6.92	2	0.031	71.10%
Sweden	1	0.039	0.000	0.927	1.93	0.00	0	.	.%
Egypt	1	8.264	3.654	12.875	1.01	7.41	2	0.025	73.00%
Poland	1	0.060	0.020	0.100	6.64	24.07	1	< 0.001	95.80%
Nigeria	1	2.690	2.060	3.330	9.21	17.06	3	0.001	82.40%
Brazil	1	0.571	0.514	0.628	3.31	0.00	0	.	.%
Romania	1	0.705	0.565	0.845	6.4	0.00	1	0.949	0.00%
USA	1	1.828	0.030	3.626	0.84	0.00	0	.	.%
Jordan	1	17.160	14.471	19.849	0.44	0.00	0	.	.%
Overall	10	2.620	2.430	2.811	100	8747.04	40	< 0.001	99.50%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO "normal condition levels" in human breast milk ($\mu\text{g}/\text{l}$): 0.8–1.5 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

can be caused by natural (migration and redistribution of soil) and human activities (mining, abandoned mining, fertilizer and pesticide application, and sewage irrigation) (Atamaleki et al., 2021). For example, Egypt has the highest concentration of Pb, As, Fe, and Mn, which can be associated with their agricultural and mining activities, sewage wastes, the nature of sediments, and anthropogenic activities (paper, pulp, ferrosilicon factories, and phosphate mining) (Al Naggar et al., 2018; El-Amier et al., 2017; Hasballah and Beheary, 2016). Waseem et al. (2014) reported that the concentrations of PTEs increased

from the south to the north in Egypt along the Nile River. They also reported that industrial activities such as sand quarry, shale mining, the nitrogen fertilizer factory at Aswan, and toxic waste drain during seasonal flash floods are the primary sources of PTEs in Egypt. The highest concentrations for Cd, Hg, and Zn in Pakistan were directly associated with environmental challenges such as unbalanced economic and social development (Khalid et al., 2020; Nighat et al., 2016; Peixoto et al., 2019; Rehman et al., 2021). Accidental urban congestion is the main reason for the deterioration of air, water, and soil quality resources

Table 8
Meta-analysis concentration of Zn ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Pakistan	1	7272.063	6796.897	7747.228	0.36	204.15	22	< 0.001	89.20%
Greece	1	512.916	268.950	756.882	9.64	144.15	1	< 0.001	99.30%
Turkey	3	410.252	355.928	464.577	18.16	57.39	6	< 0.001	89.50%
Sweden	1	451.230	419.027	483.433	4.24	0.00	0	.	.%
Egypt	1	132.000	62.810	200.000	11.32	18.79	2	< 0.001	89.40%
Poland	1	769.000	28.740	1510.000	3.36	181.61	1	< 0.001	99.40%
Nigeria	2	12.576	2.052	23.100	43.24	5858.57	4	< 0.001	99.90%
Romania	1	963.116	803.083	1123.149	0.33	0.96	1	0.328	0.00%
USA	1	2.088	1.691	2.485	8.66	0.00	0	.	.%
Jordan	1	484.900	377.052	592.748	0.68	0.00	0	.	.%
Overall	13	205.156	195.875	214.436	100	20,892.84	46	< 0.001	99.80%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO "normal condition levels" in human breast milk ($\mu\text{g}/\text{l}$): 0.7–2 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

Table 9
Meta-analysis concentration of Mn ($\mu\text{g}/\text{Kg}$ -wet weight) in human milk based on country*.

Country	Number of studies	ES	Lower	Upper	Weight (%)	Heterogeneity statistic	df ¹	P value ²	I ² (%) ³
Pakistan	1	9.193	8.363	10.023	38.96	385.16	22	< 0.001	94.30%
Greece	1	0.514	0.303	0.726	6.23	27.54	1	< 0.001	96.40%
India	1	0.525	0.426	0.625	32.94	420.91	11	< 0.001	97.40%
Turkey	1	14.589	7.951	21.227	0.88	5.80	2	0.055	65.50%
Sweden	1	0.390	0.340	0.440	3.12	0.00	0	.	.%
Egypt	1	20.400	5.380	35.500	2.19	19.60	2	< 0.001	89.80%
Poland	1	0.892	0.785	1.000	6.19	2.57	1	0.109	61.00%
Brazil	2	7.908	0.000	22.418	3.31	47.15	1	< 0.001	97.90%
Romania	1	2.114	1.487	2.741	4.13	1.33	1	0.249	24.80%
USA	1	0.933	0.000	1.921	1.7	0.00	0	.	.%
Jordan	1	10.478	7.447	13.509	0.35	0.00	0	.	.%
Overall	12	4.304	4.113	4.495	100	10,386.81	51	< 0.001	99.50%

Degrees of freedom

P value of heterogeneity

Estimates of heterogeneity

* WHO "normal condition levels" in human breast milk ($\mu\text{g}/\text{l}$): 3–4 (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991)

(Waseem et al., 2014).

3.2.2. The rank order of PTEs in the human milk

The rank order of PTEs in the breast milk samples was Fe (258.44 $\mu\text{g}/\text{kg}$) > Zn (205.16 $\mu\text{g}/\text{kg}$) > Cu (32.29 $\mu\text{g}/\text{kg}$) > Mn (4.30 $\mu\text{g}/\text{kg}$) > Cr (2.62 $\mu\text{g}/\text{kg}$) > Hg (0.44 $\mu\text{g}/\text{kg}$) > As (0.21 $\mu\text{g}/\text{kg}$) > Cd (0.16 $\mu\text{g}/\text{kg}$) > Pb (0.03 $\mu\text{g}/\text{kg}$). Compare the WHO recommendation commonly known as "normal condition levels" in human breast milk (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991), concentrations of all PTEs are below the allowable level, except for Zn and Mn. These results are significant for neonatal intensive care units and high-risk neonates, especially preterm infants.

The primary sources of PTEs in the environment are natural, agricultural, industrial solid waste, inland effluent, and atmospheric sources. Human activities (mining, electroplating, metallurgical smelting actions, and using pesticides and fertilizers) also significantly aggravate environmental pollution (Medfu Tarekegn et al., 2020). The three main pathways of PTEs entering the body are ingestion, inhalation, and dermal contact absorption; some studies have reported that ingestion is the primary pathway for human health risks (Tong Sh et al., 2020).

Fe has the highest PTEs concentration in breast milk as Fe is the fourth most abundant element in the Earth's crust and the most abundant transition metal, and most commonly found in nature in its oxides (Bhateria and Singh, 2019; Faust, 2018; Hassanien and Akl, 2018). It can be absorbed into the human body from different environmental matrices, such as air, water, soil, and food (Abbaspour et al., 2014; Ezeonu et al., 2002; Kamble et al., 2013; Kenzhebayeva et al., 2019). Furthermore, Fe plays a crucial role in several biological reactions in the

human body and has an important part in an array of enzymes, systems of energy transduction, and oxygen carriers. Its deficiency in infants is significant and can impair brain development and cause anemia (Peixoto et al., 2019; Sigel et al., 2013). In most cases, pregnant women cannot change their dietary habits. Therefore, human milk fortification with oral Fe is necessary (Idemili-Aronu et al., 2022; Mahundi, 2021; Mattila et al., 2021; Milman et al., 2016).

Zn is part of more than three thousand proteins found in the human body, also playing regulatory functions in mammalian cells and plays an important role during embryogenesis and embryo development (Boskabadi et al., 2021; Dumrongwongsiri et al., 2022; Peixoto et al., 2019). Zn concentrations (205.16 $\mu\text{g}/\text{kg}$) was higher than WHO "normal condition levels" (0.7–2 $\mu\text{g}/\text{kg}$). For Zn, like Fe, the two ways of environmental exposure from different pathways and supplements can effectively increase its concentration. However, determining each scenario's effectiveness is impossible and requires long-term interventions and cohort studies. Iqbal and Ali (Iqbal and Ali, 2021) reported that although Zn supplements reduce the risk of preterm birth, no substantial effect was found for other fetomaternal outcomes. They also emphasized that further studies need to decide the recommended Zn dose or intake during pregnancy. On the other hand, according to (Castillo-Castañeda et al., 2019), the presence of Zn ions can affect Cu absorption in breast milk.

Cu is the 25th most abundant element in the Earth's crust and the third most used metal globally (Shabbir et al., 2020). Cu is also essential for several enzymes (ceruloplasmin, cytochrome oxidase, catechol oxidase), which present electron transport, oxidation, and melanin synthesis (Khalid et al., 2020; Nighat et al., 2016; Rehman et al., 2021).

Based on the results of (de Oliveira Trinta et al., 2020), Cu correlates with immunological increases and antioxidant processes (activity versus reactive oxygen species, ROS) in infants.

Mn is an essential enzymatic cofactor for carbohydrate and lipid metabolism and a micronutrient for bone growth (Kumar et al., 2017). Although several studies reported that Fe and Mn content in mothers' milk is not enough for the nutritional requirements of preterm infants (Domellöf, 2021; Oliveira et al., 2020), the results of the present study showed that their concentration in the breast milk is higher than WHO "normal condition levels." This high concentration, similar to the above, is likely due to supplements during pregnancy. Therefore, it is suggested that in future studies, drugs used by mothers during pregnancy be recorded in order to be able to analyze their effect in this regard. In the environment, Mn is present in rocks, soil, water, and food, and its exposure can be started before birth from maternal exposure. According to (Conley et al., 2022), although bone Mn levels were strongly associated with its blood concentration, it may not be a suitable biomarker of cumulative long-term exposure (Milatovic et al., 2022).

Cr is an essential micronutrient present in supplements used during pregnancy and alters the antioxidant status in the brain. Although Cr(III) is an essential microelement for normal human metabolic activity, Excessive exposure can harm health (Wang et al., 2018). In contrast, Cr (VI) has high toxicity and carcinogenesis (Kaur et al., 2020; Yoshinaga et al., 2018). Some major Cr (VI) mechanisms that cause cellular damage are high levels of oxidative stress, chromosome breaks, and DNA adduct formation (DesMarias and Costa, 2019; Kariminejada et al., 2021). It is also classified as a Group 'A' human carcinogen (Wise et al., 2022). Some studies report that Cr is secreted significantly in breast milk (Cardoso et al., 2014; Elaridi et al., 2020; Samiee et al., 2019). Although one of the most critical exposure pathways is anthropogenic sources and industrial activities (Ukhurebor et al., 2021); but (Motas et al., 2021) reported that the highest concentrations of Cr, Mn, and Fe in human milk were detected in women living in the agricultural zone.

Hg is a potentially toxic element that has received much attention due to its toxicity, even in low concentrations (Raj and Maiti, 2019). WHO "normal condition levels" in human breast milk for Hg were 1.4–1.7 (µg/l) (1 g of human breast milk is equivalent to 0.970 mL) (Bansa et al., 2017; Parr et al., 1991). Exposure of pregnant women, lactating mothers, and weaning offspring to Hg is a tremendous public health concern, especially in the high consumption of carnivorous fish or sea mammals (Sakamoto et al., 2022). Cherkani-Hassani et al. (2021) reported a significant relationship between Hg concentration in breast milk and the occurrence of previous miscarriages, anemia before pregnancy, taking supplements during pregnancy, consumption of cereals, and the use of lipstick (Örün et al., 2021) stated there is a relationship between high Hg concentration and maternal vitamin use. It may have been related to the Hg contamination of vitamin supplements, or these mothers are prone to take medicine for various complaints due to subtle Hg exposure.

As is a metalloid that occurs in many minerals. Exposure to As is usually due to the consumption of water and seafood, especially shellfish. Exposure to Pb and Hg can cause neurotoxic problems. The organic forms of As are relatively nontoxic compared to the inorganic forms (Bassil et al., 2018). Although (Samiee et al., 2019) reported that the transfer of As to the mammary glands is limited, and this issue protects infants from exposure to As, there is an information gap in the discrimination between intrauterine postnatal effects (Rebello and Caldas, 2016).

Cd is reported among the top 10 toxic metals on the Agency for Toxic Substances and Disease Registry (Huang et al., 2019). During pregnancy, Cd can be transferred to the fetus from placental transfer. This process continues after birth in the lactation period through the transport from the maternal blood to the mammary glands (Cherkani-Hassani et al., 2017). According to (Cherkani-Hassani et al., 2020) study, Cd concentration in human milk has a significant association with the area of residence (low in mothers living in urban areas), gestational duration

(high in mothers who delivered preterm), mode of delivery (high in mothers who delivered by cesarean), level of maternal education (high in mothers by low level of education), use of cosmetic powders (high when high frequency using), dietary habits (high when the frequency of consumption of milk, dairy products and dry fruits and low in a high frequency of consumption of wheat bread).

Pb is a PTEs naturally occurring in the earth's crust without known biological activity (Yurdakök, 2015). Pb can cause central nervous system disorders, encephalitis, anemia, hepatitis, kidney, liver, heart, and blood vessel failure, damage to the immune system, the genital system, the digestive tract, and the development of cancer. LetiniL et al. (2016) reported that Pb concentrations were almost 2-fold higher in the colostrum of non-smokers and about 2.5 times higher who continued smoking during pregnancy. According to (Dursun et al., 2016) study, There was a negative correlation between the level of Pb and Hg and a positive correlation between the level of Pb and Cd in breast milk. However, As, Cd, Cr, Hg, and Pb, even in small amounts, are hazardous and can accumulate in the mother's skeleton and release during the lactation period (Lorenzetti et al., 2021). Due to the competitive effect between calcium absorption and Cd, Hg, and Pb, adequate calcium intake during pregnancy and lactation reduces their mobilization in bones and releases Pb during lactation (Lorenzetti et al., 2021; Rădulescu and Lundgren, 2019; Shiyu et al., 2020). Also, exposure of infants to toxic metals during pregnancy was more than through breastfeeding (Bansa et al., 2017).

Ultimately, it is necessary to compare our results with a similar systematic review study (Ghane et al., 2022). Based on the results obtained, PTE concentrations in the breast milk were Cu (1.84 mg/kg) > Zn (1.80 mg/kg) > Fe (1.03 mg/kg) > Ni (0.60 mg/kg) > Pb (0.10 mg/kg) > As (0.15 mg/kg) ≈ Cd (0.15 mg/kg) and the non-carcinogenic risk assessment of the PTEs in breast milk indicated different patterns in various countries, and the calculated TTHQ level of metals in infants was < 1 value. As it is known, the results of the two studies are different from each other. One of the most important reasons for this difference in the results of the two studies can be related to the literature search strategy. In the present study, collected literature studies were published between 2000 and 2021, and in Ghane et al., articles were published between 1982 and 2020. Given the current policy momentum on universal health coverage and primary health care reform globally, there is an urgent need for high-quality and new studies to evaluate the data available from existing studies. The selection of more recent studies makes meta-analysis results more real and valuable. The next difference is in the calculation of health risk assessment. In Ghane et al., standard deviation, the main input variable for calculating health risk assessment, was considered for the ingestion rate of human breast milk, EF, Ed, and BW, and the average life expectancy. However, in the present study, the standard deviation was considered for per capita consumption and body weight. In addition, based on studies conducted by (Fakhri et al., 2022; Mahdavi et al., 2022), BW does not have log-normal distribution, but in Ghane et al., BW has log-normal distribution.

3.3. Analytical methods and instruments for measuring PTEs in human milk

Choosing suitable analysis methods and instruments for PTEs is a severe challenge due to the complexity of the human milk matrix (Draghici et al., 2011; Zeiner et al., 2007). Accordingly, many analytical techniques have been used for measuring PTEs, such as Thin layer chromatography (TLC) (Jumde and Gurnule, 2015; Kholghi et al., 2020), UV-VIS spectrometry (Hasankola et al., 2020; Zhou et al., 2019), Atomic absorption spectrometry (AAS) by Flame Atomic Absorption (FAA) (Muhib et al., 2016; Wang et al., 2015), Graphite Furnace Absorption (GFA) (Forero-Mendieta et al., 2022; Zhong et al., 2016) and Vapor Generation Absorption (VGA) (Mostafavi and Ebrahimi, 2019; Perelonia et al., 2021), X-ray fluorescence (XRF) (Byers et al., 2019; Zhou et al.,

Table 10
Non-carcinogenic (THQ) risk in the infants due to ingestion of human milk.

Country	AS	Cd	Total Cr	Cu	Fe	Methyl Hg	Mn	Pb	Zn
Austria	NC	NC	NC	NC	NC	2.61E-01	NC	6.80E-03	NC
Brazil	1.65E+ 00	5.95E-01	NC	NC	1.54E-02	8.24E-01	7.00E-03	1.20E+ 00	NC
Cyprus	3.90E-02	7.10E-03	NC	NC	NC	NC	NC	5.30E-03	NC
Egypt	9.37E+ 00	2.02E-01	NC	9.73E-02	7.02E-01	NC	1.80E-02	1.33E+ 00	5.42E-02
Greece	NC	2.70E-03	NC	1.56E-01	1.15E-02	NC	5.00E-04	1.40E-03	2.11E-01
Hungary	2.20E-02	3.10E-03	NC	NC	NC	NC	NC	8.70E-03	NC
India	7.40E-02	1.39E-02	NC	NC	NC	3.01E-01	5.00E-04	1.12E-02	NC
Iran	4.70E-02	3.68E-02	NC	NC	NC	1.48E-01	NC	1.79E-01	NC
Iraq	NC	1.23E-01	NC	4.09E-02	NC	NC	NC	5.16E-02	NC
Jordan	1.68E+ 00	1.01E-01	NC	3.76E-01	7.41E-02	NC	9.20E-03	3.44E-01	1.99E-01
Nigeria	NC	3.64E-01	NC	1.46E-02	1.10E-03	NC	NC	1.62E-01	5.20E-03
Pakistan	1.07E-01	8.48E-01	NC	NC	6.93E-02	1.24E+ 00	8.10E-03	3.90E-01	2.99E+ 00
Poland	9.30E-02	1.30E-03	NC	1.69E-01	1.25E-01	NC	8.00E-04	3.70E-03	3.16E-01
Romania	2.44E-01	6.40E-03	NC	2.35E-01	NC	NC	1.90E-03	9.00E-04	3.96E-01
Saudi Arabia	NC	2.66E-02	NC	NC	NC	5.03E-01	NC	1.38E-01	NC
Sweden	3.00E-02	1.40E-03	NC	1.89E-01	7.80E-03	NC	3.00E-04	6.70E-03	1.85E-01
Taiwan	3.20E-02	1.02E-02	NC	NC	NC	NC	NC	4.21E-02	NC
Turkey	5.30E-02	4.29E-01	NC	1.87E-01	2.72E-02	2.92E-01	1.28E-02	1.05E-02	1.69E-01
USA	NC	4.90E-03	NC	4.00E-03	NC	NC	8.00E-04	3.84E-02	9.00E-04

References: Bansa et al. (2017); Parr et al. (1991)

2018), Electrochemical Methods(Cui et al., 2015; Ding et al., 2021), Inductively Coupled Plasma Optical Emission Spectroscopy (ICP-OES) (Fathabad et al., 2018; Yamini and Safari, 2018) and Inductively coupled plasma mass spectrometry (ICP-MS)(Ashrit et al., 2018; Zhang et al., 2020). In some studies, combining at least two analytical methods helps achieve the best results in different matri(Kwon et al., 2018; Li et al., 2021). Among them, AAS and ICP-MS/AES, due to high sensitivity, low levels of detail(LOD), and simultaneous determination of multiple elements, are suitable for measuring PTEs in many samples(Jin et al., 2020).

Among the studies reviewed in the present study (32 articles), analytical selected methods for measuring PTEs in human milk in 58.97%(n = 23) of the studies were AAS(10.26%(n = 4)FAA, 15.38% (n = 6)VGA, 20.51%(n = 8)GFA and 12.82%(n = 5) Blind), 30.77% (n = 12) of the studies was ICP/MS, 2.56%(n = 1) of the studies was Gamma Spectrometry and 7.69%(n = 3) of the studies was ICP/OES. It should be noted that in some studies (n = 8), a combination of two methods was used. Additional information is also presented in Table 3S. These methods have strengths and limitations, which are reviewed in previous studies (Arjomandi and Shirkanloo, 2019; Llaver et al., 2021; Tyler and Jobin Yvon, 1995; Yuan et al., 2011). Among them, AAS and ICP-MS/AES, due to high sensitivity, low levels of detail(LOD), and simultaneous determination of multiple elements, are more often used for analyzing and detecting many samples (Jin et al., 2020). Significantly, the ICP-MS, also known as a green analytical method, have preferences such as high sensitivity, wide linear range, and strong anti-interference (Chojnacka and Mikulewicz, 2019; He et al., 2013).

However, different parameters should be considered for selecting the best method, such as the costs of the analyses, the time required, the required amount of samples, the limit of detection, sensitivity, and selectivity (Hung et al., 2010; Zeiner et al., 2007). In the first step, the PTEs concentration ranges should be estimated qualitatively. Then, the method for an exact quantification may be chosen depending on the analytical task (Zeiner et al., 2007).

3.4. Health risk assessment

Human health risk assessment integrates exposure and exposure-response data to predict the hazard potential for infants from PTEs exposures. Health risks caused by PTEs that enter the infant's body from different exposure pathways are divided into carcinogenic and non-carcinogenic risks (Shams et al., 2022). For this purpose, both non-carcinogenic risks (Eq. 4 and 5) and carcinogenic risks (Eq. 6) were estimated.

The Target Hazard Quotient (THQ) assessed non-carcinogenic health risks. When THQ > 1, infants are threatened (Atamaleki et al., 2021). In present study, The Lowest and highest THQ in infants due to AS was Hungary (2.20E-02) and Egypt (9.37E+00); Cd, Poland (0.13E-02) and Pakistan (0.84E-2); Cu, USA (40E-4) and Jordan(0.37E-2); Fe, Nigeria (11.00E-4) and Egypt(0.70E-2); Methyl Hg, Iran(0.14E-2) and Pakistan(1.23E00); Mn, Sweden(3.00E-4) and Egypt(0.01E-2); Pb, Greece (14.00E-4) and Egypt (1.33E0.00); and Zn, USA(9.00E-4) and Pakistan(2.96E0.00), respectively (Table 10). The high value of THQ for As, Hg, Pb, and Zn can be associated with a high concentration of PTEs, and continuous and effective monitoring is required to reduce the amount of PTEs contaminants.

Also, Total Target Health Quotient (TTHQ) values of PTEs in the human milk were computed. TTHQ shows the sum of each THQ for the whole PTEs. If TTHQ > 1, infants are considered carcinogenic health risks (Atamaleki et al., 2021). Accordingly, in the present study the rank order of country based on TTHQ was Egypt (11.778) > Pakistan (5.643) > Brazil (4.293) > Jordan (2.780) > Turkey (1.179) > Romania (0.883) > Poland (0.707) > Saudi Arabia (0.667) > Nigeria (0.546) > Sweden (0.419) > Iran (0.410) > India (0.400) > Greece (0.382) > Austria (0.268) > Iraq (0.215) > Taiwan (0.084) > Cyprus (0.051) > USA (0.049) > Hungary (0.033) (Fig. 1). TTHQ in Egypt, Pakistan, Brazil, Jordan, and Turkey was higher than 1; infants are not at safe, non-carcinogenic risk (Fig. 1).

In the following, Carcinogenic Risk (CR) was used as the cumulative probability of individual developing cancer. Following US-EPA guidance, a CR < 1E-06 represents negligible levels, values CR < 1E-4–1E-6 are acceptable levels, while CR > 1E-4 signifies a high cancer risk to humans (Atamaleki et al., 2021). The rank order of country based on CR of inorganic As was Egypt (4.22E-03) > Jordan (7.54E-04) > Brazil (7.42E-04) > Romania (1.10E-04) > Pakistan (4.82E-05) > Poland (4.19E-05) > India (3.34E-05) > Turkey (2.40E-05) > Iran (2.11E-05) > Cyprus (1.76E-05) > Taiwan (1.46E-05) > Sweden (1.33E-05) > Hungary (9.98E-06). CR in Egypt, Jordan, Brazil, and Romania was an unsafe cancer risk (CR>1E-4), and CR in other countries was in threshold cancer risk (CR <1E-4–1E-6) (Fig. 2).

4. Conclusion

After a systematic search among Scopus, PubMed, and Web of Science databases, 32 studies measured the concentration of PTEs in human milk included in this work. The extracted data were meta-analyzed, and their health effects were assessed via the health risk models (carcinogenic and non-carcinogenic). According to the meta-analysis results, Fe




Fig. 3. TTHQ in infant due to consumption of mother milk content of heavy metals.




Fig. 4. CR in infant due to consumption of mother milk content of inorganic As.

had the most concentration in human milk, followed by Zn, Cu, Mn, Cr, Hg, As, Cd, and Pb, respectively. The total non-carcinogenic risk showed a considerable effect (TTHQ > 1) for Egypt, Pakistan, Brazil, Jordan, and Turkey. The counties, including Egypt, Jordan, Brazil, and Romania, also were at an unsafe carcinogenic risk (CR > 1E-4). As discussed above, lactating mothers can intake PTEs from different sources. However, their diet can be the main reason directly affecting the milk's content. Therefore, monitoring their environment, feeding behavior (especially supplements taken during pregnancy), and the quality of food consumption is necessary to reduce the exposure. Periodic bio-monitoring surveys in each country can also help estimate the level of exposure of lactating women and newborns. The reason remains unclear without developing our knowledge about the screened mothers' diet. (Figs. 3 and 4).

Ethical approval

Not applicable.

Consent to participate

The authors declare their Consent to Participate in this article.

Consent to publish

The authors declare their Consent to Publish this article.

CRedit authorship contribution statement

Aliasghar Neshat: Investigation, Data curation, Resources, Conceptualization, Methodology, Writing – original draft. **Ali Oghazyan:** Investigation, Data curation, Resources, Conceptualization, Methodology, Writing – original draft. **Fatemeh Kariminejad,** Investigation, Data curation, Resources, Conceptualization, Methodology, Writing – original draft. **Trias Mahmudiono:** Literature searching, Writing – review & editing. **Yadolah Fakhri:** Literature searching, Writing – review & editing. **Amir Mohammad Sheikh Asadi:** Literature searching, Writing – review & editing. **Amir Mohammad Sheikh Asadi:** Supervision, Writing – review & editing. **Ali Atamaleki:** Supervision, Writing – review & editing. **Amin Mousavi Khaneghah:** Supervision, Writing – review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

No data was used for the research described in the article.

Acknowledgments

This study was supported by Esfarayen Faculty of Medical Sciences [Ethical code: IR.ESFARAYENUMS.REC.1400.020].

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.jfca.2022.104933.

References

- Abbaspour, N., Hurrell, R., Kelishadi, R., 2014. Review on iron and its importance for human health. *J. Res. Med. Sci. Off. J. Isfahan Univ. Med. Sci.* 19 (2), 164.
- Al Nagggar, Y., Khalil, M.S., Ghorab, M.A., 2018. Environmental pollution by heavy metals in the aquatic ecosystems of Egypt. *Open Acc. J. Toxicol.* 3, 555603.
- Al-Saleh, I., Shinwari, N., Mashhour, A., 2003. Heavy metal concentrations in the breast milk of Saudi women. *Biol. Trace Elem. Res.* 96 (1), 21–37.
- Aranega, J.P., Oliveira, C.A., 2022. Occurrence of mycotoxins in pastures: a systematic review. *Qual. Assur. Saf. Crops Foods* 14 (3), 135–144.
- Arjomandi, M., Shirkhanloo, H., 2019. A review: analytical methods for heavy metals determination in environment and human samples. *Anal. Methods Environ. Chem. J.* 2 (03), 97–126.
- Ashrit, S., Chatti, R.V., Sarkar, S., Venugopal, R., Nair, G.U., 2018. Application of ICP-MS technique for analysis of heavy metals in LD slag fines. *Curr. Sci.* 115 (5), 973–977.
- Atamaleki, A., Yazdanbakhsh, A., Gholizadeh, A., Naimi, N., Karimi, P., Thai, V.N., Fakhri, Y., 2021. Concentration of potentially harmful elements (PHEs) in eggplant vegetable (*Solanum melongena*) irrigated with wastewater: a systematic review and meta-analysis and probabilistic health risk assessment. *Int. J. Environ. Health Res.* 1–13.
- Bansa, D.K., Awua, A.K., Boatun, R., Adom, T., Brown-Appiah, E.C., Amewosina, K.K., Diaba, A., Datoghe, D., Okwabi, W., 2017. Cross-sectional assessment of infants' exposure to toxic metals through breast milk in a prospective cohort study of mining communities in Ghana. *BMC Public Health* 17 (1), 1–12.
- Bassil, M., Daou, F., Hassan, H., Yamani, O., Abi Kharma, J., Attieh, Z., Elaridi, J., 2018. Lead, cadmium and arsenic in human milk and their socio-demographic and lifestyle determinants in Lebanon. *Chemosphere* 191, 911–921.
- Bastos, W.R., Vieira, S.M., Manzatto, A.G., Dórea, J.G., Rubira, M.C., de Souza, V.F.P., da Costa Junior, W.A., Souza Bastos, M.T., 2018. Heterogeneity of multimedia exposures to neurotoxic elements (Al, As, Cd, Pb, Mn, and Hg) in breastfed infants from Porto Velho, Brazil. *Biol. Trace Elem. Res.* 184 (1), 7–15.
- Bhateria, R., Singh, R., 2019. A review on nanotechnological application of magnetic iron oxides for heavy metal removal. *J. Water Process Eng.* 31, 100845.
- Borenstein, M., Hedges, L.V., Higgins, J.P.T., Rothstein, H.R., 2011. *Introduction to Meta-Analysis*. Wiley.
- Boskabadi, H., Maamouri, G., Akhondian, J., Ashrafzadeh, F., Boskabadi, A., Faramarzi, R., Heidari, E., Pourbadakhsan, N., Shojaei, S.R.H., Zakerihamidi, M., 2021. Comparison of birth weights of neonates of mothers receiving vs. not receiving zinc supplement at pregnancy. *BMC Pregnancy Childbirth* 21 (1), 1–7.
- Boumar, A., Boukaka, K., Leghouchi, E., 2020. Determination of heavy metals in tomatoes cultivated under green houses and human health risk assessment. *Qual. Assur. Saf. Crops Foods* 12 (1), 76–86.
- Byers, H.L., McHenry, L.J., Grundl, T.J., 2019. XRF techniques to quantify heavy metals in vegetables at low detection limits. *Food Chem. X* 1, 100001.
- Cardoso, O.O., Julião, F.C., Alves, R.I., Baena, A.R., Díez, I.G., Suzuki, M.N., Celere, B.S., Nadal, M., Domingo, J.L., Segura-Muñoz, S.I., 2014. Concentration profiles of metals in breast milk, drinking water, and soil: relationship between matrices. *Biol. Trace Elem. Res.* 160 (1), 116–122.
- Castillo-Castañeda, P.C., García-González, A., Bencomo-Alvarez, A.E., Barros-Núñez, P., Gaxiola-Robles, R., Méndez-Rodríguez, L.C., Zenteno-Savín, T., 2019. Micronutrient content and antioxidant enzyme activities in human breast milk. *J. Trace Elem. Med. Biol.* 51, 36–41.
- Chen, F., Aqeel, M., Maqsood, M.F., Khalid, N., Irshad, M.K., Ibrahim, M., Akhter, N., Afzaal, M., Ma, J., Hashem, M., 2022. Mitigation of lead toxicity in *Vigna radiata* genotypes by silver nanoparticles. *Environ. Pollut.* 308, 119606.
- Cherkani-Hassani, A., Ghanname, I., Mouane, N., 2017. Assessment of cadmium levels in human breast milk and the affecting factors: a systematic review, 1971–2014. *Crit. Rev. Food Sci. Nutr.* 57 (11), 2377–2391.
- Cherkani-Hassani, A., Ghanname, I., Benitez-Rexach, A.M., Mouane, N., 2019. Systematic review of the literature of factors affecting the exposure and the levels of lead in human breast milk. *Rev. Environ. Contam. Toxicol. Volume* 252, 97–129.
- Cherkani-Hassani, A., Slaoui, M., Ghanname, I., Mojemmi, B., Belhaj, A., Kers, B., Flayou, M., Mouane, N., 2020. Cadmium contamination in breast milk of Moroccan lactating women and the associated factors: contamilk study. *Biol. Trace Elem. Res.* 196 (1), 47–59.
- Cherkani-Hassani, A., Slaoui, M., Ghanname, I., Mojemmi, B., Belhaj, A., Kers, B., Flayou, M., Mouane, N., 2021. Levels of mercury in Moroccan breast milk and the affecting factors: CONTAMILK study. *Regul. Toxicol. Pharmacol.* 127, 105065.
- Chojnacka, K., Mikulewicz, M., 2019. Green analytical methods of metals determination in biosorption studies. *TrAC Trends Anal. Chem.* 116, 254–265.
- Conley, T.E., Richardson, C., Pacheco, J., Dave, N., Jursa, T., Guazzetti, S., Lucchini, R. G., Fendorf, S., Ritchie, R.O., Smith, D.R., 2022. Bone manganese is a sensitive biomarker of ongoing elevated manganese exposure, but does not accumulate across the lifespan. *Environ. Res.* 204, 112355.
- Cui, L., Wu, J., Ju, H., 2015. Electrochemical sensing of heavy metal ions with inorganic, organic and bio-materials. *Biosens. Bioelectron.* 63, 276–286.
- De Souza, C., Khaneghah, A.M., Oliveira, C.A.F., 2021. The occurrence of aflatoxin M1 in industrial and traditional fermented milk: a systematic review study. *Ital. J. Food Sci.* 33 (SP1), 12–23.
- DesMarias, T.L., Costa, M., 2019. Mechanisms of chromium-induced toxicity. *Curr. Opin. Toxicol.* 14, 1–7.
- Ding, Q., Li, C., Wang, H., Xu, C., Kuang, H., 2021. Electrochemical detection of heavy metal ions in water. *Chem. Commun.* 57 (59), 7215–7231.
- Domellöf, M., 2021. Microminerals: iron, zinc, copper, selenium, manganese, iodine, chromium and molybdenum. *Nutr. Care Preterm Infants* 122, 140–148.
- Draghici, C., Jelescu, C., Dima, C., Coman, G., Chirila, E., 2011. Heavy metals determination in environmental and biological samples. *Environmental Heavy Metal Pollution and Effects on Child Mental Development*. Springer, pp. 145–158.
- Dumrongwongsi, O., Winichagoon, P., Chongviriyaphan, N., Suthutvoravut, U., Grote, V., Koletzko, B., 2022. Zinc and iron adequacy and relative importance of zinc/iron storage and intakes among breastfed infants. *Matern. Child Nutr.* 18 (1), e13268.
- Dursun, A., Yurdakok, K., Yalcin, S.S., Tekinalp, G., Aykut, O., Orhan, G., Morgil, G.K., 2016. Maternal risk factors associated with lead, mercury and cadmium levels in umbilical cord blood, breast milk and newborn hair. *J. Matern. Fetal Neonatal Med.* 29 (6), 954–961.
- Ecsedi-Angyal, M., Tatár, E., Óvári, M., Kurin-Csörgei, K., Záray, G., Mihucz, V.G., 2020. Determination of low-level arsenic, lead, cadmium and mercury concentration in breast milk of Hungarian women. *Int. J. Environ. Anal. Chem.* 100 (5), 549–566.
- Ekeanyanwu, C.L., Alisi, C.S., Ekeanyanwu, R.C., 2020. Levels of Aflatoxin M1 and selected heavy metals (Pb, Cd, Cr, Cu, Zn, Fe, As, and Hg) in the breast milk of lactating mothers in South Eastern, Nigeria. *Food Control* 112, 107150.
- El-Amier, Y.A., Elnaggar, A.A., El-Alfy, M.A., 2017. Evaluation and mapping spatial distribution of bottom sediment heavy metal contamination in Burullus Lake. *Egypt. J. Basic Appl. Sci.* 4 (1), 55–66.
- Elaridi, J., Dimassi, H., Estephan, M., Hassan, H.F., 2020. Determination of aluminum, chromium, and barium concentrations in infant formula marketed in Lebanon. *J. Food Prot.* 83 (10), 1738–1744.
- Ezeonu, F.C., Musa, A., Udedi, S.C., Edeogu, O.C., 2002. Iron and zinc status in soils, water and staple food cultivars in Itakpe, Kogi State of Nigeria. *Environmentalist* 22 (3), 237–240.
- Fakhri, Y., Rahmani, J., Oliveira, C.A.F., Franco, L.T., Corassin, C.H., Saba, S., Rafique, J., Khaneghah, A.M., 2019. Aflatoxin M1 in human breast milk: a global systematic review, meta-analysis, and risk assessment study (Monte Carlo simulation). *Trends Food Sci. Technol.* 88, 333–342.
- Fakhri, Y., Hoseinvandtabar, S., Heidarinejad, Z., Borzoei, M., Bagheri, M., Dehbandi, R., Khaneghah, A.M., 2021. The concentration of potentially hazardous elements (PHEs) in the muscle of blue crabs (*Callinectes sapidus*) and associated health risk. *Chemosphere* 279, 130431.
- Fakhri, Y., Omar, S.S., Mehri, F., Hoseinvandtabar, S., Mahmudiono, T., 2022. Global systematic review and meta-analysis on prevalence and concentration of aflatoxins in peanuts oil and probabilistic risk assessment. *Rev. Environ. Health.*
- Fathabad, A.E., Shariatifar, N., Moazzen, M., Nazmara, S., Fakhri, Y., Alimohammadi, M., Azari, A., Khaneghah, A.M., 2018. Determination of heavy

- metal content of processed fruit products from Tehran's market using ICP-OES: a risk assessment study. *Food Chem. Toxicol.* 115, 436–446.
- Faust, B.C., 2018. A review of the photochemical redox reactions of iron (III) species in atmospheric, oceanic, and surface waters: influences on geochemical cycles and oxidant formation. *Aquat. Surf. Photochem.* 3–38.
- Forero-Mendieta, J.R., Varón-Calderón, J.D., Varela-Martínez, D.A., Riaño-Herrera, D.A., Acosta-Velásquez, R.D., Benavides-Piracón, J.A., 2022. Validation of an analytical method for the determination of manganese and lead in human hair and nails using graphite furnace atomic absorption spectrometry. *Separations* 9 (7), 158.
- Gao, L., Huang, X., Wang, P., Chen, Z., Hao, Q., Bai, S., Tang, S., Li, C., Qin, D., 2022. Concentrations and health risk assessment of 24 residual heavy metals in Chinese mitten crab (*Eriocheir sinensis*). *Qual. Assur. Saf. Crops Foods* 14 (1), 82–91.
- Ghane, E.T., Khanverdilo, S., Mehri, F., 2022. The concentration and health risk of potentially toxic elements (PTEs) in the breast milk of mothers: a systematic review and meta-analysis. *J. Trace Elem. Med. Biol.*, 126998.
- Guo, G., Li, K., Zhang, D., Lei, M., 2022. Quantitative source apportionment and associated driving factor identification for soil potential toxicity elements via combining receptor models, SOM, and geo-detector method. *Sci. Total Environ.*, 154721.
- Hasankola, Z.S., Rahimi, R., Shayegan, H., Moradi, E., Safarifard, V., 2020. Removal of Hg²⁺ heavy metal ion using a highly stable mesoporous porphyrinic zirconium metal-organic framework. *Inorg. Chim. Acta* 501, 119264.
- Hasballah, A.F., Beheary, M.S., 2016. Detection of heavy metals in breast milk and drinking water in damietta governorate, Egypt. *Asian J. Biol.* 1 (2), 1–7.
- Hassanien, A.S., Akl, A.A., 2018. Optical characteristics of iron oxide thin films prepared by spray pyrolysis technique at different substrate temperatures. *Appl. Phys. A* 124 (11), 1–16.
- He, B., Yun, Z., Shi, J., Jiang, G., 2013. Research progress of heavy metal pollution in China: sources, analytical methods, status, and toxicity. *Chin. Sci. Bull.* 58 (2), 134–140.
- Heshmati, A., Mehri, F., Karami-Momtaz, J., Khaneghah, A.M., 2020. The concentration and health risk of potentially toxic elements in black and green tea—both bagged and loose-leaf. *Qual. Assur. Saf. Crops Foods* 12 (3), 140–150.
- Higgins, J.P.T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M.J., Welch, V.A., 2019. *Cochrane Handbook for Systematic Reviews of Interventions*. Wiley.
- Hu, L., Wang, X., Zou, Y., Wu, D., Gao, G., Zhong, Z., Liu, Y., Hu, S., Fan, H., Zhang, B., 2022. Effects of inorganic and organic selenium intervention on resistance of radish to arsenic stress. *Ital. J. Food Sci.* 34 (1), 44–58.
- Huang, S., Kuang, J., Zhou, F., Jia, Q., Lu, Q., Feng, C., Yang, W., Fan, G., 2019. The association between prenatal cadmium exposure and birth weight: a systematic review and meta-analysis of available evidence. *Environ. Pollut.* 251, 699–707.
- Hung, Y.-L., Hsiung, T.-M., Chen, Y.-Y., Huang, Y.-F., Huang, C.-C., 2010. Colorimetric detection of heavy metal ions using label-free gold nanoparticles and alkanethiols. *J. Phys. Chem. C* 114 (39), 16329–16334.
- Idemili-Aronu, N., Igweonu, O., Onyehoro, N., 2022. Uptake of iron supplements and anemia during pregnancy in Nigeria. *J. Public Health* 44 (1), 111–120.
- Iqbal, S., Ali, I., 2021. Effect of maternal zinc supplementation or zinc status on pregnancy complications and perinatal outcomes: an umbrella review of meta-analyses. *Heliyon* 7 (7), e07540.
- Jin, M., Yuan, H., Liu, B., Peng, J., Xu, L., Yang, D., 2020. Review of the distribution and detection methods of heavy metals in the environment. *Anal. Methods* 12 (48), 5747–5766.
- Jumde, M.H., Gurnule, W.B., 2015. Separation and identification of heavy metal ions by thin layer chromatography on silica gel-G. *J. Chem. Pharm. Res.* 7 (4), 889–895.
- Kamble, R., Thakare, M., Ingle, A., 2013. Iron in the environment. *Indian J. Environ. Prot.* 33, 881–888.
- Kariminejada, F., Ghadimib, S.B., Rahmanc, F., Haghghid, M., Senec, R.A., Zazoulie, M. A., Heydaria, E.S., 2021. Kinetic and isotherm study of Cr (VI) biosorption from industrial effluents by biomass of dried sludge. *Desalination Water Treat.* 209, 91–104.
- Kaur, H., Sinha, S., Krishnan, V., Koner, R.R., 2020. Photocatalytic reduction and recognition of Cr (VI): new Zn (II)-based metal–organic framework as catalytic surface. *Ind. Eng. Chem. Res.* 59 (18), 8538–8550.
- Ke, Z., Bai, Y., Bai, Y., Chu, Y., Gu, S., Xiang, X., Ding, Y., Zhou, X., 2022. Cold plasma treated air improves the characteristic flavor of Dry-cured black carp through facilitating lipid oxidation. *Food Chem.* 377, 131932.
- Kenzhebayeva, S., Abekova, A., Atabayeva, S., Yernazarova, G., Omirbekova, N., Zhang, G., Turasheva, S., Asrandina, S., Sarsu, F., Wang, Y., 2019. Mutant lines of spring wheat with increased iron, zinc, and micronutrients in grains and enhanced bioavailability for human health. *BioMed. Res. Int.* 2019.
- Khalid, S., Shahid, M., Shah, A.H., Saeed, F., Ali, M., Qaisrani, S.A., Dumat, C., 2020. Heavy metal contamination and exposure risk assessment via drinking groundwater in Vehari, Pakistan. *Environ. Sci. Pollut. Res.* 27 (32), 39852–39864.
- Khanjani, N., Jafari, M., Ahmadi Mousavi, E., 2018. Breast milk contamination with lead and cadmium and its related factors in Kerman, Iran. *J. Environ. Health Sci. Eng.* 16 (2), 323–335.
- Kholghi, N., Amani, H., Malekmahmoodi, S., Amiri, A., 2020. Investigation on heavy metal removal from a crude oil contaminated soil using rhamnolipid biosurfactant as a new eco-friendly method. *Tenside Surfactants Deterg.* 57 (6), 515–520.
- Kılıç Altun, S., Dinç, H., Temamogulları, F.K., Paksoy, N., 2018. Analyses of essential elements and heavy metals by using ICP-MS in maternal breast milk from Şanlıurfa, Turkey. *Int. J. Anal. Chem.* 2018.
- Kim, S.-Y., Park, J.H., Kim, E.A.-R., Lee-Kim, Y.C., 2012. Longitudinal study on trace mineral compositions (selenium, zinc, copper, manganese) in Korean human preterm milk. *J. Korean Med. Sci.* 27 (5), 532–536.
- Kumar, R.K., Singhal, A., Vaidya, U., Banerjee, S., Anwar, F., Rao, S., 2017. Optimizing nutrition in preterm low birth weight infants—consensus summary. *Front. Nutr.* 4, 20.
- Kunter, İ., Hürer, N., Gülcan, H.O., Öztürk, B., Doğan, İ., Şahin, G., 2017. Assessment of aflatoxin M1 and heavy metal levels in mothers breast milk in Famagusta, Cyprus. *Biol. Trace Elem. Res.* 175 (1), 42–49.
- Kwon, J.-Y., Kim, B.-G., Lim, H.-J., Seo, J.-W., Kang, M.-K., Kim, Y.-M., Hong, Y.-S., 2018. Comparison of human blood cadmium concentrations using graphite furnace atomic absorption spectrometry (GF-AAS) and inductively coupled plasma-mass spectrometry (ICP-MS). *J. Environ. Health Sci.* 44 (5), 491–501.
- Lei, J., Dong, Y., Hou, Q., He, Y., Lai, Y., Liao, C., Kawamura, Y., Li, J., Zhang, B., 2022. Intestinal microbiota regulate certain meat quality parameters in chicken. *Front. Nutr.* 9.
- LetiniĹ, J.G., SariĹ, M.M., Piasek, M., JurasoviĹ, J., Varnai, V.M., Grgec, A.S., Orct, T., 2016. Use of human milk in the assessment of toxic metal exposure and essential element status in breastfeeding women and their infants in coastal Croatia. *J. Trace Elem. Med. Biol.* 38, 117–125.
- Li, W., Niu, N., Guo, N., Zhou, H., Bu, J., Ding, A., 2021. Comparative study on the determination of heavy metals in soil by XRF and ICP-MS. *J. Phys. Conf. Ser. IOP Publ.* 012075.
- Llaver, M., Oviedo, M.N., Quintas, P.Y., Wuilloud, R.G., 2021. *Analytical Methods for the Determination of Heavy Metals in Water, Remediation of Heavy Metals*. Springer, pp. 1–50.
- Lorenzetti, S., Plösch, T., Teller, I.C., 2021. Antioxidative molecules in human milk and environmental contaminants. *Antioxidants* 10 (4), 550.
- Luo, C., Sun, J., Tan, Y., Xiong, L., Peng, B., Peng, G., Bai, X., 2022. Comparison of the health risks associated with exposure to toxic metals and metalloids following consumption of freshwater catches in China. *Qual. Assur. Saf. Crops Foods* 14 (4), 1–12.
- Mahdavi, V., Eslami, Z., Omidvari, Z., Rezadoost, H., Fakhri, Y., 2022. Carcinogenic and non-carcinogenic risk assessment induced by pesticide residues in honey of Iran based on Monte Carlo simulation. *J. Food Compos. Anal.* 109, 104521.
- Mahundi, P., (2021). **Knowledge, acceptance and barriers to optimal use of iron supplements amongst pregnant women attending Mutare city clinic in Manicaland, Zimbabwe.**
- Malakootian, M., Mohammadi, A., Nasiri, A., Asadi, A.M.S., Conti, G.O., Faraji, M., 2021. Spatial distribution and correlations among elements in smaller than 75 µm street dust: ecological and probabilistic health risk assessment. *Environ. Geochem. Health* 43 (1), 567–583.
- Mattila, M., Hakola, L., Niinistö, S., Tapanainen, H., Takkinen, H.-M., Ahonen, S., Ilonen, J., Toppari, J., Veijola, R., Kniip, M., 2021. Maternal vitamin C and iron intake during pregnancy and the risk of islet autoimmunity and type 1 diabetes in children: a birth cohort study. *Nutrients* 13 (3), 928.
- Medfu Tarekegn, M., Zewdu Salilih, F., Ishetu, A.I., 2020. Microbes used as a tool for bioremediation of heavy metal from the environment. *Cogent Food Agric.* 6 (1), 1783174.
- Milatovic, D., Gupta, R.C., Yin, Z., Zaja-Milatovic, S., Brockett, M.M., Aschner, M., 2022. Manganese, Reproductive and developmental toxicology. Elsevier, pp. 587–602.
- Milman, N., Paszkowski, T., Cetin, I., Castelo-Branco, C., 2016. Supplementation during pregnancy: beliefs and science. *Gynecol. Endocrinol.* 32 (7), 509–516.
- Mostafavi, S.M., Ebrahimi, A., 2019. Mercury determination in work place air and human biological samples based on dispersive liquid-liquid micro-extraction coupled with cold vapor atomic absorption spectrometry. *Anal. Methods Environ. Chem. J.* 2 (04), 49–58.
- Motas, M., Jiménez, S., Oliva, J., Cámara, M.Á., Pérez-Cárceles, M.D., 2021. Heavy metals and trace elements in human breast milk from industrial/mining and agricultural zones of southeastern Spain. *Int. J. Environ. Res. Public Health* 18 (17), 9289.
- Muhib, M.I., Chowdhury, M.A.Z., Easha, N.J., Rahman, M.M., Shammi, M., Fardous, Z., Bari, M.L., Uddin, M.K., Kurasaki, M., Alam, M.K., 2016. Investigation of heavy metal contents in cow milk samples from area of Dhaka, Bangladesh. *Int. J. Food Contam.* 3 (1), 1–10.
- Nematollahi, A., Abdi, L., Abdi-Moghadam, Z., Fakhri, Y., Borzoei, M., Tajdar-Oranj, B., Thai, V.N., Linh, N.T.T., Mousavi Khaneghah, A., 2021. The concentration of potentially toxic elements (PTEs) in sauses: a systematic review and meta-analysis study. *Environ. Sci. Pollut. Res.* 28 (39), 55186–55201.
- Night, S., Nadeem, M.S., Mahmood, T., Kayani, A.R., Mushtaq, M., ul Hassan, M.M., 2016. Estimation of heavy metals in Indian flying fox *Pteropus giganteus* (Brünnich, 1782) from Punjab, Pakistan. *Pak. J. Zool.* 48, 6.
- Oliveira, M.M., Trevilato, T.M., Segura-Muñoz, S.I., Aragon, D.C., Alves, L.G., Nadal, M., Marquês, M., Domingo, J.L., Sierra, J., Camelo Jr, J.S., 2020. Essential and toxic elements in human milk concentrate with human milk lyophilizate: a preclinical study. *Environ. Res.* 188, 109733.
- de Oliveira Trinta, V., de Carvalho Padilha, P., Petronilho, S., Santelli, R.E., Braz, B.F., Freire, A.S., Saunders, C., da Rocha, H.F., Sanz-Medel, A., Fernández-Sánchez, M.L., 2020. Total metal content and chemical speciation analysis of iron, copper, zinc and iodine in human breast milk using high-performance liquid chromatography separation and inductively coupled plasma mass spectrometry detection. In: *Food chemistry*, 326.
- Olowoyo, J.O., Macheka, L.R., Mamejta, P.M., 2021. Health risk assessments of selected trace elements and factors associated with their levels in human breast milk from pretoria, South Africa. *Int. J. Environ. Res. Public Health* 18 (18), 9754.
- Oliszowska, T., Baranowska-Bosiacka, I., Rebacz-Maron, E., Gutowska, I., JamioĹ, D., Prokopowicz, A., Goschorska, M., Chlubek, D., 2016. Cadmium concentration in mother's blood, milk, and newborn's blood and its correlation with fatty acids,

- anthropometric characteristics, and mother's smoking status. *Biol. Trace Elem. Res.* 174 (1), 8–20.
- Örün, E., Yalçın, S.S., Aykut, O., 2021. Lead, mercury, and cadmium levels in breast milk and infant hair in the late period of lactation in Ankara, Turkey. *Int. J. Environ. Health Res.* 1–12.
- Parr, R.M., DeMaeyer, E.M., Iyengar, V.G., Byrne, A.R., Kirkbright, G.F., Schöch, G., Niinistö, L., Pineda, O., Vis, H.-L., Hofvander, Y., 1991. Minor and trace elements in human milk from Guatemala, Hungary, Nigeria, Philippines, Sweden, and Zaire. *Biol. Trace Elem. Res.* 29 (1), 51–75.
- Peixoto, R.R.A., Codo, C.R.B., Sanches, V.L., Guiraldelo, T.C., da Silva, F.F., Ribessi, R.L., Marba, S.T.M., Cadore, S., 2019. Trace mineral composition of human breast milk from Brazilian mothers. *J. Trace Elem. Med. Biol.* 54, 199–205.
- Perelonia, K.B.S., Benitez, K.C.D., Banicod, R.J.S., Tadifa, G.C., Cambia, F.D., Montojo, U.M., 2021. Validation of an analytical method for the determination of cadmium, lead and mercury in fish and fishery resources by graphite furnace and Cold Vapor Atomic Absorption Spectrometry. *Food Control* 130, 108363.
- Rădulescu, A., Lundgren, S., 2019. A pharmacokinetic model of lead absorption and calcium competitive dynamics. *Sci. Rep.* 9 (1), 1–27.
- Raj, D., Maiti, S.K., 2019. Sources, toxicity, and remediation of mercury: an essence review. *Environ. Monit. Assess.* 191 (9), 1–22.
- Rebello, F.M., Caldas, E.D., 2016. Arsenic, lead, mercury and cadmium: toxicity, levels in breast milk and the risks for breastfed infants. *Environ. Res.* 151, 671–688.
- Rehman, A., Ma, H., Ozturk, I., 2021. Do industrialization, energy importations, and economic progress influence carbon emission in Pakistan. *Environ. Sci. Pollut. Res.* 28 (33), 45840–45852.
- Rezaei, M., Malekiran, A.A., Jabbari, M., Karimi-Dehkordi, M., Ghasemidehkordi, B., Teimoori, H., Fakhri, Y., Khaneghah, A.M., (2020). Essential elements in the different types of fruits, soil, and water samples collected from Markazi province, Iran: a health risk assessment study. *Quality Assurance Safety of Crops Foods* 12(3), 111–125.
- Sakamoto, M., Haraguchi, K., Tatsuta, N., Marumoto, M., Yamamoto, M., Nakamura, M., 2022. Breast milk contribution to tissue mercury levels in rat pups examined by cross-fostering at birth. *Environ. Res.*, 112772
- Salmani, M.H., Mozaffari-Khosravi, H., Rezaei, Z., 2016. The Nickel Concentration In Breast Milk During The First Month Of Lactation In Yazd, Center of Iran. *Biol. Trace Elem. Res.* 174 (1), 65–70.
- Salmani, M.H., Rezaei, Z., Mozaffari-Khosravi, H., Ehrampoush, M.H., 2018. Arsenic exposure to breast-fed infants: contaminated breastfeeding in the first month of birth. *Environ. Sci. Pollut. Res.* 25 (7), 6680–6684.
- Samiee, F., Leili, M., Faradmal, J., Torkshavand, Z., Asadi, G., 2019. Exposure to arsenic through breast milk from mothers exposed to high levels of arsenic in drinking water: Infant risk assessment. *Food Control* 106, 106669.
- Shabbir, Z., Sardar, A., Shabbir, A., Abbas, G., Shamshad, S., Khalid, S., Murtaza, G., Dumat, C., Shahid, M., 2020. Copper uptake, essentiality, toxicity, detoxification and risk assessment in soil-plant environment. *Chemosphere* 259, 127436.
- Shams, M., Tavakkoli Nezhad, N., Dehghan, A., Alidadi, H., Paydar, M., Mohammadi, A. A., Zarei, A., 2022. Heavy metals exposure, carcinogenic and non-carcinogenic human health risks assessment of groundwater around mines in Joghatai, Iran. *Int. J. Environ. Anal. Chem.* 102 (8), 1884–1899.
- Shiyu, Q., Hongen, L., Zhaojun, N., Rengel, Z., Wei, G., Chang, L., Peng, Z., 2020. Toxicity of cadmium and its competition with mineral nutrients for uptake by plants: a review. *Pedosphere* 30 (2), 168–180.
- Sigel, A., Sigel, H., Sigel, R.K., 2013. *Interrelations Between Essential Metal Ions and Human Diseases*. Springer.
- Sun, Y., Li, J., Zhu, L., Jiang, L., 2022. Cooperation and competition between CRISPR and omics-based technologies in foodborne pathogens detection: a state of the art review. *Curr. Opin. Food Sci.* 44, 100813.
- Szukalska, M., Merritt, T.A., Lorenc, W., Sroczynska, K., Miechowicz, I., Komorowicz, I., Mazela, J., Baratkiewicz, D., Florek, E., 2021. Toxic metals in human milk in relation to tobacco smoke exposure. *Environ. Res.* 197, 111090.
- Tahboub, Y.R., Massadeh, A.M., Al-Sheyab, N.A., Nsserat, I.A., 2021. Levels Of Trace Elements In Human Breast Milk In Jordan: A Comparison With Infant Formula Milk Powder. *Biol. Trace Elem. Res.* 199 (11), 4066–4073.
- Tong Sh, Li.H., Wang, L., Tudi, M., Yang, L., 2020. Concentration, spatial distribution, contamination degree and human health risk assessment of heavy metals in urban soils across China between 2003 and 2019—a systematic review. *Int. J. Environ. Res. Public Health* 17 (9), 3099.
- Tyler, G., Jobin Yvon, S., 1995. ICP-OES, ICP-MS and AAS techniques compared. *ICP Opt. Emiss. Spectrosc. Tech. Note* 5.
- Ukhurebor, K.E., Aigbe, U.O., Onyancha, R.B., Nwankwo, W., Osibote, O.A., Paumo, H. K., Ama, O.M., Adetunji, C.O., Siloko, I.U., 2021. Effect of hexavalent chromium on the environment and removal techniques: a review. *J. Environ. Manag.* 280, 111809.
- Ursinyova, M., Masanova, V., 2005. Cadmium, lead and mercury in human milk from Slovakia. *Food Addit. Contam.* 22 (6), 579–589.
- Vahidinia, A., Samiee, F., Faradmal, J., Rahmani, A., Taravati Javad, M., Leili, M., 2019. Mercury, lead, cadmium, and barium levels in human breast milk and factors affecting their concentrations in Hamadan, Iran. *Biol. Trace Elem. Res.* 187 (1), 32–40.
- Wang, H., Jiang, Y., Tian, C., Pan, R., Dang, F., Feng, J., Li, M., Zhang, Y., Li, H., Man, C., 2018. Determination of the transfer of lead and chromium from feed to raw milk in Holstein cows. *Food Addit. Contam.: Part A* 35 (10), 1990–1999.
- Wang, Y., Chen, H., Tang, J., Ye, G., Ge, H., Hu, X., 2015. Preparation of magnetic metal organic frameworks adsorbent modified with mercapto groups for the extraction and analysis of lead in food samples by flame atomic absorption spectrometry. *Food Chem.* 181, 191–197.
- Waseem, A., Arshad, J., Iqbal, F., Sajjad, A., Mehmood, Z., Murtaza, G., 2014. Pollution status of Pakistan: a retrospective review on heavy metal contamination of water, soil, and vegetables. *BioMed. Res. Int.* 2014.
- Wise Jr, J.P., Young, J.L., Cai, J., Cai, L., 2022. Current understanding of hexavalent chromium [Cr (VI)] neurotoxicity and new perspectives. *Environ. Int.* 158, 106877.
- Wongsasuluk, P., Sematong, S., Robson, M., Siri Wong, W., 2020. Heavy metals levels in breast milk of lactating mothers working in heavy metals contaminated factories. *EnvironmentAsia* 13, 1.
- Xiao, Z.L., Cong, Q., Qu, J., 2010. Assessment of heavy metal pollution in orchard soil and its affections to fruit quality around molybdenum mining area. *Sci. Technol. Eng.* 10 (23), 5831–5834.
- Yamini, Y., Safari, M., 2018. Modified magnetic nanoparticles with catechol as a selective sorbent for magnetic solid phase extraction of ultra-trace amounts of heavy metals in water and fruit samples followed by flow injection ICP-OES. *Microchem. J.* 143, 503–511.
- Yang, Y., Zhu, H., Xu, X., Bao, L., Wang, Y., Lin, H., Zheng, C., 2021. Construction of a novel lanthanum carbonate-grafted ZSM-5 zeolite for effective highly selective phosphate removal from wastewater. *Microporous Mesoporous Mater.* 324, 111289.
- Yoshinaga, M., Ninomiya, H., Al Hossain, M.A., Sudo, M., Akhand, A.A., Ahsan, N., Alim, M.A., Khalequzzaman, M., Iida, M., Yajima, I., 2018. A comprehensive study including monitoring, assessment of health effects and development of a remediation method for chromium pollution. *Chemosphere* 201, 667–675.
- Yu, C., Chen, X., Li, N., Zhang, Y., Li, S., Chen, J., Yao, L., Lin, K., Lai, Y., Deng, X., 2022. Ag₃PO₄-based photocatalysts and their application in organic-polluted wastewater treatment. *Environ. Sci. Pollut. Res.* 1–17.
- Yuan, X., Chapman, R.L., Wu, Z., 2011. Analytical methods for heavy metals in herbal medicines. *Phytochem. Anal.* 22 (3), 189–198.
- Yurdakok, K., 2015. Lead, mercury, and cadmium in breast milk. *J. Pediatr. Neonatal Individ. Med.* JPNIM 4 (2) e040223-e040223.
- Zeiner, M., Rezić, I., Steffan, I., 2007. Analytical methods for the determination of heavy metals in the textile industry. *Kem. Ind.* 56 (11), 587–559.
- Zhang, J., Cao, K., Zhang, X., Zhang, Q., 2020. Se-directed synthesis of polymeric carbon nitride with potential applications in heavy metal-containing industrial sewage treatment. *Appl. Organomet. Chem.* 34 (2), e5377.
- Zhong, W.-S., Ren, T., Zhao, L.-J., 2016. Determination of Pb (Lead), Cd (Cadmium), Cr (Chromium), Cu (Copper), and Ni (Nickel) in Chinese tea with high-resolution continuum source graphite furnace atomic absorption spectrometry. *J. Food Drug Anal.* 24 (1), 46–55.
- Zhou, F., Li, C., Zhu, H., Li, Y., 2019. A novel method for simultaneous determination of zinc, nickel, cobalt and copper based on UV–vis spectrometry. *Optik* 182, 58–64.
- Zhou, S., Yuan, Z., Cheng, Q., Zhang, Z., Yang, J., 2018. Rapid in situ determination of heavy metal concentrations in polluted water via portable XRF: Using Cu and Pb as example. *Environ. Pollut.* 243, 1325–1333.
- Zoghi, A., Salimi, M., Mirmahdi, R.S., Massoud, R., Khosravi-Darani, K., Mohammadi, R., Rouhi, M., Tripathy, A.D., 2022. Effect of Pretreatments on bioremoval of metals and subsequent exposure to simulated gastrointestinal conditions. *Qual. Assur. Saf. Crops Foods* 14 (3), 145–155.