

The Role of Andrographolide in Andrographis paniculata as a Potential Analgesic for Herbal Medicine based Drug Development

by Suciati Suciati

Submission date: 28-Dec-2021 02:46PM (UTC+0800)

Submission ID: 1736039324

File name: tential_Analgesic_for_Herbal_Medicine_based_Drug_Development.pdf (226.92K)

Word count: 4551

Character count: 24776

RESEARCH ARTICLE

The Role of Andrographolide in *Andrographis paniculata* as a Potential Analgesic for Herbal Medicine based Drug Development

Lidya Tumewu¹, Irfan Rayi Pamungkas², Hilkatul Ilmi¹, Achmad Fuad Hafid^{1,3},
Indah Setyawati Tantular^{1,4}, Suciati^{1,3}, Tutik Sri Wahyuni^{1,3}, Aty Widyawaruyanti^{1,3*}

¹Natural Product Medicine Research and Development, Institute of Tropical Disease,
Universitas Airlangga, Campus C Universitas Airlangga, Mulyorejo, Surabaya 60115, Indonesia.

²Undergraduate Student of Department of Pharmaceutical Sciences, Faculty of Pharmacy,
Universitas Airlangga, Mulyorejo, Surabaya 60115, Indonesia.

³Department of Pharmaceutical Sciences, Faculty of Pharmacy,
Universitas Airlangga, Mulyorejo, Surabaya 60115, Indonesia.

⁴Department of Parasitology, Faculty of Medicine, Universitas Airlangga,
Mayjen. Prof. Dr. Moestopo 47, Surabaya 60131, Indonesia.

*Corresponding Author E-mail: aty-w@ff.unair.ac.id

ABSTRACT:

Background: *Andrographis paniculata* is a herbaceous plant in the Acanthaceae family, that is widely used as a traditional medicine in Asian countries and known to exhibit a wide range of pharmacological effects. Recent studies have provided an overview of the great potential of *A. paniculata* as an analgesic. The ethanol extract and ethyl acetate (EA) fraction of *A. paniculata* were shown to contain diterpene lactone compounds, which may be useful as a potential active ingredient in analgesic drugs. The development of a herbal medicine based drug requires an effective and high quality active ingredient. Therefore, this research was aimed to compare the analgesic activity of ethanol extract and EA fraction based on their andrographolide content and further to determine the more viable active substance for analgesic herbal medicine based drug development. **Method:** The andrographolide content in the ethanol extract and EA fraction was determined by High Pressure Liquid Chromatography (HPLC). Measurement of analgesic activity was performed by writhing test. The experimental animals were randomly divided into eight groups consisting of 5 mice in each. Group 1 (negative control) received 1% Tween-80 in normal saline. Group 2 (positive control) received a standard analgesic drug (diclofenac sodium) at a dose of 40 mg/kg body weight. Group 3, 4, and 5 received ethanol extract while Group 6, 7, and 8 received EA fraction, each at a dose of 12.5, 25, and 50 mg andrographolide/kg body weight, respectively. Each mouse was injected intraperitoneally with 1% acetic acid at a dose of 10 ml/kg body weight 30 minutes after oral administration of the treatments. The number of writhes were counted 5 min after acetic acid injection over a period of 45 min. **Results:** Andrographolide content in ethanol extract and EA fraction was 15.66±0.28 and 21.25±1.08 % w/w, respectively. Ethanol extract and EA fraction displayed analgesic activity of 67.68% and 70.91% respectively, at a dose of 50 mg andrographolide/kg body weight. The positive control at a dose of 40 mg/kg body weight showed an analgesic activity of 74.33%. Statistical analysis showed no significant differences between EA fraction at a dose of 50 mg andrographolide/kg body weight and ethanol extract at the same dose as well as the positive control ($P > 0.05$). The effective dose 50% (ED₅₀) of the ethanol extract and EA fraction was determined to be 29.49 and 25.55 mg/kg body weight, respectively. **Conclusion:** It was possible to use andrographolide content as an indicator for the analgesic activity of *A. paniculata*. Ethanol extract and EA fraction of *A. paniculata* at the same dose of andrographolide showed similar analgesic activity. The amount of ethanol extract which needed to reach similar analgesic activity was higher than EA fraction. Therefore, EA fraction likely has greater potential as an analgesic active substance due to its higher content of andrographolide; however further study is needed to develop it as a dosage form.

KEYWORDS: *Andrographis paniculata*, extract, ethyl acetate fraction, analgesic activity.

Received on 09.06.2020 Modified on 17.01.2021
Accepted on 15.04.2021 © RJPT All right reserved
Research J. Pharm. and Tech 2021; 14(12):6269-6274.
DOI: 10.52711/0974-360X.2021.01084

INTRODUCTION:

Pain and inflammation are a defense mechanism of body to initiate the tissue healing process. It is a common nonspecific manifestation of many diseases¹. Drugs

presently used for the management of pain and inflammatory conditions are narcotics (e.g., opioids), non-narcotics (e.g., salicylates) or corticosteroids (e.g., hydrocortisone). All of these drugs cause well known side effect, such as a low potency, and toxic effects, such as renal failure, allergic reactions, and occasionally hearing loss while also potentially increasing the risk of haemorrhage by affecting platelet function^{2,3}. The use of nonsteroidal substances such as aspirin is also increasing due to its utility in reducing the incidence of a number of common disorders including stroke, myocardial infarction and cancer⁴. Therefore, the discovery of alternative substances to treat pain is crucial.

Plants are important sources of new drugs. Many plants were reported to have a potential analgesic activity including *Solanum surattense*, *Plumbago indica*, *Clerodendrum inerme*, *Lagenaria siceraria*, *Cuscuta reflexa*, *Alpinia conchigera*, *Sida Spp* and *Andrographis paniculata*. These plants extracts were showed inhibition on acetic acid induced writhing on mice compared to aspirin or diclofenac sodium as a positive control⁵⁻¹².

Andrographis paniculata is a herbaceous plant in the Acanthaceae family that is widely used as a traditional medicine in Asian countries¹³. This plant is known to have pharmacological effects, such as anti-inflammatory, antibacterial, antioxidant, anticancer, antidiabetic, antimalarial, hepatoprotective, immunostimulant, allergic, analgesic, and antipyretic activity¹⁴⁻¹⁷. Several studies have been conducted to determine the analgesic activity of *A. paniculata*. Previous investigations reported that the aqueous extract of *A. paniculata* at a dose of 100 mg/kg and 500 mg/kg per orally, significantly reduced ($P < 0.01$) acetic acid induced writhing in both acute and chronic examinations. The test drug at a dose of 100 mg/kg and 500 mg/kg fared better than the standard drug (aspirin) at 150 mg/kg in acute cases¹². The ethanol extract of *A. paniculata* is known to display an analgesic activity of 34%¹⁸. Andrographolide has also been shown to exhibit analgesic activity at a dose of 4 mg/kg, when given intraperitoneally¹⁹. These studies provide an overview of the significant potential of *A. paniculata* as an alternative analgesic drug.

In consideration of these results, *A. paniculata* has been identified as a prospective plant than could be used to address the current need for alternative analgesic drugs. Although many reports on the analgesic activity of *A. paniculata* have been published, little attention has been directed towards the development of analgesic herbal medicines based on effective and high quality active ingredients. The ethanol extract and ethyl acetate (EA) fraction of *A. paniculata* have been shown to contain diterpene lactone compounds and importantly

andrographolide as a potential active substance. Therefore, this research was aimed to compare the analgesic activity of ethanol extract and EA fraction based on their andrographolide content and to further determine the more viable active substance for analgesic drug development.

MATERIALS AND METHODS:

Plant Material:

The plant material used in this research was *A. paniculata* dried powder containing 1.82% andrographolide (PT. Kimia Farma Tbk). Andrographolide standard was purchased from Sigma-Aldrich (Cat No.365645-100MG).

Preparation of the ethanol extract and ethyl acetate (EA) fraction:

A. paniculata dried powder was extracted by maceration method using ethanol 96% as a solvent. The liquid ethanol extract was then evaporated using a rotary evaporator. The concentrated extract was then further separated via liquid-liquid fractionation using ethyl acetate and water (1:1 v/v) to obtain the ethyl acetate (EA) fraction.

Determination of andrographolide content:

Analysis of andrographolide content in the extract and EA fraction was conducted by High Pressure Liquid Chromatography (HPLC) method. The analysis used an Agilent HPLC system, Poroshell RP-18 column (4.6x250 mm, 5 μ m), mobile phase methanol:water pH 3.05 (50%: 50% v/v) and a flow rate of 1 ml/min. The extract and EA fraction were weighed out (10 mg) and dissolved in 10 ml of methanol. Standard working solutions was made at concentrations of 100, 200, 400, 800, and 900 ppm in methanol, taking into account the 98% standard potential. The sample and standard solutions were filtered using a filter membrane with a pore size of 0.22 μ m. The solutions were subsequently inserted into an auto sampler and injected into the HPLC system. Andrographolide content in the extract and EA fraction was then determined based on the chromatogram peak areas.

Animals:

This study used male mice BALB/C strain (25–30 g), were maintained on standard animal pellets and water *ad libitum* at the Animal Laboratory of the Institute of Tropical Disease, Universitas Airlangga, Surabaya. Animals were kept at constant temperature (25 \pm 1 $^{\circ}$ C) and underwent a regular 12/12 h light/dark cycle, while having free access to standard laboratory feed and water. Permission and approval for animal studies were obtained from the Faculty of Veterinary Medicine, Universitas Airlangga with the approval code 753-KE.

1 Evaluation of analgesic activity by acetic acid induced writhing test:

The mice were divided into eight groups (n = 5). Group 1 (Negative control) received 1% Tween-80 in normal saline. Group 2 (Positive control) received a standard drug (diclofenac sodium) at a dose of 40 mg/kg body weight. Group 3 to Group 8 were treatment groups. Group 3 to Group 5 received ethanol extract at a dose of 12.5, 25, and 50 mg andrographolide/kg body weight respectively. Group 6 to Group 8 received EA fraction at doses of 12.5, 25, and 50 mg andrographolide/kg body weight respectively. All treatments were administered orally. Each mouse was injected with 1% acetic acid intraperitoneally at a dose of 10 ml/kg body weight, 30 minutes after administration of the standard drug and test samples. The number of writhes (constriction of abdominal muscles along with the stretching of hind limbs) were counted 5 min after acetic acid injection over a period of 45 min. The percentage of analgesic activity was calculated as follows²⁰.

$$\% \text{ Analgesic activity} = \frac{N \text{ negative control} - N \text{ test}}{N \text{ negative control}} \times 100$$

Where N is the mean number of writhes for the each group.

3 Statistical analysis:

Mean and standard error of mean (SEM) was calculated for the observed values in each experimental group (n=5). Statistical analysis was performed by one way analysis of variance (ANOVA) followed by Least Significant Difference (LSD). An effect was considered to be significant at the P<0.05 level. Effective Dose 50% (ED₅₀) was analyzed by probit analysis. SPSS 17.0 was used for the statistical analysis.

RESULTS:

Determination of andrographolide content:

The andrographolide content in extract and EA fraction was determined by HPLC method using andrographolide as an external standard. The analysis was done used a Poroshell RP-18 column (4.6x250 mm, 5µm), mobile phase methanol:water pH 3.05 (50%: 50% v/v) and a flow rate of 1 ml/min. Under these conditions, the andrographolide standard was observed at Rt 2.46 min. Meanwhile, andrographolide in extract and EA fraction was detected at Rt 2.51 min (Figure 1). The peak area of the andrographolide standard at concentrations of 100, 200, 400, 800 and 900 ppm was determined. The peak areas of the known standard concentration were used for linier regression analysis as shown in Table 1. Andrographolide content in extract and EA fraction was 15.66±0.28% and 21.25± 1.08% w/w as shown in Table 2.




Figure 1. HPLC Chromatogram of Andrographolide standard at a concentration of 200 ppm (a), ethanol extract of *Andrographis paniculata* (b), EA fraction of *Andrographis paniculata* (c).

Table 1. HPLC results of andrographolide standard (concentration vs area)

Concentration (ppm)	Area
100	209.84308
200	654.00830
400	1522.93140
800	3129.94482
900	3545.97778

Y=4.1461x-177.57, R²=0.9997

Table 2. HPLC result of andrographolide content in extract and EA fraction

Sample	Area	Sample Concentration (ppm)	Andrographolide content (% w/w)	Andrographolide content (% w/w) Average±SD
Extract	486.47580	160.14	16.01	15.66±0.28
	461.17447	154.04	15.40	
	476.88068	157.83	15.78	
	463.30511	154.55	15.45	
EA Fraction	707.80560	213.53	21.35	21.25±1.08
	655.43182	200.90	20.09	
	668.67413	208.91	20.89	
	762.83105	226.80	22.68	

Table 3. Analgesic activity of extract and EA fraction of *A. paniculata* by acetic acid induced writhing in mice

Groups	Dose (mg/kg bw)	Number of writhes in 45 min (Mean±SEM)	Inhibition (%)
Negative control	-	105.2±1.80 ^{abcd}	
Diclofenac sodium	40	27.0±1.92 ^{abc}	74.33±1.83
Ethanol extract	12.5	85.8±1.16 ^{abcd}	18.44±1.10
	25	54.6±1.94 ^{abcd}	48.10±1.84
	50	34.0±1.14 ^{abd}	67.68±1.08
EA fraction	12.5	83.0±1.61 ^{abcd}	21.10±1.53
	25	47.4±1.62 ^{abcd}	54.94±2.49
	50	30.6±1.12 ^b	70.91±1.07

Value are reported as Mean±SEM for all groups. The data was analyzed by ANOVA followed by LSD test. Letters (abcd) indicate statistically significant difference, $P<0.05$

Evaluation of analgesic activity by acetic acid induced writhing test:

The effects of the ethanol extract and EA fraction of *A. paniculata* on the acetic acid-induced abdominal constrictions in mice are presented in Table 3 and Figure 2. The results indicated that ethanol extract, EA fraction of *A. paniculata* (at a dose of andrographolide 12.5, 25, and 50 mg/kg body weight) and diclofenac sodium (40 mg/kg body weight) significantly ($P<0.05$) reduced abdominal writhing in mice when compared to the negative control group. Interestingly, the reduction occurred in a dose dependent manner. Maximum inhibition from treatment with ethanol extract (67.68%) and EA fraction (70.91%) was observed at a andrographolide dose of 50 mg/kg body weight. The effective dose 50% (ED₅₀) was analyzed by probit log analysis using SPSS. The results showed that the ED₅₀ values of extract and EA fraction were 29.49 mg/kg body weight and 25.55 mg/kg body weight, respectively. Hence, the EA fraction displayed higher activity compare to the extract based on their ED₅₀ value.




Figure 2. Percentage of inhibition of abdominal contractions of acetic acid induced writhing in mice.

DISCUSSION:

In order to facilitate the use of *A. paniculata* for its analgesic activity, this study aimed to determine the active substance involved and its potential for development into an analgesic herbal medicine based drug. This was undertaken by comparing the analgesic activity of the ethanol extract and EA fraction based on their andrographolide content.

The determination of andrographolide content in ethanol extract and EA fraction was conducted by HPLC method. Several methods to quantify the andrographolide content in extract and pharmaceutical dosage form were reported including High Performance Thin Layer Chromatography (HPTLC) and spectrophotometric method²¹⁻²³. HPLC methods was adopted in this study because of its simple, precise and accuracy. The determination result reveal that andrographolide content in extract was 15.66±0.28% w/w. Meanwhile, EA fraction was contain andrographolide 21.25± 1.08% w/w, which was higher compared to the extract.

In this study, experimental animals in the form of mice (*Mus musculus*) were divided into eight groups consisting of a positive control group, negative control group, and six treatment groups. The number of animals used for each test group was determined based on the Federer formula²⁴. The ideal number of animals per group according to the Federer formula was a minimum of four animals. The experimental animals were randomly divided into eight groups where each group consisted of five mice. Thus the total for all control and treatment groups amounted to forty male mice. The negative group was given CMC-Na 0.5%. The positive

control group was given the analgesic drug sodium diclofenac at a dose of 40 mg/kg body weight of mice. Group 3 to 5 were treated with 96% *A. paniculata* ethanol extract and Groups 6 to 8 were treated with *A. paniculata* EA fraction at an andrographolide dose of 12.5, 25, and 50 mg/kg body weight of mice.

The acetic acid-induced abdominal constriction method is widely used for the screening new analgesic drugs, because it is very simple and sensitive²⁵. The working principle of this method is to observe the stretching response that occurs due to pain stimulation by an irritant. The irritants are administered by injecting acetic acid as an intraperitoneal pain inducer in mice²⁶. Administering acetic acid can cause a local inflammatory response because of the release of arachidonic acid from phospholipid tissue via Cox, prostaglandin production (PGE2 and PGE2 α) and increased production of lipoxigenase which can increase intraperitoneal fluid. The production of prostaglandin and lipoxigenase causes swelling and release of endogenous substances that stimulate nerve endings in the visceral area, especially in the chest and abdominal cavity so that pain arises. This pain is characterized by the appearance of stretching which is classified as the foot being pulled back (stretched) and the abdomen touching the base of the form plate²⁷⁻²⁹.

Based on ANOVA analysis, it was shown that significant differences occurred between the treatment groups ($P < 0.05$). Hence, the administration of different test materials produced different amount of stretching. Regarding the results of the ethanol extract and EA fraction in the acetic acid-induced abdominal constriction assay, a prominent inhibition of the writhing reflex was observed. The ethanol extract and EA fraction of *A. paniculata* display analgesic activity as determined by significantly ($P < 0.05$) reduced abdominal writhing in mice when compared to the negative control group.

Post hoc-tests were performed using the Least Significant Difference (LSD) analysis to reveal any differences between treatment groups. LSD test results showed that EA fraction at a dose of andrographolide of 12.5 mg/kg was not significantly different to the ethanol extract treatment at the same dose. Furthermore, the EA fraction at an andrographolide dose of 50 mg/kg did not differ significantly from the ethanol extract treatment at the same dose as well as from the positive control. From this it can be concluded that the administration of sodium diclofenac at a dose of 40 mg/kg and the EA fraction at a andrographolide dose of 50 mg/kg have reduce writhing in mice to a similar degree.

Diclofenac sodium, like other non-steroidal anti-inflammatory drugs, inhibits the biogenesis of prostaglandins, thus inhibiting the writhing in experimental animals like mice. The presence of phytochemicals in *A. paniculata*, such as diterpenes, lactones, and flavonoids has been reported in previous work³⁰. Flavonoids were reported to have a role in analgesic activity primarily by targeting prostaglandins^{31,32}. Moreover, andrographolide was detected in diterpene lactone compounds which are known to have analgesic effects¹⁹. Madav et al. (1995) reported that 300 mg/kg of andrographolide, administered orally, had significant analgesic activity on acetic-induced writhing in mice³³. The results of our study showed that both ethanol extract and EA fraction at the same andrographolide dose of 50 mg/kg showed similar analgesic activity. Andrographolide, which was a major compound contained in the ethanol extract and EA fraction can be considered as a marker for analgesic activity. In agreement with previous work, *A. paniculata* analgesic activity was entirely explained by the presence of a high content of andrographolide³⁴. The presence of other secondary metabolites did not influence the analgesic activity as shown by their similar activity. The EA fraction might be chosen as potential active substance in analgesic drug development based on its higher andrographolide content. The amount of EA fraction equal to 50 mg andrographolide (235.30 mg) needed to reach the similar analgesic activity was lower than ethanol extract (319.28 mg). Nevertheless, further study is needed to increase the analgesic activity of the EA fraction to enable practical applications. Specifically, formulating a study of the EA fraction wherein the solubility of active substance is increased will be of great interest study as well as working towards an understanding of the mechanism of action.

CONCLUSION:

Ethanol extract and EA fraction of *A. paniculata* at the same dose of andrographolide showed similar analgesic activity. Andrographolide could take a role as a marker for the analgesic activity of *A. paniculata*. The EA fraction contained a higher andrographolide content compared to the ethanol extract. The amount of ethanol extract which needed to reach similar analgesic activity was higher than EA fraction. Therefore, the EA fraction has great potential as an analgesic active substance, but further study is needed to elucidate the appropriate dosage form.

CONFLICT OF INTEREST:

The authors declare there is no conflict of interest.

ACKNOWLEDGEMENT:

The authors are grateful to the Directorate General of Research Technology and Higher Education, for their

funding through Penelitian Terapan Unggulan Perguruan Tinggi contract No.004/ADD/SP2H/LT/DRPM/VIII/2017.

REFERENCE:

- Tatiya AU, Saluja AK, Kalaskar MG, Surana SJ, Patil PH. Evaluation of analgesic and anti-inflammatory activity of *Bridelia retusa* (Spreng) bark. Journal of Traditional and Complementary Medicine. 2017; 7: 441-51.
- Ahmad F, Khan RA, Rashid S. Pharmacological Evaluation of Medicinal Plants For Their Analgesic Activity In Mice. Medical Journal of the Islamic Republic of Iran. 1996; 10(2): 149-52.
- Thomas MC. Diuretics, ACE inhibitors and NSAIDs - the triple whammy. Med J Aust. 2000; 172: 184-5.
- Dubois RW, Melmed GY, Henning JM, Bernal M. Risk of upper gastrointestinal injury and event in patients treated with cyclooxygenase (COX)-1/COX-2 nonsteroidal antiinflammatory drugs (NSAIDs), COX-2 selective NSAIDs and gastroprotective cotherapy: an appraisal of the literature, J Clin Rheumatol. 2004; 10(4): 78-89.
- Sravathi K, Ramana KV, Sumalatha G, Neeraja K. Evaluation of anti-inflammatory and analgesic potency of whole plant extract of *Solanum surattense* in experimental animals. Research J Pharmacology and Pharmacodynamics 2013; 5(2):101-5.
- Paul S, Saha D. Analgesic activity of methanol extract of *Plumbago indica* (L.) by acetic acid induced writhing method. Asian J Pharm Tech 2012; 2(2):74-6.
- Vijay Amirtharaj R, Suresh V, Senthil Kumar R. Studies on anti-inflammatory and analgesic properties of methanol extract of aerial part of *Clerodendrum inerme* in experimental animal models. Research J Pharmacognosy and Phytochemistry 2010; 2(5): 421-4.
- Deshmukh DB, Sherkar MR. Evaluation of in vivo analgesic and anti-inflammatory activity of ethanolic extract of medicinal plant *Lagenaria siceraria*. Asian J Pharm Tech 2019; 9(2): 75-8.
- Jadhav GB, Udavant PB, Saudagar RB, Upasani CD. Analgesic and antiinflammatory activity of amarvel extracts on experimentally induce pain and inflammation on animals. Research Journal of Pharmacy and Pharmacodynamics 2014; 6(2): 112-7.
- Paul S, Saha D. Analgesic activity of methanol extract of *Alpinia conchigera* Griff by acetic acid induced writhing method. Asian J Res Pharm Sci 2012; 2(2): 62-5.
- Bhaskar VH, Sangameswaran B, Balakrishnan N, Panda AB, Raj NR, Sathis A. Screening of analgesic and anti-inflammatory activity of hydroalcohol extract of Sida (Indian) species root. Research J Pharm Tech 2008; 1(3): 287-9.
- Shivaprakash G, Gopalakrishna HN, Padbidri DS, Sadanand S, Sekhar SS, Nivedita RS. Evaluation of *Andrographis paniculata* leaves extract for analgesic activity. Journal of Pharmacy Research 2011; 4(10): 3375-7.
- Nanduri S, Nyavanandi VK, Thunuguntla S S R, Kasu S, Palleria MK, Ram PS et al. Synthesis and structure-activity relationships of andrographolide analogues as novel cytotoxicagents. Bioorg. Med. Chem. Lett. 2004;14(18): 4711-7.
- Akbar S. *Andrographis paniculata*: A Review of Pharmacological Activities and Clinical Effects. Alternative Medicine Review. 2011;16(1): 66-77.
- Jayakumar T, Hsieh CY, Lee JJ, Sheu JJ. Review article: Experimental and Clinical Pharmacology of *Andrographis paniculata* and Its Major Bioactive Phytoconstituent Andrographolide. Evidence-Based Complementary and Alternative Medicine. 2013. Article ID846740. <https://doi.org/10.1155/2013/846740>
- Joselin J, Jeeva S. *Andrographis paniculata*: A Review of its traditional uses, phytochemistry and pharmacology. Med Aromat Plants. 2014;3:169.
- Okhuarobo A, Falodun JE, Erharuyi O, Imieje V, Falodun A, Langer P. Harnessing the medicinal properties of *Andrographis paniculata* for diseases and beyond: a review of its phytochemistry and pharmacology. Asian Pac J Trop Dis. 2014; 4(3):213-22.
- Hassan MM, Khan SA, Shaikat AH, Hossain ME, Hoque MA, Ullah MH et al. Analgesic and anti-inflammatory effects of ethanol extracted leaves of selected medicinal plants in animal model. Vet World. 2013; 6(2):68-71.
- Suebsasana S, Pongnaratom P, Sattayasai J. Analgesic, antipyretic, antiinflammatory and toxic effects of Andrographolide derivatives in experimental animals. Arch Pharm Res. 2009; 32(9):1191-200.
- Hijazi MA, El-Mallah A, Aboul-Ela M, Ellakany A. Evaluation of analgesic activity of *Papaver libanoticum* extract in mice: involvement of opioids receptor. Evidence based complementary and alternative medicine. 2017. article ID 8935085.
- Bhope SG, Kuber VV, Patil MJ, Ghosh VK. Validated HPTLC method for the quantitation of andrographolide from raw material and pharmaceutical dosage form. Asian J Research Chem 2009; 2(3): 314-7.
- Rajasekaran A, Arivukkaran R, Linda M. Estimation of andrographolide content in aqueous extract of siddha formulation by HPTLC. Asian J Pharm Ana 2015; 5(4): 206-8.
- Suneetha A, Manasa K. Development and validation of visible spectrophotometric method for determination of andrographolide in kalmegh plant extract. Asian J Pharm Ana 2014; 4(2): 85-8.
- Federer WT. Statistics and Society: Data Collection and Interpretation, 2nd Ed. New York: Marcel Dekker. 1991.
- Collier HOJ, Dinneen LC, Johnson CA, Schneider C. The Abdominal Constriction Response and its Suppression by Analgesic Drugs in the Mouse. British J Pharmacol. 1968; 32:295-310.
- Gawade S. Acetic Acid Induced Painful Endogenous Infliction in Writhing Test on Mice. Journal of Pharmacology and Pharmacotherapeutics. 2012; 3(4):348.
- Thompson EB. Drug Bioscreening Fundamental of Drug Evaluation Technique In Pharmacology. New York Publishing Co, Inc. 1985.
- Granger D. Role Of Xanthine Oxidase and Granulocytes in Ischemia-Reperfusion Injury. American Journal of Physiology-Heart and Circulatory Physiology. 1988; 255(6):1269-75.
- Duarte I, Nakamura M, Ferreira S. Participation of the Sympathetic System in Acetic Acid-Induced Writhing In Mice. Brazilian Journal of Medical and Biological Research. 1987; 21(2):341-3.
- Das P, Srivastav AK. Phytochemical Extraction and Characterization of the Leaves of *Andrographis paniculata* for Its Antibacterial, Anti-Oxidant, Anti-Pyretic and Antidiabetic Activity. International Journal of Innovative Research in Science, Engineering and Technology. 2014; 3(8): 15176-84.
- Rajnarayana K, Reddy MS, Chaluvadi MR, Krishna DR. Biflavonoids classification, pharmacological, biochemical effects and therapeutic potential. Indian Journal of Pharmacology. 2001; 33:2-16. <http://www.ijonline.com/article.asp?issn=0253-7613>
- Rao MR, Rao YM, Rao AV, Prabhkar MC, Rao CS, Muralidhar N. Antinociceptive and antiinflammatory activity of a flavonoid isolated from *Caralluma attenuate*. Journal of Ethnopharmacology. 1998; 62: 63-6. DOI: 10.1016/S0378-8741(98)00048-8.
- Madav S, Tripathi HC, Tandan, Mishra SK. Analgesic, antipyretic and antiulcerogenic effects of andrographolide. Indian Journal of Pharmaceutical Sciences. 1995; 57(3):121-5.
- Thakur AK, Rai G, Chatterjee SS, Kumar V. *Andrographis paniculata* and andrographolide in diabetic rodents. EC Pharmaceutical Science. 2015; 1(1): 19-28.

The Role of Andrographolide in Andrographis paniculata as a Potential Analgesic for Herbal Medicine based Drug Development

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	ijpsr.com Internet Source	1%
2	www.amhsr.org Internet Source	1%
3	ecronicon.com Internet Source	1%
4	www.jcimjournal.com Internet Source	1%
5	Antara Ghosh, Sujan Banik, Mohammad Nurul Amin, Jamiuddin Ahmed. "Evaluation of antinociceptive, antihyperglycemic, and membrane stabilizing activities of <i>Garcinia lancifolia</i> Roxb", <i>Journal of Traditional and Complementary Medicine</i> , 2017 Publication	1%
6	cyberleninka.org Internet Source	1%

7	Internet Source	1 %
8	Zhou, Bin, Dayong Zhang, and Xiaoming Wu. "Biological Activities and Corresponding SARs of Andrographolide and Its Derivatives", Mini Reviews in Medicinal Chemistry, 2013. Publication	1 %
9	bmccomplementmedtherapies.biomedcentral.com Internet Source	1 %
10	ir.library.ui.edu.ng Internet Source	<1 %
11	www.ncbi.nlm.nih.gov Internet Source	<1 %
12	pubs.rsc.org Internet Source	<1 %
13	wto-cn.net/www.mobile.jci.org Internet Source	<1 %
14	Sharuti Mehta, Anil Kumar Sharma, Rajesh K. Singh. "Pharmacological Activities and Molecular Mechanisms of Pure and Crude Extract of Andrographis paniculata: An update", Phytomedicine Plus, 2021 Publication	<1 %
15	apjtc.com Internet Source	<1 %

16

www.hilarispublisher.com

Internet Source

<1 %

17

Hossain, Md Mokarram, Rumana Jahangir, SM Raquibul Hasan, Raushanara Akter, Taksim Ahmed, Md Imamul Islam, and Abdullah Faruque. "Antioxidant, analgesic and cytotoxic activity of *Michelia champaca* Linn. Leaf", Stamford Journal of Pharmaceutical Sciences, 2010.

Publication

<1 %

18

armaghanj.yums.ac.ir

Internet Source

<1 %

19

impactfactor.org

Internet Source

<1 %

20

www.ijpsonline.com

Internet Source

<1 %

21

clinphytoscience.springeropen.com

Internet Source

<1 %

22

ijrar.com

Internet Source

<1 %

23

link.springer.com

Internet Source

<1 %

24

www.i-scholar.in

Internet Source

<1 %

25

Alexander G. Panossian, Thomas Efferth, Alexander N. Shikov, Olga N. Pozharitskaya et al. "Evolution of the adaptogenic concept from traditional use to medical systems: Pharmacology of stress - and aging - related diseases", Medicinal Research Reviews, 2020

Publication

<1 %

26

CRS Prakoeswa, FR Natallya, D Harnindya, A Thohiroh et al. "The efficacy of topical human amniotic membrane-mesenchymal stem cell-conditioned medium (hAMMSC-CM) and a mixture of topical hAMMSC-CM + vitamin C and hAMMSC-CM + vitamin E on chronic plantar ulcers in leprosy: A randomized control trial", Journal of Dermatological Treatment, 2018

Publication

<1 %

27

Kilani-Jaziri Soumaya, Mhalla Dhekra, Châbane Fadwa, Ghedira Zied, Limem Ilef, Ghedira Kamel, Chekir-Ghedira Leila. "Pharmacological, antioxidant, genotoxic studies and modulation of rat splenocyte functions by Cyperus rotundus extracts", BMC Complementary and Alternative Medicine, 2013

Publication

<1 %

28

journals.plos.org

Internet Source

<1 %

29

www.phytojournal.com

Internet Source

<1 %

30

N Ibrahim, S Nuryanti, A Hasanuddin, M S Zubair. "The effect of growing place on the oil quality and antioxidant activity of kelor (*Moringa oleifera lamk*) seed", *Journal of Physics: Conference Series*, 2021

Publication

<1 %

31

Nesrin Gökhan-Kelekçi, Meriç Köksal, Songül Ünüvar, Göknur Aktay, Hakki Erdoğan. "Synthesis and characterization of some new 2()-benzoxazolones with analgesic and antiinflammatory activities ", *Journal of Enzyme Inhibition and Medicinal Chemistry*, 2008

Publication

<1 %

32

Preeti Sharma, Swapnil Kumar, Shruti Beriwal, Priyanka Sharma et al. "Comparative transcriptome profiling and co-expression network analysis reveals functionally coordinated genes associated with metabolic processes of *Andrographis paniculata*", *Plant Gene*, 2020

Publication

<1 %

33

Yan Tong. "Analgesic activity of myricetin isolated from *Myrica rubra* Sieb. et Zucc. leaves", *Archives of Pharmacal Research*, 04/2009

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Role of Andrographolide in Andrographis paniculata as a Potential Analgesic for Herbal Medicine based Drug Development

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
