

# Liver Function Characteristics of COVID-19 Patients with Obesity at Dr. Soetomo Hospita: Case Series

*by Erwin Maulana Farmanda Putra*

---

**Submission date:** 09-Apr-2023 09:58AM (UTC+0700)

**Submission ID:** 2059310377

**File name:** BMJ\_Manusricpt.docx (299.2K)

**Word count:** 3370

**Character count:** 18626

## Case Series

2

### Liver Function Characteristics of COVID-19 Patients with Obesity at Dr. Soetomo Hospita: Case Series

Erwin Maulana Farmanda Putra<sup>1</sup>, Ummi Maimunah<sup>2</sup>

1

<sup>1</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo Hospital, Surabaya, Indonesia

<sup>2</sup>Division of Gastroenterology and Hepatology, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo Hospital, Surabaya, Indonesia

Corresponding author:

Ummi Maimunah, Division of Gastroenterology and Hepatology, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo Hospital, Surabaya, 60132, Indonesia; [ummima@gmail.com](mailto:ummima@gmail.com)

#### Email IDs

Erwin M.F. Putra : [erwin.mfp@gmail.com](mailto:erwin.mfp@gmail.com)

Ummi Maimunah : [ummima@gmail.com](mailto:ummima@gmail.com)

## ABSTRACT

**Background:** Coronavirus disease 2019 (COVID-19) patients with obesity are more susceptible to liver injury. There are currently no published overview regarding COVID-19 patients with liver injury in Indonesia. Our study reported 7 cases of obese COVID-19 patients with an increase of serum glutamic oxaloacetic transaminase (SGOT) and serum glutamic pyruvic transaminase (SGPT), followed by 5 of them developing severe symptoms of COVID-19 in isolation room of Dr. Soetomo Hospital Surabaya within June-December 2020.

**Case Presentation:** A total of 7 COVID-19 patients, by the average age  $46.2 \pm 15.5$  years, were obese. The mean body mass index (BMI) was  $36.3 \pm 3.4$  kg/m<sup>2</sup>. Five of them experienced serious symptoms. There were 3 patients (patient no. 2, 4, 7) who had normal SGOT level, while the rest had an increased SGOT level for  $<2x$  upper limit of normal (ULN) at admission. During treatment, SGOT level increased in 5 patients for  $1- <3x$  ULN and normal in the rest of them (patient no. 2, 4). Meanwhile, the normal SGPT levels were shown in 3 patients (patient no. 2, 4, 7) and increased in 4 others at admission for  $1- <4x$  ULN. All of them tended to have an increased SGPT level during treatment for  $1- <5x$  ULN. However, since the increase of SGOT and/or SGPT for  $>3x$  ULN is defined as liver injury, it happened in 2 patients that occur since admission (patient no. 3) and during treatment (patient no. 5).

**Conclusion:** Obesity may impede the recovery of COVID-19 and manifest in the deterioration of liver function.

**Keywords:** COVID-19, liver, angiotensin converting enzyme 2, obesity, SARS-CoV-2

## INTRODUCTION

Coronavirus disease 2019 (COVID-19) is one of the international health concern that have caused long-term consequences<sup>1-3</sup> and have disturbed multi-sectors.<sup>4, 5</sup> Multiple diagnostic methods,<sup>6</sup> prevention measures<sup>7</sup> and treatments<sup>8, 9</sup> have been proposed; however COVID-19 waves are continuing. Obesity is one of factors that aggravates COVID-19, caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2 infection).<sup>10</sup> A study involving 382 COVID-19 patients, 32% of overweight and 10.7% of obese patients, showed that the obese patients were having more severe symptoms compared to the normal and overweight group.<sup>11</sup> In addition to respiratory symptoms, COVID-19 patients are also more susceptible to liver injury during the treatment.<sup>12</sup>

A study conducted in China revealed that 99 COVID-19 patients admitted to intensive care units experienced the improved indicators of liver function such as serum glutamic oxaloacetic transaminase (SGOT) and serum glutamic pyruvic transaminase (SGPT).<sup>13</sup> Another study confirmed that obese or overweight patients with COVID-19 might have liver injury following the lung manifestations.<sup>14</sup> Based on these exposures, this case series is expected to provide an overview of liver function characteristics in COVID-19 patients with obesity at Dr. Soetomo Hospital Surabaya, Indonesia.

## CASE PRESENTATION

A total of 7 patients (5 women, 2 men) included in this study were admitted to isolation room of Dr. Soetomo Hospital Surabaya within June-December 2020. Their laboratory examinations were carried out in the clinical pathology laboratory. The general profiles, symptoms, laboratory results, treatments and outcome of these patients were recorded as shown in Table 1.

The average age of all patients was  $46.2 \pm 15.5$  years. The COVID-19 diagnosis was confirmed by real-time reverse transcription–polymerase chain reaction (RT-PCR) with nasopharyngeal swab specimen. The degree of obesity was assessed by measuring body mass index (BMI) with an average BMI was  $36,3 \pm 3,4$  kg/m<sup>2</sup>. All patients had fever and cough symptoms. Short breath was experienced by 5 patients (71.4%). One (14.2%) complained other symptoms such as nausea, vomiting, anosmia and diarrhea. None of them had chronic liver disease. There were 5 patients (62.5%) identified as severe COVID-19 cases. In terms of preventing the likelihood of chronic hepatitis virus infection, all patients underwent hepatitis B surface antigen (HbsAg) screening.

After admission, all patients received antiviral treatment, such as lopinavir-ritonavir 400mg twice a day in 6 patients (85.7%) and oseltamivir 75mg twice a day in 1 patient (14.3%). All patients received symptomatic treatment, like paracetamol 500mg three times a day. The average length of treatment was  $11.4 \pm 4.9$  days and 4 patients died during treatment.

Table 1. Characteristics of the obese COVID-19 patients.

Category	Overall (n=7)	Obesity Class 1 (n=2)	Obesity Class 2 (n=3)	Obesity Class 3 (n=2)
BMI (kg/m <sup>2</sup> ). mean±SD	36.4±3.4	32.1±1.4	36.8±1.5	40.1±0.1
Age (years). mean±SD	46.2±15.5	37.5±10.6	53±18.1	45±19.7
Male. n (%)	2 (28.5)	2 (100)	0 (0)	0 (0)
Female. n (%)	5 (62.5)	0 (0)	3 (100)	2 (100)
<b>Symptoms. n (%)</b>				
Fever	7 (100)	2 (100)	3 (100)	2 (100)
Anosmia	1 (14.2)	1 (50)	0 (0)	0 (0)
Cough	7 (100)	2 (100)	3 (100)	2 (100)
Shortness of breath	5 (62.5)	1 (50)	2 (66.6)	2 (100)
Nausea & vomiting	1 (14.2)	0 (0)	0 (0)	1 (50)
Diarrhea	1 (14.2)	0 (0)	0 (0)	1 (50)
Malaise	1 (14.2)	0 (0)	0 (0)	1 (50)
<b>Laboratory</b>				
HbsAg (-). n (%)	7 (100)	2 (100)	3 (100)	2 (100)
SARS-CoV-2 rapid test (+). n (%)	3 (42.8)	1 (50)	2 (66.6)	1 (50)
SGOT (u/L). mean±SD	39.8±17.4	42.5±19.9	42±26.1	34±2.8
SGPT (u/L). mean±SD	62±45.5	86±73.5	64.3±45.3	34.5±14.8
Leucocyte (μL). mean±SD	14141.4±10926.1	14675±13654.2	9573.3±3686.1	20460±18964.6
Neutrophil (%). mean±SD	76.4±16.3	79.3±21.4	69.6±20.6	83.5±5.8
Lymphocyte (%). mean±SD	15.3±13.0	11.8±13.1	21.7±17.5	9.2±2.6
NLR (%). mean±SD	9.8±13.0	20.5±24.7	5±3.6	6.5±7.7
CRP (mg/dL). mean±SD	8±6.3	5.85±6.5	10±9	7.2±4.1
RBG (mg/dL). mean±SD	213.4±109.3	157±5.6	267.3±149.5	189±103.2
Chest x-ray of pneumonia. n (%)	7 (100)	2 (100)	3 (100)	2 (100)
<b>Treatment</b>				
Antivirus. n (%)	7 (100)	2 (100)	3 (100)	2 (100)
Oseltamivir	1 (14.2)	0 (0)	1 (33.3)	0 (0)
Lopinavir-Ritonavir	6 (85.8)	2 (100)	2 (66.6)	2 (100)
Other Treatment. n (%)				
Paracetamol	7 (100)	2 (100)	3 (100)	2 (100)
<b>Outcome</b>				
Length of stay (days). mean±SD	11.4±4.9	16.5±6.3	9.3±2.0	9.5±5
Death. n (%)	4 (57.1)	0 (0)	2 (66.6)	2 (100)

BMI - body mass index, HbsAg - hepatitis B surface antigen, SARS-CoV-2 - severe acute respiratory syndrome coronavirus 2, SGOT - serum glutamic oxaloacetic transaminase, SGPT - serum glutamic pyruvic transaminase, NLR - neutrophil-to-lymphocyte ratio, PTT - partial prothrombin time, APTT - activated partial thromboplastin time, CRP - c-reactive protein, RBG - random blood glucose.

Bilirubin level was only measured in 2 patients during treatment. One patient had 0.53 mg/dL of total bilirubin and 0.34 mg/dL of direct bilirubin. Meanwhile, another patient had 0.46 mg/dL of total bilirubin and 0.22 mg/dL of direct bilirubin.

The SGOT and SGPT levels were examined every 3 days during treatment, as shown in **Figure 1**. The overall mean of SGOT<sup>34</sup> was  $39.8 \pm 17.4$  U/L and SGPT was  $62 \pm 45.5$  U/L. The mean SGOT in obesity class 1 (Group a), 2 (Group b) and 3 (Group c) groups were  $42.5 \pm 19.9$  U/L,  $42 \pm 26.1$  U/L and  $34 \pm 2.8$  U/L, respectively. The mean SGPT in obesity class 1, 2 and 3 were  $86 \pm 73.5$  U/L,  $64.3 \pm 45.3$  U/L, and  $34.5 \pm 14.8$  U/L, respectively. At admission, 3 patients (patient no. 2, 4, 7) had normal SGOT level, while the rest (patient no. 1, 3, 5, 6) had an increased SGOT level for  $<2x$  upper limit of normal (ULN). During treatment, SGOT level increased in 5 patients; 2 (patient no. 1, 3) for  $<2x$  ULN and 3 (patient no. 5, 6, 7) for  $<3x$  ULN. The rest (patient no. 2, 4) showed normal SGOT level. Meanwhile, SGPT level also varied in each patient. The normal results were shown in 3 patients (patient no. 2, 4, 7) and increased in 4 others at admission; 2 (patient no. 5, 6) for  $<2x$  ULN, 1 (patient no. 1) for  $<3x$  ULN and another 1 (patient no. 3) had  $<4x$  ULN. All of them tended to have an increased SGPT level during treatment;  $<2x$  ULN in 4 patients (patient no. 2, 4, 6, 7),  $<3x$  ULN in 1 patient (patient no. 1) and  $<5x$  ULN in 2 patients (patient no. 3, 5).




Figure 1. The SGOT and SGPT levels during hospitalization of the 7 patients in the isolation room of Dr. Soetomo Hospital, (Group a) 2 male patients obesity class 1 group, (Group b) 3 female patients of obesity class 2 group and (Group c) 2 female patients of obesity class 3 group. Three patients (patient no.2, 4, 7) had normal SGOT levels at admission, while 4 patients had SGOT levels that were 2x upper limit of normal (ULN). Five patients (patient no. 1, 3, 5, 6, 7) had SGOT levels that rose during treatment. On the other hand, 3 patients (patient no.2, 4, 7) had normal SGPT results, while 4 others had increased results at admission. During treatment, they all tended to have elevated SGPT levels. The normal reference for SGOT and SGPT levels is 0-50 U/L for men and 0-35 U/L for women. All examinations were carried out in the clinical pathology laboratory of Dr. Soetomo General Hospital Surabaya. Liver injury is defined by the level of SGPT and/or SGOT >3x ULN or ALP, GGT, and/or TBIL > 2x ULN.



## DISCUSSION

The 7 cases presented here highlight a number of noteworthy insights. First, the rise in SGOT and SGPT levels upon admission and during therapy suggests that obesity in COVID-19 patients may make liver injury more likely. Second, obesity may cause COVID-19's recovery more difficult by exhibiting severe symptoms. Third, the use of paracetamol and lopinavir/ritonavir may also contribute in the occurrence of liver injury. People who are obese have a higher risk of contracting infections like COVID-19 and experience more severe symptoms than patients who are not obese.<sup>15</sup> A study explained that overweight patients were 1.84 times more likely to have a severe COVID-19 condition than obese patients, who had a 3.4-fold increased risk.<sup>11</sup> Severe COVID-19 is defined when a patient meets one of the following criteria: (1) respiratory rate  $\geq 30$  /minute; (2) blood oxygen saturation  $\leq 93\%$  at rest; (3) the partial pressure of arterial oxygen (PaO<sub>2</sub>) to the fraction of inspired oxygen (FiO<sub>2</sub>)  $\leq 300$  mmHg (1 mmHg = 0.133kPa); (4) and pulmonary infiltration  $> 50\%$  in 48-72 hours.<sup>16</sup> These criteria indicated that there were 5 out of 7 patients were categorized as severe COVID-19 sufferers in this case series.

Obese people have continuous low-grade inflammation, which affects their innate and adaptive immune responses to be dysregulated. Moreover, angiotensin-converting enzyme 2 (ACE-2) receptor is abundantly expressed in obese patients, encouraging viral internalization into adipocytes and boost TNF- $\alpha$  and IL-6 production. Given their reduced mobility, increased insulin resistance, and gut dysbiosis, obese people may have greater inflammatory reactions to SARS-CoV-2.<sup>17-19</sup> Such aberration can lead to cytokine storm, characterized by an increase in inflammatory markers such as CRP and neutrophil lymphocyte ratio (NLR), which were revealed high in the three study groups. The cytokine storm also becomes a plausible reason for the process of liver injury.<sup>20, 21</sup> Additionally, obesity causes compensatory changes in respiratory system that include weakened diaphragm and lungs, higher airway resistance, and poorer airflow.<sup>22, 23</sup>

Most of COVID-19 patients were associated with abnormal liver function especially during treatment.<sup>24</sup> Liver injury is characterized by an increase in SGPT and/or SGOT  $>3x$  ULN (both ULN values are 0-50 U/L for men and 0-35 U/L for women), or alkaline phosphatase (ALP), gamma-glutamyl transferase (GGT), and/or total bilirubin (TBIL)  $>2x$  ULN, with each ULN is 49 U/L, 135 U/L and 17.1  $\mu\text{mol/L}$ .<sup>25</sup> There were only 2 patients examined for bilirubin, but none of them having it increasing. However, the increase for  $>3x$  ULN of SGPT and/or SGOT were found in two patients that occur since admission (patient no. 3) and during treatment (patient no. 5).

Numerous investigations found a connection between obesity in COVID-19 patients and decreased liver function.<sup>26, 27</sup> Innate immune cells such as macrophages, NK cells and T cells, are abundant in the liver. Adipose cells and Kupffer cells have been shown



to produce more pro-inflammatory cytokines in obesity. Adipose tissue and free fatty acids flow to the liver and activate hepatic macrophages which are categorized as M1 and M2 that respond in two different ways. M1 macrophages start the inflammatory process, providing the opposite effect of M2 macrophage response. M2 macrophage has anti-inflammatory properties and reparative functions with different chemokine expressions. The patient's clinical condition is determined by how well the two inflammatory reactions above balance one another. The dysregulation of the hepatic innate immune system contributes to the pathogenesis.<sup>28, 29</sup>

In this case series, all patients who had fever received 500 mg of paracetamol three times a day, and 6 of 7 patients received antiviral therapy with lopinavir-ritonavir 400 mg every 12 hours. Although<sup>32</sup> it has not been sufficiently proven for lopinavir-ritonavir, the aforementioned drugs can also cause liver damage in COVID-19 patients, especially when administered at high doses as >4000<sup>26</sup> mg per day for paracetamol.<sup>30, 31</sup> One of the mechanisms underlying the occurrence of drug-induced liver injury (DILI) is the interaction of drugs with cytochrome P-450 (CYP) in the liver. For instance, CYP3A inhibition by lopinavir-ritonavir hampers the drug metabolism and clearance while CYP2E1 and CYP2A6 induction by paracetamol produces N-acetyl-p-benzo-quinone imine (NAPQI) that later becomes hepatotoxic.<sup>32, 33</sup>

<sup>21</sup> This study has several limitations. First, this study is a case series with 7 patients. Second, coronavirus is known to enter bile duct epithelial cells via the ACE2 receptor, which in turn causes liver damage, but data regarding ALP and GGT were not available in this study. Third, there are no data related to abdominal ultrasound examination, so the influence of chronic liver disease and fatty liver on liver function abnormalities cannot be completely ruled out.

## CONCLUSION

We presented 7 obese patients with COVID-19,<sup>6</sup> of them developed severe symptoms. At admission, normal SGOT values were found in 3 patients and increased in 4 patients, while normal SGPT values were found in 3 patients and increased in 4 patients. Increased SGOT values were found during treatment in 5 patients, while SGPT values increased in all patients. All patients<sup>7</sup> had elevated markers of inflammation. Obesity may hamper the recovery process of COVID-19 and manifest in the deterioration of liver function.

## PATIENT CONSENT

All patients signed informed consent form and had agreed to be published in scientific journal.

## ACKNOWLEDGMENTS

We would like to thank to all <sup>14</sup> the patients who participated in this study.

## DISCLOSURE OF CONFLICTS OF INTEREST

There is no conflict of interest.

## FUNDING

None.

## <sup>30</sup> AUTHOR CONTRIBUTION

Authors contributed significantly to the study.

## REFERENCES

1. Fahriani M, Ilmawan M, Fajar JK, Maliga HA, Frediansyah A, Masyeni S, et al. Persistence of long COVID symptoms in COVID-19 survivors worldwide and its potential pathogenesis-a systematic review and meta-analysis. *Narra J.* 2021;1(2):e36.
2. Fajar JK, Ilmawan M, Mamada S, Mutiawati E, Husnah M, Yusuf H, et al. Global prevalence of persistent neuromuscular symptoms and the possible pathomechanisms in COVID-19 recovered individuals: A systematic review and meta-analysis. *Narra J.* 2021;1(3):e48.
3. Lusida MAP, Salamah S, Jonatan M, Wiyogo IO, Asyari CH, Ali ND, et al. Prevalence of and risk factors for depression, anxiety, and stress in non-hospitalized asymptomatic and mild COVID-19 patients in East Java province, Indonesia. *PloS one.* 2022;17(7):e0270966.
4. Fahriani M, Anwar S, Yufika A, Bakhtiar B, Wardani E, Winardi W, et al. Disruption of childhood vaccination during the COVID-19 pandemic in Indonesia. *Narra J.* 2021;1(1):e7.
5. Wagner AL, Rajamoorthy Y, Taib NM. Impact of economic disruptions and disease experiences on COVID-19 vaccination uptake in Asia: a study in Malaysia. *Narra J.* 2021;1(2).
6. Sarengat R, Islam MS, Ardhi MS. Correlation of neutrophil-to-lymphocyte ratio and clinical outcome of acute thrombotic stroke in patients with COVID-19. *Narra J.* 2021;1(3):e50.
7. Dinia F, Trisnantoro L, Helmi M. Managing Human Resources for Surge Capacity in Referral Hospitals Based on WHO Hospital Readiness Checklist for Covid-19. *Folia Medica Indonesiana.* 2022;58(2):141-9.

8. Sutadji JC, Widodo ADW, Indiasuti DN. Mortality Comparison of Using Anti Interleukin-6 Therapy and Using Standard Treatment in Severe Covid-19. *Folia Medica Indonesiana*. 2021;57(2):158-65.
9. Sharun K, Tiwari R, Iqbal Yattoo M, Patel SK, Natesan S, Dhama J, et al. Antibody-based immunotherapeutics and use of convalescent plasma to counter COVID-19: advances and prospects. *Expert opinion on biological therapy*. 2020;20(9):1033-46.
10. Triyono EA, Seipalla F, Djaja N, Akbas AMI, Ar-Rahmah KA, Budiono PS, et al. Clinical characteristics of patients with COVID-19 admitted to the COVID-19 Emergency Field Hospital of Bangkalan, Indonesia. *F1000Research*. 2022;11(414):414.
11. Cai Q, Chen, F., Wang, T., et al. Obesity and COVID-19 Severity in a Designated Hospital in Shenzhen, China. *Diabetes Care*. 2020;43(7):1392–8.
12. Harapan H, Fajar JK, Supriono S, Soegiarto G, Wulandari L, Seratin F, et al. The prevalence, predictors and outcomes of acute liver injury among patients with COVID-19: A systematic review and meta-analysis. *Rev Med Virol*. 2022 May;32(3):e2304. PubMed PMID: 34643006. Pubmed Central PMCID: PMC8646502. Epub 2021/10/14. eng.
13. Chen N, Zhou, M., Dong, X., et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet*. 2020;395(10223):507–13.
14. Hu X, Pan, X., Zhou, W., et al. Clinical epidemiological analyses of overweight/obesity and abnormal liver function contributing to prolonged hospitalization in patients infected with COVID-19. *International Journal of Obesity*. 2020;44(8):1784–9.
15. Zheng KI, Gao, F., Wang, X. B., et al. Obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism: Clinical and Experimental*. 2020;108:154244.
16. Lei Liu HQ, Jun Jian Li, et al. Effectiveness of methylprednisolone therapy in patients with a high-risk common type of COVID-19 pneumonia: a retrospective cohort study. *Clin Exp Med*. 2021;2021 Oct 22:1–11.
17. Apicella M CM, Mantuano M, et al. COVID-19 in people with diabetes: understanding the reasons for worse outcomes. *Lancet Diabetes Endocrinol*. 2020;8(9):782-92.
18. Petrakis D, Margina, D., Tsarouhas, K., et al. Obesity a risk factor for increased COVID-19 prevalence, severity and lethality (Review). *Mol Med Rep*. 2020;22(1):9-19.
19. Zahra Z, Ramadhani CT, Mamfaluti T, Pamungkas SR, Firdaus S. Association between depression and HbA1c levels in the elderly population with type 2 diabetes mellitus during COVID-19 pandemic. *Narra J*. 2022;2(1).
20. Bangash M, Patel, J., Parekh, D. COVID-19 and the liver: little cause for concern. *Lancet Gastroenterol Hepatol*. 2020;5(6):529-30.
21. Xu L LJ, Lu M, et al. Liver injury during highly pathogenic human coronavirus

- infections. *Liver Int.* 2020;40(5):998e1004.
22. Bansal R GS, Muniyappa R. Metabolic Syndrome and COVID 19: Endocrine-Immune-Vascular Interactions Shapes Clinical Course. *Endocrinology.* 2020;161(10).
  23. Ritter A, Kreis, NN., Louwen, F., et al. Obesity and COVID-19: Molecular Mechanisms Linking Both Pandemics. *Int J Mol Sci.* 2020;21(16).
  24. Li X, Zhang, Z. C., Zhang, P. L. Severe COVID-19 patients with liver injury a seven-case series. *European Review for Medical and Pharmacological Sciences.* 2020;24(14):7855–60.
  25. Cai Q, Huang, D., Yu, H., et al. COVID-19: Abnormal liver function tests. *Journal of Hepatology.* 2020;73(3):566–74.
  26. Ji D, Qin, E., Xu, J., et al Non-alcoholic fatty liver diseases in patients with COVID-19: A retrospective study. *Journal of Hepatology.* 2020;73(2):451–3.
  27. Eslam M, Newsome, P. N., Sarin, S. K., et al. A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement. *Journal of Hepatology.* 2020;73(1):202–9.
  28. Praveen Sharma AK. Metabolic dysfunction associated fatty liver disease increases risk of severe Covid-19. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews.* 2020;14(5):825-7.
  29. van der Poorten D MK, Hui J. Visceral fat: a key mediator of steatohepatitis in metabolic liver disease. *Hepatology* 2008;48:449e57.
  30. Huilin Tang LZ, Xiaotong Li, et al. Drug-induced liver injury associated with lopinavir-ritonavir in patients with COVID-19: a disproportionality analysis of U.S. food and drug administration adverse event reporting system (FAERS) data. *International Journal of Clinical Pharmacy.* 2021;43(5):1116–22.
  31. Enas M.A. Mostafa AMT, Khadiga M. Abd-Elrahman. Egyptian perspectives on potential risk of Paracetamol/acetaminophen-induced toxicities: Lessons learnt during COVID-19 pandemic. *Toxicol Rep.* 2022;9:541–8.
  32. Cao B, Wang, Y., Wen, D., et al. A trial of lopinavir-ritonavir in adults hospitalized with severe COVID-19. *New England Journal of Medicine.* 2020;382(19):1787–99.
  33. Lizi Zhao GP. Paracetamol metabolism and related genetic differences. *Drug Metabolism Reviews.* 2011;43(1):41-52.

# Liver Function Characteristics of COVID-19 Patients with Obesity at Dr. Soetomo Hospita: Case Series

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

13%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	2%
2	<a href="https://daten-quadrat.de">daten-quadrat.de</a> Internet Source	1%
3	<a href="https://medvestnik.stgmu.ru">medvestnik.stgmu.ru</a> Internet Source	1%
4	<a href="https://studyres.com">studyres.com</a> Internet Source	1%
5	Heng Ge, Mingli Zhu, Jing Du, Yong Zhou, Wei Wang, Wei Zhang, Handong Jiang, Zhiqing Qiao, Zhichun Gu, Fenghua Li, Jun Pu. "Cardiac Structural and Functional Characteristics in Patients with Coronavirus Disease 2019: A Serial Echocardiographic Study", Cold Spring Harbor Laboratory, 2020 Publication	1%
6	<a href="https://ejournal.neurona.web.id">ejournal.neurona.web.id</a> Internet Source	1%

7	Dandan Tian, Qing Ye. "Hepatic complications of COVID - 19 and its treatment", Journal of Medical Virology, 2020 Publication	1 %
8	David S. Cheng. "Idiopathic refractory sideroblastic anemiaIncidence and risk factors for leukemic transformation", Cancer, 08/1979 Publication	1 %
9	Submitted to Study Group Worldwide Student Paper	1 %
10	Yun Feng, Yaping Liu, Qian Zhao, Jiao Zhu et al. "Liver Injury in Patients with COVID-19: A Retrospective Study", International Journal of Medical Sciences, 2023 Publication	1 %
11	<a href="http://www.jogh.org">www.jogh.org</a> Internet Source	1 %
12	Olivia J. Bono, Gregory W. Poorman, Norah Foster, Cyrus M. Jalai et al. "Body mass index predicts risk of complications in lumbar spine surgery based on surgical invasiveness", The Spine Journal, 2017 Publication	1 %
13	Submitted to Syiah Kuala University Student Paper	1 %



14 Modesto M Maestre-Muñiz, Ángel Arias, Laura Arias-González, Basilio Angulo-Lara, Alfredo J Lucendo. "Prognostic Factors at Admission for In-Hospital Mortality from COVID-19 Infection in an Older Rural Population in Central Spain", *Journal of Clinical Medicine*, 2021  
Publication

---

15 [bmcgeriatr.biomedcentral.com](https://bmcgeriatr.biomedcentral.com)  
Internet Source

---

16 [covidlit.spectrumhealth.org](https://covidlit.spectrumhealth.org)  
Internet Source

---

17 [downloads.hindawi.com](https://downloads.hindawi.com)  
Internet Source

---

18 [journals.lww.com](https://journals.lww.com)  
Internet Source

---

19 [sciendo.com](https://sciendo.com)  
Internet Source

---

20 [assets.researchsquare.com](https://assets.researchsquare.com)  
Internet Source

---

21 [jamanetwork.com](https://jamanetwork.com)  
Internet Source

---

22 [preprints.jmir.org](https://preprints.jmir.org)  
Internet Source

---

23 [www.scilit.net](https://www.scilit.net)  
Internet Source

---

24	<a href="https://1library.net">1library.net</a> Internet Source	<1 %
25	<a href="https://issuu.com">issuu.com</a> Internet Source	<1 %
26	<a href="https://purehost.bath.ac.uk">purehost.bath.ac.uk</a> Internet Source	<1 %
27	<a href="http://www.qeios.com">www.qeios.com</a> Internet Source	<1 %
28	Michał Kukła, Karolina Skonieczna-Żydecka, Katarzyna Kotfis, Dominika Maciejewska et al. "COVID-19, MERS and SARS with Concomitant Liver Injury—Systematic Review of the Existing Literature", <i>Journal of Clinical Medicine</i> , 2020 Publication	<1 %
29	Taylon Felipe Silva, Fernanda Tomiotto-Pellissier, Raquel Arruda Sanfelice, Manoela Daiele Gonçalves et al. "A 21st Century Evil: Immunopathology and New Therapies of COVID-19", <i>Frontiers in Immunology</i> , 2020 Publication	<1 %
30	Submitted to Universitas Airlangga Student Paper	<1 %
31	<a href="https://bariatrictimes.com">bariatrictimes.com</a> Internet Source	<1 %

32

Internet Source

<1 %

33

[globalheartjournal.com](http://globalheartjournal.com)

Internet Source

<1 %

34

[repository-tnmgrmu.ac.in](http://repository-tnmgrmu.ac.in)

Internet Source

<1 %

35

Erwin Astha Triyono, Joni Wahyuhadi, Jongky Hendro Prajitno, Hermina Novida et al.

"Clinical characteristics and outcomes of hospitalized COVID-19 patients with diabetes mellitus in East Java, Indonesia: A cross-sectional study", F1000Research, 2022

Publication

<1 %

36

Donovan A. McGrowder, Fabian Miller, Melisa Anderson Cross, Lennox Anderson-Jackson, Sophia Bryan, Lowell Dilworth. "Abnormal Liver Biochemistry Tests and Acute Liver Injury in COVID-19 Patients: Current Evidence and Potential Pathogenesis", Diseases, 2021

Publication

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On

# Liver Function Characteristics of COVID-19 Patients with Obesity at Dr. Soetomo Hospita: Case Series

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---