

AQUAT-CEK

by Muhamad Amin

Submission date: 06-Aug-2023 04:11PM (UTC+0800)

Submission ID: 2141929476

File name: Aquat_Sci_2023-2.pdf (1.91M)

Word count: 8174

Character count: 45350

RESEARCH ARTICLE



The structure, composition, and predicted microbiome functional genes in Pacific white shrimp (*Litopenaeus vannamei*) grow-out ponds with different survival rates through high-throughput sequencing

Muhamad Amin¹ · Yoga Pramujisunu² · Ni Kadek Dita Cahyani³ · Akhmad Taufiq Mukti¹ · Mirni Lamid⁴ · Muhamad Ali⁵ · Orhan Tufan Eroldoğan^{6,7}

Received: 13 October 2022 / Accepted: 19 May 2023 / Published online: 19 June 2023
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2023

Abstract

Microbiomes inhabiting rearing ponds have been frequently associated with the survival rate of cultured animals. However, specific studies reporting on the structure, composition, and functional genes of bacteria in commercial ponds with different survival rates are still scarce. Thus, using high-throughput sequencing the present study addressed this gap by investigating microbiomes in commercial ponds with different survival rates of Pacific white shrimp, *Litopenaeus vannamei*. Water samples were collected from ponds with low survival rates (LSR) and ponds with high survival rates (HSR) and profiled by high-throughput sequencing of the 16S rRNA gene. Then, functional genes of microbiomes were predicted using PICRUSt (Phylogenetic Investigation of Communities by Reconstruction of Unobserved States) software. The results showed that ponds with different survival rates had distinct bacterial communities. At the genus level, 20 genera had different relative abundance in the two pond groups (LDA > 3), of which 11 genera were in HSR ponds and nine genera were in LSR ponds. The top five most abundant bacterial genera found in LSR ponds were *Vibrio*, *Kocuria*, *Tepidiphilus*, *unidentified_Alphaproteobacteria*, and *Pseudoalteromonas*, while the five most abundant found in the HSR ponds were *Candidatus_Aquiluna*, *unidentified_Acinobacteria*, *Ilumatobacter*, *unidentified_Deltaproteobacteria*, and *Marivita*. PICRUSt analysis revealed that nine metabolic pathways were higher in HSR grow-out ponds and associated with carbohydrate metabolism, chlorophyll biosynthesis, lignin degradation, arginine and glucosamine biosynthesis, and vitamin K2 synthesis, while nine metabolic pathways were found to be significantly higher in LSR grow-out ponds including lipid IVA biosynthesis (conserved structure in diverse Gram-negative pathogens), and L-tryptophan degradation activity (proteolytic agents). To the authors' knowledge, this is the first study to report the diversity, composition, and metabolic activity of microbiomes in commercial shrimp grow-out ponds and their effect on the survival rate of Pacific white shrimp. Overall, these findings suggest that the specific microbiomes in rearing water may be linked to the impact on improved survival of Pacific white shrimp.

Keywords Functional prediction · Microbiota · Grow-out pond · Survival rate

✉ 94 hamad Amin
muhamad.amin@fpk.unair.ac.id

¹ Department of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, East Java 60115, Indonesia

² Aquaculture Study Program, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya, East Java 60115, Indonesia

³ Department of Biology, Faculty of Science and Mathematics, Universitas Diponegoro, Semarang, Central Java, Indonesia

⁴ Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, East Java 60115, Indonesia

⁵ Laboratory of Microbiology and Biotechnology, Faculty of Animal Science, University of Mataram, Mataram, Indonesia

⁶ Faculty of Fisheries, Department of Aquaculture, Çukurova University, 01330 Adana, Turkey

⁷ Biotechnology Research and Application Center, Çukurova University, 01130 Adana, Turkey

Introduction

Shrimp cultures in Asia have recently been found to suffer from diseases caused by viruses, bacteria, fungi, and protozoa (Kurniawinata et al. 2022; Shin et al. 2023). In particular, viral and bacterial diseases typically lead to mass mortality in the grow-out phase and cause great economic losses. Thus, assessment of microbiota and nutrient dynamics in the pond is imperative to avoiding rearing failure. In this sense, the composition and diversity of microbiomes, particularly pathogenic bacteria, in rearing water have been frequently reported to contribute to the water quality and the survival rate of cultured animals (Zheng et al. 2017). This has been correlated with the predominance of certain bacterial communities in the pond ecosystem such as in water and sediment. Several bacteria may have the capacity to decompose organic matter and inorganic nutrients such as ammonia (NH₃) or nitrite (NO₂) into less or non-toxic compounds such as nitrate (NO₃) (Hlordzi et al. 2020; Amin et al. 2021) or to produce antimicrobial compounds antagonistic to bacterial pathogens (Amin et al. 2020). A study by Kewcharoen and Srisapoom (2019) reported that *Bacillus* spp. can suppress the growth of pathogenic microbiota in shrimp pond ecosystems where planktonic bacteria proliferate with shrimp and thereby increase the survival rate of Pacific white shrimp (*Litopenaeus vannamei*). On the contrary, some bacteria living in the water column can be pathogens and may also infect cultured animals (Zhang et al. 2021). *Vibrio harveyi*, for instance, has been associated with luminous vibriosis diseases in Pacific white shrimp (Supono et al. 2019). Thus, a complex interaction between the host, the aquatic ecosystem, and the activity of the corresponding microbial communities affects the survival rate of cultured organisms (Chen et al. 2017).

Studies reporting how certain bacterial species inhabiting rearing ponds affect the survival of Pacific white shrimps have been performed by several authors, as the characteristics and dynamics of microorganisms surrounding shrimp are important factors (Imaizumi et al. 2022). A study by Nguyen Thi Truc et al. (2021), for instance, described a strong correlation between the presence of *Lactobacillus* spp. in rearing ponds and the survival rate of Pacific white shrimp. Additionally, a study by Chen et al. (2019) documented how bacterial communities in rearing water of a recirculating aquaculture system (RAS) affected water quality parameters and the survival of Pacific white shrimp. Similarly, Khademzade et al. (2020) reported that the presence of *Bacillus cereus* and *Proteus acidilactici* in rearing water improved the water quality and survival of Pacific white shrimp in earthen ponds. However, these studies were mostly performed in small rearing

systems or laboratory-scale experiments and focused only on certain species such as *Bacillus* spp., *Lactobacillus* spp., and *P. acidilactici*. Meanwhile, broader studies which investigate a correlation between the structure and composition of all microbiomes in large-scale grow-out shrimp ponds with different survival rates are still very limited. Thus, a better understanding of microbial communities as an initial step in screening probiotic candidates for improving water quality or suppressing bacterial pathogen presence in rearing ponds is needed to improve production in the shrimp industry.

Current technological developments such as high-throughput sequencing of the V3–V4 region of the 16S rRNA gene have been a great help for researchers to monitor and characterize potential pathogenic or beneficial bacteria (probiotics) that live and thrive in shrimp ponds (Gu et al. 2021; Amin et al. 2022). The next-generation sequencing (NGS) technique is an effective tool used to ascertain the complexity and interaction of microbes from environments with high microbial diversity such as culture ponds (Infante-Villamil et al. 2021). In addition, a metagenomics analysis can explain the relationship between microorganisms by associating genetic information extracted from environmental samples or certain environments (Yu et al. 2020). Characterization of the bacterial microbiome not only aids in monitoring potential pathogens to control disease outbreaks and increase pond productivity, but also may be used as an initial step to screen potential indigenous probiotic candidates.

Therefore, this study aimed to assess the structure and composition and to predict the metabolic activity of microbiomes that thrive in shrimp pond water with a high and low survival rate. In addition, this research is intended to determine the correlation of the types of bacteria that live in pond water considering the survival rate of shrimp reared. To the authors' knowledge, this report is the first study to report the diversity, composition, and metabolic activity of microbiomes in commercial shrimp grow-out ponds and their relation with the survival rate of Pacific white shrimp. The results of this study, therefore, are expected to provide initial information about the correlation of the bacteria community in pond water with the survival rate of cultured shrimp in intensive pond systems.

Material and methods

Sampling sites and culture condition

Water samples were taken on the same day (day 47 of culture) from four intensive shrimp ponds at Madura, East Java, Indonesia (113°01'14.7" E and 6°52'59.3" LS). The shrimp ponds were constructed with 800 m² high-density

polyethylene (HDPE). The shrimp seeds were F1 fry obtained from a certified commercial hatchery and cultured in a stocked density of 275 fish/m². Solid waste was taken out by siphoning at the beginning (after water sterilization) and once a week during the rearing period to reduce the accumulation of solid waste at the bottom of ponds. Shrimps were cultured in a semi-flow-through system and fed a commercial feed (IRAWAN, CP Proteina Tbk) at 5% body weight (bw) day⁻¹ and five times a day. Water quality parameters were measured 1 week before sampling, during sampling, and 1 week after sampling. Water quality parameters were taken in the form of acidity (pH), salinity, temperature (°C), phosphate (PO₄³⁻), ammonia (NH₃), nitrite (NO₂), dissolved oxygen (DO, mg/l), and alkalinity (Table 1).

Water sample collection

Water sampling was performed by collecting rearing water from six shrimp ponds (three ponds with high survival rates [HSR, 94.68%] and three ponds with low survival rates [LSR, 66.7%]). The results of the *t*-test showed that the two groups of ponds had significantly different survival rates ($p < 0.05$). Then, 200 ml of pond water was collected from four distant locations at 15–30 cm depth (from the left corner to the right corner) of each pond and mixed in a sterile bottle, 20 ml of which was collected and mixed with 30 ml 95% ethanol and stored in a 50 ml falcon tube. The water samples were then filtered through 0.45 and 0.22 µm pore sizes of MF-Millipore membrane Filter (Millipore), and both filter papers were stored at -20 °C until DNA extraction.

DNA extraction, amplification, purification, and sequencing

Bacterial DNA was extracted from the samples as described previously by Amin (2010), with slight modification. In brief, half of each filter paper (0.45 and 0.22 µl) was cut into

Table 1 Physicochemical water quality parameters of rearing ponds

Water quality parameters	LSR	HSR
	Average ± Std	Average ± Std
pH	7.88 ± 0.06	7.84 ± 0.16
Salinity (psu)	31.17 ± 0.87	32.00 ± 1.55
T (°C)	28.50 ± 0.58	28.50 ± 0.58
PO ₄ (mg/l)	0.20 ± 0.12	0.44 ± 0.24
NH ₃	0.86 ± 0.73	1.17 ± 0.74
NO ₂	0.38 ± 0.13	0.29 ± 0.07
DO (mg/l)	4.65 ± 0.20	4.27 ± 0.31
Alkalinity	110.32 ± 0.00	109.61 ± 1.04

LSRn is pond n with the low survival rate of Pacific white shrimp, HSR is pond n with the high survival rate of white shrimps

pieces and extracted using a commercial DNA extraction kit (ZymoBIOMICS™ DNA Miniprep Kit). The V3–V4 region of the 16S rRNA gene was amplified using a pair of specific primers (16S V4: 515F-806R, 18S V4: 528F-706R) with the barcode. All polymerase chain reactions (PCR) were carried out with Phusion® High-Fidelity PCR Master Mix (New England Biolabs). The quantity and quality of PCR products were measured by mixing the same volume of 1X loading buffer (contained SYBR Green) with PCR products and conducting electrophoresis on 2% agarose gel for detection. Samples with a bright main strip between 400 and 450 base pairs (bp) were chosen for further analysis. PCR products were mixed at equal density ratios and were purified with QIAquick Gel Extraction Kit (Qiagen, Germany). The libraries were generated with NEBNext® Ultra™ DNA Library Prep Kit for Illumina and quantified via Qubit and qPCR with the Illumina platform.

Sequence data analysis, OTU cluster, and taxonomic annotation

Paired-end reads were assigned to the samples based on their unique barcodes and identified by cutting off the barcode and primer sequences. Paired-end reads were merged using FLASH (V1.2.7) (Magoč and Salzberg 2011). Quality filtering on the raw tags was performed under specific filtering conditions to obtain high-quality clean tags according to Qiime (V1.7.0) (Caporaso et al. 2010). The tags were compared with the reference database (SILVA database) using the UCHIME algorithm to detect and remove chimeric sequences (Edgar et al. 2011). Then the effective tags were finally obtained. Sequences were analyzed with UPARSE software (UPARSE v7.0.1090) (Edgar 2013) using all the effective tags. Sequences with ≥ 97% similarity were assigned to the same operational taxonomic unit (OTUs). A representative sequence for each OTU was screened for further annotation. For each representative sequence, Qiime (Version 1.7.0) in the Mothur method was performed against the small subunit ribosomal RNA (SSU rRNA) database of the SILVA Database for species annotation at each taxonomic rank (threshold: 0.8–1) (Quast et al. 2012). To obtain the phylogenetic relationship of all OTU representative sequences, MUSCLE (version 3.8) was used to compare multiple sequences rapidly. OTU abundance information was normalized using a standard sequence number corresponding to the sample with the least sequences. Subsequent analyses of alpha diversity (Shannon index and OTU number) and beta diversity were all performed based on this output normalized data.

Profiling of predictive metagenome functions from bacterial communities

PICRUSt (Phylogenetic Investigation of Communities by Reconstruction of Unobserved States) software provided predictive metagenome functions from bacterial 16S rRNA gene data (Langille et al. 2013). PICRUSt wraps a number of tools to generate functional predictions from amplicon sequences (Mirarab et al. 2012; Barbera et al. 2019; Czech et al. 2020). These tools were used for the placement of reads (citations), predicting hidden states (Louca and Doebeli 2018), and inferring the pathway (Ye and Doak 2009). Then, metagenome functions of bacterial communities were predicted and categorized with Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. The functional diversity of microbiome data was visualized with STAMP (Parks et al. 2014).

Identification of bacterial features that discriminate groups

LEfSe (Linear discriminant analysis Effect Size) was conducted using MicrobiomeAnalyst software as described previously by Amin et al. (2022). All bacterial quantitative taxa and functional prediction estimations were applied to calculation of effect size on the survival rate of shrimps. The threshold for the logarithmic linear discriminant analysis (LDA) score was set as 3.0. The Kruskal–Wallis rank-sum statistical test was conducted during the LDA score calculation to assess whether the measurements discriminating groups were significant. Statistically different bacterial taxa and metabolic pathways were used for the analyses of the relationships between bacterial communities and ponds with different survival rates.

Results

Overall bacterial community composition

A total of 559,152 high-quality sequences (average 415 bp in sequence length) of the V3–V4 region of the 16S rRNA gene were obtained from four rearing water shrimp ponds.

The sequence number for an individual sample varied from 133,656 to 147,261. The number of OTUs from each sample ranged from 272 to 753 (Table 2).

Completeness of the sequences was estimated with Good's coverage, which ranged from 0.99 to 1.00 in all samples (Table 2). No significant difference in diversity indexes was found between the two sample groups (> 0.05). The Shannon index was 4.75 ± 0.94 and 5.74 ± 0.16 for the HSR and LSR, respectively. The Simpson index values were 0.85 ± 0.10 and 0.94 ± 0.04 for the HSR and LSR, respectively. The two indices varied greatly in ponds with both high and low survival rates. The Bray–Curtis index distance-cluster tree showed that bacterial communities in the LSR clustered together, while bacterial communities in the HSR shrimp formed a branch but separated from pond water with a low survival rate (Fig. 1A). Furthermore, the principal component analysis (PCA) plot indicated distinct bacterial communities between the two pond water groups (Fig. 1B).

The dominant population of bacterial communities

At the phylum level, Proteobacteria (48.1%) was the most dominant phylum in the HSR, followed by Cyanobacteria (19.3%), Bacteroidetes (11.4%), Firmicutes (9.46%), Actinobacteria (6.41%), and Verrucomicrobia (1.81%), while the remaining phyla including Tenericutes, Chloroflexi, and Planctomycetes were less than 1%, as shown in Fig. 2. Deeper taxonomic annotation indicated that more Firmicutes members including *Lactobacillus* and *Bacillus* were observed in the HSR (Fig. 2). On the other hand, the most abundant phylum in the LSR was Cyanobacteria (37.42%), followed by Actinobacteria (22.73%), Proteobacteria (21.55%), Bacteroidetes (9.12%) and Firmicutes (5.74%), while other phyla including Verrucomicrobia, Tenericutes, Chloroflexi, and Planctomycetes accounted for less than 1% each (Fig. 3). In general, Proteobacteria was the most abundant phylum in the HSR, but Cyanobacteria was the most abundant in LSR.

At the genus level, the plots with low survival rates were the unidentified Cyanobacteria genus (unidentified Cyanobacteria), with a value of 37.3% and 20.0%, respectively. In addition, a relative abundance of some dominant bacterial genera (e.g., *Massilia*, *Acinetobacter*, *Sphingomonas*,

Table 2 16S rRNA sequencing information and alpha diversity indexes of bacterial communities from rearing water of white shrimp pond

SR status	Effective tags	No. of OTUs	Shannon index	Simpson index	Chao_1	ACE	Good's coverage
HSR-1	115,896	464	4.086	0.774	458.118	458.550	1.00
HSR-2	103,527	753	5.413	0.918	730.128	737.334	1.00
LSR-1	111,670	618	5.620	0.908	619.550	623.801	0.99
LSR-2	120,773	272	5.850	0.963	281.474	279.713	1.00

SR survival rate; HSR pond water with high survival rate; LSR: pond water with low survival rate




Fig. 1 **A** Bray-Curtis Index distance-cluster tree showed that bacterial communities in the pond water of low survival rate (LSR) and high survival rate (HSR). **B** Principal component analysis (PCA) constructed based on OTUs indicates the distinctions of bacterial com-

munities in the pond waters with different survival rates. HSR: pond water with a high survival rate; LSR: pond water with a low survival rate




Fig. 2 Comparison of relative abundance at the phylum level. Composition of microbial communities (relative to OTU composition) in phyla stages inhabiting pond waters with different survival rates of white shrimps. LSR is a pond with a low survival rate and HSR is a high survival rate for Pacific white shrimp ponds




Fig. 3 Relative abundance of microbial communities (relative to OTU composition) in top 10 genera inhabiting pond waters with different growth rates of Pacific white shrimps. LSR is a pond with a low survival rate and HSR is a high survival rate for shrimp ponds

and *Ruegeria*) corresponded to pond water with a high survival rate of white shrimps (Fig. 3). *Massilia* accounted for 7.6% in the HSR and only 0.01% in the LSR. *Acinetobacter* accounted for 2.3% in the HSR and only 0.06% in the LSR. *Sphingomonas* accounted for 1.7% of the HSR and <0.01% in the LSR. On the contrary, the five most abundant bacterial genera identified from the LSR were *Candidatus_Aquilana*, *unidentified_Actinobacteria*, and *Ilumatobacter*, which accounted for 12.8%, 3.9%, and 2.3%, respectively. However, the present study results also showed that the majority of OTUs were categorized into the “Others” group, which was 62% and 41% in HSR and LSR, respectively.

Bacterial signatures

The most prominent differences in the pond-rearing water between the two groups led us to identify a key variable that significantly contributes to the segregation of water microbiota. Based on the LEfSe analysis tool with LDA score > 3, 45 bacterial genera had different relative abundance in both HSR and LSR which consistently explained their differences based on the effect size. The results of this analysis are shown in Fig. 4. Eleven genera were found in relative abundance in HSR. Among them, the top five dominant genera were *Candidatus_Aquilana*, *unidentified_Actinobacteria*, *Ilumatobacter*, *unidentified*




Fig. 4 Factors associated with survival variations in the pond-water microbiome. A total of two factors ($LDA > 3$) were associated with the variation in the pond-water microbiome. The bar plot indicates that HSR is a pond-water microbiome with a high survival rate, and LSR is a pond-water microbiome with a low survival rate




Fig. 5 Venn diagrams showing the OTU numbers that were shared or unique in pond water with different survival rates of white shrimps. The bubble plot represents the relative abundance of unique or shared bacterial taxa at the OTU level. HSRn is OTU microbiomes of pond water with a high survival rate at pond n. LSRn is bacterial OTUs of pond water with a low survival rate at pond n

Deltaproteobacteria, and *Marivita*. Meanwhile, microbiota in the LSR was dominated by nine genera, and the

top five dominant genera were *Vibrio*, *Kocuria*, *Tepidiphilus*, *unidentified_Alphaproteobacteria*, and *Pseudoalteromonas* (Fig. 4). Notably, *Vibrio* appeared to be the most dominant taxa, with high relative abundance in the LSR (Fig. 6).

In the lower taxa level, there were 81 OTUs with different relative abundance in the two pond water groups: 20 OTUs in LSR, and 61 OTUs in the HSR. These OTUs belonged to 20 bacterial species, including 11 species of HSR and nine species with LSR (Fig. 4). The top five abundant bacterial species found in LSR were *Vibrio*, *Kocuria*, *Tepidiphilus*, *unidentified_Alphaproteobacteria*, and *Pseudoalteromonas*. On the genus level, the top five abundant bacterial species found in the HSR were *Candidatus_Aquiluna*, *unidentified_Acinobacteria*, *Ilumatobacter*, *unidentified_Deltaproteobacteria*, and *Marivita*.


To determine the common bacteria between the examined rearing water, a Venn diagram showed 88 common OTUs that were associated with HSR and LSR pond water (Fig. 5). Among these were 30 OTUs shared between pond water that were associated mostly with *Proteobacteria* such as *Ruegeria atlantica*, *Enterobacter kobei*, *Acinetobacter junii*, and *Moraxella osloensis*. Specific species showing 10 genera in high relative abundance were selected to construct the taxonomy tree (Fig. 4).

The present study also showed that ponds with a high survival rate of white shrimp might have dominance of Firmicutes. Two genera belonging to Firmicutes (*Lactobacillus* and *Bacillus*) were more abundant in ponds with a high survival rate (Fig. 6a and b), while the number of *Vibrio* spp. was also higher in ponds with lower survival rates, 0.7% and 5% (Fig. 7a and b).

Predictive metagenome functions in grow-out ponds with different shrimp growth

Predicted KEGG pathways from 16S rRNA gene sequences were profiled via PICRUSt2 to compare between pond groups. Nine survival-rate-related metabolic pathways were identified as significantly different between the two pond groups ($p < 0.05$). Five of these pathways—PWY-7198, PWY-7210, PWY-5531, PWY-7159, and PWY-7046—were significantly higher in HSR ponds and were classified into 15 metabolic processes (Fig. 8b.) The 15 metabolic processes were associated with pyrimidine deoxyribonucleotides de novo biosynthesis, pyrimidine deoxyribonucleotides de novo biosynthesis from CTP, chlorophyllide a biosynthesis I, chlorophyllide a biosynthesis II, chlorophyllide a biosynthesis III, 6-hydroxymethyl-dihydroprotein diphenylphosphate biosynthesis, 4-coumarate degradation, L-arginine biosynthesis II, UDP-N-acetyl-D-glucosamine biosynthesis I, superpathway of menaquinol-11 biosynthesis, superpathway of menaquinol-12 biosynthesis, superpathway of menaquinol-13 (K2 vitamin) biosynthesis,

Fig. 7 The proportion of Firmicutes genera in the shrimp's ponds of HSR



transfer to lipid IVA III (chlamydia), aromatic biogenic amine degradation, and norspermidine biosynthesis (Fig. 8). These metabolic processes were associated with lipid biosynthesis (NAGLIPASYN-PWY), NAD biosynthesis, and protein degradation.

Discussion

The structure and composition of bacteria in rearing water determined the survival rate of cultured organisms (Deng et al. 2019). Several studies have reported that *Nitrosomonas* sp. and *Nitrobacter* sp. maintained good water quality by degrading organic and inorganic aquaculture waste (Nair et al. 2019; Wei et al. 2021). Other bacteria such as




Fig. 8 Predictive metagenome functions that discriminate between pond microbiomes with different survival rates. HSR is grow-out ponds with a high survival rate and LSR is grow-out ponds with a low survival rate. Extended error bar plot for two-group analysis module comparison of PICRUSt predicted KEGG function data based on the HSR and LSR using Welch's *t*-test for the two groups.

Lactobacillus sakei, *Bacillus natto*, and *Bacillus amyloliquefaciens* have been reported to suppress dominant pathogenic bacteria such as *Vibrio* spp. (Sahandi et al. 2019). However, the abovementioned findings were mostly obtained from in vitro studies. To the best of our knowledge, this article presents the first descriptions of the rearing water microbiome diversity and taxonomic composition in commercial shrimp grow-out ponds. There are no previous NGS-based studies assessing the effects of the microbiome on the survival rate of Pacific white shrimp in pond water. Overall, the findings in the present study showed that the types and composition of water microbiomes obtained from the HSR and LSR were significantly different. At the phylum level, bacteria dominating in HSR were Proteobacteria (48.1%), followed by Cyanobacteria (19.3%), Bacteroidetes (11.38%), and Firmicutes (9.5%), while LSR were dominated by Cyanobacteria (37.4%), followed by Actinobacteria (22.7%), Proteobacteria (21.6%), and Bacteroidetes (9.1%). Proteobacteria and Firmicutes appeared to be more dominant in HSR. This result might be related to the large number of probiotic species belonging to Proteobacteria and Firmicutes. A similar result was reported by Huerta-Rábago et al. (2019) where proteobacteria dominated the pond after probiotic consortia were added. Further analysis using LefSe indicated that two members of phylum Proteobacteria (*unidentified*

3 An extended error bar plot was used for the comparison between two water samples, and only predicted functions with $p < 0.05$ are shown. Bar plots on the left side display the mean proportion of each KEGG pathway while the dot plots on the right show the differences in mean proportions between two water samples using p -values

Deltaproteobacteria and *Litoricola* sp.) were found to be very dominant in HSR. However, studies which reported metabolic activity of these two genera are still scarce today. Only a few studies have reported that *unidentified Deltaproteobacteria* and *Litoricola* sp. are beneficial. A study by Langwig et al. (2022) reported that *Deltaproteobacteria* was uncultured marine bacteria which may play essential roles in global carbon, sulfur, and nutrient cycling. The other bacterial species, *Litoricola* sp., is also a relatively new species and less studied in the shrimp ponds. However, a study by Huang et al. (2018) suggested that *Litoricola* sp. might be a potential probiotic for shrimps. Other possible mechanisms could be that these bacteria have *N*-acyl homoserine lactone (AHL)-degrading activity which ultimately suppresses the presence of pathogenic bacteria such as *Vibrio* spp. as reported by Zheng et al. (2016). The results further suggested that for these species, research to investigate metabolic function is highly recommended for further studies.

The second possibility is that the high survival rate of white shrimp in HSR might be attributable to the dominance of Firmicutes. Two genera belonging to Firmicutes (*Lactobacillus* and *Bacillus*) were more abundant in the HSR pond (Fig. 8c and d). Lin et al. (2019) reported that Firmicutes, especially from the genera *Lactobacillus* and *Bacillus*, are among the most common probiotics and have been applied

not only in aquaculture species but also in terrestrial cultured animals. Among the possible mechanisms is the production of antimicrobial compounds against bacterial pathogens or stimulation of the immune function of cultured organisms. For instance, *Lactobacillus farraginis*, *Pediococcus acidilactici*, and *Pediococcus pentosaceus* have been reported to produce anti-vibrios against *Vibrio anguillarum*, *Vibrio alginolyticus*, and *Vibrio harveyi* (Amin et al. 2016). In addition, Ahmmed et al. (2020) reported that supplementation of *Lactobacillus* spp. reduced the number of *Vibrio* spp., the most common pathogen in marine species, and improved the survival rate of giant tiger shrimp, *P. monodon*. Consistent with previous studies, in the present study, Firmicutes—especially the genera *Lactobacillus* and *Bacillus*—were found to be more abundant in HSR relative to a lower number of *Vibrio* spp. (3% in LSR and only 0.08% in HSR), as shown in Fig. 6. However, the specific mechanisms by which these genera suppress *Vibrio* spp. should be further investigated to gain more comprehensive information.

Besides the two genera, LEfSe analysis showed that nine other genera—*Candidatus_Aquiluna*, *unidentified Actinobacteria*, *Ilumatobacter*, *Marivita*, *Balneola*, *Acholeplasma*, *unidentified Methylophilaceae*, *Bacteroides*, and *Faecalibacterium*—were prevalent in the water samples. Of these bacteria, only a few have been reported as probiotics. *Methylophilaceae*, for instance, has been reported as denitrification bacteria in wastewater treatment (Baytshtok et al. 2009). It was further documented that this bacterial genus can remove nitrate (NO₃), which is the end product of ammonia degradation (Srinandan et al. 2012) and is one of the most toxic compounds in shrimp farms. In addition, *Bacteroides* have been reported as probiotics in several studies. Zafar and Saier (2018) reported four species (*Bacteroides thetaiotaomicron*, *Bacteroides vulgatus*, *Bacteroides ovatus*, and *Bacteroides fragilis*) to be good probiotic candidates for humans. In addition, *Bacteroides* were revealed to contribute to the host's nutrition, especially by producing fatty acids and vitamins (Lara-Flores 2011). However, since the present study identified at the genus level, it is too preliminary to conclude that these signature bacteria can be probiotic candidates. Therefore, studies on species level and their metabolic activity should be performed to gain more comprehensive conclusion.

The remaining signature bacteria appeared to be less commonly found in either terrestrial or aquatic environments. To the author's knowledge, *Candidatus_Aquiluna*, for instance, has been reported by only three studies. The first was by Alfiansah (2019), who found the bacterium from the intestine of white shrimp cultured in Rembang Regency, Central Java, Indonesia. Other studies reported *Candidatus_Aquiluna* sp. from marine water in Toulon Bay, France (Coclet et al. 2019), and marine sediment in the Adriatic Sea, Croatia (Kajan et al. 2022). The other study

was reported by Gutiérrez et al. (2018), who found *Candidatus_Aquiluna* sp. from seawater in Puyuhuapi Fjord, Chile. However, none of the studies reported the metabolic activity of the bacterium; therefore, further studies which isolate and characterize the metabolic activity of the bacterium are highly recommended. Similarly, although *Ilumatobacter* has also been reported in several studies including shrimp guts (Amin et al. 2022) and marine sediments (Conte et al. 2018), none of these studies also reported the metabolic activity of this bacterium. These signify that bacteria identified from the shrimp ponds may be good candidates for indigenous probiotics, especially to improve the survival rate of Pacific white shrimp. Further studies to investigate the metabolic activity of these bacteria in their capacity to degrade aquaculture wastes or produce antimicrobial compounds against opportunistic or obligate pathogens are still required.

Further analyses by predicting metabolic function using PICRUSt showed that 24 metabolic processes were significantly higher in the HSR grow-out ponds ($p < 0.05$), which nine metabolic processes were associated with pyrimidine deoxyribonucleotide de novo biosynthesis, pyrimidine deoxyribonucleotide de novo biosynthesis from CTP, chlorophyllide a biosynthesis I, chlorophyllide a biosynthesis II, chlorophyllide a biosynthesis III, 6-hydroxymethyl-dihydroprotein diphosphate biosynthesis, 4-coumarate degradation, L-arginine biosynthesis II, UDP-N-acetyl-D-glucosamine biosynthesis I, superpathway of menaquinol-11 biosynthesis, superpathway of menaquinol-12 biosynthesis, superpathway of menaquinol-13 (K2 vitamin) biosynthesis, superpathway of menaquinol-6 biosynthesis I, superpathway of menaquinol-10 biosynthesis, and chondroitin sulfate degradation I. These nine metabolic processes were associated with carbohydrate metabolism, chlorophyll biosynthesis, lignin degradation, arginine and glucosamine biosynthesis, and vitamin K2 synthesis (Schröder et al. 2005; Yang et al. 2011). Likewise, nine metabolic processes were found to be significantly higher in LSR grow-out ponds including lipid IVA biosynthesis, NAD biosynthesis II (from tryptophan), L-tryptophan degradation to 2-amino-3-carboxymucon, superpathway of pyrimidine deoxyribonuclease degradation, superpathway of thiamin diphosphate biosynthesis I, TCA cycle VIII, Kdo transfer to lipid IVA III (chlamydia), aromatic biogenic amine degradation, and norspermidine biosynthesis. Some of these metabolic pathways have been described as signals for pathogenic environments. For instance, the lipid IVA biosynthesis is one of the main conserved structures in diverse Gram-negative pathogens (Raetz et al. 2009), and the chondroitin sulfate degradation was a cell wall degrading enzyme (Wang et al. 2020). In addition, 2-amino-3-carboxymucon has been described as a marker for constant exposure to a wide range of endogenous caustic or proteolytic agents (Toribara et al. 1997), and Kdo transfer to lipid IVA III (chlamydia), which is common to most

Gram-negative bacteria, is the most toxic part (Schomburg and Schomburg 2013). These data might suggest that most bacterial species dominating the LSR ponds are pathogens.

Overall, these findings point to the fact that the specific microbiomes in rearing water may be linked to the impact on improved survival of Pacific white shrimp. Some bacteria were identified to be signature in the pond with high survival rate including *Candidatus_Aquiluna*, unidentified Actinobacteria, *Ilumatobacter*, unidentified Deltaproteobacteria, *Marivita*, *Lactobacillus* spp., and *Bacillus* spp. These bacterial genera can be indigenous potential probiotics, and therefore should be isolated for further study. As previously reported by Yang et al. (2018), *Candidatus_Aquiluna* has been identified as keystone, suggestive of a vital role in driving bacterioplankton succession in white shrimp aquaculture ponds and maintaining healthy aquaculture environments. Meanwhile, the number of *Vibrio* spp. was also lower in HSR ponds, which may indicate that the two bacterial genera can produce antimicrobial compounds which are antagonistic to the *Vibrio* spp. However this speculative hypothesis should be further studied in order to obtain a more comprehensive conclusion. On the other hand, the five most abundance microbiota in the LSR were *Vibrio*, *Kocuria*, *Tepidiphilus*, unidentified Alphaproteobacteria, and *Pseudoalteromonas*. Among these genera, *Vibrio* has been regarded as an opportunistic pathogen in shrimp aquaculture (Zhang et al. 2020). Meanwhile, the other three genera might be indigenous pathogens, which should be further studied. Thus, suppressing the growth of these bacteria in aquaculture ponds might improve the survival of white shrimp.

In conclusion, the study results revealed that the structure, composition, and predicted metabolic functions of the microbiome in rearing ponds appeared to correlate strongly with the survival rate of Pacific white shrimps. LSR grow-out ponds were dominated by *Vibrio*, *Kocuria*, *Tepidiphilus*, unidentified Alphaproteobacteria, and *Pseudoalteromonas*, while the HSR ponds were dominated by *Candidatus_Aquiluna*, unidentified Acinobacteria, *Ilumatobacter*, unidentified Deltaproteobacteria, and *Marivita*. In addition, PICRUSt analysis revealed that nine metabolic pathways were higher in HSR grow-out ponds and associated with carbohydrate metabolism, chlorophyll biosynthesis, lignin degradation, arginine and glucosamine biosynthesis, and vitamin K2 synthesis. Likewise, nine metabolic pathways were significantly higher in LSR grow-out ponds including lipid IVA biosynthesis (conserved structure in diverse Gram-negative pathogens) and L-tryptophan degradation activity (proteolytic agents). Overall, these findings suggest that the specific microbiomes in rearing water may be linked to the impact on survival improvements of Pacific white shrimp. Higher OTUs belonging to Firmicutes in HSR, especially from the genera *Bacillus* and

Lactobacillus, might play an important role in suppressing the vibrios in ponds. Nevertheless, further research is still required to confirm this preliminary conclusion.

Acknowledgements This study was funded by Universitas Airlangga, the Ministry of Education, Culture, Science, and Technology, the Republic of Indonesia, with grant number 309/UN3.15/PT/2023.

Author contributions MA: Conceptualization, Funding acquisition, Data curation, Formal analysis, Methodology, Software, Writing—original draft, Validation, Submission. YP: Data curation, Formal analysis, Methodology, Software, Writing—original draft, Validation. NKDC: Conceptualization, Investigation, Data analysis. ATM: Conceptualization, Investigation, Writing & editing. ML: Conceptualization, Investigation, Writing & editing, Supervision. MA: Project administration, Methodology, Formal analysis, Validation. OTE: Formal analysis, Methodology, Validation, Supervision, Writing.

Data availability Data will be made available by request to the authors.

Declarations

Conflict of interest The authors declare no conflict of interest.

References

- Ahmed F, Ahmed MK, Khushi SS, Sumon MS, Karamcheti SS, Sarower M (2020) Host gut-derived probiotic *Lactobacillus* sp. improves resistance of giant freshwater prawn *Macrobrachium rosenbergii* against *Vibrio harveyi*. *Aquac Int* 28:1709–1724
- Alfiansah YR 2019. Microbiome in shrimp *Litopenaeus vannamei* aquaculture: dynamic changes and bacterial lifestyles. Universitas Bremen.
- Amin M 2010. Bacterial variation in gastrointestinal tract of Nile tilapia, *Oreochromis niloticus*, reared in recirculating aquaculture system and active suspension tank. M. Sc.
- Amin M, Adams M, Bolch CJS, Burke CM (2016) *In vitro* screening of lactic acid bacteria isolated from gastrointestinal tract of Atlantic Salmon (*Salmo salar*) as probiotic candidates. *Aquac Int* 23:1–14
- Amin M, Liliyanti M, Nufus N, Ali M (2020) Screening of antivibrio-producing lactic acid bacteria originated from aquatic animals as probiotic candidates. IOP conference series: earth and environmental science. IOP Publishing, p 012092
- Amin M, Ago A, Prayugo P, Ali M, Hum NNMF (2021) Comparison of total nutrient recovery in aquaponics and conventional aquaculture systems. *Open Agriculture* 6:682–688
- Amin M, Kumala RRC, Mukti AT, Lamid M, Nindarwi DD (2022) Metagenomic profiles of core and signature bacteria in the guts of white shrimp, *Litopenaeus vannamei*, with different growth rates. *Aquaculture* 550:737849
- Barbera P, Kozlov AM, Czech L, Morel B, Darriba D, Flouri T, Stamatakis A (2019) EPA-ng: massively parallel evolutionary placement of genetic sequences. *Syst Biol* 68:365–369
- Baytshok V, Lu H, Park H, Kim S, Yu R, Chandran K (2009) Impact of varying electron donors on the molecular microbial ecology and biokinetics of methylotrophic denitrifying bacteria. *Biotechnol Bioeng* 102:1527–1536
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7:335–336

- Chen W-Y, Ng TH, Wu J-H, Chen J-W, Wang H-C (2017) Microbiome dynamics in a shrimp grow-out pond with possible outbreak of acute hepatopancreatic necrosis disease. *Sci Rep* 7:1–12
- Chen Z, Chang Z, Zhang L, Jiang Y, Ge H, Song X, Chen S, Zhao F, Li J (2019) Effects of water recirculation rate on the microbial community and water quality in relation to the growth and survival of white shrimp (*Litopenaeus vannamei*). *BMC Microbiol* 19:1–15
- Coclet C, Garnier C, Durrieu G, Omanović D, D'Onofrio S, Le Poupon C, Mullot J-U, Briand J-F, Misson B (2019) Changes in bacterioplankton communities resulting from direct and indirect interactions with trace metal gradients in an urbanized marine coastal area. *Front Microbiol* 10:257
- Conte A, Papale M, Amalfitano S, Mikkonen A, Rizzo C, De Domenico E, Michaud L, Giudice AL (2018) Bacterial community structure along the subtidal sandy sediment belt of a high Arctic fjord (Kongsfjorden, Svalbard Islands). *Sci Total Environ* 619:203–211
- Czech L, Barbera P, Stamatakis A (2020) Genesis and Gappa: processing, analyzing and visualizing phylogenetic (placement) data. *Bioinformatics* 36:3263–3265
- Deng Y, Xu X, Yin X, Lu H, Chen G, Yu J, Ruan Y (2019) Effect of stock density on the microbial community in biofloc water and Pacific white shrimp (*Litopenaeus vannamei*) gut microbiota. *Appl Microbiol Biotechnol* 103:4241–4252
- Edgar RC (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 10:996–998
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194–2200
- Gu W, Deng X, Lee M, Sucu YD, Arevalo S, Stryke D, Federman S, Gopez A, Reyes K, Zorn K (2021) Rapid pathogen detection by metagenomic next-generation sequencing of infected body fluids. *Nat Med* 27:115–124
- Gutiérrez MH, Narváez D, Daneri G, Montero P, Perez-Santos I, Pantoja S (2018) Linking seasonal reduction of microbial diversity to increase in winter temperature of waters of a Chilean Patagonia fjord. *Front Mar Sci* 5:277
- Hlorzdi V, Kuebutornye FKA, Afriyie G, Abarike ED, Lu Y, Chi S, Anokyewaa MA (2020) The use of *Bacillus* species in maintenance of water quality in aquaculture: a review. *Aquac Res* 18:100503
- Huang F, Pan L, Song M, Tian C, Gao S (2018) Microbiota assemblages of water, sediment, and intestine and their associations with environmental factors and shrimp physiological health. *Appl Microbiol Biotechnol* 102:8585–8598
- Huerta-Rábago JA, Martínez-Porchas M, Miranda-Baeza A, Nieves-Soto M, Rivas-Vega ME, Martínez-Córdova LR (2019) Addition of commercial probiotic in a biofloc shrimp farm of *Litopenaeus vannamei* during the nursery phase: effect on bacterial diversity using massive sequencing 16S rRNA. *Aquaculture* 502:391–399
- Imaizumi K, Molex W, Jitnavee C, Direkbusarakom S, Kondo H, Hirono I (2022) Bacterial and eukaryotic communities in pond water of whiteleg shrimp *Litopenaeus vannamei* and the bacterial communities of their stomach and midgut. *Aquaculture* 554:738139
- Infante-Villamil S, Huerlimann R, Jerry DR (2021) Microbiome diversity and dysbiosis in aquaculture. *Rev Aquac* 13:1077–1096
- Kajan K, Cukrov N, Cukrov N, Bishop-Pierce R, Orlić S (2022) Micro-eukaryotic and prokaryotic diversity of anchialine caves from Eastern Adriatic Sea Islands. *Microb Ecol* 83:257–270
- Kewcharoen W, Srisapoom P (2019) Probiotic effects of *Bacillus* spp. from Pacific white shrimp (*Litopenaeus vannamei*) on water quality and shrimp growth, immune responses, and resistance to *Vibrio parahaemolyticus* (AHPND strains). *Fish Shellfish Immunol* 94:175–189
- Khademzade O, Zakeri M, Haghi M, Mousavi SM (2020) The effects of water additive *Bacillus cereus* and *Pediococcus acidilactici* on water quality, growth performances, economic benefits, immunohematology and bacterial flora of whiteleg shrimp (*Penaeus vannamei* Boone, 1931) reared in earthen ponds. *Aquac Res* 51:1759–1770
- Kurniawinata MI, Sukenda S, Wahjuningrum D, Widanarni W (2022) Bacterial diversity and community composition in the gut and rearing water of Pacific White shrimp *Penaeus vannamei* during an outbreak of white feces disease. *Aquaculture* 559:738431
- Langille MG, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkepile DE, Vega Thurber RL, Knight R (2013) Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nat Biotechnol* 31:814–821
- Langwig MV, De Anda V, Dombrowski N, Seitz KW, Rambo IM, Greening C, Teske AP, Baker BJ (2022) Large-scale protein level comparison of Deltaproteobacteria reveals cohesive metabolic groups. *ISME J* 16:307–320
- Lara-Flores M (2011) The use of probiotic in aquaculture: an overview. *Int Res J Microbiol* 2:471–478
- Lin T-L, Shu C-C, Lai W-F, Tzeng C-M, Lai H-C, Lu C-C (2019) Investiture of next generation probiotics on amelioration of diseases—Strains do matter. *Med Microecol* 1:100002
- Louca S, Doebeli M (2018) Efficient comparative phylogenetics on large trees. *Bioinformatics* 34:1053–1055
- Magoč T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27:2957–2963
- Mirarab S, Nguyen N, Warnow T (2012) SEPP: SATé-enabled phylogenetic placement. *Biocomputing 2012*. World Scientific, pp 247–258
- Nair RR, Boobal R, Vrinda S, Bright Singh I, Valsamma J (2019) Ammonia-oxidizing bacterial and archaeal communities in tropical bioaugmented zero water exchange shrimp production systems. *J Soils Sed* 19:2126–2142
- Nguyen Thi Truc L, Nguyen Thanh T, Tran Thi Hong T, Van Pham D, Vo Thi Tuyet M, Nguyen Trong N, Phan Cong M, Cao Ngoc D, Truong Quoc P (2021) Effects of feed mixed with lactic acid bacteria and carbon, nitrogen, phosphorus supplied to the water on the growth and survival rate of white leg shrimp (*Penaeus vannamei*) infected with acute hepatopancreatic necrosis disease caused by *Vibrio parahaemolyticus*. *Biology* 10:280
- Parks DH, Tyson GW, Hugenholtz P, Beiko RG (2014) STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics* 30:3123–3124
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO (2012) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 41:D590–D596
- Raetz CR, Guan Z, Ingram BO, Six DA, Song F, Wang X, Zhao J (2009) Discovery of new biosynthetic pathways: the lipid a story. *J Lipid Res* 50(Suppl):S103–108
- Sahandi J, Sorgeloos P, Xiao H, Wang X, Qi Z, Zheng Y, Tang X (2019) The use of selected bacteria and yeasts to control *Vibrio* spp. in live food. *Antibiotics* 8:95
- Schomburg D, Schomburg I (2013) (KDO)-lipid IV A 3-deoxy-d-manno-octulosonic acid transferase 2.4.99.13, Class 2–3.2 transferases, hydrolases. Springer, pp 217–223
- Schröder M, Giermann N, Zrenner R (2005) Functional analysis of the pyrimidine de novo synthesis pathway in solanaceous species. *Plant Physiol* 138:1926–1938
- Shin J, Yun K-S, Gunathilaka BE, Hasanthi M, Ko D, Lim H, Lim J, Eom G, Kim H-S, Lee K-J (2023) Piperine supplementation in diet improves growth, feed efficiency, innate immunity,

- digestibility and disease resistance of Pacific white shrimp (*Litopenaeus vannamei*). *Aquac Rep* 29:101490
- Srinandan C, D'souza G, Srivastava N, Nayak BB, Nerurkar AS (2012) Carbon sources influence the nitrate removal activity, community structure and biofilm architecture. *Bioresour Technol* 117:292–299
- Supono S, Wardiyanto W, Harpeni E (2019) Identification of *Vibrio* sp. as a cause of white feces diseases in white shrimp *Penaeus vannamei* and handling with herbal ingredients in East Lampung Regency, Indonesia. *AACL Bioflux* 12:417–425
- Toribara NW, Ho SB, Gum E, Gum JR, Lau P, Kim YS (1997) The carboxyl-terminal sequence of the human secretory mucin, MUC6: analysis of the primary amino acid sequence*. *J Biol Chem* 272:16398–16403
- Wang W, Shi L, Qin Y, Li F (2020) Research and application of chondroitin sulfate/dermatan sulfate-degrading enzymes. *Front Cell Dev Biol* 8:560442
- Wei D, Zeng S, Hou D, Zhou R, Xing C, Deng X, Yu L, Wang H, Deng Z, Weng S (2021) Community diversity and abundance of ammonia-oxidizing archaea and bacteria in shrimp pond sediment at different culture stages. *J Appl Microbiol* 130:1442–1455
- Yang J, Chen F, Yu O, Beachy RN (2011) Controlled silencing of 4-coumarate:CoA ligase alters lignocellulose composition without affecting stem growth. *Plant Physiol Biochem* 49:103–109
- Yang W, Zhu J, Zheng C, Qiu H, Zheng Z, Lu K (2018) Succession of bacterioplankton community in intensive shrimp (*Litopenaeus vannamei*) aquaculture systems. *Aquaculture* 497:200–213
- Ye Y, Doak TG (2009) A parsimony approach to biological pathway reconstruction/inference for genomes and metagenomes. *PLoS Comput Biol* 5:e1000465
- Yu LH, Teh CSJ, Yap KP, Thong KL (2020) Diagnostic approaches and contribution of next-generation sequencing technologies in genomic investigation of *Vibrio parahaemolyticus* that caused acute hepatopancreatic necrosis disease (AHPND). *Aquac Int* 28:2547–2559
- Zafar H, Saier MH Jr (2018) Comparative genomics of transport proteins in seven *Bacteroides* species. *PLoS ONE* 13:e0208151
- Zhang X-H, He X, Austin B (2020) *Vibrio harveyi*: a serious pathogen of fish and invertebrates in mariculture. *Mar Life Sci Technol* 2:231–245
- Zhang X, Sun J, Han Z, Chen F, Lv A, Hu X, Sun X, Qi H, Guo Y (2021) *Vibrio parahaemolyticus* alters the community composition and function of intestinal microbiota in Pacific white shrimp, *Penaeus Vannamei*. *Aquaculture* 544:737061
- Zheng Y, Yu M, Liu Y, Su Y, Xu T, Yu M, Zhang X-H (2016) Comparison of cultivable bacterial communities associated with Pacific white shrimp (*Litopenaeus vannamei*) larvae at different health statuses and growth stages. *Aquaculture* 451:163–169
- Zheng X, Tang J, Ren G, Wang Y (2017) The effect of four microbial products on production performance and water quality in integrated culture of freshwater pearl mussel and fishes. *Aquac Res* 48:4897–4909

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

AQUAT-CEK

ORIGINALITY REPORT

16%

SIMILARITY INDEX

10%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Han, Li, Huang Shaobin, Wei Zhendong, Chen Pengfei, and Zhang Yongqing. "Performance of a new suspended filler biofilter for removal of nitrogen oxides under thermophilic conditions and microbial community analysis", *The Science of The Total Environment*, 2016.

Publication

1 %
- 2 www.ukm.my

Internet Source

<1 %
- 3 Alexander Dietrich, Monica Steffi Matchado, Maximilian Zwiebel, Benjamin Ölke et al. "Namco: A microbiome explorer", *Cold Spring Harbor Laboratory*, 2021

Publication

<1 %
- 4 Qi Chen, Bo Zhu, Dong Sun, Weijie Liu, Xian Sun, Shunshan Duan. "The effect of protocatechuic acid on the phycosphere in harmful algal bloom species *Scrippsiella trochoidea*", *Aquatic Toxicology*, 2020

Publication

<1 %

5

Awik P.D Nurhayati, Enny Zulaika, Muhamad Amin, Edwin Setiawan, Zaki Muhammad Wijaya. "Isolation and screening of lactic acid bacteria producing anti-edwardsiella from the gastrointestinal tract of wild-caught catfish (*Clarias gariepinus*) for probiotic candidates", Research Square Platform LLC, 2022

Publication

<1 %

6

David Allaway, Richard Haydock, Zoe N. Lonsdale, Oliver D. Deusch, Ciaran O'Flynn, Kevin R. Hughes. "Rapid Reconstitution of the Faecal Microbiome after Extended Diet-Induced Changes Indicate a Stable Gut Microbiome in Healthy Adult Dogs", Applied and Environmental Microbiology, 2020

Publication

<1 %

7

Bin Wang, Shu-qin Zhang, Jia-li Dong, Yuan Li, Yu-xiao Jin, Hui-wen Xiao, Hai-chao Wang, Sai-jun Fan, Ming Cui. "Ambient temperature structures the gut microbiota of zebrafish to impact the response to radioactive pollution", Environmental Pollution, 2022

Publication

<1 %

8

jurnal-iktiologi.org

Internet Source

<1 %

9

Nuril Azhar, Ervia Yudiati. "Outbreak simulation of *Litopenaeus vannamei* recovery rate with oral alginate and spirulina diet

<1 %

supplementation against *Vibrio*
parahaemolyticus AHPND", Aquaculture
International, 2023

Publication

10

biotechnologyforbiofuels.biomedcentral.com

Internet Source

<1 %

11

www.pure.ed.ac.uk

Internet Source

<1 %

12

Liu Ruizhe, Wang Shan, Huang Dongliang,
Huang Yulu, He Tianliang, Chen Xinhua. " The
probiotic roles of E2 as a dietary supplement
in growth promotion and disease resistance
of juvenile large yellow croaker () ", Cold
Spring Harbor Laboratory, 2023

Publication

<1 %

13

pureadmin.uhi.ac.uk

Internet Source

<1 %

14

Yuxin Hu, Hanyue Chen, Yihang Jiang, Jianguo
Zhang, Xiongqing Zhang. "Effects of forest age
and seasonal changes on soil microbial
community diversity in Chinese fir
plantations", Research Square Platform LLC,
2023

Publication

<1 %

15

public.thinkonweb.com

Internet Source

<1 %

bkms.brenda-enzymes.org

16

Internet Source

<1 %

17

rupress.org

Internet Source

<1 %

18

Kaouther Mejri Kousri, Amel Belaaj Zouari, Marouan Meddeb, Oumayma Chkili et al. "Structure of planktonic food web in the Gulf of Gabès (Southeastern Mediterranean): potential importance of heterotrophic and mixotrophic microzooplankton", Aquatic Sciences, 2023

Publication

<1 %

19

Yuan Yuan, Li Lu, Nan Bo, Yang Chaoyue, Yan Haiyang. "Allicin Ameliorates Intestinal Barrier Damage via Microbiota-Regulated Short-Chain Fatty Acids-TLR4/MyD88/NF-κB Cascade Response in Acrylamide-Induced Rats", Journal of Agricultural and Food Chemistry, 2021

Publication

<1 %

20

patents.google.com

Internet Source

<1 %

21

repository.lppm.unila.ac.id

Internet Source

<1 %

22

Shujie Xu, Fei Wang, Peng Zou, Xiang Li, Qian Jin, Qi Wang, Baikui Wang, Yuanhao Zhou, Li Tang, Dongyou Yu, Weifen Li. "Bacillus

<1 %

amyloliquefaciens SC06 in the diet improves egg quality of hens by altering intestinal microbiota and the effect is diminished by antimicrobial peptide", *Frontiers in Nutrition*, 2022

Publication

23

Xinzhu Chen, Yao Zhang, Jianguo Zhang, Xiusheng Huang, Qinlou Huang, Jing Liu, Wen Yue. "Dynamics of microbial community composition on grass surfaces in response to glucose and sucrose spraying for enhanced silage quality", *Research Square Platform LLC*, 2023

Publication

<1 %

24

www.hindawi.com

Internet Source

<1 %

25

www.nwipb.cas.cn

Internet Source

<1 %

26

www.verticalfarmdaily.com

Internet Source

<1 %

27

Gi Ppeum Han, Kyu-Chan Lee, Hwan Ku Kang, Han Na Oh, Woo Jun Sul, Dong Yong Kil. "Analysis of excreta bacterial community after forced molting in aged laying hens", *Asian-Australasian Journal of Animal Sciences*, 2019

Publication

<1 %

28

Hao Guan, Yanhong Yan, Xiaoling Li, Xiaomei Li, Yang Shuai, Guangyan Feng, Qifan Ran, Yimin Cai, Ying Li, Xinquan Zhang. "Microbial communities and natural fermentation of corn silages prepared with farm bunker-silo in Southwest China", *Bioresource Technology*, 2018

Publication

<1 %

29

Sitong Dong, Jiao jiao, Shuangshuo Jia, Gaoyu Li, Wei Zhang, Kai Yang, Zhen Wang, Chao Liu, Da Li, Xiuxia Wang. "16S rDNA Full-Length Assembly Sequencing Technology Analysis of Intestinal Microbiome in Polycystic Ovary Syndrome", *Frontiers in Cellular and Infection Microbiology*, 2021

Publication

<1 %

30

Yustian Rovi Alfiansah, Sonja Peters, Jens Harder, Christiane Hassenrück, Astrid Gärdes. "Structure and co-occurrence patterns of bacterial communities associated with white faeces disease outbreaks in Pacific white-leg shrimp *Penaeus vannamei* aquaculture", *Scientific Reports*, 2020

Publication

<1 %

31

escholarship.org

Internet Source

<1 %

32

jjbs.hu.edu.jo

Internet Source

<1 %

33

proceeding.unram.ac.id

Internet Source

<1 %

34

Jennifer G. Jankowiak, Christopher J. Gobler. "The Composition and Function of Microbiomes Within Microcystis Colonies Are Significantly Different Than Native Bacterial Assemblages in Two North American Lakes", *Frontiers in Microbiology*, 2020

Publication

<1 %

35

Ludovica Rolando, Paola Grenni, Jasmin Rauseo, Tanita Pescatore et al. "Isolation and Characterization in a Soil Conditioned With Foaming Agents of a Bacterial Consortium Able to Degrade Sodium Lauryl Ether Sulfate", *Frontiers in Microbiology*, 2020

Publication

<1 %

36

Wen Yang, Jinyong Zhu, Cheng Zheng, Haijie Qiu, Zhongming Zheng, Kaihong Lu. "Succession of bacterioplankton community in intensive shrimp (*Litopenaeus vannamei*) aquaculture systems", *Aquaculture*, 2018

Publication

<1 %

37

Young-Sam Kim, Sang-Eon Kim, Seong-Jin Kim, Hyun-Kyoung Jung et al. "Effects of wheat flour and culture period on bacterial community composition in digestive tracts of *Litopenaeus vannamei* and rearing water in

<1 %

biofloc aquaculture system", Aquaculture, 2021

Publication

38

github.com

Internet Source

<1 %

39

m.moam.info

Internet Source

<1 %

40

www.ijbs.com

Internet Source

<1 %

41

Cecília de Souza Valente, Alex H.L. Wan. "Vibrio and major commercially important vibriosis diseases in decapod crustaceans", *Journal of Invertebrate Pathology*, 2021

Publication

<1 %

42

Muhammad Usama Marghoob, Alejandro Rodriguez-Sanchez, Asma Imran, Fathia Mubeen, Lori Hoagland. "Diversity and functional traits of indigenous soil microbial flora associated with salinity and heavy metal concentrations in agricultural fields within the Indus Basin region, Pakistan", *Frontiers in Microbiology*, 2022

Publication

<1 %

43

Ramya R. Nair, R. Boobal, S. Vrinda, I. S. Bright Singh, Joseph Valsamma. "Ammonia-oxidizing bacterial and archaeal communities in tropical bioaugmented zero water exchange shrimp

<1 %

production systems", Journal of Soils and Sediments, 2018

Publication

44

Jianchun Shao, Keyong Jiang, Lei Wang. " fed diets with different replacement levels of fish meal by fish silage: A molecular approach on intestinal microbiota ", Aquaculture Nutrition, 2019

Publication

45

Jiung-Wen Chen, Wei-Fan Chiang, Jer-Horng Wu, Yuh-Ling Chen, Wei-Sheng Wu, Li-Wha Wu. "Functional Profiling of Saliva Microbiome is Essential for Oral Cancer Prediction", Research Square, 2020

Publication

46

Sebastian VadilloGonzalez, Sofie Vranken, Melinda A. Coleman, Thomas Wernberg, Peter D. Steinberg, Ezequiel M. Marzinelli. " Host genotype and microbiome associations in co - occurring clonal and non - clonal kelp, ", Molecular Ecology, 2023

Publication

47

Shan Cong, Lingrui Wang, Yan Meng, Xuanlin Cai, Chenxi Zhang, Yanqin Gu, Xiumin Ma, Li Luo. " oral liquid regulates gut microbiota and serum metabolism during alleviation of collagen - induced arthritis in rats ", Phytotherapy Research, 2022

Publication

<1 %

<1 %

<1 %

<1 %

48 sites.utexas.edu Internet Source <1 %

49 wjgnet.com Internet Source <1 %

50 www.leibniz-zmt.de Internet Source <1 %

51 Carolina Ramírez, María Soledad Gutiérrez, Lucas Venegas, Catalina Sapag et al. "Microbiota composition and susceptibility to florfenicol and oxytetracycline of bacterial isolates from mussels (*Mytilus* spp.) reared on different years and distance from salmon farms", Environmental Research, 2022
Publication <1 %

52 Catarina Leal, Ales Eichmeier, Kateřina Štůsková, Josep Armengol et al. "Biocontrol agents establishment and their impact on rhizosphere microbiome and induced grapevine defenses is highly soil-dependent", Research Square Platform LLC, 2023
Publication <1 %

53 Darong Yang, Zhenhua Wang, Xixi Dai, Minggang Liu, Dongmei Zhang, Yan Zeng, Dong Zeng, Xueqin Ni, Kangcheng Pan. "Addition of *Brevibacillus laterosporus* to the rearing water enhances the water quality, growth performance, antioxidant capacity,

and digestive enzyme activity of crucian carp
Carassius auratus", Fisheries Science, 2023

Publication

54

GBOLAGA OLUWASEGUN OLANREWAJU,
ABIOLA O AREMU, David D Sarpong,
OMOBOLANLE E ADE-ADEMILUA. "Growth
performance and nutritional composition of
Clarias gariepinus in flood-and-drain
aquaponics coupled with three species of
leafy vegetable.", Cold Spring Harbor
Laboratory, 2022

Publication

<1 %

55

Mohsen Abdel-Tawwab, Hossam A.M.
Mounes, Sherien H.H. Shady, Kareem M.
Ahmed. "Effects of yucca, *Yucca schidigera*,
extract and/or yeast, *Saccharomyces*
cerevisiae, as water additives on growth,
biochemical, and antioxidants/oxidant
biomarkers of Nile tilapia, *Oreochromis*
niloticus", Aquaculture, 2021

Publication

<1 %

56

Parisa Kheirandish, Renee Maxine Petri, Arife
Sener - Aydemir, Heidi Elisabeth Schwartz -
Zimmermann et al. " Characterization of
Microbial Intolerances and Ruminal Dysbiosis
Towards Different Dietary Carbohydrate
Sources Using an Model ", Journal of Applied
Microbiology, 2022

Publication

<1 %

57 Peng Sheng, Qiang Chen, Qingxi Han, Jiaqi Lu, Bo Zhang, Maowang Jiang, Ruibing Peng, Xiamin Jiang. "Bacterial community in gut, shell sediment, and surrounding water of *Hyriopsis cumingii* fed three different microalgal diets", *Aquaculture*, 2021
Publication

58 Sameeha Misriya Shroff, chandrashekhar T Sreeramare. "Marketing claims, promotional strategies, and product information on Malaysian e-cigarette retailer websites - a systematic content analysis", *Research Square Platform LLC*, 2023
Publication

59 Zidan Liu, Linglin Qiuqian, Zhiyuan Yao, Xin Wang, Lei Huang, Jialai Zheng, Kai Wang, Laiguo Li, Demin Zhang. "Effects of a Commercial Microbial Agent on the Bacterial Communities in Shrimp Culture System", *Frontiers in Microbiology*, 2018
Publication

60 etd.auburn.edu
Internet Source

61 lipidworld.biomedcentral.com
Internet Source

62 mbio.asm.org
Internet Source

63

soybase.org

Internet Source

<1 %

64

www.medrxiv.org

Internet Source

<1 %

65

Estefanía Garibay-Valdez, Luis Rafael Martínez-Córdova, Marco A. López-Torres, F. Javier Almendariz-Tapia et al. "The implication of metabolically active *Vibrio* spp. in the digestive tract of *Litopenaeus vannamei* for its post-larval development", *Scientific Reports*, 2020

Publication

<1 %

66

Ken Cullings, Shilpa R Bhardwaj, Michael Spector. "Variation in the Microbiomes of the Basidiomycete Fungi *Scleroderma citrinum* (Pers.) and *Pisolithus arhizus* (Pers.): a tale of two saprotrophs.", *Cold Spring Harbor Laboratory*, 2023

Publication

<1 %

67

Kentaro Imaizumi, Wanlapha Molex, Chakrit Jitnavee, Sataporn Direkbusarakom, Hidehiro Kondo, Ikuo Hirono. "Bacterial and eukaryotic communities in pond water of whiteleg shrimp *Litopenaeus vannamei* and the bacterial communities of their stomach and midgut", *Aquaculture*, 2022

Publication

<1 %

68

Kui Li, Shuang Zhao, Weibing Guan, Kejun Li. " Planktonic bacteria in white shrimp () and channel catfish () aquaculture ponds in a salt - alkaline region ", Letters in Applied Microbiology, 2021

Publication

<1 %

69

Linh Nguyen Thi Truc, Tuu Nguyen Thanh, To Tran Thi Hong, Day Pham Van et al. "Effects of Feed Mixed with Lactic Acid Bacteria and Carbon, Nitrogen, Phosphorus Supplied to the Water on the Growth and Survival Rate of White Leg Shrimp (Penaeus vannamei) Infected with Acute Hepatopancreatic Necrosis Disease Caused by Vibrio parahaemolyticus", Biology, 2021

Publication

<1 %

70

Neyssa M. Marina, Qi Liu, Sarah S. Donaldson, Charles A. Sklar et al. "Longitudinal follow-up of adult survivors of Ewing sarcoma: A report from the Childhood Cancer Survivor Study", Cancer, 2017

Publication

<1 %

71

Ning Kong, Shuo Han, Qiang Fu, Zichao Yu, Lingling Wang, Linsheng Song. "Impact of ocean acidification on the intestinal microflora of the Pacific oyster Crassostrea gigas", Aquaculture, 2022

Publication

<1 %

72

Paulina Średnicka, Edyta Juszcuk-Kubiak, Michał Wójcicki, Monika Akimowicz, Marek Ł. Roszko. "Probiotics as a biological detoxification tool of food chemical contamination: A review", *Food and Chemical Toxicology*, 2021

Publication

<1 %

73

Qiufen Li. "Chapter 9 Probiotics for Biofloc System and Water Quality", Springer Science and Business Media LLC, 2022

Publication

<1 %

74

Surapun Tapaamorndech, Kanittha Chantarasakha, Yutthana Kingcha, Sage Chaiyapechara et al. "Effects of *Bacillus aryabhattai* TBRC8450 on vibriosis resistance and immune enhancement in Pacific white shrimp, *Litopenaeus vannamei*", *Fish & Shellfish Immunology*, 2018

Publication

<1 %

75

Vanessa Oliveira, Newton C.M. Gomes, Daniel F.R. Cleary, Adelaide Almeida et al. "Halophyte plant colonization as a driver of the composition of bacterial communities in salt marshes chronically exposed to oil hydrocarbons", *FEMS Microbiology Ecology*, 2014

Publication

<1 %

76

Wen Yang, Shi-Zhan Zheng, Shou-Heng Zhou, Li Zhao et al. "Structure and Functional Diversity of Surface Bacterioplankton Communities in an Overwintering Habitat for Large Yellow Croaker, *Pseudosciaena crocea*, of the Southern East China Sea", *Frontiers in Marine Science*, 2020

Publication

<1 %

77

Yihui Ding, Nuo Chen, Jiangdong Ke, Zhenyu Qi, Wenjie Chen, Sizhi Sun, Zhongming Zheng, Jilin Xu, Wen Yang. "Response of the rearing water bacterial community to the beneficial microalga *Nannochloropsis oculata* cocultured with Pacific white shrimp (*Litopenaeus vannamei*)", *Aquaculture*, 2021

Publication

<1 %

78

Yuanfeng Liu, Xiang Li, Yudie Yang, Ye Liu, Shijun Wang, Boyang Ji, Yongjun Wei. "Exploring Gut Microbiota in Patients with Colorectal Disease Based on 16S rRNA Gene Amplicon and Shallow Metagenomic Sequencing", *Frontiers in Molecular Biosciences*, 2021

Publication

<1 %

79

hal.science

Internet Source

<1 %

80

ijabe.org

Internet Source

<1 %

81	medcraveonline.com Internet Source	<1 %
82	pubag.nal.usda.gov Internet Source	<1 %
83	researchmgt.monash.edu Internet Source	<1 %
84	vetmedmosul.com Internet Source	<1 %
85	www.bioflux.com.ro Internet Source	<1 %
86	www.scielo.cl Internet Source	<1 %
87	Farapti Farapti, Chusnul Fadilla, Niwanda Yogiswara, Merryana Adriani. "Effects of vitamin D supplementation on 25(OH)D levels and blood pressure in the elderly: a systematic review and meta-analysis", F1000Research, 2020 Publication	<1 %
88	Mohamed I. Kord, Sahya Maulu, Tarek M. Srour, Eglal A. Omar et al. "Impacts of water additives on water quality, production efficiency, intestinal morphology, gut microbiota, and immunological responses of Nile tilapia fingerlings under a zero-water-exchange system", Aquaculture, 2022	<1 %

89

Nicolas Layglon, Benjamin Misson, Gaël Durieu, Clément Coclet et al. "Long-term monitoring emphasizes impacts of the dredging on dissolved Cu and Pb contamination along with ultraplankton distribution and structure in Toulon Bay (NW Mediterranean Sea, France)", *Marine Pollution Bulletin*, 2020

Publication

<1 %

90

Peng Yu, Hongwei Shan, Yu Cheng, Jingjing Ma, Kai Wang, Hongyang Li. "Translucent disease outbreak in *Penaeus vannamei* post-larva accompanies the imbalance of pond water and shrimp gut microbiota homeostasis", *Aquaculture Reports*, 2022

Publication

<1 %

91

Rooks, Michelle G, Patrick Veiga, Leslie H Wardwell-Scott, Timothy Tickle, Nicola Segata, Monia Michaud, Carey Ann Gallini, Chloé Beal, Johan ET van Hylckama-Vlieg, Sonia A Ballal, Xochitl C Morgan, Jonathan N Glickman, Dirk Gevers, Curtis Huttenhower, and Wendy S Garrett. "Gut microbiome composition and function in experimental colitis during active disease and treatment-induced remission", *The ISME Journal*, 2014.

Publication

<1 %

92

Clément Coclet, Cédric Garnier, Gaël Durrieu, Dario Omanović et al. "Changes in Bacterioplankton Communities Resulting From Direct and Indirect Interactions With Trace Metal Gradients in an Urbanized Marine Coastal Area", *Frontiers in Microbiology*, 2019

Publication

<1 %

93

Dae-Young Kim, Surendra Krushna Shinde, Avinash Ashok Kadam, Rijuta Ganesh Saratale et al. "Advantage of Species Diversification to Facilitate Sustainable Development of Aquaculture Sector", *Biology*, 2022

Publication

<1 %

94

M Ali, M Zubair, A Rosyidi, M Amin. "Screening of ammonia-degrading bacteria to reduce ammonia content in the manure of laying hens", *IOP Conference Series: Earth and Environmental Science*, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

AQUAT-CEK

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13
