

Clinical Findings in Biliary Atresia A Literature Review

by Bagus Setyoboedi

Submission date: 03-Feb-2024 10:28PM (UTC+0800)

Submission ID: 2285263616

File name: 08._Clinical_Findings_in_Biliary_Atresia_A_Literature_Review.pdf (7.18M)

Word count: 2787

Character count: 16254

Clinical Findings in Biliary Atresia: A Literature Review

Daryn Ikrima Mumtaz Zahravi^a, Bagus Setyoboedi^{*}, Ariandi Setiawan^c, Alpha Fardah Athiyyah^b

17

^{*}Corresponding E-mail: bagus.setyoboedi@fk.unair.ac.id

^a Medical Programme, Faculty of medicine, Universitas Airlangga, Surabaya, Indonesia

^bDepartment of Pediatric, Dr. Soetomo General Hospital, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

^cDepartment of Surgery, Dr. Soetomo General Hospital, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

2 Abstract

Biliary atresia is an obstructive disease affecting the biliary tract. Patients with biliary atresia often appear similar to normal infants, leading to delayed detection in many cases. There are several clinical symptoms and typical laboratory findings that typically manifest in patients with biliary atresia. As the age of the patient advances, the manifestations worsen, potentially leading to the development of cirrhosis, liver failure by the age of 1, and other complications. The aim of this review is to understand the clinical finding of biliary atresia patients, facilitating early detection and intervention to reduce the likelihood of the disease progressing to a more severe stage. Delay in the diagnosis of biliary atresia can result in a poor prognosis.

Keywords: Biliary Atresia; Clinical sign; Laboratory result; Treatment

2 1. Introduction

Biliary atresia is an obstructive disease of the biliary tract that results in the inability of bile to reach the intestinal tract [1]. Biliary atresia is found in 1 in every 8000 to 18000 births and is more prevalent in Asian and African populations compared to Europeans [2]. Biliary atresia has some common symptoms include jaundice, pale stools, and hepatomegaly. Untreated biliary atresia can lead to liver cirrhosis, liver failure by the first year of life, and other complications [3,4].

Biliary atresia presents in two forms based on associated anomalies: perinatal and embryonic. The embryonic type has an earlier onset and is often accompanied by abnormal developments. The perinatal type is suspected to result from postnatal cholangiopathy obstruction and presents with alcoholic stools [3,5]. The Japanese Association of Pediatric Surgeons categorizes biliary atresia into three main types based on the location of obstruction. Type I involves atresia in the common bile duct, while the gallbladder and hepatic

ducts remain patent (referred to as distal biliary atresia). Type II further divides into IIA and IIb. In Type IIA, the gallbladder and common bile duct are patent, whereas Type IIb involves obstruction in the gallbladder, cystic duct, and common bile duct. Type III, the most prevalent condition at 73%, is characterized by obliteration of all extrahepatic bile ducts (complete biliary atresia) [6].

Biliary atresia results from inflammation and fibrous tissue formation, leading to obstruction of the extrahepatic bile ducts, disrupting bile flow [7]. Simultaneously, intrahepatic bile ducts undergo hyperplasia. There is substantial evidence indicating that viruses, toxins, and genetic variations contribute to the etiology of biliary atresia, eliciting an inflammatory response affecting the epithelial layer. Immune cells activate the expression of type 2 cytokines, inducing epithelial cell proliferation and the formation of matrix cells from non-parenchymal cells [1,8].

2. Clinical Manifestation

The most prevalent clinical manifestation in patients with Biliary Atresia is jaundice. This occurs due to the accumulation of bile fluid in the body, resulting in a yellowish appearance in the sclera, skin, and mucous membranes. Biliary atresia patients experience post-hepatic jaundice, a condition where the blocked bile ducts affect the amount of conjugated bilirubin [9]. This leads to the buildup of bilirubin in the circulation, soluble in fat beneath the skin, causing the appearance of jaundice [10]. Many infants exhibit jaundice in the early stages of life, making it challenging to differentiate if the jaundice is physiological or not. [11]. Jaundice in biliary atresia patients differs from physiological jaundice. Physiological jaundice typically improves with exposure to UV light, whereas pathological jaundice, as in biliary atresia, does not improve even with sunlight exposure and worsens over time. [12].

Additionally, biliary atresia patients typically present with pale stools. This is because bilirubin, which should reach the intestines, fails to do so. Bilirubin in the digestive tract serves as a colorant for feces. Some countries have started implementing fecal color detection cards for early biliary atresia detection,

including at RS DR. Soetomo Surabaya. These cards display various stool colors ranging from pale to normal, aiding parents in identifying alarming signs of their child's stool color. [13,14].

Another common symptom in biliary atresia patients is dark-colored urine. Normally, urine does not contain bilirubin. However, in biliary atresia cases, ⁵ conjugated bilirubin is excreted through the kidneys, giving the urine a dark color [15]. Another commonly observed symptom is hepatomegaly [16]. Jaundice, pale stools, and dark-colored urine are an early signs of biliary atresia, so there is no difference between early and late diagnosis symptoms in biliary atresia. [14].

3. Laboratory Result

Patients with biliary atresia typically exhibit elevated levels of conjugated bilirubin, a slight increase in serum gamma-glutamyl transferase (GGT), ALP, and elevation in serum transaminases [17]. The increase in conjugated bilirubin is indicate of liver cell damage or cholestasis. Although the measurement of conjugated bilirubin is effective in detecting liver diseases in neonates, it is not entirely accurate for biliary atresia, necessitating further examination. Several studies have explored early detection strategies for infants, with commonly used methods worldwide including the measurement of conjugated bilirubin levels and the use of color stool cards provided to parents [18].

Patients also tend to exhibit elevated levels of GGT [19]. Elevated GGT levels can serve as a crucial parameter to rule out other diagnoses related to neonatal cholestasis. ¹³ Infants with biliary atresia tend to have higher GGT levels compared to those without biliary atresia. However, it's important to note that GGT measurements are most effective when performed in infants aged less than 120 days [20].

Various enzymes, including SGPT, SGOT, ALP, among others, can indicate liver function. In this context, liver function tests aim to identify the source of liver damage. An increase in the SGPT and SGOT ratio to alkaline phosphatase (ALP) and bilirubin indicates hepatocellular disease. Conversely, disproportionate increases in SGPT and SGOT to ALP and bilirubin suggest a cholestatic condition,

prompting the consideration of differential diagnoses. SGPT and SGOT are part of serum aminotransferases, which are markers of liver cell damage [21,22].

4. Diagnosis

Early diagnosis, ideally within the first 3 months, is crucial. This is because cirrhosis can begin to develop at around 3 months of age, leading to poorer outcomes. Hepatosplenomegaly and ascites tend to appear in patients older than 3 months, indicating cirrhosis [23]. Late diagnosis is considered when the patient is over 90 days old [24].

Abdominal ultrasonography is the first and non-invasive imaging examination of choice for suspected biliary atresia patients [25]. One of the findings in the ultrasonography (USG) of patients with biliary atresia is the presence of the triangular cord sign [26]. Another diagnostic option is percutaneous cholecysto-cholangiography, which is more accurate as it directly observes the progressive filling of iodine in the bile ducts, confirming biliary atresia in challenging cases. The absence of retrograde contrast filling in the intrahepatic bile ducts is also indicative of biliary atresia. In patients without biliary atresia, clarity in both intrahepatic and extrahepatic bile ducts is always observed [27].

Histopathological examination findings in biliary atresia patients reveal bile duct proliferation (BDP), bile duct obstruction, portal infiltration with lymphocytes, neutrophils, and eosinophils, hepatocyte transformation, and liver cell swelling [28].

5. Treatment

Infants with biliary atresia can have a better prognosis if detected and treated early, ideally within the first 8 weeks of life. The procedures used to manage biliary atresia include the Kasai Procedure (KP) and liver transplantation.

5.1 Kasai Procedure

The Kasai Procedure aims to restore bile flow between the liver and the intestine, helping to reduce the progression of liver disease using a Roux-en-Y jejunal limb. Kasai Procedure can successfully restore bile ducts from the liver to the intestine especially when performed within the first 60 days of life. The success of the Kasai Procedure is determined by the restoration of bile flow, the return of normal stool color, and the disappearance of jaundice [25,29]. It is advisable to perform the Kasai Procedure before the age of 3 months to prevent further liver damage or progression to cirrhosis [23].

5.2 Liver Transplantation

Liver transplantation (LT) is considered in various conditions, including Kasai Procedure failure, significant malnutrition, recurrent cholangitis, portal hypertension, hepatopulmonary syndrome, and malignancies. Unfortunately, patients with biliary atresia often face long waiting times for liver transplantation [23]. The Pediatric End-stage Liver Disease (PELD) score is used to allocate liver transplants in children, considering factors such as albumin levels, bilirubin levels, prothrombin time, and growth failure [30]. Early detection and appropriate intervention significantly impact the prognosis and quality of life for patients with biliary atresia.

6. Conclusions

Biliary atresia is a neonatal disease that requires early diagnosis. Commonly observed symptoms include jaundice, pale stools, dark urine, and hepatomegaly. Laboratory test results in patients with biliary atresia tend to show elevated levels of conjugated bilirubin, GGT, and serum transaminases. The later the diagnosis, the more likely patients are to experience clinical deterioration, and the efficacy of treatment is compromised. Ultrasonography (USG) is routinely performed in the early stages of diagnosis to assess the patient's biliary tract condition. A more accurate diagnostic procedure is percutaneous cholecysto-cholangiography. Treatment options for biliary atresia include the Kasai Procedure and Liver Transplant. Therefore, it is imperative to prioritize early diagnosis for biliary atresia.

References

1. Asai, A., Miethke, A., & Bezerra, J. A. (2015). Pathogenesis of biliary atresia: defining biology to understand clinical phenotypes. *Nature reviews. Gastroenterology & hepatology*, 12(6), 342–352. <https://doi.org/10.1038/nrgastro.2015.74>
2. Bezerra, J. A., Wells, R. G., Mack, C. L., Karpen, S. J., Hoofnagle, J. H., Doo, E., & Sokol, R. J. (2018). Biliary Atresia: Clinical and Research Challenges for the Twenty-First Century. *Hepatology (Baltimore, Md.)*, 68(3), 1163–1173. <https://doi.org/10.1002/hep.29905>
3. Siddiqui AI, Ahmad T. Biliary Atresia. [Updated 2023 Jun 26]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK537262/>
4. Fanna, M., Masson, G., Capito, C., Girard, M., Guerin, F., Hermeziu, B., Lachaux, A., Roquelaure, B., Gottrand, F., Broue, P. and Dabadie, A., 2019. Management of biliary atresia in France 1986 to 2015: long-term results. *Journal of pediatric gastroenterology and nutrition*, 69(4), pp.416-424.
5. Ghazy, R.M., Adawy, N.M., Khedr, M.A. and Tahoun, M.M., (2018). Biliary atresia recent insight. *Egyptian Pediatric Association Gazette*, 66(1), pp.1-8
6. Wildhaber B. E. (2012). Biliary atresia: 50 years after the first kasai. *ISRN surgery*, 2012, 132089. <https://doi.org/10.5402/2012/132089>
7. Vij, M., & Rela, M. (2020). Biliary atresia: pathology, etiology and pathogenesis. *Future science OA*, 6(5), FSO466. <https://doi.org/10.2144/fsoa-2019-0153>
8. Ortiz-Perez, A., Donnelly, B., Temple, H., Tiao, G., Bansal, R., & Mohanty, S. K. (2020). Innate Immunity and Pathogenesis of Biliary Atresia. *Frontiers in immunology*, 11, 329. <https://doi.org/10.3389/fimmu.2020.00329>
9. Joseph, A., & Samant, H. (2022). Jaundice. In *StatPearls*. StatPearls Publishing.
10. Ansong-Assoku B, Shah SD, Adnan M, et al. Neonatal Jaundice. [Updated 2023 Feb 20]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK532930/>
11. Lianti, H., Kurnia, N., Rinaldhy, K., Aji, A.S., Ismet, M.F. and Amaliah, R., (2020). Analysis of knowledge regarding Biliary Atresia among healthcare providers and laypersons in East Jakarta after educational intervention. *ASEAN Journal of Community Engagement*, 4(1), p.13.
12. Setyoboedi, B., Prihaningtyas, R.A., Utomo, M.T. and Arief, S., (2022). Early detection of biliary atresia in primary health care: still a problem. *F1000Research*, 11, p.1245.
13. Borgeat, M., Korff, S., & Wildhaber, B. E. (2018). Newborn biliary atresia screening with the stool colour card: a questionnaire survey of parents. *BMJ paediatrics open*, 2(1), e000269. <https://doi.org/10.1136/bmjpo-2018-000269>
14. Couturier, L., Jarvis, C., Rousseau, H., & Jimenez, V. (2015). Biliary atresia. *Canadian family physician Medecin de famille canadien*, 61(11), 965–968.
15. Coucke EM, Akbar H, Kahloon A, et al. Biliary Obstruction. [Updated 2022 Nov 26]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK539698/#>
16. He, Lin, Patrick Ho Yu Chung, Vincent Chi Hang Lui, Clara Sze Man Tang, and Paul Kwong Hang Tam. 2022. "Current Understanding in the Clinical Characteristics and Molecular Mechanisms in Different Subtypes of Biliary Atresia" *International Journal of Molecular Sciences* 23, no. 9: 4841. <https://doi.org/10.3390/ijms23094841>
17. Abdel-Aziz, S. A., Sira, M. M., Gad, E. H., Ayoub, I., & Soltan, M. (2019). Preoperative alkaline phosphatase is a potential predictor of short-term outcome of surgery in infants with biliary atresia. *Clinical and experimental hepatology*, 5(2), 155–160. <https://doi.org/10.5114/ceh.2019.85072>
18. Wang, K. S., Section on Surgery, Committee on Fetus and Newborn, & Childhood Liver Disease Research Network (2015). Newborn Screening for Biliary Atresia. *Pediatrics*, 136(6), e1663–e1669. <https://doi.org/10.1542/peds.2015-3570>
19. Shankar, S., Bolia, R., Foo, H. W., D'Arcy, C. E., Hardikar, N., Wensing, M., & Hardikar, W. (2020). Normal Gamma Glutamyl Transferase Levels at Presentation Predict Poor Outcome in Biliary Atresia. *Journal of pediatric gastroenterology and nutrition*, 70(3), 350–355. <https://doi.org/10.1097/MPG.0000000000002563>
20. Govindarajan K. K. (2016). Biliary atresia: Where do we stand now?. *World journal of hepatology*, 8(36), 1593–1601. <https://doi.org/10.4254/wjh.v8.i36.1593>
21. Lala V, Zubair M, Minter DA. Liver Function Tests. [Updated 2023 Jul 30]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482489/>
22. Rafeey, M., Saboktakin, L., Shoa Hassani, J., Farahmand, F., Aslanabadi, S., Ghorbani-Haghjoui, A., & Poorebrahim, S. (2014). Diagnostic value of procalcitonin and apo-e in extrahepatic biliary atresia. *Iranian journal of pediatrics*, 24(5), 623–629.
23. Sundaram, S.S., Mack, C.L., Feldman, A.G. and Sokol, R.J., 2017. Biliary atresia: Indications and timing of liver transplantation and optimization of pretransplant care. *Liver Transplantation*, 23(1), pp.96-109.
24. Fuchs, J., Mrad, C., Gonzales, E., Ndiaye, D., Fouquet, V., Héry, G., Baujard, C., Guérin, F. and Branchereau, S., 2023. Biliary drainage surgery before or after 3 months of life versus primary liver transplantation in children with biliary atresia: comparative cohort study. *BJS open*, 7(2), p.zrac175.
25. Harpavat, S., Hawthorne, K., Setchell, K. D. R., Rivas, M. N., Henn, L., Beil, C. A., Karpen, S. J., Ng, V. L., Alonso, E. M., Bezerra, J. A., Guthery, S. L., Horslen, S., Loomes, K. M., McKiernan, P., Magee, J. C., Merion, R. M., Molleston, J. P., Rosenthal, P., Thompson, R. J., Wang, K. S., ... for Childhood Liver Disease Research Network (ChILDRen) (2023). Serum

- bile acids as a prognostic biomarker in biliary atresia following Kasai portoenterostomy. *Hepatology (Baltimore, Md.)*, 77(3), 862–873. <https://doi.org/10.1002/hep.32800>
26. Zhou, W., & Zhou, L. (2021). Ultrasound for the Diagnosis of Biliary Atresia: From Conventional Ultrasound to Artificial Intelligence. *Diagnostics (Basel, Switzerland)*, 12(1), 51. <https://doi.org/10.3390/diagnostics12010051>
 27. Napolitano, M., Franchi-Abella, S., Damasio, B.M. *et al.* (2021). Practical approach for the diagnosis of biliary atresia on imaging, part 2: magnetic resonance cholangiopancreatography, hepatobiliary scintigraphy, percutaneous cholecysto-cholangiography, endoscopic retrograde cholangiopancreatography, percutaneous liver biopsy, risk scores and decisional flowchart. *Pediatr Radiol* 51, 1545–1554. <https://doi.org/10.1007/s00247-021-05034-7>
 28. El-Araby, H. A., Saber, M. A., Radwan, N. M., Taie, D. M., Adawy, N. M., & Sira, A. M. (2021). Temporal histopathological changes in biliary atresia: A perspective for rapid fibrosis progression. *Annals of hepatology*, 21, 100263. <https://doi.org/10.1016/j.aohep.2020.09.007>
 29. Wildhaber B. E. (2012). Biliary atresia: 50 years after the first kasai. *ISRN surgery*, 2012, 132089. <https://doi.org/10.5402/2012/132089>
 30. Chang, C. H., Bryce, C. L., Shneider, B. L., Yabes, J. G., Ren, Y., Zenarosa, G. L., Tomko, H., Donnell, D. M., Squires, R. H., & Roberts, M. S. (2018). Accuracy of the Pediatric End-stage Liver Disease Score in Estimating Pretransplant Mortality Among Pediatric Liver Transplant Candidates. *JAMA pediatrics*, 172(11), 1070–1077. <https://doi.org/10.1001/jamapediatrics.2018.2541>

Clinical Findings in Biliary Atresia A Literature Review

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	link.springer.com Internet Source	1%
2	www.mdpi.com Internet Source	1%
3	Marcello Napolitano, Stéphanie Franchi-Abella, Beatrice Maria Damasio, Thomas Angell Augdal et al. "Practical approach for the diagnosis of biliary atresia on imaging, part 2: magnetic resonance cholecystopancreatography, hepatobiliary scintigraphy, percutaneous cholecysto-cholangiography, endoscopic retrograde cholangiopancreatography, percutaneous liver biopsy, risk scores and decisional flowchart", <i>Pediatric Radiology</i> , 2021 Publication	1%
4	Submitted to Prince of Songkla University Student Paper	1%
5	www.ncbi.nlm.nih.gov Internet Source	1%

6	pubs.rsna.org Internet Source	1 %
7	Juniastuti, Abdul Hadi Furqoni, Mochamad Amin, Yanna Debby Restifan et al. "The evaluation results of proposed antigen rapid diagnostic tests for COVID-19: some possible factors might influence", Infection, 2023 Publication	1 %
8	Mary Ayers, Patrick McKiernan. "Biliary tract malformations of infancy", Paediatrics and Child Health, 2021 Publication	1 %
9	Submitted to The University of Wolverhampton Student Paper	1 %
10	jamanetwork.com Internet Source	1 %
11	www.frontiersin.org Internet Source	1 %
12	www.researchgate.net Internet Source	1 %
13	www.wjgnet.com Internet Source	1 %
14	Masahiro Takeda, Hajime Takei, Mitsuyoshi Suzuki, Takafumi Tsukui et al. "Bile acid profiles in adult patients with biliary atresia"	1 %

who achieve native liver survival after portoenterostomy", Scientific Reports, 2024

Publication

15	psasir.upm.edu.my Internet Source	1 %
16	scholarhub.ui.ac.id Internet Source	1 %
17	www.obgynia.com Internet Source	1 %
18	Ronald J. Sokol. "Pathogenesis and Outcome of Biliary Atresia: Current Concepts :", Journal of Pediatric Gastroenterology & Nutrition, 07/2003 Publication	<1 %
19	patient.info Internet Source	<1 %
20	portlandpress.com Internet Source	<1 %
21	pubmed.ncbi.nlm.nih.gov Internet Source	<1 %
22	pure.rug.nl Internet Source	<1 %
23	www.ejmanager.com Internet Source	<1 %

24

Ali Islek, Gokhan Tumgor. "Biliary atresia and congenital disorders of the extrahepatic bile ducts", World Journal of Gastrointestinal Pharmacology and Therapeutics, 2022

Publication

<1 %

25

Hanaa A El-Araby, Magdy A Saber, Noha M Radwan, Doha M Taie, Nermin M Adawy, Ahmad M Sira. "Temporal histopathological changes in biliary atresia: A perspective for rapid fibrosis progression", Annals of Hepatology, 2020

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Clinical Findings in Biliary Atresia A Literature Review

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
