

A prospective randomized study with a 5-year minimum follow-up evaluation of transanal endoscopic microsurgery versus laparoscopic total mesorectal excision after neoadjuvant therapy

**G. Lezoche · M. Baldarelli · Mario Guerrieri · A. M. Paganini ·
A. De Sanctis · S. Bartolacci · E. Lezoche**

Published online: 20 November 2007
© Springer Science+Business Media, LLC 2007

Erratum to: Surg Endosc
DOI 10.1007/s00464-007-9596-y

The surname of the third author was inadvertently omitted.
The Given Name and Family Name should read Mario
Guerrieri.

Springer regrets the error.

The online version of the original article can be found under
doi: [10.1007/s00464-007-9596-y](https://doi.org/10.1007/s00464-007-9596-y)

G. Lezoche · A. M. Paganini · E. Lezoche (✉)
Department of Surgery “Paride Stefanini,” II Clinica Chirurgica,
University of Rome “La Sapienza”, Viale del Policlinico,
00161 Rome, Italy
e-mail: emanuele.lezoche@uniroma1.it

M. Baldarelli · M. Guerrieri · A. De Sanctis
Department of General Surgery, University “Politecnica delle
Marche”, Ancona, Italy

S. Bartolacci
Department of Epidemiology and Statistics, University
“Politecnica delle Marche”, Ancona, Italy