

MECHANISMS IN ENDOCRINOLOGY

White, brown and pink adipocytes: the extraordinary plasticity of the adipose organ

Antonio Giordano, Arianna Smorlesi, Andrea Frontini, Giorgio Barbatelli and Saverio Cinti¹

Section of Neuroscience and Cell Biology, Department of Experimental and Clinical Medicine, Università Politecnica delle Marche, Ancona, Italy and ¹Center of Obesity, Università Politecnica delle Marche-United Hospitals, Ancona, Italy

Correspondence should be addressed to S Cinti

Email
s.cinti@univpm.it

Abstract

In mammals, adipocytes are lipid-laden cells making up the parenchyma of the multi-depot adipose organ. White adipocytes store lipids for release as free fatty acids during fasting periods; brown adipocytes burn glucose and lipids to maintain thermal homeostasis. A third type of adipocyte, the pink adipocyte, has recently been characterised in mouse subcutaneous fat depots during pregnancy and lactation. Pink adipocytes are mammary gland alveolar epithelial cells whose role is to produce and secrete milk. Emerging evidence suggests that they derive from the transdifferentiation of subcutaneous white adipocytes. The functional response of the adipose organ to a range of metabolic and environmental challenges highlights its extraordinary plasticity. Cold exposure induces an increase in the 'brown' component of the organ to meet the increased thermal demand; in states of positive energy balance, the 'white' component expands to store excess nutrients; finally, the 'pink' component develops in subcutaneous depots during pregnancy to ensure litter feeding. At the cell level, plasticity is provided not only by stem cell proliferation and differentiation but also, distinctively, by direct transdifferentiation of fully differentiated adipocytes by the stimuli that induce genetic expression reprogramming and through a change in phenotype and, consequently function. A greater understanding of adipocyte transdifferentiation mechanisms would have the potential to shed light on their biology as well as inspire novel therapeutic strategies against metabolic syndrome (browning) and breast cancer (pinkening).

*European Journal of
Endocrinology*
(2014) **170**, R159–R171

The adipose organ is made up of white and brown adipocytes...

In experimental animals, gross anatomy demonstrates that the adipose organ has a multi-depot organisation (1, 2, 3), consisting of two large subcutaneous depots and of numerous visceral depots (intended here as fat in close

apposition to viscera, regardless of their portal or caval venous drainage) located in the visceral cavities of the trunk (Fig. 1). Dissection studies show that the depots have a fairly consistent shape throughout life and that similar shapes are found in different mouse strains. The subcutaneous depots are located in the upper portion of the

Invited Author's profile

Prof. Saverio Cinti is a specialist in internal medicine and anatomic pathology. He has been Professor of Anatomy, School of Medicine, Ancona University, Italy since 1984. Having started work in the field of adipose tissues since 1980 in the lab of Prof. Björntorp (Gothenburg University), he developed the concept of the adipose organ focusing on the plastic properties of adipose tissues and outlined the importance of this plasticity for the future treatment of obesity and related disorders. He described the CLS as the cause of low-grade chronic inflammation in the obese adipose organ. He has authored more than 250 peer reviewed articles. In 2008, he received the Blaise Pascal medal of the European Academy of Science for biology and in 2013 the Wasserman Prize for senior scientist of the European Society of Obesity.






Figure 1

Gross anatomy of the adipose organ of adult female 129Sv mice. The subcutaneous and visceral depots were dissected and positioned on a mouse template to show their respective location in the body. The mouse on the left was maintained at temperatures close to thermoneutrality (28 °C for 10 days), whereas the mouse on the right was acclimated to cold (6 °C for 10 days). Browning of the adipose organ is visually evident in the cold-acclimated mouse. The adipose organ is made up of two subcutaneous depots: (A) anterior (deep cervical, superficial cervical, interscapular, subscapular, axillo-thoracic) and (F) posterior (dorso-lumbar, inguinal, gluteal), and of several visceral depots: (B) mediastinal, (C) mesenteric, (D) retro-peritoneal and (E) abdomino-pelvic (perirenal, periovarian, parametrial, perivescical). Scale bar: 1 cm. Reproduced with permission. Murano I, Zingaretti CM & Cinti S. The adipose organ of Sv129 mice contains a prevalence of brown adipocytes and shows plasticity after cold exposure. *Adipocytes* 2005 1 121–130.

thorax close to the forelimbs and the neck (anterior) and in the lower part of the abdomen, close to the hind legs (posterior). In animals kept at an environmental temperature close to thermoneutrality (about 28 °C), the adipose organ is predominantly white, with a few brownish areas in the anterior subcutaneous depot at the interscapular, subscapular, axillary, perisusclavian and pericarotid levels; brownish areas in the visceral depots are found mainly around the heart, the aorta and its main branches (mediastinal and perirenal sites). By light microscopy the adipose organ contains two different cell types: brown and white adipocytes (4, 5, 6). ‘Brownish’ areas are those where brown adipocytes are the predominant parenchymal cell type: they correspond to brown adipose tissue (BAT), which is richly innervated and vascularised. ‘White’ areas, where white adipocytes are the predominant cell type, are white adipose tissue (WAT) and show fewer nerves and a lower number of blood vessels. Depot colour is determined by the

relative amount of the two cell types and the degree of vascularisation (7). The relative amount of BAT and WAT in the adipose organ is variable, depending on several factors of which age, diet and environmental temperature are the most important. Quantitative histological studies have shown that the vast majority of depots have a mixed composition also in warm-acclimated mice (see below).


White and brown adipocytes display distinctive features by transmission electron microscopy (1, 2, 3). White adipocytes contain a single large lipid droplet occupying about 90% of the cell volume. The nucleus is squeezed to the cell periphery and the cytoplasm forms a very thin rim. The organelles are poorly developed; in particular mitochondria are small, elongated and have short, randomly organised cristae (Fig. 2). Because of these ultrastructural characteristics, these cells are also called unilocular adipocytes.

Brown adipocytes are smaller than white adipocytes, their cytoplasm contains several lipid droplets, a roundish nucleus and numerous, large, generally spherical mitochondria with lamellar cristae (Fig. 3). These mitochondria contain a unique protein, uncoupling protein 1 (UCP1), that supports the thermogenic function of brown adipocytes (8, 9). These cells are also called multilocular adipocytes.

The different morphology of white and brown adipocytes underpins their different functional roles. The white adipocyte stores energy (in the form of lipids) that is released between meals: its lipid droplet is spherical because this is the geometrical shape maximising volume and minimising space occupation. The brown adipocyte burns lipids to produce heat: its multilocularity maximises the cytoplasmic–lipid interface, making large amounts of fatty acids available quickly for mitochondrial uncoupling and consequently thermogenesis (8).

Notably, WAT and BAT do not exhibit distinct anatomical boundaries, but rather are found as a seamless continuum in all depots at both the macroscopic and microscopic levels. Indeed, in the areas between WAT and BAT, we described adipocytes with an intermediate morphology between white and brown adipocytes and designated them as paucilocular adipocytes (Fig. 4). Recent data from our laboratory have suggested that paucilocular adipocytes, which are found in all adipose depots, are the population showing the greatest proneness to transdifferentiate into brown adipocytes upon environmental or pharmacological stimulation (see below).

In conclusion, both white and brown adipocytes harbour a large amount of lipids in their cytoplasm. However, they serve two almost opposite functions that are essential for survival, to store and to dissipate energy respectively.

**Figure 2**

Transmission electron microscopy of an epididymal WAT white adipocyte from a 3-week-old rat. N, nucleus; m, mitochondria; L, lipid droplet. Scale bar: 3 μ m. Reproduced with permission. Reproduced with permission. Cinti S. *The Adipose Organ*. Milan: Kurtis, 1999.


...but 'pink' adipocytes arise during pregnancy

During pregnancy and lactation the anterior and posterior subcutaneous depots of the female adipose organ turn into an organ whose function is to produce and secrete milk: the mammary gland (1, 10) (Fig. 5). Such transformation involves mainly the parenchyma through the development of milk-secreting lobulo-alveolar glandular structures. This dynamic process is generally referred to as alveologensis.


Epithelial ducts ending in symmetrical nipples (three in anterior and two in posterior subcutaneous fat) develop before puberty and branch through the subcutaneous depot parenchyma, where in normal conditions white and brown adipocytes account for about 90% of depot volume. In terms of anatomical organisation, the mouse mammary glands are five symmetrical structures, each ending with a nipple; in terms of adipose organ organisation, the subcutaneous depot contains two glands, one anterior and one posterior (corresponding to the anterior and posterior

subcutaneous depots respectively), each provided with symmetrical nipples (three anterior and two posterior). During pregnancy, which in mice lasts 21 days, alveoli gradually replace adipose tissue. Well-developed alveoli formed by epithelial cells devoid of lipid droplets, likely deriving from the proliferation of ductal stem cells, are visible already 12–15 days from conception (11, 12). Alveologensis continues with the appearance of lipid-laden epithelial cells and culminates on days 18–21. Our data suggest that these cells derive from transdifferentiation of subcutaneous white adipocytes (13). We have proposed the name of 'pink adipocytes' for these adipocyte-derived milk-producing cells, because: i) they meet the definition of adipocyte, i.e. a parenchymal cell capable of storing large amounts of lipids; ii) they arise exclusively in female subcutaneous depots during pregnancy and lactation; and iii) the pregnant mammary gland is pink at the macroscopic level.

Thus, the adipose organ parenchyma contains three cell types characterised by a discrete morphology and function: i) the white adipocyte stores and secretes lipids; ii) the brown adipocyte produces heat; and iii) the pink adipocyte produces milk (Fig. 6). Each function is critical for individual and species survival. Interestingly, despite

**Figure 3**

Transmission electron microscopy of an interscapular BAT brown adipocyte from a 10-day-old rat. Note the numerous, large and generally spherical mitochondria with lamellar cristae (m). N, nucleus; L, lipid droplet; CAP, capillary. Scale bar: 3 μ m. Reproduced with permission. Cinti S. *The Adipose Organ*. Milan: Kurtis, 1999.

**Figure 4**

Paucilocular adipocytes. (A) Transmission electron microscopy of subcutaneous fat of a cold acclimated adult mouse (6 °C for 5 days) showing a paucilocular adipocyte with an intermediate morphology between white and brown adipocytes. Note the predominant large central lipid droplet (L) and several small cytoplasmic lipid droplets (l). Mitochondria are numerous and exhibit an intermediate morphology between those typical of white and brown adipocytes (inset: enlargement of squared area in (A)). Scale bar: A = 5 μ m and inset = 0.5 μ m. (B) UCP1-immunoreactive paucilocular adipocyte found in omental fat from a patient suffering for pheochromocytoma. Note the morphology corresponding to that described in (A). Surrounding (upper and right) white adipocytes are unilocular and UCP1 negative. Scale bar: 10 μ m.

their different morphology and physiology, the three cell types nonetheless share the expression of some genes. For example, most of the genes related to lipid metabolism are expressed in all three cells: leptin is expressed in both white (14, 15) and pink adipocytes (16), S-100b is expressed in white (17) and pink adipocytes (18) and perilipin A is expressed in white and brown adipocytes (19). From a physiological point of view, all three types of adipose cells have endocrine properties. White adipocytes secrete a number of adipokines that affect eating behaviour (leptin) (20) and metabolism (e.g. adiponectin, resistin, adipsin) (21). Brown adipocytes also secrete hormones and growth factors (e.g. betatrophin and FGF21) (22, 23, 24). Pink adipocytes, besides milk

**Figure 5**

Gross anatomy of the adipose organ of a lactating female mouse. Both anterior and posterior subcutaneous depots are transformed into mammary glands. Scale bar: 1.5 cm. Reproduced with permission. Cinti S. The Adipose Organ. Milan: Kurtis, 1999.




Figure 6

Scheme of the three adipocytes forming the parenchyma of the adipose organ.

components, also secrete leptin, which seems to have an important role in preventing obesity in pups (25, 26).

Plasticity of the adipose organ

The anatomical, cytological and physiological aspects reviewed above prompt the question of why three different cell types, each playing discrete physiological roles, should all be found in the same organ, the adipose organ. Similarities in morphology and gene expression can to some extent explain their coexistence. Yet distinctive phenotypic aspects underlie striking morphological and functional differences. For instance, some important genes appear to be specific and crucial for the function exerted by each cell type: *UCP1* underpins thermogenesis and is unique to brown adipocytes (8, 9), leptin is not found in classic multilocular brown adipocytes (14) and perilipin B is found in pink adipocytes along with a number of epithelial and milk-related genes (27, 28).

White–brown plasticity

Over the last three decades our and other laboratories have collected a large body of evidence documenting that fully differentiated adipocytes have the outstanding physiological ability to transdifferentiate. In particular, mature adipocytes undergo genome reprogramming and turn into a different cell type, serving different physiological roles (15, 29, 30, 31, 32, 33, 34, 35); crucially, the process is reversible. White-to-brown transdifferentiation is essential to meet increased heat production requirements during

chronic cold exposure. Cold exposure activates BAT by acting on the sympathetic nervous fibres that directly innervate brown adipocytes at the parenchymal level (36, 37, 38, 39); chronic cold exposure results in branching of noradrenergic parenchymal fibres, significantly increasing BAT sympathetic innervation, a phenomenon that appears to be closely related to white-to-brown transdifferentiation (40, 41). β 3-adrenoceptors (AR) are specifically expressed by brown adipocytes. When activated by noradrenaline they drive brown adipocyte thermogenic activation, but are also likely responsible for white-to-brown transdifferentiation. ‘Browning’, i.e. an increase in the brown component of the organ, is detectable even at the macroscopic level in the adipose organ of a mouse kept at 6 °C compared with one acclimated to 28 °C (Fig. 1). The tissue remodelling is partly due to recruitment of precursor cells, especially in interscapular and inguinal subcutaneous depots (42), and partly to direct conversion of a subpopulation of unilocular/paucilocular adipocytes (6, 31, 32, 33, 35, 43). The two processes, which most likely coexist, are driven by the same physiological stimulus through β -AR activation. Cold-exposed mice lacking β 3-AR do not undergo browning (32, 43), but precursor development, probably driven by β 1-AR, is not hampered in these animals, because pre-adipocyte development has been documented after administration of β 1-AR agonists and lack of development has been demonstrated after administration of β 3-AR agonists (43). As also suggested by *in vitro* findings (44), β 3-AR could thus be responsible for white-to-brown transdifferentiation and β 1-AR for precursor proliferation and differentiation.

Browning is of remarkable pathophysiological interest, because it could be harnessed to tackle obesity and metabolic syndrome (45, 46, 47, 48). Indeed, ectopic *UCP1* expression (49) and expression in white adipocytes of key molecules involved in brown adipocyte differentiation, such as *Prdm16* (50), induce obesity resistance and ameliorate insulin sensitivity. Obesity-prone mouse strains have less BAT than obesity-resistant strains (3, 4, 51); mice lacking brown fat (52) and β 3-AR (34) are prone to diet-induced obesity; and specific β 3-AR agonists curb obesity in obese rats (53, 54, 55).

Warm exposure, ageing and obesity lead to ‘whitening’, which involves a significant reduction in the density of parenchymal noradrenergic nerve fibres in the adipose organ (1, 2, 3). Different gene expression profiles are found in brown adipocytes from different depots (56) and even in *UCP1*-expressing adipocytes (57). Furthermore, a different gene expression has been found during development in interscapular

(anterior subcutaneous depot) and perirenal (visceral depot) brown adipocytes as compared with those found predominantly in white depots, such as the posterior subcutaneous depot (58). These findings have prompted terms such as 'brite' and 'beige' for a population of UCP1-containing multilocular adipocytes (57, 59). However, we feel that the notion of 'brown' should be maintained until different functions are demonstrated for these cells. The different gene expression profiles described in interscapular brown adipocytes and in other depots could merely depend on differences in the extracellular hormonal milieu, the degree and type of innervation, and/or the cell development stage. As mentioned earlier, paucilocular adipocytes, i.e. adipocytes with an intermediate morphology, are found in the areas between BAT and WAT. We documented a varied morphology and degree of UCP1 expression in these adipocytes (43). Some were UCP1-negative, others were UCP1-positive; accordingly, mitochondrial morphology spanned from that typical of white adipocytes to that typical of brown adipocytes, thus accounting for the variable immunoreactivity of these cells for the brown marker UCP1. It is conceivable that UCP1 immunoreactivity is acquired only upon achievement of a given degree of mitochondrial differentiation. The distinctive gene expression profile of brite/beige adipocytes may reflect the density of multilocular adipocytes and their level of differentiation. A true, functional difference between interscapular brown adipocytes and those found in other areas of the organ has not yet been documented *in vivo* and at single cell level.

A recent paper seems to confirm the direct, reversible transdifferentiation of white adipocytes into brown adipocytes (35). Cold-induced UCP1-expressing adipocytes from posterior subcutaneous fat turned into unilocular adipocytes, expressing genuine white phenotype genes when the animals were exposed to a warm environment, and re-exposure to cold involved a return to the multilocular UCP1 phenotype. Not all white adipocytes seem to have the ability to turn into brown adipocytes, and some white adipocytes might never be able to undergo the phenotype change, possibly because of their distance from noradrenergic fibres. In this connection, we found a positive correlation between the density of brown adipocytes and the density of noradrenergic fibres in the adipose organ of two different mouse strains (5).


The plasticity of the adipose organ raises the question of the origin of adipocytes. Despite contrasting findings from different laboratories (60) our data, including lineage-tracing studies, seem to indicate that endothelial cells

of adipose tissue capillaries are able to turn into pericytes and then into either white and brown adipocytes (61, 62, 63, 64, 65, 66). Such unique origin could support their ability to undergo multiple changes before achievement of the adult phenotype.

White-pink plasticity

During pregnancy and lactation, all subcutaneous depots of the adipose organ turn into mammary glands (1, 10). Our morphological studies of the transforming subcutaneous depots seem to suggest that mammary gland alveoli develop in two stages through two different mechanisms. In the first stage of pregnancy, alveoli are constituted of epithelial cells lacking cytoplasmic lipid droplets that could derive from stem cell proliferation (Fig. 7), but in the second stage of pregnancy they are constituted of lipid-laden epithelial cells while subcutaneous fat shrinks progressively (Fig. 8). Ultrastructural data support the astonishing possibility that in the second part of pregnancy subcutaneous adipocytes progressively acquire epithelial-like features, likely under hormonal stimuli, aggregating with similarly committed pink adipocytes and with myoepithelial cells to form adipose tissue-derived milk-secreting alveoli (13) (Fig. 9). To document the striking transdifferentiation of adipocytes into milk-producing glands and establish whether the opposite process occurs during mammary gland involution, we carried out lineage-tracing studies using aP2-cre/R26R and WAP-cre/R26R double transgenic mice respectively. In aP2-cre/R26R mice, about 70% of alveolar epithelial cells expressed the reporter gene in late pregnancy, whereas in WAP-cre/R26R mice most of the adipocytes found in subcutaneous fat post-lactation were positive for the reporter gene. Notably, about 30% of alveolar epithelial cells never stained for the gene during pregnancy, in line with the origin of a large number of alveolar cells from the well-characterised ductal alveolar progenitor cells (10, 11, 12). The transdifferentiation of white adipocytes into pink cells was also confirmed by explants experiments, where both adipose tissue and isolated adipocytes from *Rosa26* (*Gt(ROSA)26Sor*) mice implanted in pregnant WT female mice gave rise to donor-derived glands (68).

Reversible white-to-pink transdifferentiation could shed light on breast cancer biology, as suggested by recent data showing that loss of *PPAR* γ expression by mammary secretory epithelial cells creates a pro-breast tumourigenic environment (69). Notably, *PPAR* γ seems to be a key factor for pink-to-white transdifferentiation *in vitro* (70).

**Figure 7**


Haematoxylin–eosin staining of the subcutaneous adipose depot (mammary gland) from a female mouse at day 10 of pregnancy. In the first stage of pregnancy, the alveoli (asterisk) are constituted of epithelial cells lacking cytoplasmic lipid droplets. D, duct; V, blood vessel. Scale bar: 60 μ m.

The obese adipose organ

In 2003, two independent groups in the United States showed that the adipose organ of obese animals and humans is infiltrated by macrophages; the infiltration was found to relate to adipocyte size and to the development of insulin resistance (71, 72). The majority of cytokines with key roles in inducing insulin resistance are expressed by the stroma-vascular fraction of fat (including macrophages) and a minority by the floating fraction formed by mature adipocytes, reflecting the importance of macrophage infiltration in the development of insulin resistance and subsequently type 2 diabetes. Our group found that most of the macrophages infiltrating obese fat are arranged around dead adipocyte remnants into distinctive figures that we denominated crown-like-structures (CLS) (73). In a subsequent paper, we described ultrastructural abnormalities (such as calcium build-up and cholesterol crystals), signs of oxidative stress and *NLRP3* inflammasome activation with formation of active caspase 1 in hypertrophic adipocytes from obese mice and suggested that these cells die of pyroptosis, a proinflammatory programmed cell death (74).

Resorption of dead adipocyte remnants, especially the large lipid droplets, is an extended process characterised by a chronic low-grade inflammation similar to that seen in foreign body reactions. Accordingly, CLS may also

contain syncytial giant macrophages. To assess whether CLS originate from adipocyte debris, we used Philip Scherer's transgenic model, where white adipocyte apoptotic death is specifically induced by administration of a dimeriser that activates caspase 8 (75). All dead adipocytes gave rise to CLS, in line with our hypothesis (76). The time course of fat histopathology in this model disclosed that CLS form after adipocyte death, demonstrating that this event may be sufficient to recruit macrophages and induce CLS formation (76). Altogether, we think that the death and degeneration of hypertrophic adipocytes, with the consequent exposure to the extracellular milieu of nuclear and cytoplasmic (mainly, lipid droplets) determinants that are normally segregated into the cell, represent the primary events triggering the inflammatory and immune reactions in the obese adipose tissue. This view has been recently reinforced by data from Xu *et al.* (77), showing the importance of lipid catabolism in the macrophages infiltrating the obese adipose tissue. Interestingly, this

**Figure 8**

Subcutaneous adipose depot (mammary gland) from a female mouse at day 17 of pregnancy. Haematoxylin–eosin staining (A and B) shows the appearance of alveoli constituted of lipid-laden epithelial cells (pink adipocytes). B is the enlargement of the area framed in A: most of the alveolar cells are pink adipocytes with large cytoplasmic lipid droplets. Pink adipocytes of well-developed alveoli (with central lumen: L) and of early alveoli (without central lumen: *) show strongly immunoreactive nuclei for the transcription factor *Elf5* (C), a master regulator of alveologenesis (67), and are immunoreactive for the milk whey acidic protein (WAP) (D). CAP, capillary; WAT, white adipose tissue. Scale bar: A = 50 μ m; B = 15 μ m; C and D = 30 μ m.




Figure 9

Transmission electron microscopy of pink adipocytes. (A) Pink adipocytes contain large cytoplasmic lipid droplets (L). (B) Enlargement of the pink adipocyte marked by * in A. Abundant stacked rough endoplasmic reticulum (RER) and a distinct basal membrane (BM) are visible. (C) Enlargement of the framed area in (A). Milk granules (G) and microvilli projecting in the early lumen are visible. Scale bars: 2 μm .

view may also predict that autoimmune reactions could develop in obesity, and be possibly involved in some physiopathological aspects of the metabolic syndrome.

Hormone-sensitive lipase knockout mice created by Grant Michell are lean but their fat is characterised by hypertrophic adipocytes (78). These animals exhibited the same CLS density as obese animals (73). Notably, we found a positive correlation between CLS density and adipocyte size both in subcutaneous and visceral fat depots; their density was lower in subcutaneous fat containing larger adipocytes (79). Collectively, these data suggest that visceral adipocytes have a smaller death critical size

(size-triggering death) (41), in line with the well-known greater morbidity due to accumulation of visceral fat (80). Interestingly, we failed to detect CLS in either mice or humans with hyperplastic obesity, which is characterised by small adipocytes and the absence of secondary metabolic disorders (73, 81). The positive correlation between adipocyte size and insulin resistance has recently been confirmed in non-obese humans (82).

The plasticity of the adipose organ could be the basis for future treatment, or prevention, of obesity and type 2 diabetes. As mentioned above, white-to-brown transdifferentiation involves a reduction in adipocyte size and an increase in their mitochondrial content. Thus, 'mild' white-to-brown transdifferentiation could make white adipocytes less prone to death and turns the adipose organ parenchyma into a 'healthier' tissue.

The human adipose organ

As in experimental animals, also in humans the adipose organ is made up of subcutaneous and visceral depots (83, 84, 85, 86, 87). Whereas in rodents, dermal and subcutaneous adipose tissues are separated by a layer of skeletal muscle cells, in humans they are continuous with one another; moreover, subcutaneous adipose tissue is not confined to some areas, but forms an uninterrupted layer throughout the body with the exception of hands and feet. Importantly, mammary and gluteo-femoral subcutaneous adipose tissues are more developed in females than in males. The distribution of the visceral depots is very similar to that described in rodents, but the omental depot is particularly well developed in humans. In lean adults, the human adipose organ accounts for 8–18% of body weight in males and for 14–28% in females (~5% in monkeys) (88, 89).

Independent of total body fat, body fat distribution is a well-known important risk factor for obesity-associated diseases, with visceral obesity displaying greater morbidity (80). It should be noted, however, that the vast majority of free fatty acids, that are especially involved in the cardiovascular complications of the metabolic syndrome (90), are released by the upper body subcutaneous fat depots (91, 92).

As in small rodents, the human adipose organ contains brown adipocytes organised into typical BAT. The lower surface:volume ratio, hence thermal dispersion, of the human compared with the rodent's body involves lower heat production demands, at least in adults. On the other hand, newborns are characterised by a greater surface:volume ratio and a considerable amount of BAT

(93, 94). Human BAT thus seems to undergo an age-related morphofunctional involution (likely, brown-to-white transdifferentiation). The histological and electron microscopic features of human adipose tissue are identical to those of their murine counterpart (95). In particular, UCP1-positive brown adipocytes are found among white adipocytes (96, 97, 98). In human newborns, BAT is found in almost all the areas described in rodents, and *UCP1* gene expression has been documented in visceral adipose tissue of lean and obese adult patients. The brown:white adipocyte ratio in the visceral adipose tissue of lean adult humans has been put at 1/100–200 (99).

As in experimental animals, also in humans the adipose organ displays outstanding plasticity. An increased amount of BAT has been described in outdoor workers in northern Europe (100) and in patients with pheochromocytoma, a tumour derived from the cells of the adrenal medullary and characterised by catecholamine secretion (101, 102). Furthermore hibernoma, a rare BAT tumour, has been described in several anatomical areas, including subcutaneous and visceral fat (103).

Brown adipocytes have a strong oxidative metabolism and incorporate high levels of fluorodeoxyglucose, the tracer used in positron emission tomography (PET); this has enabled unexpectedly large amounts of BAT to be detected in adult humans (83, 84, 85, 86, 104, 105). In normal adults, BAT depots are found at the base of the neck, the root of the upper limbs, and the intercostal spaces (106). We found UCP1-positive brown adipocytes in perithyroid fat from adult biopsies. These specimens also contained parenchymal noradrenergic fibres in direct contact with brown adipocytes (Fig. 10) (87, 107), documenting a similar parenchymal innervation of the adipose organ in humans and small mammals. Electron microscopy has disclosed preadipocytes in close proximity to capillary walls in human BAT (87); interestingly, the density of preadipocytes was about five times higher in a case of hibernoma than in normal BAT (108).

Human adipose organ plasticity and therapeutic prospects

As the physiological role of BAT in adult humans continues to be explored, the possibility to expand this energy-dissipating tissue through pharmacological interventions is being hailed as a possible approach to treat obesity and related disorders (109). Functional BAT has clearly been demonstrated in adult humans and proved to contribute to the overall energy balance (83, 84, 85, 86, 110). Cold exposure can recruit BAT to produce




Figure 10

Double immunostaining and confocal microscopy of human BAT. The sympathetic nerve fibres (detected by tyrosine hydroxylase immunofluorescence: green) are closely apposed to the surface of *UCP1*-immunoreactive (red) brown adipocytes. Nuclei are stained by TOTO3 (blue). Scale bar: 10 μ m.

non-shivering thermogenesis (111). Moreover, poor BAT activity correlates with ageing, BMI and measures of metabolic disease (112, 113, 114).

It is interesting to note that humans with a reduced brown phenotype of abdominal subcutaneous adipose tissue have reduced insulin sensitivity (115), and human white adipocyte precursors can be induced *in vitro* to express UCP1 through administration of drugs (116, 117). We recently reported that a white visceral depot (the omentum), which normally contains only unilocular UCP1-negative (white) adipocytes, showed multilocular UCP1-positive (brown) adipocytes in six of 12 patients with pheochromocytoma (118). In these six patients, we also detected several UCP1-positive paucilocular adipocytes, the intermediate phenotype preceding white-to-brown transdifferentiation. In such adipocytes, electron microscopy documented mitochondria with an intermediate morphology between typical white and typical brown. Similar features have been detected in transdifferentiating human brown adipocytes *in vitro* (119). Taken together, these data suggest that white-to-brown transdifferentiation also occurs in humans and might be harnessed for therapeutic purposes. The master

molecular pathways could be the noradrenergic stimulation via the β 3-AR signalling, despite the unsuccessful clinical trials performed with β 3-AR agonists (120, 121) before PET, and bioptic studies renewed the interest in the topic in 2009. A range of new molecular mechanisms to induce browning have recently been proposed (reviewed in (107)). Among these, secreted factors such as ANP (NPPA), BMP8B, irisin and FGF21 that affect brown adipocyte activation and recruitment seem to be particularly promising for the development of new anti-obesity drugs in the near future (22, 122, 123, 124, 125).

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

This work was supported by DIABAT Collaborative Project of the European Community's FP7, grant agreement number HEALTH-F2-2011-278373 to S Cinti.

References

- Cinti S. *The Adipose Organ*. Milan: Kurtis, 1999.
- Cinti S. The adipose organ. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* 2005 **73** 9–15. (doi:10.1016/j.plefa.2005.04.010)
- Frontini A & Cinti S. Distribution and development of brown adipocytes in the murine and human adipose organ. *Cell Metabolism* 2010 **11** 253–256. (doi:10.1016/j.cmet.2010.03.004)
- Murano I, Zingaretti CM & Cinti S. The adipose organ of Sv129 mice contains a prevalence of brown adipocytes and shows plasticity after cold exposure. *Adipocytes* 2005 **1** 121–130.
- Murano I, Barbatelli G, Giordano A & Cinti S. Noradrenergic parenchymal nerve fiber branching after cold acclimatisation correlates with brown adipocyte density in mouse adipose organ. *Journal of Anatomy* 2009 **214** 171–178. (doi:10.1111/j.1469-7580.2008.01001.x)
- Vitali A, Murano I, Zingaretti MC, Frontini A, Ricquier D & Cinti S. The adipose organ of obesity-prone C57BL/6J mice is composed of mixed white and brown adipocytes. *Journal of Lipid Research* 2012 **53** 619–629. (doi:10.1194/jlr.M018846)
- Nechad M. Structure and development of brown adipose tissue. In *Brown Adipose Tissue*. Eds P Trayhurn & D Nicholls. London: Edward Arnold, 1986.
- Cannon B & Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiological Reviews* 2004 **84** 277–359. (doi:10.1152/physrev.00015.2003)
- Ricquier D. Respiration uncoupling and metabolism in the control of energy expenditure. *Proceedings of the Nutrition Society* 2005 **64** 47–52. (doi:10.1079/PNS2004408)
- Richert MM, Schwertfeger KL, Ryder JW & Anderson SM. An atlas of mouse mammary gland development. *Journal of Mammary Gland Biology and Neoplasia* 2000 **5** 227–241. (doi:10.1023/A:1026499523505)
- Ercan C, van Diest PJ & Vooijs M. Mammary development and breast cancer: the role of stem cells. *Current Molecular Medicine* 2011 **11** 270–285. (doi:10.2174/156652411795678007)
- Bussard KM & Smith GH. The mammary gland microenvironment directs progenitor cell fate *in vivo*. *International Journal of Cell Biology* 2011 **2011** 451676. (doi:10.1155/2011/451676)
- Morrioni M, Giordano A, Zingaretti MC, Boiani R, De Matteis R, Kahn BB, Nisoli E, Tonello C, Pisoschi C, Luchetti MM *et al*. Reversible transdifferentiation of secretory epithelial cells into adipocytes in the mammary gland. *PNAS* 2004 **101** 16801–16806. (doi:10.1073/pnas.0407647101)
- Cinti S, Frederich RC, Zingaretti MC, De Matteis R, Flier JS & Lowell BB. Immunohistochemical localization of leptin and uncoupling protein in white and brown adipose tissue. *Endocrinology* 1997 **138** 797–804. (doi:10.1210/endo.138.2.4908)
- Cancello R, Zingaretti MC, Sarzani R, Ricquier D & Cinti S. Leptin and UCP1 genes are reciprocally regulated in brown adipose tissue. *Endocrinology* 1998 **139** 4747–4750. (doi:10.1210/endo.139.11.6434)
- Smith-Kirwin SM, O'Connor DM, de Johnston J, Lancey ED, Hassink SG & Funanage VL. Leptin expression in human mammary epithelial cells and breast milk. *Journal of Clinical Endocrinology and Metabolism* 1998 **83** 1810–1813. (doi:10.1210/jcem.83.5.4952)
- Cinti S, Cigolini M, Morrioni M & Zingaretti MC. S-100 protein in white preadipocytes: an immunoelectronmicroscopic study. *Anatomical Record* 1989 **224** 466–472. (doi:10.1002/ar.1092240403)
- Barracrough R & Rudland PS. The S-100-related calcium-binding protein, p9Ka, and metastasis in rodent and human mammary cells. *European Journal of Cancer* 1994 **30A** 1570–1576. (doi:10.1016/0959-8049(94)00320-5)
- Blanchette-Mackie EJ, Dwyer NK, Barber T, Coxey RA, Takeda T, Rondinone CM, Theodorakis JL, Greenberg AS & Londos C. Perilipin is located on the surface layer of intracellular lipid droplets in adipocytes. *Journal of Lipid Research* 1995 **36** 1211–1226.
- Zhang Y, Proenca R, Maffei M, Barone M, Leopold L & Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature* 1994 **372** 425–432. (doi:10.1038/372425a0)
- Trayhurn P. Hypoxia and adipose tissue function and dysfunction in obesity. *Physiological Reviews* 2013 **93** 1–21. (doi:10.1152/physrev.00017.2012)
- Fisher FM, Kleiner S, Douris N, Fox EC, Mepani RJ, Verdeguer F, Wu J, Kharitonov A, Flier JS, Maratos-Flier E *et al*. FGF21 regulates PGC-1 α and browning of white adipose tissues in adaptive thermogenesis. *Genes and Development* 2012 **26** 271–281. (doi:10.1101/gad.177857.111)
- Yi P, Park JS & Melton DA. Betatrophin: a hormone that controls pancreatic β cell proliferation. *Cell* 2013 **153** 747–758. (doi:10.1016/j.cell.2013.04.008)
- Villarroya J, Cereijo R & Fillarroya F. An endocrine role for brown adipose tissue? *American Journal of Physiology. Endocrinology and Metabolism* 2013 **305** E567–E572. (doi:10.1152/ajpendo.00250.2013)
- Oliver P, Picò C, De Matteis R, Cinti S & Palou A. Perinatal expression of leptin in rat stomach. *Developmental Dynamics* 2002 **223** 148–154. (doi:10.1002/dvdy.1233)
- Palou A, Sánchez J & Picò C. Nutrient-gene interactions in early life programming: leptin in breast milk prevents obesity later in life. *Advances in Experimental Medicine and Biology* 2009 **646** 95–104. (doi:10.1007/978-1-4020-9173-5_10)
- Brasaemle DL, Barber T, Wolins NE, Serrero G, Blanchette-Mackie EJ & Londos C. Adipose differentiation-related protein is a ubiquitously expressed lipid storage droplet-associated protein. *Journal of Lipid Research* 1997 **38** 2249–2263.
- Russell TD, Palmer CA, Orlicky DJ, Fischer A, Rudolph MC, Neville MC & McManaman JL. Cytoplasmic lipid droplet accumulation in developing mammary epithelial cells: roles of adipophilin and lipid metabolism. *Journal of Lipid Research* 2007 **48** 1463–1475. (doi:10.1194/jlr.M600474-JLR200)
- Barbatelli G, Morrioni M, Vinesi P, Cinti S & Michetti F. S-100 protein in rat brown adipose tissue under different functional conditions: a morphological, immunocytochemical, and immunochemical study. *Experimental Cell Research* 1993 **208** 226–231. (doi:10.1006/excr.1993.1241)
- Cousin B, Bascands-Viguerie N, Kassis N, Nibbelink M, Ambid L, Casteilla L & Pénicaud L. Cellular changes during cold acclimatation in

- adipose tissues. *Journal of Cellular Physiology* 1996 **167** 285–289. (doi:10.1002/(SICI)1097-4652(199605)167:2<285::AID-JCP12>3.0.CO;2-7)
- 31 Himms-Hagen J, Melnyk A, Zingaretti MC, Ceresi E, Barbatelli G & Cinti S. Multilocular fat cells in WAT of CL-316243-treated rats derive directly from white adipocytes. *American Journal of Physiology. Cell Physiology* 2000 **279** C670–C681.
 - 32 Jimenez M, Barbatelli G, Allevi R, Cinti S, Seydoux J, Giacobino JP, Muzzin P & Preitner F. β 3-adrenoceptor knockout in C57BL/6J mice depresses the occurrence of brown adipocytes in white fat. *European Journal of Biochemistry* 2003 **270** 699–705. (doi:10.1046/j.1432-1033.2003.03422.x)
 - 33 Granneman JG, Li P, Zhu Z & Lu Y. Metabolic and cellular plasticity in white adipose tissue I: effects of β 3-adrenergic receptor activation. *American Journal of Physiology. Endocrinology and Metabolism* 2005 **289** E608–E616. (doi:10.1152/ajpendo.00009.2005)
 - 34 Bachman ES, Dhillon H, Zhang CY, Cinti S, Bianco AC, Kobilka BK & Lowell BB. BAR signaling required for diet-induced thermogenesis and obesity resistance. *Science* 2002 **297** 843–845. (doi:10.1126/science.1073160)
 - 35 Rosenwald M, Perdikari A, Rulicke T & Wolfrum C. Bi-directional interconversion of brite and white adipocytes. *Nature Cell Biology* 2013 **15** 659–667. (doi:10.1038/ncb2740)
 - 36 Giordano A, Morroni M, Santone G, Marchesi GF & Cinti S. Tyrosine hydroxylase, neuropeptide Y, substance P, calcitonin gene-related peptide and vasoactive intestinal peptide in nerves of rat periovarian adipose tissue: an immunohistochemical and ultrastructural investigation. *Journal of Neurocytology* 1996 **25** 125–136. (doi:10.1007/BF02284791)
 - 37 Giordano A, Morroni M, Carle F, Gesuita R, Marchesi GF & Cinti S. Sensory nerves affect the recruitment and differentiation of rat periovarian brown adipocytes during cold acclimation. *Journal of Cell Science* 1998 **111** 2587–2594.
 - 38 Giordano A, Frontini A, Castellucci M & Cinti S. Presence and distribution of cholinergic nerves in rat mediastinal brown adipose tissue. *Journal of Histochemistry and Cytochemistry* 2004 **52** 923–930. (doi:10.1369/jhc.3A6246.2004)
 - 39 Foster MT & Bartness TJ. Sympathetic but not sensory denervation stimulates white adipocytes proliferation. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* 2006 **291** R1630–R1637. (doi:10.1152/ajpregu.00197.2006)
 - 40 Cinti S. Transdifferentiation properties of adipocytes in the adipose organ. *American Journal of Physiology. Endocrinology and Metabolism* 2009 **297** E977–E986. (doi:10.1152/ajpendo.00183.2009)
 - 41 Cinti S. Reversible physiological transdifferentiation in the adipose organ. *Proceedings of the Nutrition Society* 2009 **68** 340–349. (doi:10.1017/S0029665109990140)
 - 42 Wang QA, Tao C, Gupta RK & Scherer PE. Tracking adipogenesis during white adipose tissue development, expansion and regeneration. *Nature Medicine* 2013 **19** 1338–1344. (doi:10.1038/nm.3324)
 - 43 Barbatelli G, Murano I, Madsen L, Hao Q, Jimenez M, Kristiansen K, Giacobino JP, De Matteis R & Cinti S. The emergence of cold-induced brown adipocytes in mouse white fat depots is determined predominantly by white to brown adipocyte transdifferentiation. *American Journal of Physiology. Endocrinology and Metabolism* 2010 **298** E1244–E1253. (doi:10.1152/ajpendo.00600.2009)
 - 44 Bronnikov G, Houstek J & Nedergaard J. β -adrenergic, cAMP-mediated stimulation of proliferation of brown fat cells in primary culture. Mediation via β 1 but not β 3 adrenoceptors. *Journal of Biological Chemistry* 1992 **267** 2006–2013.
 - 45 Guerra C, Navarro P, Valverde AM, Arribas M, Bruning J, Kozak LP, Khan CR & Benito M. Brown adipose tissue-specific insulin receptor knockout shows diabetic phenotype without insulin resistance. *Journal of Clinical Investigation* 2001 **108** 1205–1213. (doi:10.1172/JCI13103)
 - 46 Nedergaard J, Bengtsson T & Cannon B. New powers of brown fat: fighting the metabolic syndrome. *Cell Metabolism* 2011 **13** 238–240. (doi:10.1016/j.cmet.2011.02.009)
 - 47 Bartel A, Bruns OT, Reimer R, Hohenberg H, Ittrich H, Peldschus K, Kaul MG, Tromsdorf UI, Weller H, Waurisch C *et al.* Brown adipose tissue activity controls triglyceride clearance. *Nature Medicine* 2011 **17** 200–205. (doi:10.1038/nm.2297)
 - 48 Stanford KI, Middelbeek RJ, Townsend KL, An D, Nygaard EB, Hitchcox KM, Markan KR, Nakano KR, Hirshman MF, Tseng YH *et al.* Brown adipose tissue regulates glucose homeostasis and insulin sensitivity. *Journal of Clinical Investigation* 2013 **123** 215–223. (doi:10.1172/JCI62308)
 - 49 Kopecky J, Hodny Z, Rossmeisl M, Syrový I & Kozak LP. Reduction of dietary obesity in aP2-Ucp transgenic mice: physiology and adipose tissue distribution. *American Journal of Physiology* 1996 **270** E768–E775.
 - 50 Seale P, Conroe HM, Estall J, Kajimura S, Frontini A, Ishibashi J, Cohen P, Cinti S & Spiegelman BM. Prdm16 determines the thermogenic program of subcutaneous white adipose tissue in mice. *Journal of Clinical Investigation* 2011 **121** 96–105. (doi:10.1172/JCI44271)
 - 51 Almind K, Manieri M, Sivitz WI, Cinti S & Kahn CR. Ectopic brown adipose tissue in muscle provides a mechanism for differences in risk of metabolic syndrome in mice. *PNAS* 2007 **104** 2366–2371. (doi:10.1073/pnas.0610416104)
 - 52 Lowell BB, S-Susulic V, Hamann A, Lawitts JA, Himms-Hagen J, Boyer BB, Kozak LP & Flier JS. Development of obesity in transgenic mice after genetic ablation of brown adipose tissue. *Nature* 1993 **366** 740–742. (doi:10.1038/366740a0)
 - 53 Ghorbani M, Claus TH & Himms-Hagen J. Hypertrophy of brown adipocytes in brown and white adipose tissues and reversal of diet-induced obesity in rats treated with a β 3-adrenoceptor agonist. *Biochemical Pharmacology* 1997 **54** 121–131. (doi:10.1016/S0006-2952(97)00162-7)
 - 54 Ghorbani M & Himms-Hagen J. Appearance of brown adipocytes in white adipose tissue during CL 316,243-induced reversal of obesity and diabetes in Zucker fa/fa rats. *International Journal of Obesity and Related Metabolic Disorders* 1997 **21** 465–475. (doi:10.1038/sj.ijo.0800432)
 - 55 Ghorbani M & Himms-Hagen J. Treatment with CL 316,243, a β 3-adrenoceptor agonist, reduces serum leptin in rats with diet- or aging-associated obesity, but not in Zucker rats with genetic (fa/fa) obesity. *International Journal of Obesity and Related Metabolic Disorders* 1998 **22** 63–65. (doi:10.1038/sj.ijo.0800544)
 - 56 Waldén TB, Hansen IR, Timmons JA, Cannon B & Nedergaard J. Recruited vs. nonrecruited molecular signatures of brown, "brite," and white adipose tissues. *American Journal of Physiology. Endocrinology and Metabolism* 2012 **302** E19–E31. (doi:10.1152/ajpendo.00249.2011)
 - 57 Petrovic N, Walden TB, Shabalina IG, Timmons JA, Cannon B & Nedergaard J. Chronic peroxisome proliferator-activated receptor γ (PPAR γ) activation of epididymally derived white adipocyte cultures reveals a population of thermogenically competent, UCP1-containing adipocytes molecularly distinct from classic brown adipocytes. *Journal of Biological Chemistry* 2010 **285** 7153–7164. (doi:10.1074/jbc.M109.053942)
 - 58 Seale P, Bjork B, Yang W, Kajimura S, Chin S, Kuang S, Scimè A, Devarakonda S, Conroe HM, Erdjument-Bromage H *et al.* PRDM16 controls a brown fat/skeletal muscle switch. *Nature* 2008 **454** 961–967. (doi:10.1038/nature07182)
 - 59 Wu J, Bostrom P, Sparks LM, Ye L, Choi JH, Giang AH, Khandekar M, Virtanen KA, Nuutila P, Schaart G *et al.* Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell* 2012 **150** 366–376. (doi:10.1016/j.cell.2012.05.016)
 - 60 Berry R & Rodeheffer MS. Characterization of the adipocyte cellular lineage *in vivo*. *Nature Cell Biology* 2013 **15** 302–308. (doi:10.1038/ncb2696)
 - 61 Iyama K, Ohzono K & Usuku G. Electron microscopical studies on the genesis of white adipocytes: differentiation of immature pericytes into adipocytes in transplanted preadipose tissue. *Virchows Archiv* 1979 **31** 143–155. (doi:10.1007/BF02889932)

- 62 Slavin BG. Fine structural studies on white adipocyte differentiation. *Anatomical Record* 1979 **195** 63–72. (doi:10.1002/ar.1091950106)
- 63 Cinti S, Cigolini M, Bosello O & Bjorntorp P. A morphological study of the adipocyte precursor. *Journal of Submicroscopic Cytology and Pathology* 1984 **16** 243–251.
- 64 Tang W, Zeve D, Suh JM, Bosnakovski D, Kyba M, Hammer RE, Tallquist MD & Graff JM. White fat progenitor cells reside in the adipose vasculature. *Science* 2008 **322** 583–586. (doi:10.1126/science.1156232)
- 65 Tran KV, Gealekman O, Frontini A, Zingaretti MC, Morroni M, Giordano A, Smorlesi A, Perugini J, De Matteis R, Sbarbati A *et al.* The vascular endothelium of the adipose tissue gives rise to both white and brown fat cells. *Cell Metabolism* 2012 **15** 222–229. (doi:10.1016/j.cmet.2012.01.008)
- 66 Gupta RK, Mepani RJ, Kleiner S, Lo JC, Khandekar MJ, Cohen P, Frontini A, Bhowmic DC, Ye L, Cinti S *et al.* Zfp423 expression identifies committed preadipocytes and localizes to adipose endothelial and perivascular cells. *Cell Metabolism* 2012 **15** 230–239. (doi:10.1016/j.cmet.2012.01.010)
- 67 Lee HJ & Ormandi CJ. Elf5, hormones and cell fate. *Trends in Endocrinology and Metabolism* 2012 **23** 292–298. (doi:10.1016/j.tem.2012.02.006)
- 68 De Matteis R, Zingaretti MC, Murano I, Vitali A, Frontini A, Giannulis I, Barbatelli G, Marcucci F, Bordicchia M, Sarzani R *et al.* *In vivo* physiological transdifferentiation of adult adipose cells. *Stem Cells* 2009 **27** 2761–2768. (doi:10.1002/stem.197)
- 69 Apostoli AJ, Skelhorne-Gross GE, Rubino RE, Peterson NT, Di Lena MA, Schneider MM, Senqupta SK & Nicol CJ. Loss of PPAR γ expression in mammary secretory epithelial cells creates a pro-breast tumorigenic environment. *International Journal Cancer* 2014 **134** 1055–1066. (doi:10.1002/ijc.28432)
- 70 Yin Y, Yuan H, Wang C, Pattabiraman N, Rao M, Pestell RG & Glazer RI. 3-phosphoinositide-dependent, protein kinase-1 activates the peroxisome proliferator-activated receptor- γ and promotes adipocyte differentiation. *Molecular Endocrinology* 2006 **20** 268–278. (doi:10.1210/me.2005-0197)
- 71 Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA *et al.* Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *Journal of Clinical Investigation* 2003 **112** 1821–1830. (doi:10.1172/JCI200319451)
- 72 Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL & Ferrante AW Jr. Obesity is associated with macrophage accumulation in adipose tissue. *Journal of Clinical Investigation* 2003 **112** 1796–1808. (doi:10.1172/JCI200319246)
- 73 Cinti S, Mitchell G, Barbatelli G, Murano I, Ceresi E, Faloia E, Wang S, Fortier M, Greenberg AS & Obin MS. Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. *Journal of Lipid Research* 2005 **46** 2347–2355. (doi:10.1194/jlr.M500294-JLR200)
- 74 Giordano A, Murano I, Mondini E, Perugini J, Smorlesi A, Severi I, Barazzoni R, Scherer PE & Cinti S. Obese adipocytes show ultrastructural features of stressed cells and die of pyroptosis. *Journal of Lipid Research* 2013 **54** 2423–2436. (doi:10.1194/jlr.M038638)
- 75 Pajvani UB, Trujillo ME, Combs TP, Iyengar P, Jelicks L, Roth KA, Kitsis RN & Scherer PE. Fat apoptosis through targeted activation of caspase 8: a new mouse model of inducible and reversible lipoatrophy. *Nature Medicine* 2005 **11** 797–803. (doi:10.1038/nm1262)
- 76 Murano I, Rutkowski JM, Wang QA, Cho YR, Scherer PE & Cinti S. Time course of histomorphological changes in adipose tissue upon acute lipoatrophy. *Nutrition, Metabolism, and Cardiovascular Diseases* 2013 **23** 723–731. (doi:10.1016/j.numecd.2012.03.005)
- 77 Xu X, Grijalva A, Skowronski A, van Eijk M, Serlie MJ & Ferrante AW Jr. Obesity activates a program of lysosomal-dependent lipid metabolism in adipose tissue macrophages independently of classic activation. *Cell Metabolism* 2013 **18** 816–830. (doi:10.1016/j.cmet.2013.11.001)
- 78 Wang SP, Laurin N, Himms-Hagen J, Rudnicki MA, Levy E, Robert MF, Pan L, Oligny L & Mitchell GA. The adipose tissue phenotype of hormone-sensitive lipase deficiency in mice. *Obesity Research* 2001 **9** 119–128. (doi:10.1038/oby.2001.15)
- 79 Murano I, Barbatelli G, Parisani V, Latini C, Muzzonigro G, Castellucci M & Cinti S. Dead adipocytes, detected as crown-like structures, are prevalent in visceral fat depots of genetically obese mice. *Journal of Lipid Research* 2008 **49** 1562–1568. (doi:10.1194/jlr.M800019-JLR200)
- 80 Bjorntorp P. Metabolic abnormalities in visceral obesity. *Annals of Medicine* 1992 **24** 3–5. (doi:10.3109/07853899209164137)
- 81 Valet P, Grujic D, Wade J, Ito M, Zingaretti MC, Soloveva V, Ross SR, Graves RA, Cinti S, Lafontan M *et al.* Expression of human α 2-adrenergic receptors in adipose tissue of β 3-adrenergic receptor-deficient mice promotes diet-induced obesity. *Journal of Biological Chemistry* 2000 **275** 34797–34802. (doi:10.1074/jbc.M005210200)
- 82 Arner E, Westermark PO, Spalding KL, Britton T, Rydén M, Frisén J, Bernard S & Arner P. Adipocyte turnover: relevance to human adipose tissue morphology. *Diabetes* 2010 **59** 105–109. (doi:10.2337/db09-0942)
- 83 Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, Kuo FC, Palmer EL, Tseng YH, Doria A *et al.* Identification and importance of brown adipose tissue in adult humans. *New England Journal of Medicine* 2009 **360** 1509–1517. (doi:10.1056/NEJMoa0810780)
- 84 Virtanen KA, Lidell ME, Orava J, Heglind M, Niemi T, Taittonen M, Laine J, Savisto NJ, Enerback S & Nuutila P. Functional brown adipose tissue in healthy adults. *New England Journal of Medicine* 2009 **360** 1518–1525. (doi:10.1056/NEJMoa0808949)
- 85 van Marken Lichtenbelt WD, Vanhomerig JW, Smulders NM, Drossaerts JM, Kemerink GJ, Bouvy ND, Schrauwen P & Teule GJ. Cold-activated brown adipose tissue in healthy men. *New England Journal of Medicine* 2009 **360** 1500–1508. (doi:10.1056/NEJMoa0808718)
- 86 Saito M, Okamatsu-Ogura Y, Matsushita M, Watanabe K, Yoneshiro T, Nio-Kobayashi J, Iwanaga T, Miyagawa M, Kameya T, Nakada K *et al.* High incidence of metabolically active brown adipose tissue in healthy adult humans: effects of cold exposure and adiposity. *Diabetes* 2009 **58** 1526–1531. (doi:10.2337/db09-0530)
- 87 Zingaretti MC, Crosta F, Vitali A, Guerrieri M, Frontini A, Cannon B, Nedergaard J & Cinti S. The presence of UCP1 demonstrates that metabolically active adipose tissue in the neck of adult humans truly represents brown adipose tissue. *FASEB Journal* 2009 **23** 3113–3120. (doi:10.1096/fj.09-133546)
- 88 Pond CM & Mattacks CA. The anatomy of adipose tissue in captive *Macaca* monkeys and its implications for human biology. *Folia Primatologica* 1987 **48** 164–185. (doi:10.1159/000156293)
- 89 Prins JB & O’Rahilly S. Regulation of adipose cell number in man. *Clinical Science* 1997 **92** 3–11.
- 90 McBride P. Triglycerides and risk for coronary artery disease. *Current Atherosclerosis Reports* 2008 **10** 386–390. (doi:10.1007/s11883-008-0060-9)
- 91 Jensen MD. Gender differences in regional fatty acid metabolism before and after meal ingestion. *Journal of Clinical Investigation* 1995 **96** 2297–2303. (doi:10.1172/JCI118285)
- 92 Ebbert JO & Jensen MD. Fat depots, free fatty acids, and dyslipidemia. *Nutrients* 2013 **5** 498–508. (doi:10.3390/nu5020498)
- 93 Merklin RJ. Growth and distribution of human fetal brown fat. *Anatomical Record* 1974 **178** 637–645. (doi:10.1002/ar.1091780311)
- 94 Lidell ME, Betz MJ, Dahlqvist Leinhard O, Heglind M, Elander L, Slawik M, Mussack T, Nilsson D, Romu T, Nuutila P *et al.* Evidence for two types of brown adipose tissue in humans. *Nature Medicine* 2013 **19** 631–634. (doi:10.1038/nm.3017)
- 95 Cinti S. The role of brown adipose tissue in human obesity. *Nutrition, Metabolism, and Cardiovascular Diseases* 2006 **16** 569–574. (doi:10.1016/j.numecd.2006.07.009)
- 96 Lean ME, James WP, Jennings G & Trayhurn P. Brown adipose tissue uncoupling protein content in human infants, children and adults. *Clinical Science* 1986 **71** 291–297.

- 97 Kortelainen ML, Pelletier G, Ricquier D & Bukowiecki LJ. Immunohistochemical detection of human brown adipose tissue uncoupling protein in an autopsy series. *Journal of Histochemistry and Cytochemistry* 1993 **41** 759–764. (doi:10.1177/41.5.8468458)
- 98 Garruti G & Ricquier D. Analysis of uncoupling protein and its mRNA in adipose tissue deposits of adult humans. *International Journal of Obesity and Related Metabolic Disorders* 1992 **16** 383–390.
- 99 Oberkofler H, Dallinger G, Liu YM, Hell E, Krempler F & Patsch W. Uncoupling protein gene: quantification of expression levels in adipose tissues of obese and non-obese humans. *Journal of Lipid Research* 1997 **38** 2125–2133.
- 100 Huttunen P, Hirvonen J & Kinnula V. The occurrence of brown adipose tissue in outdoor workers. *European Journal of Applied Physiology and Occupational Physiology* 1981 **46** 339–345. (doi:10.1007/BF00422121)
- 101 Lean ME, James WP, Jennings G & Trayhurn P. Brown adipose tissue in patients with pheochromocytoma. *International Journal of Obesity* 1986 **10** 219–227.
- 102 Kuji I, Imabayashi E, Minagawa A, Matsuda H & Miyauchi T. Brown adipose tissue demonstrating intense FDG uptake in a patient with mediastinal pheochromocytoma. *Annals of Nuclear Medicine* 2008 **22** 231–235. (doi:10.1007/s12149-007-0096-x)
- 103 Zancanaro C, Pelosi G, Accordini C, Balercia G, Sbabo L & Cinti S. Immunohistochemical identification of uncoupling protein in human hybernoma. *Biology of the Cell* 1994 **80** 75–78. (doi:10.1016/0248-4900(94)90021-3)
- 104 Gelfand MJ, O'Hara SM, Curtwright LA & Maclean JR. Premedication to block [¹⁸F] FDG uptake in the brown adipose tissue of pediatric and adolescent patients. *Pediatric Radiology* 2005 **35** 984–990. (doi:10.1007/s00247-005-1505-8)
- 105 Hany TF, Gharehpapagh E, Kamel EM, Buck A, Himms-Hagen J & von Schulthess GK. Brown adipose tissue: a factor to consider in symmetrical tracer uptake in the neck and upper chest region. *European Journal of Nuclear Medicine and Molecular Imaging* 2002 **29** 1393–1398. (doi:10.1007/s00259-002-0902-6)
- 106 Nedergaard J, Bengtsson T & Cannon B. Unexpected evidence for active brown adipose tissue in adult humans. *American Journal of Physiology. Endocrinology and Metabolism* 2007 **293** E444–E452. (doi:10.1152/ajpendo.00691.2006)
- 107 Smorlesi A, Frontini A, Giordano A & Cinti S. The adipose organ: white-brown adipocyte plasticity and metabolic inflammation. *Obesity Reviews* 2012 **13** (Suppl 2) 83–96. (doi:10.1111/j.1467-789X.2012.01039.x)
- 108 Manieri M, Murano I, Fianchini A, Brunelli A & Cinti S. Morphological and immunohistochemical features of brown adipocytes and preadipocytes in a case of human hybernoma. *Nutrition, Metabolism, and Cardiovascular Diseases* 2010 **20** 567–574. (doi:10.1016/j.numecd.2009.04.020)
- 109 Carruba M, Tonello C, Briscini L & Nisoli E. Advances in pharmacotherapy for obesity. *International Journal of Obesity and Related Metabolic Disorders* 1998 **22** (Suppl 1) S13–S16.
- 110 Ouellet V, Labbé SM, Blondin DP, Phoenix S, Guérin B, Haman F, Turcotte EE, Richard D & Carpentier AC. Brown adipose tissue oxidative metabolism contributes to energy expenditure during acute cold exposure in humans. *Journal of Clinical Investigation* 2012 **122** 545–552. (doi:10.1172/JCI60433)
- 111 van der Lans AA, Hoeks J, Brans B, Vijgen GH, Visser MG, Vosselman MJ, Hansen J, Jörgensen JA, Wu J, Mottaghy FM *et al.* Cold acclimation recruits human brown fat and increases nonshivering thermogenesis. *Journal of Clinical Investigation* 2013 **123** 3395–3403. (doi:10.1172/JCI68993)
- 112 Ouellet V, Routhier-Labadie A, Bellemare W, Lakhil-Chaieb L, Turcotte E, Carpentier AC & Richard D. Outdoor temperature, age, sex, body mass index, and diabetic status determine the prevalence, mass, and glucose-uptake activity of ¹⁸F-FDG-detected BAT in humans. *Journal of Clinical Endocrinology and Metabolism* 2011 **96** 192–199. (doi:10.1210/jc.2010-0989)
- 113 Yoneshiro T, Aita S, Matsushita M, Kameya T, Nakada K, Kawai Y & Saito M. Brown adipose tissue, whole-body energy expenditure, and thermogenesis in healthy adult men. *Obesity* 2011 **19** 13–16. (doi:10.1038/oby.2010.105)
- 114 Yoneshiro T, Aita S, Matsushita M, Kayahara T, Kameya T, Kaway Y, Iwanaga T & Saito M. Recruited brown adipose tissue as an antiobesity agent in humans. *Journal of Clinical Investigation* 2013 **123** 3404–3408. (doi:10.1172/JCI67803)
- 115 Yang X, Enerback S & Smith U. Reduced expression of FOXC2 and brown adipogenic genes in human subjects with insulin resistance. *Obesity Research* 2003 **11** 1182–1191. (doi:10.1038/oby.2003.163)
- 116 Elabd C, Chiellini C, Carmona M, Galitzky J, Cochet O, Petersen R, Pénicaud L, Kristiansen K, Bouloumié A, Casteilla L *et al.* Human multipotent adipose-derived stem cells differentiate into functional brown adipocytes. *Stem Cells* 2009 **27** 2753–2760. (doi:10.1002/stem.200)
- 117 Beranger GE, Karbiener M, Barquissau V, Pisani DF, Scheideler M & Amri EZ. *In vitro* brown and "brite"/"beige" adipogenesis: human cellular models and molecular aspects. *Biochimica et Biophysica Acta* 2013 **1831** 905–914. (doi:10.1016/j.bbali.2012.11.001)
- 118 Frontini A, Vitali A, Perugini J, Murano I, Romiti C, Ricquier D, Guerrieri M & Cinti S. White-to-brown transdifferentiation of omental adipocytes in patients affected by pheochromocytoma. *Biochimica et Biophysica Acta* 2013 **1831** 950–959. (doi:10.1016/j.bbali.2013.02.005)
- 119 Cigolini M, Cinti S, Brunetti L, Bosello O, Osculati F & Bjorntorp P. Human brown adipose cells in culture. *Experimental Cell Research* 1985 **159** 261–266. (doi:10.1016/S0014-4827(85)80056-2)
- 120 Larsen TM, Toubro S, van Baak MA, Gottesdiener KM, Larson P, Saris WH & Astrup A. Effect of a 28-d treatment with L-796568, a novel β (3)-adrenergic receptor agonist, on energy expenditure and body composition in obese men. *American Journal of Clinical Nutrition* 2002 **76** 780–788.
- 121 van Baak MA, Hul GB, Toubro S, Astrup A, Gottesdiener KM, DeSmet M & Saris WH. Acute effect of L-796568, a novel β 3-adrenergic receptor agonist, on energy expenditure in obese men. *Clinical Pharmacology and Therapeutics* 2002 **71** 272–279. (doi:10.1067/mcp.2002.122527)
- 122 Hondares E, Rosell M, Gonzalez FJ, Giral M, Iglesias R & Villarroya F. Hepatic FGF21 expression is induced at birth via PPAR α in response to milk intake and contributes to thermogenic activation of neonatal brown fat. *Cell Metabolism* 2010 **11** 206–212. (doi:10.1016/j.cmet.2010.02.001)
- 123 Bordicchia M, Liu D, Amri EZ, Ailhaud G, Dessi-Fulgheri P, Zhang C, Takahashi N, Sarzani R & Collins S. Cardiac natriuretic peptides act via p38 MAPK to induce the brown fat thermogenic program in mouse and human adipocytes. *Journal of Clinical Investigation* 2012 **122** 1022–1036. (doi:10.1172/JCI59701)
- 124 Bostrom P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, Rasbach KA, Bostrom EA, Choi JH, Long JZ *et al.* A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 2012 **481** 463–468. (doi:10.1038/nature10777)
- 125 Whittle AJ, Carobbio S, Martins L, Slawik M, Hondares E, Vázquez MJ, Morgan D, Csikasz RI, Gallego R, Rodriguez-Cuenca S *et al.* BMP8B increases brown adipose tissue thermogenesis through both central and peripheral actions. *Cell* 2012 **149** 871–885. (doi:10.1016/j.cell.2012.02.066)

Received 22 November 2013

Revised version received 24 January 2014

Accepted 27 January 2014