



P112 Influence of Cuff Blood Pressure Accuracy on Identification of Isolated Systolic Hypertension

Dean Picone^{1,*}, Martin Schultz¹, Matthew Armstrong¹, Willem Bos^{2,3}, Nathan Dwyer^{4,5}, Peter Lacy⁶, Esben Laugesen⁷, Stefano Omboni^{8,9}, Giacomo Pucci¹⁰, Philip Roberts-Thomson^{5,1}, George Stouffer¹¹, Kenji Takazawa¹², Thomas Weber¹³, Berend Westerhof¹⁴, James Sharman¹⁵

¹Menzies Institute for Medical Research, College of Health and Medicine, University of Tasmania, Australia

²St Antonius Hospital, Department of Internal Medicine, Nieuwegein, The Netherlands

³Department of Internal Medicine, Leiden University Medical Center, Leiden, The Netherlands

⁴Menzies Institute for Medical Research, College of Health and Medicine, University of Tasmania, Hobart, Australia

⁵Royal Hobart Hospital, Hobart, Australia

⁶UCL Institute of Cardiovascular Science

⁷Department of Endocrinology and Internal Medicine, Aarhus University Hospital, Aarhus, Denmark

⁸Clinical Research Unit, Italian Institute of Telemedicine, Varese, Italy

⁹Scientific Research Department of Cardiology, Science and Technology Park for Biomedicine, Sechenov First Moscow State Medical University, Moscow, Russian Federation

¹⁰Unit of Internal Medicine at Terni University Hospital, Department of Medicine, University of Perugia, Perugia, Italy

¹¹Division of Cardiology, University of North Carolina at Chapel Hill, Chapel Hill, United States

¹²Center for Health Surveillance and Preventive Medicine, Tokyo Medical University Hospital, Tokyo, Japan

¹³Cardiology Department, Klinikum Wels-Grieskirchen, Wels, Austria

¹⁴Department of Pulmonary Diseases, VU University Medical Center, Amsterdam, The Netherlands

¹⁵Menzies Institute for Medical Research, College of Health and Medicine, University of Tasmania, Hobart, Australia

ABSTRACT

Introduction: Isolated systolic hypertension (ISH) is the most common form of hypertension in older people. However, accurate identification of ISH may be hindered because cuff blood pressure (BP) underestimates systolic BP (SBP) and overestimates diastolic BP (DBP). This study aimed to determine the influence of cuff BP accuracy on the identification of ISH.

Methods: Cuff BP and invasive aortic BP were measured simultaneous (or near-simultaneously) in 1737 subjects (63 ± 12 years, 68% male) during coronary angiography. Data was derived from 32 studies, using 20 different cuff BP devices, from the Invasive Blood Pressure Consortium (INSPECT). ISH was defined as $\geq 140 / < 90$ mmHg according to cuff BP and invasive aortic BP.

Results: According to cuff BP, 25% of subjects ($n = 430$) had ISH, however, 37% ($n = 648$) were identified with ISH from invasive aortic BP. There was 77% concordance between cuff and invasive BP for identifying ISH. Among the 23% ($n = 408$) of subjects misclassified by cuff BP, 38% ($n = 155$) of misclassification was from SBP underestimation (mean: -16.6 , 95% CI: -18.9 to -13.9 mmHg), whereas 35% ($n = 143$) was from DBP overestimation (15.6, 11.9 to 19.0 mmHg) and 20% ($n = 83$) from SBP overestimation (17.6, 14.4 to 20.5 mmHg). Subjects misclassified were on average 2.7 years older and had greater body mass index (0.8 kg/m^2) than those correctly classified.

Conclusion: Approximately one quarter of older subjects have ISH misclassified, mostly because of underestimation of cuff SBP and overestimation of cuff DBP. This demonstrates a need to improve the accuracy of cuff BP methods for greater precision in identifying ISH.

© 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).